Frequentist and Bayesian Analysis of the Matter–Antimatter Asymmetry in $B^{\pm} \to K^{\pm}K^{+}K^{-}$ Decays at LHCb

Final project for the *Statistics and Data Analysis* course
Academic Year 2025–2026

Marco Belotti and Giacomo Moretti
Università degli Studi di Milano-Bicocca

We measure the raw charge asymmetry $A_{\text{raw}} = [N(B^-) - N(B^+)]/[N(B^-) + N(B^+)]$ of $B^{\pm} \to K^{\pm}K^{+}K^{-}$ decays in the 2011 LHCb Open Data record and compare two inference frameworks on the same data and the same question: an extended unbinned maximum-likelihood fit with profile likelihood and a Bayesian analysis sampled with Markov chain Monte Carlo (MCMC). After a common preselection (19,731 candidates in the fit window), a common mass model gives $A_{\text{raw}} = -0.0441 \pm 0.0085\,(\text{stat}) \pm 0.0072\,(\text{syst})$ in the frequentist fit and $-0.0443 \pm 0.0085 \pm 0.0070$ in the Bayesian one, 5.2 standard deviations from zero (Bayes factor $\approx 9 \times 10^3$ against no asymmetry). The two methods agree in value, uncertainty and coverage, which was verified on toy-MC experiments; the dominant systematic uncertainty is the particle-identification requirement. The asymmetry is not uniform over the Dalitz plane: in the region of the published analysis of the same data we find $A_{\text{raw}} = -0.234 \pm 0.025$, reproducing -0.239 ± 0.020. A machine-learning selection trained on weak labels, described in Appendix I, leaves A_{raw} unchanged within its statistical spread. A_{raw} is not corrected for production and detection asymmetries and is therefore not a measurement of A_{CP}.

CONTENTS

I. Introduction 1

II. Data 2
A. Dataset, Preselection and Mass 2
B. Data Exploration 2

III. Mass Model and Likelihood 3

IV. Frequentist Analysis 4
A. Mass Model and Fit Strategy 4
B. Separate vs. Simultaneous Fit 4
C. Profile Likelihood and Confidence Interval 4
D. Coverage of the Wilks Intervals 5

V. Bayesian Analysis 5
A. Priors and Posterior 5
B. MCMC Sampling and Convergence Diagnostics 5
C. Posterior Results and Credible Intervals 6
D. Bayes Factors 6

VI. Systematic Uncertainties and Result 6
A. Stability Checks of the Frequentist Fit 7

VII. Validation of the Estimator and of the Intervals 8
A. Bias, Efficiency and Coverage 8

VIII. Local Asymmetry in the Dalitz Plane 9

IX. Comparison and Discussion 10
A. Where the Approaches Differ 10
B. Comparison with the Published Analysis 11
C. Machine-Learning Cross-Check 12

X. Conclusions 12

A. Dataset and Preselection 12

B. Mass Model and Likelihood 13

C. Frequentist Analysis 14

D. Bayesian Analysis 16

E. Systematic Uncertainties 23

F. Validation of the Estimator and of the Intervals 28

G. Local Asymmetry in the Dalitz Plane 30

H. Comparison of the Two Approaches 32

I. Machine-Learning Analysis 35

J. Conclusions, Scope and Limitations 44

K. Code and Data Availability 46

L. Notation 47

References 47

I. INTRODUCTION

Charge-parity (CP) violation – the small observed asymmetry between the behavior of matter and antimatter – is one of the necessary ingredients for the matter-dominated universe we observe, yet the CP violation built into the Standard Model is far too small to account for it.

Studying CP violation in the decays of B mesons, where it is comparatively large and experimentally accessible, is one of the main ways to test the Standard Model's Cabibbo-Kobayashi-Maskawa (CKM) description of the effect and search for physics beyond it. Charmless three-body decays such as $B^\pm \to h^\pm h^+ h^-$ ($h = K, \pi$) are especially interesting in this respect: unlike a two-body decay, the three-body phase space (the Dalitz plot) allows the asymmetry to be studied not only as a single global number, but also *locally*, since interference between intermediate resonances can concentrate CP-violating effects in specific kinematic regions rather than spreading them uniformly [1, 2].

The quantity measured in this work is the raw asymmetry between the signal yields of the two charges,

$$A_{\rm raw} = \frac{N(B^-) - N(B^+)}{N(B^-) + N(B^+)}. \tag{1}$$

It differs from the physical asymmetry A_{CP} by the production asymmetry A_P of $B^\pm$ mesons in pp collisions and by the detection asymmetry A_D of the final state; to first order in these small quantities,

$$A_{CP} \approx A_{\rm raw} - A_P - A_D, \tag{2}$$

where A_D arises mainly from the different interaction cross sections of K^+ and K^- with the detector material and depends on the kaon momenta [1]. Neither correction is applied here, so $A_{\rm raw}$ is not a measurement of A_{CP}. The corrections applied in the published analysis are discussed in Sec. IX B.

In this work we use the LHCb Open Data record for $B^\pm \to K^\pm K^+ K^-$ [3, 4], a subset of the 2011 data on which the first evidence of an inclusive CP asymmetry in this decay was published [5], to ask: *to what extent do frequentist likelihood fits and Bayesian inference produce consistent, well-controlled estimates of the raw asymmetry $A_{\rm raw}$?* The aim is a methodological comparison of the two frameworks on the same data and, by construction, on the same likelihood, so that any difference between the two results comes from the inference and not from the model; it is not a competitive measurement of A_{CP}, which cannot be obtained from this record because it lacks the control channel for A_P and A_D (Appendix J.5); the other three channels of the record need a different selection and a model of peaking backgrounds and are not analysed. Secs. II–III describe the data and the common model, Secs. IV–V the two fits, and Secs. VI–IX the systematic study, the validation, the Dalitz plane and the comparison; a machine-learning selection is in Appendix I. The main results are in Table I.

II. DATA

A. Dataset, Preselection and Mass

In short: 8,556,118 *candidates are reduced to* 23,895 *by a kaon-identification and muon-veto requirement common to the analyses; the fit window* $[5100, 5600]$ *MeV contains* 19,731 *candidates (*10,292 B^+, 9439 B^-*).* The dataset is the CERN Open Data record 4900 [3]: $B^\pm \to h^\pm h^+ h^-$ candidates collected by LHCb in 2011 at $\sqrt{s} = 7\,\text{TeV}$, already through trigger and offline selection, distributed as ROOT `TTree`s with the most relevant branches and split by magnet polarity, for 8,556,118 candidates. The record also contains a phase-space simulation of $B^\pm \to K^\pm K^+ K^-$ (50,000 decays, no asymmetry), used only to test the local-asymmetry procedure (Sec. VIII).

For each of the three final-state tracks ($H1$, $H2$, $H3$) the analysis reads the momentum components, the particle-identification (PID) likelihoods (`ProbK`, `ProbPi`), a muon flag and the charge. The same preselection is used in all analyses,

$$\texttt{H*_ProbPi} < 0.5 \ \wedge\ \texttt{H*_ProbK} > 0.5 \ \wedge\ \neg\texttt{H*_isMuon},$$

and retains 23,895 of the 8,556,118 candidates (0.28%), the PID requirement being the dominant reduction (Table V of Appendix A).

$H1$ is always the kaon of charge opposite to the B, so the Dalitz variables are $m^2(H1, H2)$ and $m^2(H1, H3)$.

The three final-state tracks are assigned the charged-kaon mass hypothesis ($M_K = 493.677$ MeV) regardless of their PID likelihoods, and their four-momenta are summed to reconstruct the B candidate, $m(KKK) = \sqrt{(\sum_i E_i)^2 - |\sum_i \vec{p}_i|^2}$. The nominal fit window, common to the frequentist and Bayesian analyses, is $[5100, 5600]$ MeV and contains 19,731 candidates (10,292 B^+, 9439 B^-; 11,636 magnet down, 8095 magnet up). Below ∼5150 MeV the background rises, consistent with a partially reconstructed contribution that a single exponential cannot describe; the window includes the start of this rise, which is described by a second exponential in the background model (Secs. IV–V). The machine-learning selection (Appendix I) uses the same window.

B. Data Exploration

In short: the spectrum peaks at 5284 *MeV with a similar shape for the two charges and the two magnet polarities, and the counting asymmetry of all candidates is* -0.043 ± 0.007.

Figure I shows the spectrum in the fit window, split by B charge (a) and by magnet polarity (b). The peak at $m \approx 5284$ MeV over a falling combinatorial background has the same shape for the two charges and for the two polarities, which is what allows a mass and a resolution common to B^+ and B^- in the model (Sec. III). The asymmetry itself is not visible by eye: the counting asymmetry of all candidates, signal and background together, is $A_{\rm count} = (N^- - N^+)/(N^- + N^+) = -0.043 \pm 0.007$ (binomial uncertainty), and the fits of Secs. IV and V refine it by separating signal and background.

Further material is in Appendix A.

Method	Sample	A_{raw}	Statistical	Systematic	Section
Counting (no fit)	fit window, 19,731	-0.0432	0.0071	–	II B
Frequentist fit	fit window, 19,731	-0.0441	0.0085	0.0072	IV, VI
Bayesian fit	fit window, 19,731	-0.0443	0.0085	0.0070	V, VI
Bayesian fit, D^0 veto	18,744	-0.0496	0.0088	–	D.15
Bayesian fit, ML selection	score > 0.5, 15,709	-0.0451	0.0087	–	App. I.9
Frequentist fit, ML selection	score > 0.5, 15,709	-0.0447	0.0086	–	App. I.9
Bayesian fit, Dalitz region of Ref. [5]	2146	-0.234	0.025	–	VIII
Published analysis (same data) [5]	different selection and model	-0.060	0.007	–	–

TABLE I. Summary of the raw asymmetry A_{raw} (not corrected for production and detection asymmetries). The Bayesian value is the posterior median and its standard deviation; the systematic uncertainty is given only for the nominal fits (Sec. VI). The published A_{raw} comes from a different selection and a model with additional background components, so it is not directly comparable with the first rows.




FIG. I. Reconstructed $B \to KKK$ invariant mass in the nominal fit window $[5100, 5600]$ MeV, split by (a) B charge and (b) magnet polarity. Both splits show a consistent peak at $m \approx 5284$ MeV over a falling combinatorial background, which rises below ∼5150 MeV.

III. MASS MODEL AND LIKELIHOOD

In short: a Gaussian signal and a double-exponential background for each charge, with the raw asymmetry A as a parameter of the fit; both inference methods use the same likelihood. The mass spectrum of each charge sample is described by a signal-plus-background mixture on the nominal fit window $[5100, 5600]$ MeV (Sec. II A),

$$f^{\pm}(m) = N_s^{\pm}\, S(m;\mu,\sigma) + N_b^{\pm}\, B^{\pm}(m), \qquad (3)$$

where S is a Gaussian truncated to the window, with mean μ and width σ shared by the two charges, and the background is a double exponential with a shape of its own for each charge,

$$B^{\pm}(m) = f_{\pm}\, E(m;\lambda_1^{\pm}) + (1 - f_{\pm})\, E(m;\lambda_2^{\pm}),$$

with $E(m;\lambda) \propto e^{-\lambda m}$ a truncated exponential normalized on the window. The steep component (λ_1) describes the rise of the background towards the lower edge of the window, the slow one (λ_2) the combinatorial background under and above the peak. The ordering $\lambda_1 > \lambda_2$ removes the label-switching symmetry of the mixture. Signal and background yields are split between the charges as

$$N_s^{\pm} = \tfrac{N_s}{2}(1 \mp A), \qquad N_b^{\pm} = \tfrac{N_b}{2}(1 \mp A_b),$$

so that $A = (N_s^- - N_s^+)/(N_s^- + N_s^+)$, the raw asymmetry A_{raw} of Eq. (1), is itself a parameter of the fit, the parameter of interest (POI, all the others being nuisance parameters), and the background asymmetry A_b is free. The parameter vector has twelve components, $\theta = (N_s, N_b, A, A_b, \mu, \sigma, \lambda_1^{\pm}, \lambda_2^{\pm}, f_{\pm})$.

Given the data $D = \{(m_i, q_i)\}_{i=1}^{N}$ (mass and charge of each candidate), the extended unbinned likelihood of the two charge samples is

$$\mathcal{L}(D \mid \theta) \propto e^{-(N_s+N_b)} \prod_{i:\, q_i=+1} \Big[N_s^+ S(m_i) + N_b^+ B^+(m_i)\Big] \times \prod_{i:\, q_i=-1} \Big[N_s^- S(m_i) + N_b^- B^-(m_i)\Big], \qquad (4)$$

where $S(m_i) \equiv S(m_i;\mu,\sigma)$. The likelihood is extended (the yields are parameters, hence the Poisson term, Appendix B); it is the common ground of the two mass

analyses: Sec. IV maximizes it and Sec. V multiplies it by a prior. Since both use this model unchanged, any difference between their results can only come from the inference.

IV. FREQUENTIST ANALYSIS

The frequentist analysis estimates $A_{\rm raw}$ by maximizing the extended unbinned likelihood of Eq. (4), i.e. the same model and the same window $[5100, 5600]$ MeV as the Bayesian analysis, with `iminuit` [6]. The B^+ and B^- samples were fitted simultaneously with $A_{\rm raw}$ as a fit parameter, and its uncertainty and the test of $A = 0$ were derived from the profile likelihood, the other parameters being treated as nuisance parameters (Secs. IV A–IV C).

A. Mass Model and Fit Strategy

In short: the common nominal model of Sec. III, fitted by extended unbinned maximum likelihood with a background started from the sidebands, gives $N_s = 15{,}328 \pm 133$ *and* $N_b = 4{,}403 \pm 82$*; a single-exponential background is rejected.*

The frequentist fit uses the common mass model of Sec. III (Eq. (3)) on the same window $[5100, 5600]$ MeV: a Gaussian signal, with mean μ and width σ shared by the two charges, and a double-exponential background with a shape and a yield of its own for each charge. The twelve parameters are the same as in the Bayesian fit, the two background yields being left free (equivalent to the pair N_b, A_b). They were estimated by maximizing the extended unbinned likelihood of Eq. (4) through `iminuit.cost.ExtendedUnbinnedNLL`: the extended form is needed because the signal and background yields are fit parameters. The minimization started from a background description obtained independently of the signal, with an unbinned fit of the sidebands (the fit window without the region $[5200, 5350]$ MeV) for each charge. The fit gives $N_s = 15{,}328 \pm 133$ signal and $N_b = 4{,}403 \pm 82$ background candidates, to be compared with the Bayesian posterior medians $15{,}328 \pm 133$ and 4404 ± 82 (Sec. V C).

The shapes were not tuned on the frequentist fit: the nominal model is the common one of Sec. III, and the alternatives are systematic variants (Sec. VI). The nominal fit has $\chi^2/\mathrm{ndof} = 160/88$ ($p = 4.8 \cdot 10^{-6}$) in 2×50 bins of 10 MeV. The double exponential is required by the data: the background rises towards the lower edge of the window (Sec. II A), and a single exponential is rejected ($\chi^2/\mathrm{ndof} = 294/92$ ($p = 5.7 \cdot 10^{-23}$)), so for the systematic study only the double exponential is considered.

B. Separate vs. Simultaneous Fit

In short: fitting the two charges separately gives $A_{\rm raw} = -0.0442 \pm 0.0087$ *and fitting them simultaneously with* A *as a parameter gives* -0.0441 ± 0.0085*: the two parametrizations agree.*

The asymmetry was estimated in two ways. In the *simultaneous* fit the two charge samples share the signal parametrization $N_s^+ = N_s(1 - A)/2$, $N_s^- = N_s(1 + A)/2$ (the same one used on the Bayesian side, Sec. III) and the signal mean and width, so that A is a parameter of the likelihood; mass, resolution and background-shape parameters are nuisance parameters, profiled out when reporting uncertainty on A. In the *separate* fit each charge was fitted on its own with the same model and no parameter in common (seven free parameters per charge), and $A_{\rm raw} = (N_s^- - N_s^+)/(N_s^- + N_s^+)$ was computed afterwards from the two signal yields, with the uncertainty propagated from those of the yields, which are independent because the two samples are. The simultaneous fit gives $A_{\rm raw} = -0.0441 \pm 0.0085$ and the separate fits $A_{\rm raw} = -0.0442 \pm 0.0087$.

The two estimates agree, because sharing μ and σ between B^+ and B^-, as expected for a particle and its antiparticle and a symmetric detector, is adequate. Therefore the simultaneous fit, which estimates A directly and gives the profile-likelihood interval of the next subsection, is taken as the nominal one.

C. Profile Likelihood and Confidence Interval

In short: $\hat{A} = -0.0441$ *with 68% and 95% Minos intervals* $[-0.0526, -0.0356]$ *and* $[-0.0607, -0.0275]$*;* $A = 0$ *is rejected with* $q(0) \simeq 27$*,* $p = 2.05 \times 10^{-7}$ *(*5.20σ*).*

The maximum-likelihood estimate is $\hat{A} = -0.0441$, with a Hesse uncertainty of 0.0085, and the 68% and 95% Confidence Interval, CI, obtained with `MINOS` [7] are the following:

$$A_{\rm raw} = -0.0441,$$
$$68\%\ \mathrm{CI}\ [-0.0526, -0.0356],$$
$$95\%\ \mathrm{CI}\ [-0.0607, -0.0275].$$

The uncertainties are statistical only; the systematic uncertainty is given in Sec. VI.

The profile likelihood ratio,

$$q(A) = -2 \log \frac{L(A, \hat{\hat{\theta}})}{L(\hat{A}, \hat{\theta})},$$

where $\hat{\hat{\theta}}$ are the values of the other eleven parameters that maximize the likelihood at fixed A, was evaluated by re-minimizing them at each value of A (the `mnprofile` of `iminuit`). By Wilks' theorem [8], $q(A)$ follows a χ^2 distribution with one degree of freedom at the true value of A, so the 68% and 95% confidence intervals are the

sets of A with $q(A) \leq 1$ and $q(A) \leq 3.84$ (the MINOS intervals [7]), which need not be symmetric.

For the null hypothesis $A = 0$ it gives $q(0) \simeq 27$, i.e. a p-value of 2.05×10^{-7} corresponding to a significance of 5.20σ: the hypothesis of no asymmetry is rejected. This is the frequentist counterpart of the Savage-Dickey test of Sec. V D, where $A = 0$ lies at 5.2 posterior standard deviations.The two quantities do not measure the same thing (the p-value is not the probability of H_0, and the Bayes factor depends on the width of the prior), but they lead to the same conclusion.

The result agrees with the Bayesian one (Sec. V C: $A_{\text{raw}} = -0.0443$, 68% CI $[-0.0528, -0.0358]$). This is expected: with flat priors and $\sim 2 \times 10^4$ events the posterior is dominated by the likelihood and nearly Gaussian, so the credible and confidence intervals coincide numerically although their interpretations differ.

D. Coverage of the Wilks Intervals

In short: in 1000 *bootstrap replicas the* 68% *interval contains the fitted value in* $68.4 \pm 1.5\%$ *of the cases, the nominal coverage.*

Wilks' theorem [8] is an asymptotic result, so the coverage of the profile-likelihood interval on A was checked directly on the data with a bootstrap. 1000 replicas of the 19,731 candidates of the fit window were drawn with replacement, which reproduces the statistical fluctuations of the sample, including those of the charge counts, and the nominal fit was repeated on each of them. The reference value of the asymmetry in the replicas is the best-fit value of the original data, $A_{\text{true}} = -0.0441$. For each replica the statistic $q(A_{\text{true}})$ was computed as the difference of $-2 \log L$ between the fit with A fixed to A_{true} and the fit with A free and the other eleven parameters profiled. The 68% interval of a replica contains A_{true} if and only if $q(A_{\text{true}}) \leq 1$; replicas in which a fit did not converge were discarded.

The fraction of replicas with $q(A_{\text{true}}) \leq 1$ is $68.4 \pm 1.5\%$ (binomial uncertainty), against the nominal 68.3%: the interval has the nominal coverage (Fig. II). This agrees with the toy-MC check of the Bayesian credible interval (Sec. D.16, $65.0 \pm 3.3\%$ for the 68% interval); the two values differ by less than one standard deviation, and the uncertainty here is smaller because of the larger number of replicas (1000 against 200). The two checks are not equivalent. The Bayesian pseudo-datasets are generated from the fitted model, which makes that check a closure test, while the replicas here are resampled from the data and therefore carry the features of the spectrum that the model does not describe, such as the shape of the signal tails (Sec. D.13).




FIG. II. Coverage check, 1000 toys. Left: the statistics $q(A_{\text{true}})$ following a $\chi^2_{dof=1}$ distribution as expected from Wilks theorem. Right: pulls $(\hat{A} - A_{\text{true}})/\sigma(\hat{A})$, with $\hat{A}$ and $\sigma(\hat{A})$ the fit results of each toy, compared with a standard normal.

V. BAYESIAN ANALYSIS

The Bayesian analysis takes the likelihood of Sec. III as it is and adds only a prior and a sampler; it changes the question asked of the data, not the model.

A. Priors and Posterior

In short: uniform bounded priors on twelve parameters; with $\sim 2 \times 10^4$ *candidates the posterior is dominated by the likelihood.* Bayes' theorem combines the likelihood of Eq. (4) (Sec. III) with a prior $\pi(\theta)$ into the posterior

$$p(\theta \mid D) = \frac{\mathcal{L}(D \mid \theta)\, \pi(\theta)}{p(D)}, \tag{5}$$

The priors are uniform, with independent ranges except for the ordering of the background slopes: $N_s, N_b \in (0, 5N)$ with $N = 19{,}731$, $A, A_b \in (-1, 1)$, $\mu \in (5260, 5310)$ MeV, $\sigma \in (3, 40)$ MeV, $\lambda_1^{\pm} \in (10^{-5}, 0.1)$ MeV^{-1}, $\lambda_2^{\pm} \in (10^{-5}, \lambda_1^{\pm})$, $f_{\pm} \in (0, 1)$. The prior density is therefore a constant on the allowed region Θ,

$$\pi(\theta) \propto \mathbf{1}\left[\theta \in \Theta\right], \tag{6}$$

and inside it the posterior is proportional to the likelihood. With $\sim 2 \times 10^4$ events the posterior is dominated by the likelihood, as the prior-sensitivity check of Appendix D.15 confirms for A. The evidence $p(D)$ is a constant: the MCMC samples $\mathcal{L}\,\pi$ without it, and the Savage-Dickey ratio of Sec. V D avoids it as well; the marginal posterior of A is the histogram of the A component of the chain.

B. MCMC Sampling and Convergence Diagnostics

In short: 32 walkers for 25,000 *steps; the chain is at least* 113 *autocorrelation times long for every parameter and independent runs from dispersed starting points agree.*

The posterior was sampled with the affine-invariant ensemble sampler of Goodman and Weare [9] as implemented in `emcee` [10] (Appendix D.1): 32 walkers started around the maximum of the posterior, 25,000 steps with the first 1/6 discarded as burn-in (6.7×10^5 samples, acceptance 0.38). The integrated autocorrelation time is $\tau \in [147, 183]$ steps, so the chain is $N/\tau \geq 113$ autocorrelation times long for every parameter, above the threshold of 50 [10] ($\gtrsim 3600$ effectively independent samples). Two runs started uniformly over the whole prior box converge to the same single mode ($A_{\mathrm{raw}} = -0.0446$ and -0.0442; trace plot in Fig. XIII, Appendix D.5).

C. Posterior Results and Credible Intervals

In short: $A_{\mathrm{raw}} = -0.0443$*, 68% credible interval* $[-0.0528, -0.0358]$ *and 95%* $[-0.0608, -0.0276]$*; the posterior is Gaussian.*

Figure III compares the nominal model with the data of the two charges: it follows the spectrum over two orders of magnitude, and the pulls show signal tails heavier than a Gaussian, whose effect on A_{raw} is negligible (Appendix D.13). The marginal posterior on the raw asymmetry gives (median and equal-tailed credible intervals, CI)

$$A_{\mathrm{raw}} = -0.0443,$$
$$68\%\ \mathrm{CI}\ [-0.0528, -0.0358],$$
$$95\%\ \mathrm{CI}\ [-0.0608, -0.0276].$$

The posterior is very close to Gaussian (standard deviation 0.0085), so equal-tailed and highest-posterior-density intervals coincide; with uniform priors its mode, -0.0441, is the maximum-likelihood estimate and the result can be compared directly with the frequentist fit (Sec. IV, Appendix C.1). The other physics parameters (Fig. IV) are $N_s = 15{,}328 \pm 133$, $N_b = 4404 \pm 82$, $A_b = -0.040 \pm 0.018$, $\mu = 5284.39 \pm 0.16$ MeV and $\sigma = 18.00 \pm 0.14$ MeV. A is almost uncorrelated with them, which is why profiling (frequentist) and marginalizing (Bayesian) the nuisance parameters give the same result; the largest correlation, -0.17, is with A_b: conditioning on $A_b = 0$ moves A to -0.0475, as the fit with a charge-symmetric background does (-0.0471, Sec. VI).

D. Bayes Factors

In short: decisive evidence against $A = 0$ *(Bayes factor* $BF_{10} \approx 9 \times 10^3$*); the evidence for a background asymmetry is not significant, and depends on the prior width.*

The Bayes factor compares two hypotheses through their evidences, $BF_{10} = P(D \mid H_1)/P(D \mid H_0)$, and requires proper priors. For the nested hypothesis H_0: $A = 0$ against H_1: A free, with shared priors on the other parameters, it reduces to the Savage-Dickey ratio [11] (Appendix C.1),

$$BF_{01} = \frac{p(A = 0 \mid D)}{\pi(A = 0)}, \qquad \pi(A = 0) = \tfrac{1}{2}.$$

$A = 0$ lies 5.2 posterior standard deviations from the median, where the chain has essentially no samples, so $p(A = 0 \mid D)$ was taken from the Gaussian approximation of the marginal posterior: $BF_{01} \approx 1.1 \times 10^{-4}$ ($BF_{10} \approx 9 \times 10^3$), decisive evidence against $A = 0$ on Jeffreys' scale [12]. The value scales with the prior width (Ockham factor): with $A \in (-0.3, 0.3)$, BF_{01} drops by a factor 3.3 and the conclusion is unchanged (Fig. V; the Savage-Dickey densities are in Fig. XIV of Appendix D, further checks in Appendix H.3).

The same test on the background asymmetry shows the Ockham factor at work: $A_b = -0.040 \pm 0.018$ is 2.3 standard deviations from zero, yet $BF_{01}(A_b = 0) \approx 3.5$, because the likelihood gain does not compensate for spreading the prior over $(-1, 1)$; unlike for A, the preference changes sign with the prior width (Fig. V, crossing at $W \approx 0.3$). Of the Bayesian results only the Bayes factor depends on the prior; the interval of A does not (Sec. V C). A negative A_b is plausible given the misidentified $B^{\pm} \to K^{\pm}\pi^{+}\pi^{-}$ and $\pi^{\pm}K^{+}K^{-}$ decays; A_b is kept free in the nominal model and fixing it to zero is one of the systematic variants (Sec. VI).

Further material is in Appendix D. *Model Adequacy: Goodness of Fit and Posterior Predictive Check*: the model does not reach $\chi^2/\mathrm{ndof} \approx 1$ (164.1/88), mainly because the signal tails are heavier than a Gaussian, but A_{raw} changes by only -0.0002 with a Crystal Ball signal.

VI. SYSTEMATIC UNCERTAINTIES AND RESULT

In short: thirteen variants in four groups give a systematic uncertainty of 0.0070 *(Bayesian) and* 0.0072 *(frequentist), dominated by the particle-identification requirement; the final result is* $A_{\mathrm{raw}} = -0.0443 \pm 0.0085\,(\mathit{stat}) \pm 0.0070\,(\mathit{syst})$. The systematic study asks how much the result depends on choices of the analysis that the data do not fix. The frequentist and Bayesian fits share one study, so that the two uncertainties are comparable: the same thirteen variants of the nominal choices, fitted by both methods on exactly the same candidates, and the same rule to combine them. The variants (listed in Table II) change one choice at a time and fall in four groups: the *window*, the *background* model, the *signal* model and the *preselection* cuts.

The shift $\Delta = A_{\mathrm{raw}}^{\mathrm{var}} - A_{\mathrm{raw}}^{\mathrm{nom}}$ of a variant is not necessarily a systematic effect: when a variant changes the sample (window, preselection) the two samples are nested and part of Δ is a statistical fluctuation of expected size $\sigma_B = \sqrt{|\sigma^2_{\mathrm{var}} - \sigma^2_{\mathrm{nom}}|}$ [13], and such a variant contributes only if $|\Delta| > \sigma_B$. A variant that the data reject ($\Delta\chi^2 > 25$ on the same data) is excluded: this removes the single exponential ($\Delta\chi^2 = 134$, Appendix D.13). The




FIG. III. Mass distributions of (a) B^+ and (b) B^- candidates with the nominal model (posterior medians, logarithmic scale): total, signal and background components. Bottom: pulls, $(\text{obs} - \text{exp})/\sqrt{\text{exp}}$.

largest remaining $|\Delta|$ of each group is its uncertainty, since the variants of a group are alternative descriptions of the same choice, and the groups, being independent choices, are added in quadrature.

The results are in Table II, and all the Bayesian fits in Table XII of Appendix E.1. The two methods give the same shifts to within ∼0.0005 and the same group uncertainties, with totals of 0.0070 (Bayesian) and 0.0072 (frequentist):

$$\text{Bayesian:}$$
$$A_{\text{raw}} = -0.0443 \pm 0.0085\,(\text{stat}) \pm 0.0070\,(\text{syst}),$$
$$\text{frequentist:}$$
$$A_{\text{raw}} = -0.0441 \pm 0.0085\,(\text{stat}) \pm 0.0072\,(\text{syst}).$$

The preselection dominates, through the single variant `ProbK` > 0.6 $(+0.0058,\ 1.2\,\sigma_B)$. The scan of each cut in steps of 0.05 (Fig. XX of Appendix E) shows that this value is a fluctuation: on the tight side the shifts scatter around zero within $1.3\,\sigma_B$ without a trend, whereas loosening `ProbK` moves A_{raw} monotonically, as expected if misidentified decays not in the mass model enter the sample. The nominal requirement is therefore well placed and the preselection uncertainty conservative. The background group is second, driven by the charge dependence of the background, and the window third.

A. Stability Checks of the Frequentist Fit

While the systematic uncertainties study is shared with the Bayesian approach – and discussed in detail in Sec. VI – three specific stability checks were performed within the frequentist one, to evaluate the robustness of the fitted raw asymmetry A_{raw}. Each test consists of a two-dimensional grid scan in which two selection or fit parameters are varied simultaneously, performing a simultaneous fit with the nominal double-exponential background model at each point and observing the stability of A_{raw}.

1. **Fit Window Scan:** The lower and upper boundaries of the invariant mass fit window, nominally set to $[5100, 5600]$ MeV, were scanned over the ranges $[5000, 5200]$ MeV and $[5350, 6100]$ MeV, respectively, with a 50 MeV step. This test evaluates the sensitivity of A_{raw} to the exponential background parametrization, and is useful to choose the nominal window: too small or too asymmetric fit window produces unstable fit, due to worse background estimation.

2. **Signal Region Scan:** Changing the extremes of the signal region not significant fluctation in A_{raw} are observed, suggesting that the sideband preliminar fit could be avoided.

3. **Particle Identification Cut Scan:** The preselection cuts on the final-state kaon PID likelihoods (`ProbK` and `ProbPi`) were varied simultaneously towards a better kaon purity (larger than 0.5 for kaons, smaller than 0.5 for pions). This influences significantly the A_{raw} estimation, but always within the 68% CI, confirming the cut selection as reasonable.

Across all scan configurations, the maximum observed variation in A_{raw} remains below the statistical uncertainty ($\Delta A_{\text{raw}} < \sigma_{\text{stat}}$), demonstrating the stability of the nominal fit result.

The numbers of the three scans are given in Appendix E.9.




FIG. IV. Posterior of the six physics parameters of the nominal fit (the six background-shape parameters are marginalized). Diagonal: marginal posteriors with the 16th, 50th and 84th percentiles (dashed lines and titles). Off-diagonal: 2D marginals, contours enclosing the regions of the 1σ and 2σ levels of a 2D Gaussian (39% and 86%).




FIG. V. BF_{01} of $A = 0$ and of $A_b = 0$ as a function of the half-width W of the uniform prior $(-W, W)$ (Gaussian approximation of the posterior, Savage-Dickey). The vertical line marks the nominal prior.

VII. VALIDATION OF THE ESTIMATOR AND OF THE INTERVALS

In short: toy-MC experiments generated from the fitted model show that the maximum-likelihood estimate of A is unbiased and attains the minimum variance bound, that the Wilks interval has the nominal coverage, and that the test of $A = 0$ has the expected size and a power that is known as a function of the injected asymmetry.

The asymptotic properties of the maximum-likelihood estimator and of the profile-likelihood ratio were also verified on toy MC experiments generated from the fitted model of Sec. III (Poisson yields, masses drawn from the model), as done for the Bayesian interval in Appendix D.16: 1000 experiments at the fitted value $A_{\text{true}} = -0.0441$ and 300 at each of -0.1, -0.02, 0 and $+0.05$, each fitted with the nominal frequentist procedure (1–3% of the fits are not flagged valid by `MIGRAD` and are excluded).

A. Bias, Efficiency and Coverage

In short: no bias larger than 0.0008, *an estimator spread equal to the Hesse uncertainty within* 6%, *and coverage of* 0.70 ± 0.01 *and* 0.95 ± 0.01 *at the fitted value; the test of* $A = 0$ *has size* 0.051 ± 0.013 *and power* 0.68 *at* $A = -0.02$. Table III summarizes the estimator over the five injected values: the bias of $\hat{A}$ is at most 0.0008 (1.6 standard errors), its spread equals the mean Hesse uncertainty within 6% and the pulls have width 0.94–1.05, so the estimator is unbiased and reaches the minimum variance bound. At the fitted value the coverage of $t(A_{\text{true}}) \leq 1$ is 0.696 ± 0.015 (68.3% level) and

Group	Variant	n	Δ Bayes	σ_B Bayes	Δ freq.	σ_B freq.
Window	$[5050, 5600]$	21164	-0.0003	0.0009	-0.0004	0.0005
	$[5150, 5600]$	18566	-0.0001	0.0007	-0.0005	0.0021
	$[5100, 5450]$	19004	$+0.0002$	0.0009	-0.0001	0.0003
	$[5100, 5700]$	20191	$\mathbf{-0.0014}$	0.0010	$\mathbf{-0.0014}$	0.0001
Background	single exponential (rejected)	19731	-0.0037	–	-0.0039	–
	shape shared between charges	19731	$\mathbf{-0.0037}$	–	$\mathbf{-0.0040}$	–
	symmetric, $A_b = 0$	19731	-0.0028	–	-0.0031	–
Signal	Crystal Ball	19731	$\mathbf{-0.0002}$	–	$\mathbf{-0.0006}$	–
	μ, σ per charge	19731	-0.0000	–	-0.0001	–
Preselection	`ProbK` > 0.4	26553	-0.0032	0.0038	-0.0032	0.0037
	`ProbK` > 0.6	13981	$\mathbf{+0.0058}$	0.0048	$\mathbf{+0.0058}$	0.0050
	`ProbPi` < 0.4	17984	$+0.0022$	0.0022	$+0.0020$	0.0023
	`ProbPi` < 0.6	20953	-0.0004	0.0020	-0.0006	0.0017
Group uncertainties: window / background / signal / preselection			0.0014/0.0037/0.0002/0.0058		0.0014/0.0040/0.0006/0.0058	
Total (quadrature)			0.0070		0.0072	

TABLE II. Systematic study common to the Bayesian and frequentist fits: shift of A_{raw} for each variant and, for the variants that change the sample, the expected statistical spread σ_B of the shift. The largest contributing shift of each group is in bold. The single exponential is rejected by the data and excluded. For the window $[5150, 5600]$ the frequentist uncertainty is scaled from the number of events, since the Hesse matrix of that fit is not accurate.

0.954 ± 0.007 (95.4% level); at $A_{\text{true}} = +0.05$ the 68% coverage is 0.614 ± 0.028, 2.5 standard deviations low, not unexpected among five values.

A_{true}	n	bias	$\text{sd}(\hat{A})/\langle\sigma_{\text{Hesse}}\rangle$	cov. 68%	cov. 95%
-0.1	297	-0.0005 ± 0.0005	0.94	0.71	0.96
-0.0441	985	-0.0001 ± 0.0003	0.97	0.70	0.95
-0.02	290	-0.0008 ± 0.0005	1.02	0.69	0.94
0	293	$+0.0002 \pm 0.0005$	1.01	0.69	0.95
$+0.05$	293	-0.0004 ± 0.0005	1.05	0.61	0.96

TABLE III. Toy-MC experiments at five injected values of the asymmetry: number of valid fits, bias of $\hat{A}$ with its standard error, spread of the estimates relative to the mean Hesse uncertainty (0.0085), and coverage of the intervals $t(A_{\text{true}}) \leq 1$ and ≤ 3.84.

Size and power. Under $A_{\text{true}} = 0$ the test $q(0) > 3.84$ rejects in 0.051 ± 0.013 of the toy experiments, the nominal size, and $q(0)$ is compatible with χ^2_1 (Kolmogorov-Smirnov $p = 0.79$). Figure VI gives the power: 1.00 at the fitted value and 0.68 (0.27 at 3σ) at $A_{\text{true}} = -0.02$, so with 19,731 candidates an asymmetry below ~ 0.02 is not resolved. Together, the estimator is unbiased and efficient, the interval has its nominal coverage and the test its nominal size; the 5.2σ of the data is an extrapolation of the χ^2_1 tail, which the 293 experiments check only down to the 3σ level (Appendix F.4).

Further material is in Appendix F.

VIII. LOCAL ASYMMETRY IN THE DALITZ PLANE

In short: the asymmetry is not uniform over the Dalitz plane ($\chi^2 = 80.8$ for 31 degrees of freedom); it is concentrated at low m^2_{lo} outside the $\phi(1020)$ and, in the region of the published analysis of the same data, it reproduces the published value. To test whether the asymmetry is uniform over phase space or concentrated in specific regions, each candidate was placed in the Dalitz plane using the two opposite-sign K^+K^- invariant masses $m^2(H1, H2)$ and $m^2(H1, H3)$ (Sec. II A), sorted into m^2_{lo} and m^2_{hi} to remove the $H2/H3$ labeling ambiguity. The plane $[0, 16] \times [0, 30]$ GeV^2 in $(m^2_{\text{lo}}, m^2_{\text{hi}})$ was divided into an 8×8 grid, fixed before looking at the asymmetries. In each bin with at least 20 candidates (32 of 64), the local asymmetry $A_i = (N_i^- - N_i^+)/N_i$, with binomial uncertainty $\sigma(A_i) = \sqrt{(1 - A_i^2)/N_i}$, was compared with the global counting asymmetry $A_{\text{glob}} = -0.043$ through $S_i = (A_i - A_{\text{glob}})/\sigma(A_i)$.




FIG. VI. Probability to reject $A = 0$ as a function of the injected asymmetry, with the test $q(0) > 3.84$ (size 5%, dotted line) and $q(0) > 9$ (3σ); 1000 toy experiments at -0.0441, about 300 at each other value.

Figure VII shows the occupancy of the bins (a), the




FIG. VII. Fixed 8×8 Dalitz-plot binning, all candidates in the fit window: (a) number of candidates per bin, (b) local asymmetry A_i for bins with ≥ 20 candidates, and (c) local significance with respect to the global asymmetry, $(A_i - A_{\rm glob})/\sigma(A_i)$. The two bins with $|S_i| > 3$ are outlined. Dashed lines: the squared masses of the D^0, J/ψ and χ_{c0}, in either opposite-sign pair.

local asymmetry (b) and its significance (c). Two of the 32 bins have $|S_i| > 3$, adjacent at $m^2_{\rm lo} < 4$ GeV^2 and $7.5 < m^2_{\rm hi} < 11.25$ GeV^2 ($S_i = -3.7$ and -3.0); about 0.09 such bins would be expected if the asymmetry were uniform, and two or more with probability 3×10^{-3}, a look-elsewhere number over any two of the 32 bins. The global homogeneity test, $\chi^2 = \sum_i S_i^2 = 80.8$ for 31 degrees of freedom ($p = 3 \times 10^{-6}$), confirms that the local asymmetry is not uniform; it also holds in the signal window alone, with the charm vetoes and for other grids (Appendix G).

The structure is the same with both magnet polarities: in the region of the two significant bins the counting asymmetry is -0.129 ± 0.023 (magnet down) and -0.126 ± 0.028 (magnet up), and the two maps agree bin by bin ($\chi^2 = 23.0$ for 29 bins, $p = 0.78$). The split tests the part of the detection asymmetry that changes sign with the polarity, not the different interaction of K^+ and K^- with the detector material.

Where is the asymmetry in terms of the decay dynamics? The ordered Dalitz plot (Fig. VIIIa) shows the $\phi(1020)$ band at the kinematic edge ($m^2_{\rm lo} \approx 1.04$ GeV^2) and the charm bands of the D^0, J/ψ and χ_{c0}. The negative asymmetry is not produced by the $\phi(1020)$: its band (1964 candidates) has $A_{\rm count} = -0.004 \pm 0.023$, while the significant bins without it have -0.145 ± 0.019 (Fig. XXVI of Appendix G). It sits just above the ϕ, where the published analysis located it, defining the region $1.2 < m^2_{\rm lo} < 2.0$ GeV^2, $m^2_{\rm hi} < 15$ GeV^2 with $A_{\rm raw} = -0.239 \pm 0.020$ [5]. The region is fixed by the paper, not chosen on our data, so the comparison is unbiased: the nominal model fitted to its 2146 candidates gives $A_{\rm raw} = -0.234 \pm 0.025$ (Bayesian) and -0.237 ± 0.025 (frequentist, 9.4σ from zero), reproducing the published value (spectra of the two charges in Fig. VIIIb, where the B^- peak is visibly smaller).

Three caveats limit the interpretation: the binning is coarse and was not optimized, the asymmetries are counts without sWeight background subtraction (Appendix J.5), and local variations can be partly instrumental, since the $K^\pm$ interaction asymmetry depends on momenta that are correlated with the Dalitz position. In summary, the global asymmetry is not uniform over phase space, and the region identified by the published analysis carries it.

Further material is in Appendix G.

IX. COMPARISON AND DISCUSSION

In short: the frequentist and Bayesian analyses give the same value, uncertainty and coverage for $A_{\rm raw}$ because they use the same likelihood and the data constrain it well; they differ in what the numbers mean and in the hypothesis tests, and a machine-learning selection (Appendix I) confirms the result without improving it.

The frequentist and Bayesian analyses agree on every number and differ in what each number means. Table IV puts the two approaches side by side, from the question each one asks to how it was validated; the subsections explain the rows where the numbers coincide but the statements do not.

A. Where the Approaches Differ

In short: the numbers coincide but the statements differ: the meaning of the interval, the test of $A = 0$, the goodness of fit, the treatment of the nuisance parameters and what has been validated, as summarized in Table IV. Meaning of the interval. The credible interval states that, given the data and the prior, A lies in $[-0.0528, -0.0358]$ with 68% probability; the confidence interval $[-0.0526, -0.0356]$ is the set of values that the data do not reject at the 32% level, and the 68% is the fraction of repeated experiments in which such an interval contains the true value, a property of the procedure that the bootstrap and toy checks verified. Fig-




FIG. VIII. (a) Ordered Dalitz plot of the candidates in the fit window, with the squared masses of the $\phi(1020)$, D^0, J/ψ and χ_{c0} (dashed) and the region of the published analysis of the same data [5] (box). (b) Mass spectra of B^+ and B^- candidates in that region, same binning: the B^- peak is visibly smaller.

	Frequentist (Sec. IV)	Bayesian (Sec. V)
Question	Which value of A makes the data most probable, and which values are not rejected?	What is the distribution of A given the data and the prior?
Estimate	Maximum likelihood, -0.0441	Posterior median, -0.0443 (mode -0.0441)
Interval (68%)	$[-0.0526, -0.0356]$ (profile likelihood, Minos)	$[-0.0528, -0.0358]$ (credible, equal-tailed)
Nuisance parameters (11)	profiled: re-minimized at each fixed A	marginalized: integrated over the prior
Test of $A = 0$	$q(0) \simeq 27$, $p = 2.05 \times 10^{-7}$, 5.2σ	$BF_{01} \approx 1.1 \times 10^{-4}$ (Savage-Dickey)
Validation of the interval	1000 bootstrap replicas: $68.4 \pm 1.5\%$; 985 toys: $70 \pm 1\%$ (68%), $95 \pm 1\%$ (95%)	200 toys: 0.65 ± 0.03 (68%), 0.93 ± 0.02 (95%)
Goodness of fit	χ^2/ndof $\approx$ 160/88 (10 MeV bins); 441.7/388, $p = 0.031$ (2.5 MeV bins)	164.1/88 ($p = 1.6 \times 10^{-6}$); posterior predictive check
Computing time	one joint fit with Minos $\simeq 19$ s; 1000 bootstrap fits $\simeq 49$ min	one chain (25,000 steps, 32 walkers) $\simeq 8$ min; 200 toys of 12,000 steps

TABLE IV. Frequentist and Bayesian analyses compared on the same candidates ($A_{\rm raw}$, statistical uncertainties only). Computing times are indicative, on one laptop.

ure IX shows the two for A: the profile likelihood and the marginal posterior coincide (a), and the profile curves of the two magnet polarities add up to that of the joint fit (b).

Hypothesis test. Three tests of $A = 0$ are available. The exact binomial test of $N^- = N^+$ on all candidates gives 6.1σ ($A_{\rm count} = -0.0432 \pm 0.0071$); it sees signal and background together and does not use the mass. The frequentist test of the fit gives $p = 2 \times 10^{-7}$ ($Z = \sqrt{q(0)} = 5.2\sigma$ in the two-sided convention, 5.06σ one-sided, Appendix C.1), independent of the prior and not the probability of $A = 0$; its size and power were verified with toy MC (Appendix F.4).

The Bayes factor, $BF_{10} \approx 9 \times 10^3$, depends on the prior width (Sec. V D). For A the frequentist and Bayesian tests give the same conclusion; for the background asymmetry, $A_b = -0.040 \pm 0.018$, a frequentist test would call 2.3σ a hint while $BF_{01} \approx 3.5$ mildly favors $A_b = 0$: the p-value asks how improbable the data are under $A_b = 0$, the Bayes factor which hypothesis the data prefer after the prior volume is accounted for.

B. Comparison with the Published Analysis

In short: the published value (-0.060 ± 0.007) differs from this one (-0.0441 ± 0.0085) by the sample and the model; with the same D^0 veto the Bayesian fit gives -0.0496 ± 0.0088, and $A_{\rm raw}$ is not A_{CP}. The published analysis [5] finds $A_{\rm raw} = -0.060 \pm 0.007$ with $22{,}119 \pm 164$ signal candidates, against -0.0441 ± 0.0085 and $15{,}328 \pm 133$ here: the Open Data record is a preselected subset and the published model has a Cruijff signal, partially reconstructed four-body decays and a misidentified peaking background from simulation. With the same D^0 veto the Bayesian fit gives -0.0496 ± 0.0088, within 0.93σ of the published value.

The raw asymmetry is not the CP asymmetry: $A_{CP} \approx$




FIG. IX. (a) Frequentist profile likelihood, $\exp[-q(A)/2]$ normalized, and Bayesian marginal posterior of A. (b) Profile curves $q(A)$ of the two magnet polarities (parabolic fits to the points), their sum and the joint fit.

$A_{\rm raw} - A_P - A_D$ (Eq. (2)), and in the published analysis the corrections move the result from -0.060 to $-0.043 \pm 0.009 \pm 0.003 \pm 0.007$ (3.7σ) [5], by $+0.017$, about two statistical uncertainties of this work. Such a correction cannot be assumed for a different selection, and the 5.2σ here is the significance of a nonzero raw asymmetry, not of CP violation. A later measurement on a larger data set gives $A_{CP} = -0.037 \pm 0.002 \pm 0.002 \pm 0.003$ (8.5σ) [1, 2]; our result is consistent with these in sign and size.

C. Machine-Learning Cross-Check

In short: a classifier trained on weak labels from the mass spectrum selects candidates without changing $A_{\rm raw}$ beyond its statistical spread; it does not improve the precision and cannot remove the misidentified decays under the signal (Appendix I). A cut at 0.5 on the classifier score keeps 92% of the fitted signal and 36% of the background and gives $A_{\rm raw} = -0.0451 \pm 0.0087$ (Bayesian) and -0.0447 ± 0.0086 (frequentist), against -0.0443 and -0.0441 for the full window: both shifts are within the statistical spread expected for a subsample (Appendix I.9, Fig. XXXIX). The precision does not improve because the nominal sample has little background to remove; the classifier is described in Appendix I.

Further material is in Appendix H.

X. CONCLUSIONS

In short: two inference frameworks give the same raw asymmetry, $A_{\rm raw} = -0.044 \pm 0.009\,(stat) \pm 0.007\,(syst)$, 5.2 standard deviations from zero, with validated intervals; what separates them is the meaning of the numbers and the tests, not their values; a machine-learning selection (Appendix I) confirms the result without improving it.

The answer to the question of the Introduction is that, on this problem, the frequentist profile-likelihood fit and the Bayesian posterior are equivalent in value, uncertainty and coverage. With a common selection and mass model on the 2011 LHCb Open Data of $B^\pm \to K^\pm K^+ K^-$ they give $A_{\rm raw} = -0.0441 \pm 0.0085 \pm 0.0072$ and $-0.0443 \pm 0.0085 \pm 0.0070$ (statistical, systematic), a nonzero asymmetry at 5.2 standard deviations, stable under the thirteen variants of the fit and selection, compatible in sign and size with the published analysis, and concentrated in its Dalitz region outside the $\phi(1020)$. It is a raw asymmetry, not a measurement of A_{CP} and not a significance of CP violation (Sec. IX B).

The methods differ in what they add: the frequentist fit an unbiased estimator at the minimum variance bound with intervals of nominal coverage; the Bayesian analysis the same interval with flat priors, plus sampling diagnostics, a Bayes factor that depends on the prior width and a posterior predictive check. The machine-learning selection leaves $A_{\rm raw}$ unchanged. The dominant systematic contribution is the particle-identification requirement, in both methods. Three limitations bound the result (Appendix J.5): no tested model reaches $\chi^2/{\rm ndof} \approx 1$, although $A_{\rm raw}$ does not depend on the signal shape; no production or detection correction is applied, so $A_{\rm raw} \neq A_{CP}$; and the local Dalitz-plane asymmetries are counts without background subtraction by event weights.

Appendix A: Dataset and Preselection

In short: the record and its branches, the cut flow of the preselection, and the candidates and charge combinations that the main text treats in one line.

1. Description of the Record

In short: the record holds 8,556,118 *candidates of* $B^{\pm} \to h^{\pm}h^{+}h^{-}$ *collected in 2011, in two files by magnet polarity, plus a detector-free phase-space simulation of* $B^{\pm} \to K^{\pm}K^{+}K^{-}$ *used only to test the local-asymmetry procedure.* The dataset is the CERN Open Data record 4900, "Matter Antimatter Differences – B meson decays to three hadrons" [3]: candidates for $B^{\pm} \to h^{\pm}h^{+}h^{-}$ decays collected by LHCb in 2011 at $\sqrt{s} = 7\,\text{TeV}$, reconstructed with the LHCb software and already passed through trigger and offline selection. The data are distributed as ROOT `TTree`s (`DecayTree`) containing only the most relevant reconstructed branches, split across two files by magnet polarity (magnet up/down), for a combined 8,556,118 candidates. The same record also includes a simplified, detector-response-free phase-space simulation of $B^{\pm} \to K^{\pm}K^{+}K^{-}$ (50,000 decays, half of each charge); it is used only as a sample without asymmetry to test the local-asymmetry procedure (Sec. VIII).

2. Branches and Preselection

In short: the analysis reads momentum, particle-identification likelihoods, muon flag and charge of each track; the common preselection keeps 23,895 *candidates (*0.28%*), mostly through the kaon-identification requirement.* For each of the three final-state tracks ($H1$, $H2$, $H3$) the analysis reads momentum components (`PX`, `PY`, `PZ`), particle-identification (PID) likelihoods (`ProbK`, `ProbPi`), a muon flag (`isMuon`), and the track charge. The same preselection is used in all three analyses: a kaon-PID requirement and a muon veto on every track,

$$\texttt{H*_ProbPi} < 0.5 \ \wedge \ \texttt{H*_ProbK} > 0.5 \ \wedge \ \neg\texttt{H*_isMuon},$$

which retains 23,895 of the 8,556,118 candidates (0.28%), the PID requirement being by far the dominant reduction (Table V of Appendix A). The files contain no event identifier; as a proxy for multiple candidates from the same collision, only 42 of the candidates in the fit window (0.2%) share an identical track with another candidate, a negligible effect on the statistical uncertainties, and all are kept.

3. Cut Flow of the Preselection

In short: the kaon-identification requirement on the three tracks is by far the most selective.

Requirement	alone	cumulative	down	up
All candidates	–	8,556,118	5,135,823	3,420,295
¬`isMuon`, three tracks	6,311,517	6,311,517	3,806,297	2,505,220
`ProbPi` < 0.5, three tracks	975,724	481,101	288,908	192,193
`ProbK` > 0.5, three tracks	28,812	23,895	14,178	9717
Mass window [5100, 5600] MeV	–	19,731	11,636	8095

TABLE V. Number of candidates passing each requirement alone and after the cumulative selection, in the order of the table, and by magnet polarity. The kaon-identification requirement on the three tracks is by far the most selective.

4. Candidates Sharing a Track

In short: only 42 *candidates of the fit window (*0.2%*) share a track with another candidate, a negligible effect, and all are kept.* The files have no event identifier; only 42 candidates of the fit window (0.2%) share a track with another candidate, a negligible effect, and all are kept.

5. Charge Combinations

In short: $H1$ is always the kaon of charge opposite to the B and $H2$, $H3$ the same-sign pair, so the Dalitz variables are the two opposite-sign invariant masses. In this dataset only the charge combinations $(+1, -1, -1)$ and $(-1, +1, +1)$ occur for $(H1, H2, H3)$: $H1$ is always the kaon of charge opposite to the B, and $H2$, $H3$ are the interchangeable same-sign pair. The Dalitz-plot variables are therefore the invariant masses of the two opposite-sign pairs, $m^2(H1, H2)$ and $m^2(H1, H3)$.

6. Invariant Mass and Fit Window

In short: the mass is computed with the kaon hypothesis on all tracks; the fit window $[5100, 5600]$ *MeV holds* 19,731 *candidates and includes the start of the rise of the background below* $\sim$ 5150 *MeV.* The three final-state tracks are assigned the charged-kaon mass hypothesis ($M_K = 493.677$ MeV) regardless of their PID likelihoods, and their four-momenta are summed to reconstruct the B candidate, $m(KKK) = \sqrt{(\sum_i E_i)^2 - |\sum_i \vec{p}_i|^2}$. The nominal fit window, common to the frequentist and Bayesian analyses, is $[5100, 5600]$ MeV. Below $\sim$5150 MeV the background rises, consistent with a partially reconstructed contribution that a single exponential cannot describe; the window includes the start of this rise, which is described by a second exponential in the background model (Secs. IV–V). This leaves 19,731 preselected candidates in the fit window: 10,292 with B-charge $+1$ and 9439 with B-charge -1, split 11,636 (magnet down) / 8095 (magnet up). The machine-learning selection of Appendix I uses the same window, with the region below 5150 MeV excluded from its background-like training sample (Appendix I.4).

Appendix B: Mass Model and Likelihood

In short: the extended likelihood and its relation to a mixture, the label switching of the two background slopes,

and the six-parameter model that preceded the nominal one.

1. Extended Likelihood

In short: with a Poisson total number of candidates the likelihood is $e^{-(N_s+N_b)}\prod_i[N_sS+N_bB]$ *up to a constant, which constrains the yields and A together.* If the total number of candidates N is Poisson-distributed with mean $\nu = N_s + N_b$ and each candidate is drawn independently from the normalized density $f(m,q) = [N_sS + N_bB]/\nu$, the likelihood of the sample is $\frac{\nu^N e^{-\nu}}{N!}\prod_i f(m_i,q_i) = \frac{e^{-\nu}}{N!}\prod_i[N_sS(m_i) + N_bB(m_i)]$. The factor $1/N!$ does not depend on the parameters and is dropped, which gives the extended likelihood of Eq. (4). Unlike the likelihood with a fixed total, it constrains the yields N_s and N_b through N itself, and the asymmetry A is estimated together with them.

2. Extended Likelihood versus a Mixture with Fixed Total

In short: the extended form differs from a mixture with fixed total only by the Poisson term of the total yield, which carries no information on A; the purity of the signal window is about 96%. The extended form differs from a mixture of signal and background with a fixed total number of candidates and a signal fraction only through the Poisson term for the total yield, which carries no information on A and does not change its estimate; it is used because the yields of the two charges are the natural parameters of the model. Bayes' theorem applied to the fitted yields gives the probability that a candidate in the signal window $|m-\mu| < 2\sigma$ is a signal decay, about 96%: a purity, not a property of the single candidate, which depends on the signal fraction of the sample (here 78% of the candidates in the fit window).

3. Label Switching of the Background Components

In short: the mixture is invariant under the exchange of the two exponentials, which would give two identical modes; the ordering $\lambda_1 > \lambda_2$ *removes the symmetry.* The mixture is invariant under the exchange $(\lambda_1, f) \leftrightarrow (\lambda_2, 1-f)$, which would give the posterior two identical modes (label switching); the ordering $\lambda_1 > \lambda_2$ removes this symmetry.

4. Minimal Model as a Cross-Check

In short: a six-parameter model on a narrower window gives $A_{\rm raw} = -0.0497$ *(68% CI* $[-0.058, -0.041]$*), compatible with the nominal result within its uncertainty; it is the model used for the low-statistics comparison.* Before the nominal model was fixed, the same analysis was carried out with a minimal six-parameter model: a narrower window, $[5150, 5450]$ MeV, which excludes most of the rising background below the peak, a single exponential background with a shape and a yield shared by the two charges, and the same Gaussian signal. On its 17,839 candidates it gives $A_{\rm raw} = -0.0497$ (68% CI $[-0.058, -0.041]$) with $\chi^2/\text{ndof} = 127.4/54$, and -0.0457 when the background yields of the two charges are left free. Its toy validation, with 500 toy-MC datasets, gives coverage consistent with nominal (0.664 and 0.962 for the 68% and 95% intervals). The difference from the nominal result, 0.005, is driven by the background treatment, as in the systematic study of Sec. VI, and is smaller than the statistical uncertainty.

Appendix C: Frequentist Analysis

In short: what the Hesse and Minos uncertainties and the test of $A = 0$ *are, the Neyman construction, the uncertainties of the twelve parameters by three methods, and the behavior of the fit from different starting values.*

1. Hesse Uncertainties, Test of $A = 0$ and Significance

In short: the Hesse matrix is the inverse of the Fisher information, the test statistic is the profile likelihood ratio, and the significance is $Z = \sqrt{q(0)}$.

a. Hesse uncertainties. For n independent candidates the likelihood contains the information on the parameters θ through the Fisher information matrix, $I_{ij} = -E\big[\partial^2 \ln L/\partial\theta_i\partial\theta_j\big]$. The information inequality (minimum variance bound) states that the variance of an estimator cannot be smaller than the inverse of I (for an unbiased one), and the maximum-likelihood estimate reaches the bound asymptotically, where it is also unbiased and Gaussian [14]. The covariance matrix of the fit is therefore $V \simeq I^{-1}$ evaluated at the best-fit point, with the second derivatives of $-\ln L$ computed numerically: this is the Hesse matrix of `iminuit`, and $\sigma(A) = 0.0085$ is the square root of the diagonal element of V for A, which includes the effect of the eleven nuisance parameters. It assumes that $-\ln L$ is parabolic around the minimum. The Minos intervals do not: they follow the profile likelihood on each side of the minimum, and their agreement with the Hesse error (-0.0085, $+0.0085$) shows that the parabola is a good description here.

b. Test of $A = 0$. The null hypothesis H_0: $A = 0$ is tested against the alternative H_1: A free, the other parameters being free in both. The test statistic is the profile likelihood ratio, $q(0) = -2\ln\big[L(0,\hat{\hat{\theta}})/L(\hat{A},\hat{\theta})\big] = 2\Delta\text{NLL}$ (NLL: negative log-likelihood), large when the data disfavor H_0. For two simple hypotheses the likelihood ratio is the most powerful test of a given size,

i.e. the one with the highest probability of rejecting H_0 when H_1 is true (Neyman-Pearson lemma [15]); the profile likelihood ratio is its extension to hypotheses with unspecified parameters, with no optimality guarantee in general. By Wilks' theorem [8], under H_0 the statistic follows a χ^2 distribution with one degree of freedom, the number of parameters fixed by H_0, whatever the number of nuisance parameters, so the p-value is $P(\chi^2_1 \geq q_{\rm obs})$; this is the probability, under H_0, of a result at least as discrepant as the data, not the probability of H_0.

c. Significance. A χ^2_1 variable is the square of a standard normal one, so $p = P(|N| \geq \sqrt{q})$ and the significance quoted is $Z = \sqrt{q(0)}$ ($q \simeq 27$, $p = 2.05 \times 10^{-7}$, $Z = 5.20$). This is a two-sided convention: the one-sided one, $Z = \Phi^{-1}(1-p)$, applied to the same p would give 5.06. For a parabolic likelihood $\sqrt{q}$ coincides with the Hesse significance $|\hat{A}|/\sigma(A) = 5.19$.

2. Neyman Construction and Dependence on the Momentum

In short: inverting the test of $A_{\rm true}$ with toy-MC experiments gives the same interval as the likelihood ratio, and $A_{\rm raw}$ does not depend on the momentum of the B. The confidence interval is, by definition, the set of $A_{\rm true}$ not rejected by the data. It was constructed directly (Fig. X): for 11 values of $A_{\rm true}$ between -0.1 and 0, 300 toy-MC experiments give the central 68% and 95% ranges of $\hat{A}$ (the belt), and the interval is the set of $A_{\rm true}$ whose 68% range contains the observed $\hat{A} = -0.0441$: $[-0.0528, -0.0351]$, against $[-0.0526, -0.0356]$ from the profile likelihood (Minos), which is the content of Wilks' theorem for this problem.




FIG. X. Confidence belt of $\hat{A}$ from toy-MC experiments (central 68% and 95% ranges) as a function of $A_{\rm true}$; the vertical line is the observed $\hat{A}$ and the dashed lines the resulting 68% interval.

The detection asymmetry of kaons depends on their momentum; as a direct check, the joint fit was repeated in three bins with equal numbers of candidates of the transverse momentum p_T and of the momentum p of the B candidate (Table VI): the three values are compatible with the global one, with a largest deviation of 1.4σ.

Bin	range [GeV]	n	$A_{\rm raw}$
p_T, 1	1.0–4.7	6577	-0.0563 ± 0.0169
p_T, 2	4.7–8.2	6577	-0.0362 ± 0.0141
p_T, 3	8.2–30.5	6577	-0.0452 ± 0.0134
p, 1	26.7–78.1	6577	-0.0391 ± 0.0139
p, 2	78.1–110.1	6577	-0.0475 ± 0.0149
p, 3	110.1–288.5	6577	-0.0478 ± 0.0153

TABLE VI. $A_{\rm raw}$ (joint frequentist fit) in terciles of the transverse momentum and of the momentum of the B candidate.

The values of Table VI are plotted in Fig. XI against the global result: no trend with p_T or p is visible.

3. Uncertainties of the Twelve Parameters by Three Methods

In short: the inverse of the Fisher information (Hesse), the bootstrap of the data and the posterior give the same standard deviations for A, A_b, μ and σ; the bootstrap differs for the two yields, because it keeps the total number of candidates fixed, and for the steep slope λ_1^+, which is weakly constrained. The covariance of the fit parameters was estimated in three ways (Table VII): the inverse of the matrix of second derivatives of $-\ln L$ at the minimum (Hesse), 300 bootstrap replicas of the data, each refitted, and the posterior sample. The posterior agrees with Hesse to 10% for all the parameters, and the bootstrap does for A, A_b, μ, σ and the other shape parameters. The exceptions of the bootstrap are the steep slope of the B^+ background, λ_1^+ (0.0057 against 0.0077, a weakly constrained and non-parabolic direction, Appendix C.4), and the signal and background yields, $\sigma(N_s) = 133$ (Hesse) and 72 (bootstrap): a resampling of N candidates keeps the total yield fixed, which the extended likelihood does not, and the yields of the bootstrap show the anticorrelated fluctuation of the binomial split; the asymmetry A is not affected. The correlation between A and A_b is -0.16, -0.17 and -0.17 in the three methods.

Figure XII shows the same comparison as the ratio to the Hesse uncertainty. The posterior agrees with Hesse within 10% for all parameters; the bootstrap does too, except for the yields (the bootstrap of the data keeps the total number of candidates fixed, which removes the Poisson fluctuation of N_s) and λ_1^+.




FIG. XI. A_{raw} (joint frequentist fit) in terciles of (a) the transverse momentum and (b) the momentum of the B candidate, with the global value and its $\pm 1\sigma$ band; each bin has 6577 candidates.




FIG. XII. Standard deviation of the twelve parameters of the nominal fit from the bootstrap of the data (300 replicas) and from the posterior, divided by the Hesse uncertainty; the band is ±10%.

Parameter	Hesse	Bootstrap	Posterior
A	0.0085	0.0085	0.0085
A_b	0.0177	0.0183	0.0176
N_s	133	72	133
N_b	82	72	82
μ [MeV]	0.159	0.159	0.161
σ [MeV]	0.141	0.150	0.138
λ_1^+	0.0077	0.0057	0.0078
λ_2^+	0.00028	0.00030	0.00029
f_+	0.028	0.029	0.030
λ_1^-	0.0047	0.0043	0.0047
λ_2^-	0.00040	0.00038	0.00039
f_-	0.039	0.036	0.037

TABLE VII. Standard deviation of the parameters of the nominal fit from the Hesse matrix, from the bootstrap of the data and from the posterior.

4. Starting Values and Weakly Constrained Parameters

In short: 29 of 30 fits from random starting points reach the same minimum; the steep-background fraction is the weakly constrained direction, as the long chain of the Bayesian fit already showed. The nominal fit was repeated from 30 starting points drawn at random (yields and slopes multiplied by a log-normal factor of width 0.5, the fractions uniform, the asymmetry uniform in $(-0.2, 0.2)$): 29 reach the minimum of the nominal fit ($\hat{A} = -0.0441$); the remaining one stops in a local minimum 96 units of $-2\ln L$ above, at $\hat{A} = -0.052$, which would be rejected by comparing the likelihood values. The profile likelihood of the fraction of the steep component for B^+ (the parameter whose chain mixes slowly in the narrow window, Sec. D.13) is asymmetric: $q(f_+) = 1$ at 0.085 and 0.140 around the best value 0.112, with a longer tail towards large fractions where the two exponentials exchange their roles. The uncertainty of A does not depend on it (correlation below 0.1).

Appendix D: Bayesian Analysis

In short: the sampling and the posterior summaries behind Sec. V, the Bayes factors with their derivation, the model-adequacy and robustness checks, the toy-MC coverage of the credible interval and the alternative sampler.

1. Sampling the Posterior

In short: Metropolis-Hastings accepts a proposal with a ratio that satisfies detailed balance; the affine-invariant ensemble move samples correlated posteriors without tuning, and N/τ is the effective length of a chain. The Metropolis-Hastings algorithm builds a Markov chain whose stationary distribution is the target $\pi(\theta)$: from the state x a point y is proposed from $q(y \mid x)$ and accepted with probability $\min\big[1, \pi(y)\,q(x \mid y)/(\pi(x)\,q(y \mid x))\big]$, which satisfies detailed balance. Its efficiency depends on how the proposal is matched to the (here correlated) posterior. The ensemble sampler used for the nominal fit [9, 10] evolves K walkers; to update the walker X_k it picks another walker X_j and proposes $Y = X_j + Z\,(X_k - X_j)$, with $Z \in [1/a, a]$ drawn from $g(z) \propto z^{-1/2}$ ($a = 2$), and accepts it with probability $\min\big[1, Z^{d-1}\pi(Y)/\pi(X_k)\big]$ (d the number of parameters). The move is invariant under affine transformations of the parameters, so a strongly correlated, elongated posterior is as easy to sample as a spherical one, and it needs no hand-tuned proposal covariance.

Successive states of a chain are correlated. The integrated autocorrelation time of a parameter is $\tau = 1 + 2\sum_{t\geq 1}\rho(t)$, with $\rho(t)$ the autocorrelation of the chain at lag t, and a chain of N steps contains $N_{\text{eff}} = N/\tau$ effectively independent samples, so that the Monte Carlo error of a posterior mean is $\sigma/\sqrt{N_{\text{eff}}}$. The estimate of τ is itself reliable only for $N \gtrsim 50\,\tau$ [10].

2. Posterior Summaries and Model Comparison

In short: equal-tailed and HPD intervals, the Jeffreys scale for Bayes factors, the Ockham effect of the prior width and the definition of the posterior predictive p-value. The credible intervals quoted are the equal-tailed ones (the 16%–84% quantiles for 68%); the highest-posterior-density (HPD) interval is the shortest interval with the same probability and coincides with it for a symmetric posterior. A Bayes factor $BF_{ij} = P(D \mid H_i)/P(D \mid H_j)$ is interpreted with the Jeffreys scale [12, 16]: values between 1 and 3 are barely worth mentioning, 3–10 substantial, 10–30 strong, 30–100 very strong and above 100 decisive. The evidence automatically penalizes unnecessary freedom (Ockham's razor): for a Gaussian posterior much narrower than a uniform prior of width $2W$ the Savage-Dickey ratio of $A = 0$ is proportional to W, so widening the prior of A favors the null hypothesis by construction, which is why the dependence on W is reported.

The posterior predictive check compares the data y with replicas y^{rep} simulated from the model at parameters drawn from the posterior, through a discrepancy statistic T: the posterior predictive p-value is $p_B = P\big(T(y^{\text{rep}}, \theta) \geq T(y, \theta) \mid y\big)$ [17]. Values close to 0 or 1 signal a feature of the data that the model does not reproduce; a statistic that the fit adjusts directly (for example the total asymmetry, when A is a free parameter) is reproduced by construction and tests nothing.

3. Flat Priors and the Evidence

In short: a flat prior is not invariant under reparametrization, but with $\sim 2 \times 10^4$ events the posterior is dominated by the likelihood; the evidence is a constant that neither the MCMC nor the Savage-Dickey ratio needs. A flat prior is not uninformative in an absolute sense, since it is not invariant under reparametrization (a prior uniform in λ is not uniform in $1/\lambda$); with $\sim 2 \times 10^4$ events, however, the posterior is dominated by the likelihood, as the prior-sensitivity check of Sec. D.15 confirms for A. The evidence $p(D)$ is a constant normalization: the MCMC samples $\mathcal{L}\,\pi$ without computing it, and the Savage-Dickey ratio of Sec. V D avoids computing it as well. The marginal posterior of the asymmetry, $p(A \mid D) = \int p(\theta \mid D)\, d\theta_{\neq A}$, is the histogram of the A component of the chain.

4. Sampling and Convergence Diagnostics

In short: 32 walkers for 25,000 steps give $N/\tau \geq 113$ autocorrelation times for every parameter, and runs started over the whole prior box converge to the same single mode. The posterior was sampled with the affine-invariant ensemble sampler of Goodman and Weare [9] as implemented in `emcee` [10] (the stretch move, the autocorrelation time and the effective sample size are described in Appendix D.1). The maximum of the posterior was first located numerically (Nelder-Mead), and 32 walkers were initialized in a narrow ball around it and run for 25,000 steps, with the first 4166 (1/6) discarded as burn-in, leaving 6.7×10^5 samples; the mean acceptance fraction is 0.38. Convergence was checked through the integrated autocorrelation time of each parameter, $\tau = 1 + 2\sum_{t\geq 1}\rho(t)$, with $\rho(t)$ the autocorrelation of the chain at lag t. We find $\tau \in [147, 183]$ steps, the longest for the background-shape parameters: the post-burn-in chain is $N/\tau \geq 113$ autocorrelation times long for every parameter, above the threshold of 50 recommended for a reliable estimate of τ [10], and corresponds to $\gtrsim 3600$ effectively independent samples per parameter. To check that the chain is not trapped near its starting point, the fit was repeated twice with the walkers initialized uniformly over the whole prior box and the first half of the chain discarded: both runs converge to the same single mode, with $A_{\text{raw}} = -0.0446$ and -0.0442, the same intervals and the same values of all other parameters as the nominal fit. The trace plot of such a run (Fig. XIII, Appendix D.5) shows all walkers reaching the bulk of the posterior within $\sim$2000 steps, the background fraction f_+ last, well inside the discarded half of the chain.

5. Trace Plots of the Walkers

In short: walkers started over the whole prior box converge to the same mode.

6. Posterior of the Raw Asymmetry

In short: $A_{\rm raw} = -0.0443$ *with* 68% *and* 95% *intervals* $[-0.0528, -0.0358]$ *and* $[-0.0608, -0.0276]$*; the posterior is Gaussian, so equal-tailed and highest-density intervals coincide and its mode is the maximum-likelihood estimate.* The marginal posterior on the raw asymmetry gives (median and equal-tailed credible intervals, CI)

$$\begin{aligned} A_{\rm raw} &= -0.0443, \\ 68\%\ \text{CI}\ &[-0.0528, -0.0358], \\ 95\%\ \text{CI}\ &[-0.0608, -0.0276]. \end{aligned}$$

The posterior is very close to Gaussian, with standard deviation 0.0085; its 68% interval is symmetric about the median to within 2×10^{-5}, so the equal-tailed intervals coincide with the highest-posterior-density ones at the quoted precision. Since the priors are uniform, the posterior mode (the maximum a posteriori, MAP, estimate, -0.0441) coincides with the maximum-likelihood estimate, so this result can be compared directly with the frequentist fit of the same model (Sec. IV; the quantities it uses are summarized in Appendix C.1). The published analysis of the same 2011 data, with its own selection (which includes a D^0 veto) and a mass model with partially reconstructed and misidentified backgrounds, finds $A_{\rm raw} = -0.060 \pm 0.007$ [5]; with the D^0 veto our fit gives -0.0496 ± 0.0088 (Sec. D.15), compatible within the statistical uncertainties, although the two selections overlap and differ. After the corrections of Eq. (2), the published asymmetry is $A_{CP} = -0.043 \pm 0.009 \pm 0.003 \pm 0.007$ [5], and the most precise later measurement, on a larger data set, is $A_{CP} = -0.037 \pm 0.002 \pm 0.002 \pm 0.003$ [1, 2]. The other physics parameters are $N_s = 15{,}328 \pm 133$, $N_b = 4404 \pm 82$, $A_b = -0.040 \pm 0.018$, $\mu = 5284.39 \pm 0.16$ MeV and $\sigma = 18.00 \pm 0.14$ MeV (Fig. IV). A is almost uncorrelated with the other parameters: its largest correlation, -0.17, is with A_b: conditioning on $A_b = 0$ shifts A by $\rho\,\sigma_A/\sigma_{A_b}\,(0 - \bar{A}_b) = -0.0033$, to -0.0475, which is what the fit with a charge-symmetric background gives (-0.0471, Sec. VI).

7. Equal-Tailed and Highest-Posterior-Density Intervals

In short: for A *and* A_b *the two kinds of* 68% *interval coincide; they differ for the skewed parameters, whose highest-posterior-density interval is shifted towards the mode.* Table VIII compares the equal-tailed interval (percentiles 15.9%–84.1%) with the interval of highest posterior density, the shortest containing 68.3%, for the parameters of interest and for three skewed nuisance parameters.

Parameter	median	equal-tailed	HPD	skewness
A	-0.0442	$[-0.0529, -0.0358]$	$[-0.0525, -0.0355]$	$+0.01$
A_b	-0.0400	$[-0.0575, -0.0224]$	$[-0.0575, -0.0225]$	$+0.01$
f_+	0.114	$[0.089, 0.145]$	$[0.084, 0.138]$	$+0.90$
f_-	0.209	$[0.174, 0.248]$	$[0.168, 0.241]$	$+0.33$
λ_1^+ [MeV^{-1}]	0.0350	$[0.0281, 0.0431]$	$[0.0274, 0.0423]$	$+0.52$

TABLE VIII. Equal-tailed and highest-posterior-density 68% intervals and skewness of the marginal posteriors.

8. Savage–Dickey Ratio

In short: the Bayes factor of $A = 0$ *is the ratio of the posterior and prior densities at* $A = 0$*, with no integral over the twelve parameters.* The Bayes factor of H_0 ($A = 0$) against H_1 (A free) is the ratio of the evidences, $BF_{01} = P(D \mid H_0)/P(D \mid H_1)$. The null hypothesis is H_1 with A fixed to 0 and the same priors on the other parameters, so $P(D \mid H_0) = P(D \mid A = 0)$. Bayes' theorem for the marginal posterior of A at $A = 0$ reads $p(A = 0 \mid D) = P(D \mid A = 0)\,\pi(A = 0)/P(D \mid H_1)$. Solving for the ratio gives

$$BF_{01} = \frac{P(D \mid H_0)}{P(D \mid H_1)} = \frac{p(A = 0 \mid D)}{\pi(A = 0)},$$

the density of the posterior at $A = 0$ divided by that of the prior: how much the data have lowered the probability density of the null value. No integral over the twelve parameters is needed, only the marginal posterior of A from the chain.

9. Bayes Factor of $A = 0$

In short: the Savage-Dickey ratio with the Gaussian approximation of the posterior gives $BF_{01} \approx 1.1 \times 10^{-4}$ *(*$BF_{10} \approx 9 \times 10^3$*), decisive evidence against* $A = 0$ *that scales with the prior width.* The Bayes factor compares two hypotheses through their evidences (marginal likelihoods), $BF_{10} = P(D \mid H_1)/P(D \mid H_0)$ with $P(D \mid H_i) = \int \mathcal{L}(D \mid \theta, H_i)\,\pi(\theta \mid H_i)\,d\theta$: it is the factor by which the data update the prior odds of the two hypotheses, and it requires proper priors. Computing the evidence of each hypothesis by direct numerical integration is impractical for a 12-parameter posterior. When the null hypothesis fixes one parameter, H_0: $A = 0$, nested in H_1 (A free), and the priors on the other parameters are shared, the Bayes factor reduces to the Savage-Dickey density ratio [11] (Appendix C.1),

$$BF_{01} = \frac{p(A = 0 \mid D)}{\pi(A = 0)}, \qquad \pi(A = 0) = \tfrac{1}{2}.$$




FIG. XIII. Trace plot of the nominal fit with the 32 walkers initialized uniformly over the prior box (one line per walker, every 10th step), for A, N_s, σ and the steep-background fraction f_+. Left: first 3000 steps, full range; right: whole chain, zoomed on the posterior. The first half of the chain (dotted line) is discarded as burn-in.

With equal prior probabilities for the two hypotheses, BF_{01} is also the posterior odds of H_0 against H_1. $A = 0$ lies 5.2 posterior standard deviations from the median, in a tail where the chain contains essentially no samples, so a kernel density estimate of $p(A = 0 \mid D)$ would only reflect the kernel tails of the few most extreme samples. We therefore use the Gaussian approximation of the marginal posterior (Sec. V C), with the mean and standard deviation of the chain, which gives $BF_{01} \approx 1.1 \times 10^{-4}$ ($BF_{10} \approx 9 \times 10^3$, $\ln BF_{10} \approx 9.1$): decisive evidence against $A = 0$ on Jeffreys' scale [12] ($BF_{10} > 100$). As for any Bayes factor of a point null against a free parameter, the value scales with the prior width – the Ockham factor built into the Bayes factor, which penalizes a free parameter in proportion to the prior volume it is allowed to explore, and the mechanism behind Lindley's paradox: with the narrower prior $A \in (-0.3, 0.3)$ of Sec. D.15, $\pi(A = 0)$ grows to $1/0.6$ and BF_{01} drops by a factor 3.3, which leaves the conclusion unchanged. Figure XIV of Appendix D shows the two densities of the ratio; the dependence on the prior width, the evidence computed by direct integration and the equality of the background shapes of the two charges are in Appendix F.

10. Bayes Factor of the Background Asymmetry

In short: $A_b = -0.040 \pm 0.018$ *is* 2.3 *standard deviations from zero, yet* $BF_{01} \approx 3.5$ *favors* $A_b = 0$ *(Ockham factor);* A_b *is kept free in the nominal model.* The same test on the background asymmetry shows the Ockham factor at work. $A_b = -0.040 \pm 0.018$ lies 2.3 standard deviations from zero, yet the Savage-Dickey ratio gives $BF_{01}(A_b = 0) \approx 3.5$: the data mildly *favor* a charge-symmetric background, because the gain in likelihood does not compensate for spreading the prior of A_b over $(-1, 1)$. A negative background asymmetry is plausible: besides combinatorial background, the sample contains misidentified $B^\pm \to K^\pm\pi^+\pi^-$ and $B^\pm \to \pi^\pm K^+K^-$ decays, with A_{CP} values of their own, whose contribution grows when the kaon identification is loosened (Sec. VI). A_b is nevertheless kept free in the nominal model, which avoids imposing an assumption on the background that the frequentist analysis does not make; fixing $A_b = 0$ is one of the variants of the systematic study (Sec. VI).

11. Savage–Dickey Densities of the Parameters

In short: the posterior and prior densities at the null value, which give the Bayes factors of the main text.




FIG. XIV. Savage-Dickey ratio for (a) A and (b) A_b: marginal posterior (histogram of the chain and Gaussian approximation) and prior density, logarithmic scale. The Bayes factor BF_{01} is the ratio of the two densities at zero (dots). For A the posterior density at zero is 10^{-4} times the prior one; for A_b, 2.3 standard deviations from zero, it is higher than the prior density, and the data favor $A_b = 0$.

12. Priors and Bayes Factors

In short: no change of the prior on the nuisance parameters or of its parametrization moves A by more than 0.0002; the Bayes factor of $A = 0$ scales with the prior width, and the evidence computed by direct integration agrees in order of magnitude with the Savage-Dickey ratio. The posterior was reweighted, without new sampling, to other priors: logarithmically uniform yields and slopes, a prior uniform in the lifetimes $1/\lambda$, and a prior flat in (N_s^+, N_s^-) and (N_b^+, N_b^-) instead of (N_s, A) and (N_b, A_b), which differ by the Jacobian $N_s/2$ (Table IX). The largest shift is 0.0002, and the effective sample size of the reweighted chain is stated.

Prior	A (median)	68% interval	ESS
Nominal (uniform)	-0.0442	$[-0.0529, -0.0358]$	20000
Log-uniform N_s, N_b	-0.0442	$[-0.0529, -0.0358]$	19993
Log-uniform slopes	-0.0440	$[-0.0528, -0.0356]$	12306
Uniform in $1/\lambda$	-0.0444	$[-0.0526, -0.0360]$	153
Flat in (N_s^+, N_s^-)	-0.0442	$[-0.0529, -0.0358]$	19999
Flat in $(N_s^\pm)$ and $(N_b^\pm)$	-0.0442	$[-0.0529, -0.0358]$	19993

TABLE IX. Posterior of A under alternative priors, by importance reweighting of the chain; ESS: effective sample size of the 20,000 thinned samples.

The Bayes factor $BF_{01} = p(A = 0 \mid D)/\pi(0)$ is proportional to the density of the prior at zero, $\pi(0) = 1/(2W)$ for a uniform prior on $(-W, W)$ (Fig. V): for A it stays below 10^{-2} for every width up to $W = 1$ (decisive evidence for $A \neq 0$ whatever the prior), while for A_b it crosses 1 at $W \simeq 0.3$, which is the content of the Lindley paradox. The same evidence was computed by direct numerical integration over A on a grid, $Z_1 = \int L\,\pi\,dA$ with the nuisance parameters fixed at the maximum of the posterior, the method that is not feasible for twelve parameters: it gives $K_{01} = 7.0e-05$, against $1.3e-04$ with the profile likelihood and $1.1e-04$ for the Savage-Dickey ratio, which marginalizes the nuisance parameters; the three values are of the same order and the conclusion is the same.

Model	$A_{\rm raw}$	χ^2/ndof	p
Nominal	-0.0443	164.1/88	2×10^{-6}
Crystal Ball signal	-0.0445	131.0/86	1×10^{-3}
μ, σ per charge	-0.0443	164.2/86	8×10^{-7}
Shared bkg. shape	-0.0480	174.1/91	4×10^{-7}
Symmetric bkg. ($A_b = 0$)	-0.0471	169.8/89	5×10^{-7}
Single exponential	-0.0480	297.7/92	2×10^{-23}

TABLE X. $A_{\rm raw}$ (posterior median) and goodness of fit of the nominal model (Gaussian signal, double exponential per charge) and of the alternative signal and background models, same data and window.

The Savage-Dickey ratio also tests the equality of the background shapes of the two charges, one parameter at a time, with the difference $\delta = \theta^+ - \theta^-$: $BF_{01} = 3.1$ for the steep slope λ_1 (difference at 0.8σ), 2.2 for λ_2 (2.4σ) and 1.1 for the fraction f (2.0σ). The data are not able to distinguish the shapes, with a weak preference for equal ones: this is why the variant with a shared shape moves A by -0.0037 and is part of the systematic uncertainty, without being preferred.

13. Model Adequacy: Goodness of Fit and Posterior Predictive Check

In short: the model does not reach χ^2/ndof ≈ 1 (164.1/88), mainly because the signal tails are heavier than a Gaussian, but $A_{\rm raw}$ changes by only -0.0002 with a Crystal Ball signal.

A binned comparison of data and posterior-median model ($\chi^2 = \sum(\text{obs} - \text{exp})^2/\text{exp}$ over 50 bins of 10 MeV per charge, $100 - 12 = 88$ degrees of freedom) gives $\chi^2 = 164.1/88$ ($p = 1.6 \times 10^{-6}$), with 95.4 for B^+ and 68.7

for B^-. The largest pulls, in the B^+ sample, sit on the two flanks of the peak, about $\pm 3\sigma$ from μ (Fig. III): the observed signal shape has heavier tails than a Gaussian. Replacing the Gaussian by a Crystal Ball shape [18], a Gaussian core with a power-law low-mass tail,

$$S_{\mathrm{CB}}(x) \propto \begin{cases} e^{-x^2/2}, & x > -\beta, \\ \left(\frac{\kappa}{\beta}\right)^{\kappa} e^{-\beta^2/2} \left(\frac{\kappa}{\beta} - \beta - x\right)^{-\kappa}, & x \le -\beta, \end{cases} \tag{D1}$$

with $x = (m-\mu)/\sigma$, β the point where the tail starts (in units of σ) and κ its power, both free, improves the fit to $\chi^2 = 131.0/86$ ($p = 1.3 \times 10^{-3}$, $\beta = 1.71$, $\kappa = 35$) but moves A_{raw} by only -0.0002 (Table X). The asymmetry does not depend on the signal shape even though none of the models reaches $\chi^2/\mathrm{ndof} \approx 1$. The Gaussian is nevertheless kept as the nominal signal model, common to the two analyses: the Crystal Ball adds two tail parameters that are poorly constrained (its power κ runs to large values), and the effect of the signal shape on A_{raw} is covered by the systematic study (Sec. VI).

The background model is constrained by the data much more strongly. A single exponential per charge is rejected ($\chi^2 = 297.7/92$, $\Delta\chi^2 = 134$ for four fewer parameters), and so it is not used even as a systematic variant (Sec. VI). A shape shared between the two charges ($\Delta\chi^2 = 10.0$ for three fewer parameters) and a charge-symmetric background ($\Delta\chi^2 = 5.7$ for one fewer) are acceptable descriptions and shift A_{raw} by -0.0037 and -0.0028: the background treatment is the largest model dependence of the fit. The lower edge of the window is a compromise between two failures. Extending the window to 5050 MeV degrades the fit ($\chi^2 = 217.4/98$), because the double exponential no longer follows the background further below the peak; raising it to 5150 MeV removes most of the region that constrains the steep component, which becomes poorly identified – the chain for its fraction f_+ mixes slowly ($N/\tau = 10$) – while A_{raw} itself is unchanged (-0.0444).

The binned χ^2 uses the posterior medians only. A posterior predictive check [17] accounts for the posterior uncertainty and tests specific features of the data: for 1000 parameter sets θ_j drawn from the posterior, a dataset y_j^{rep} is generated from the model, and a discrepancy $T(y, \theta_j)$ is computed both on the replica and on the data; the posterior predictive p-value is $p_B = P\big(T(y^{\mathrm{rep}}, \theta) \geq T(y, \theta) \mid y\big)$. For the binned χ^2 the replicas give 104 ± 15 and the data 175 (median over θ_j): none of the replicas reaches the observed value ($p_B < 10^{-3}$), with the misfit concentrated in the B^+ sample ($p_B < 10^{-3}$, against 0.03 for B^-). The counting asymmetries of the whole window ($p_B = 0.996$) and of the signal window (0.99) are reproduced, but this is expected and not informative: A is a free parameter, so the replicas are centred on the observed value by construction. The asymmetry of the sidebands (0.73), which the fit constrains less directly, is also reproduced (Fig. XV). The check therefore shows that the model misdescribes the shape of the spectrum, mainly the signal tails, and does not reveal a charge-dependent distortion of the sidebands; that A_{raw} does not depend on the signal shape is shown directly by its stability under the Crystal Ball model (Table X).

14. Posterior Predictive Check with Statistics Not Fitted by the Model

In short: the asymmetry in slices of the mass, which the model predicts, is reproduced ($p_B = 0.67$), and the difference between the χ^2 of the two charges is at the edge of the replicas ($p_B = 0.08$). Section D.13 noted that the counting asymmetries of the window are fitted by construction. Two statistics that are not adjusted directly were therefore used (Fig. XVI): the asymmetry in ten slices of 50 MeV of the mass against the one predicted by the model from the same posterior draw, which tests that the charge asymmetry follows the mixture of signal and background, $T = \sum_s (A_s - \bar{A}_s)^2/\sigma_s^2$, and the difference $\chi^2(B^+) - \chi^2(B^-)$ of the binned fit. The slice asymmetries are reproduced; the difference of the χ^2 is 27 in the data against -0 ± 15 in the replicas, a 1.8σ excess that restates that the misfit is concentrated in the B^+ sample.

15. Robustness Checks: Polarity, Prior and Charm Vetoes

In short: A_{raw} agrees between the two magnet polarities (0.7σ) and does not depend on the prior; the D^0 veto moves it by 2.3 times the expected statistical spread.

The fit was repeated separately on the two magnet polarities, whose left-right detection asymmetries are opposite and partially cancel in the combined sample: $A_{\mathrm{raw}}(\mathrm{down}) = -0.0497 \pm 0.0112$ ($n = 11{,}636$) and $A_{\mathrm{raw}}(\mathrm{up}) = -0.0378 \pm 0.0129$ ($n = 8095$); the frequentist fits give -0.0492 ± 0.0111 and -0.0374 ± 0.0131, and the sum of their profile likelihoods -0.0443 ± 0.0084 (Appendix F). They differ by 0.7σ, so there is no indication that the asymmetry is an artifact of one polarity (Fig. XVII). A narrower prior, $A \in (-0.3, 0.3)$, leaves the posterior unchanged ($A_{\mathrm{raw}} = -0.0443$), as expected for a posterior far from the prior bounds; the prior width matters only for the Bayes factor (Sec. V D).

The sample contains $B^\pm$ decays through intermediate charm states, $B^\pm \to D^0 K^\pm$, $J/\psi K^\pm$ and $\chi_{c0} K^\pm$ with D^0, J/ψ, $\chi_{c0} \to K^+K^-$, which are visible as peaks in the opposite-sign $m(K^+K^-)$ spectra. They are part of the $B^\pm \to K^\pm K^+ K^-$ final state and are kept in the nominal sample; removing them changes the observable, from the asymmetry of all $B^\pm \to K^\pm K^+ K^-$ decays to that of the charmless ones, so the vetoes are reported as an informative check rather than as a systematic variation. A candidate is vetoed if either opposite-sign pair, $m(H1, H2)$ or $m(H1, H3)$, lies within ± 30 MeV of the D^0 or J/ψ mass or within ± 60 MeV of the χ_{c0} mass (world-average masses; the wider window for the χ_{c0} reflects its natural




FIG. XV. Posterior predictive check: distributions of three discrepancies over 1000 datasets replicated from posterior draws, compared with the observed values. (a) binned χ^2; (b) and (c) counting asymmetry of the fit window and of the sidebands.




FIG. XVI. (a) Counting asymmetry in slices of 50 MeV of the mass, data and model. (b) Distribution of $T = \sum_s (A_s - \bar{A}_s)^2/\sigma_s^2$ over 1000 replicas and observed value. (c) The same for $\chi^2(B^+) - \chi^2(B^-)$.




FIG. XVII. Posterior of A_{raw} for the magnet-down, magnet-up and combined samples.

width). The D^0 veto alone keeps 18,744 candidates and gives $A_{\text{raw}} = -0.0496 \pm 0.0088$; the three vetoes together keep 16,367 and give -0.0423 ± 0.0094. The shifts with respect to the nominal fit, -0.0053 and $+0.0020$, compare with expected statistical fluctuations of 0.0023 and 0.0040 for a subsample of the nominal one (Sec. VI); the D^0 veto moves A_{raw} by 2.3 times this amount, i.e. the vetoed $D^0 K^\pm$ candidates have a less negative asymmetry than the rest of the sample. The analysis was also carried out with a minimal six-parameter model, with a narrower window and a single exponential background, before the nominal model was fixed; it gives a compatible result (Appendix B.4).

16. Toy-MC Coverage Validation

In short: 200 *toy-MC experiments give coverage* 0.65 ± 0.03 *(68%) and* 0.93 ± 0.02 *(95%), consistent with nominal, with unbiased estimates and pulls of width* 1.04.

A credible interval is a probability statement on A given the data and the prior, whereas the nominal coverage of a confidence interval is a property of the procedure (Sec. IV). With flat priors and $\sim 2 \times 10^4$ events the two intervals coincide numerically (Sec. V C), but nothing guarantees the coverage of the credible interval, so it was checked: 200 toy-MC datasets were generated from the fitted model (parametric bootstrap, with the posterior




FIG. XVIII. Toy-MC validation, 200 toy-MC datasets. (a) Pulls $(\hat{A} - A_{\rm true})/\sigma(\hat{A})$, with $\hat{A}$ and $\sigma(\hat{A})$ the posterior median and standard deviation of each replica, compared with a standard normal. (b) Fraction of replicas whose 68% and 95% credible intervals contain the injected value, with binomial uncertainties, and the nominal coverage.

medians as ground truth and Poisson-fluctuated yields for each charge and component; the truncated Gaussian and exponential masses are drawn with the transformation method, i.e. by inverting their cumulative distributions) and each was refit with the nominal procedure and a 12,000-step chain. The fraction of replicas whose interval contains the injected A is 0.650 ± 0.033 for the 68% CI and 0.930 ± 0.015 for the 95% CI (binomial uncertainties), 1.0σ and 1.3σ below nominal: the coverage is consistent with nominal. The estimates themselves are unbiased and their spread matches the reported uncertainty (Fig. XVIII): their mean differs from the injected value by -0.0002 ± 0.0006, their standard deviation, 0.0088, agrees with the posterior standard deviation of the data fit, 0.0085, and the pulls have mean -0.03 ± 0.07 and standard deviation 1.04.

The toy chains are shorter than the nominal one, and in 124 of the 200 replicas the slowest-mixing parameter, always a background-shape parameter, has $N/\tau < 50$ (median 48, minimum 22). The criterion on the slowest parameter is conservative for the quantity tested here: for A itself the median autocorrelation time is 156 steps, so each chain is at least 45 autocorrelation times long in A and provides $\gtrsim 1400$ effectively independent samples of it. The Monte Carlo error on the percentiles of A is then a few per cent of $\sigma(A)$, negligible against the binomial uncertainty of the coverage, and the replicas with $N/\tau < 50$ do not show a lower coverage than the others (0.69 for the 68% interval, against 0.58 ± 0.06 for the 76 replicas with $N/\tau \geq 50$, whose coverage is 1.9 standard deviations below nominal: the split was made after the fact and 200 replicas cannot tell a trend from a fluctuation, and the check on the data of Sec. IV D gives the nominal coverage). This is a closure test of the inference procedure: since the toys are generated from the same model that is fitted, it does not probe the model mismatch of Sec. D.13, which is instead addressed by the systematic study.

17. Metropolis-Hastings Sampler and Published Likelihood

In short: an own Metropolis-Hastings sampler reproduces the posterior of the ensemble sampler, and the profile likelihood is given in a table, so that the result can be combined with others. The algorithm of the ensemble sampler was replaced by the basic Metropolis-Hastings one: a Gaussian proposal with the covariance of the posterior scaled by $2.38^2/d$, 120,000 steps with the first 1/6 discarded. The acceptance is 0.24, the integrated autocorrelation time of A is 36 steps (2813 effective samples), and $A = -0.0443$ with 68% interval $[-0.0526, -0.0357]$, against -0.0443 and $[-0.0528, -0.0358]$ from the ensemble sampler: the two samplers give the same posterior.

The profile likelihood of A is given in Table XI, together with the Minos intervals, so that other measurements of the same quantity can be combined with this one by adding $q(A)$.

A	−0.08	−0.07	−0.06	−0.05	−0.04	−0.03	−0.02	−0.01	+0.00
$q(A)$	17.9	9.3	3.5	0.5	0.2	2.8	8.1	16.1	27.0

TABLE XI. Profile likelihood ratio $q(A) = -2\Delta \ln L$ of the nominal fit (other parameters profiled); minimum at -0.0441, Minos intervals $[-0.0526, -0.0356]$ (68%) and $[-0.0607, -0.0275]$ (95%).

Appendix E: Systematic Uncertainties

In short: every fit of the nominal model and of its variants with both methods, the variants and the rule that combines them, and the validation of that rule on nested samples.

1. All Fits of the Nominal Model and of Its Variants

In short: the Bayesian and frequentist results of every variant and cross-check side by side.

2. List of the Variants

In short: thirteen variants of the nominal choices, in four groups (window, background, signal, preselection), fitted by both methods on exactly the same candidates. Each variant is fitted by both methods on exactly the same candidates.

- *Window*: lower edge at 5050 or 5150 MeV, upper edge at 5450 or 5700 MeV.
- *Background*: a single exponential per charge; a double exponential with the shape shared between charges; a charge-symmetric background ($A_b = 0$).

Fit	n	$A_{\rm raw}$ Bayesian (median)	68% CI	χ^2/ndof	$A_{\rm raw}$ frequentist
Nominal	19731	-0.0443	$[-0.0528, -0.0358]$	164.1/88	-0.0441 ± 0.0085
Systematic variants (Sec. VI)					
Window $[5050, 5600]$	21164	-0.0446	$[-0.0531, -0.0362]$	217.4/98	-0.0445 ± 0.0085
Window $[5150, 5600]$	18566	-0.0444	$[-0.0528, -0.0359]$	141.5/78	$-0.0446 \pm 0.0088^{\dagger}$
Window $[5100, 5450]$	19004	-0.0440	$[-0.0524, -0.0357]$	136.9/58	-0.0442 ± 0.0085
Window $[5100, 5700]$	20191	-0.0456	$[-0.0540, -0.0373]$	200.2/108	-0.0455 ± 0.0085
Single exponential	19731	-0.0480	$[-0.0564, -0.0395]$	297.7/92	-0.0481 ± 0.0085
Background shape shared	19731	-0.0480	$[-0.0564, -0.0397]$	174.1/91	-0.0481 ± 0.0083
Symmetric background	19731	-0.0471	$[-0.0552, -0.0388]$	169.8/89	-0.0472 ± 0.0084
Crystal Ball signal	19731	-0.0445	$[-0.0529, -0.0363]$	131.0/86	-0.0447 ± 0.0085
μ, σ per charge	19731	-0.0443	$[-0.0527, -0.0356]$	164.2/86	-0.0442 ± 0.0087
`ProbK` > 0.4	26553	-0.0474	$[-0.0550, -0.0399]$	175.6/88	-0.0473 ± 0.0076
`ProbK` > 0.6	13981	-0.0385	$[-0.0482, -0.0289]$	145.0/88	-0.0384 ± 0.0099
`ProbPi` < 0.4	17984	-0.0421	$[-0.0506, -0.0333]$	150.1/88	-0.0421 ± 0.0088
`ProbPi` < 0.6	20953	-0.0446	$[-0.0529, -0.0366]$	159.0/88	-0.0448 ± 0.0083
Cross-checks (Secs. V B, D.15)					
Dispersed start, run 1	19731	-0.0446	$[-0.0529, -0.0360]$	164.1/88	–
Dispersed start, run 2	19731	-0.0442	$[-0.0528, -0.0356]$	164.1/88	–
Magnet down	11636	-0.0497	$[-0.0607, -0.0386]$	153.1/88	–
Magnet up	8095	-0.0378	$[-0.0507, -0.0251]$	150.3/88	–
Narrow prior, $A \in (-0.3, 0.3)$	19731	-0.0443	$[-0.0528, -0.0358]$	164.1/88	–
D^0 veto	18744	-0.0496	$[-0.0584, -0.0409]$	159.1/88	-0.0493 ± 0.0087
D^0, J/ψ, χ_{c0} vetoes	16367	-0.0423	$[-0.0516, -0.0329]$	140.1/88	-0.0421 ± 0.0094

TABLE XII. All fits of the nominal model and of its variants: number of candidates, Bayesian posterior median and 68% credible interval of $A_{\rm raw}$, binned goodness of fit (10 MeV bins), and frequentist estimate with its Hesse uncertainty (Sec. IV; fitted on the same candidates; the cross-checks of the Bayesian fits that have no frequentist counterpart in this work are marked –). † Uncertainty scaled from the number of events, since the Hesse matrix of that fit is not accurate. Every fit uses 32 walkers and 25,000 steps.

- *Signal*: a Crystal Ball shape with free tail parameters, Eq. (D1); μ and σ separate for the two charges.
- *Preselection*: one cut at a time loosened or tightened, `ProbK` > 0.4 or > 0.6, `ProbPi` < 0.4 or < 0.6, on all three tracks.

3. Combination Rule

In short: a variant that changes the sample counts only if its shift exceeds σ_B; variants rejected by the data are excluded, the largest shift of each group is its uncertainty, and the groups are added in quadrature. The shift $\Delta = A_{\rm raw}^{\rm var} - A_{\rm raw}^{\rm nom}$ of a variant is not necessarily a systematic effect. When a variant changes the sample (window, preselection), the two samples are nested and part of the shift is a statistical fluctuation, with expected size $\sigma_B = \sqrt{|\sigma_{\rm var}^2 - \sigma_{\rm nom}^2|}$, σ being the statistical uncertainty of each fit [13]; such a variant contributes only if $|\Delta| > \sigma_B$. A variant that the data reject, with a χ^2 worse than the nominal one by more than 25 on the same data, is not a plausible alternative and is excluded: this removes the single exponential ($\Delta\chi^2 = 134$, Sec. D.13). The largest remaining $|\Delta|$ of each group is taken as its uncertainty, since the variants of a group are alternative descriptions of the same choice, and the four groups are added in quadrature.

4. Results and Particle-Identification Scan

In short: the total systematic uncertainties are 0.0070 *(Bayesian) and* 0.0072 *(frequentist), dominated by the preselection through the single variant* `ProbK` > 0.6 *(*$+0.0058$*,* $1.2\,\sigma_B$*).* The results are in Table II, and all the Bayesian fits, with their intervals and goodness of fit, in Table XII of the Appendix. The two methods give the same shifts to within $\sim$0.0005 and the same group uncertainties, with total systematic uncertainties of 0.0070 (Bayesian) and 0.0072 (frequentist). The final results are

Bayesian:

$$A_{\rm raw} = -0.0443 \pm 0.0085\,(\text{stat}) \pm 0.0070\,(\text{syst}),$$

frequentist:

$$A_{\rm raw} = -0.0441 \pm 0.0085\,(\text{stat}) \pm 0.0072\,(\text{syst}).$$

The preselection dominates, through the single variant `ProbK` > 0.6 ($+0.0058$, $1.2\,\sigma_B$). To tell a dependence on the PID requirement from a fluctuation of that single value, the frequentist fit was repeated in steps of 0.05 of each cut, from `ProbK` > 0.30 to > 0.85 and from `ProbPi` < 0.15 to < 0.70. On the tight side of the kaon requirement the shifts scatter around zero without a trend ($+0.0009$, $+0.0058$, -0.0017, -0.0112, -0.0054, $+0.0010$, $+0.0022$ for `ProbK` > 0.55–0.85, all within $1.3\,\sigma_B$), and the pion requirement gives a flat scan within $1.3\,\sigma_B$: the `ProbK` > 0.6 value is a statistical fluctuation, and




FIG. XIX. Shift $\Delta = A_{\rm raw}^{\rm var} - A_{\rm raw}^{\rm nom}$ of every variant of the systematic study, Bayesian and frequentist fits; bars: expected spread σ_B of a nested subsample (variants that change the sample only). Open symbols: variant rejected by the data. The dotted lines separate the groups (window, background, signal, preselection).

the preselection uncertainty is a conservative estimate. Loosening the kaon requirement, instead, moves $A_{\rm raw}$ monotonically (-0.0019, -0.0032, -0.0067, -0.0111 for ProbK $>$ 0.45–0.30, the last at $2.5\,\sigma_B$), as expected if misidentified $B \to K\pi\pi$ and $B \to \pi KK$ decays, with asymmetries of their own and not included in the mass model, enter the sample; this supports the nominal requirement. The combination rule was fixed before this scan and is not changed by it. The background group is the second contribution and is driven by the assumptions on the charge dependence of the background, the window the third.

5. Shifts of All Variants

In short: the shifts of the Bayesian and frequentist fits coincide variant by variant; the preselection group dominates, and the single exponential, rejected by the data, is drawn open. The 13 variants of the systematic study (Sec. VI) are shown in Fig. XIX. For the variants that change the sample the bar is the expected statistical spread σ_B of the shift (Appendix E.8); the others have no bar because the sample is the same. The Bayesian (purple) and frequentist (green) shifts agree variant by variant, the preselection group is the largest, and the single exponential, which the data reject, is drawn open and excluded from the uncertainty.

6. Scan of the Particle-Identification Cuts

In short: on the tight side the shift scatters around zero within $1.3\,\sigma_B$ and the pion requirement is flat; loosening ProbK moves $A_{\rm raw}$ monotonically, which supports the nominal requirement. The preselection dominates, through the single variant ProbK > 0.6 ($+0.0058$, $1.2\,\sigma_B$). A scan in steps of 0.05 of each cut (ProbK $>$ 0.30–0.85, ProbPi $<$ 0.15–0.70) shows that on the tight side the shifts scatter around zero without a trend, all within $1.3\,\sigma_B$, and the pion requirement is flat: the ProbK > 0.6 value is a statistical fluctuation and the preselection uncertainty is conservative. Loosening the kaon requirement moves $A_{\rm raw}$ monotonically (-0.0019 to -0.0111 for ProbK $>$ 0.45–0.30, the last at $2.5\,\sigma_B$), as expected if misidentified $B \to K\pi\pi$ and $B \to \pi KK$ decays, not in the mass model, enter the sample; this supports the nominal requirement. The background group is second, driven by the charge dependence of the background, and the window third.

The scan is shown in Fig. XX.

7. Sensitivity of σ_B to Differences between the Methods

In short: σ_B is the difference of two nearly equal uncertainties and is sensitive to small differences between the methods; a variant at the threshold can be classified differently without changing the result. Each method computes σ_B from its own uncertainties (posterior standard deviation, Hesse error). Since σ_B is the difference of two nearly equal uncertainties, it is sensitive to small differences between them, including the Monte Carlo noise of the posterior standard deviation (1–3%), and a variant close to the threshold can be classified differently by the two methods: ProbPi < 0.4 counts for the Bayesian fit ($|\Delta| = 0.00223$, $\sigma_B = 0.00220$) and not for the frequentist one (0.00204 vs 0.00229). This does not affect the result, since the uncertainty of each group is set by its largest shift, the same in both methods.

8. Validation of the Rule for Nested Samples

In short: the spread of the shift of $\hat{A}$ between a sample and its tighter-identification subsample is the one given by the quadrature difference of the uncertainties. The systematic study counts a variant that changes the sample only if its shift exceeds $\sigma_B = \sqrt{|\sigma_{\rm var}^2 - \sigma_{\rm nom}^2|}$ (Sec. VI). To verify the rule, 300 bootstrap replicas of the nominal sample were fitted before and after the tighter selections ProbK > 0.6 and ProbPi < 0.4 (Fig. XXI): the standard deviation of the shift is 0.0051 and 0.0025, against median σ_B of 0.0050 and 0.0023 (ratios 1.02 and 1.09), and the shifts follow a Gaussian of width σ_B. The mean shift of the replicas (0.0060 and 0.0020) is that of the data,




FIG. XX. Shift of A_{raw} (frequentist fit) as a function of (a) the `ProbK` lower cut and (b) the `ProbPi` upper cut, in steps of 0.05, with the expected statistical spread of the nested sample ($\pm\sigma_B$ dark, $\pm2\sigma_B$ light; zero at the nominal cut, dash-dotted line). On the tight side of `ProbK` the points scatter around zero without trend and the pion requirement is flat; loosening `ProbK` moves A_{raw} monotonically.

which is preserved by the resampling: it is a property of the sample and not a fluctuation of the rule.

9. Stability Scans of the Frequentist Fit: Numbers

In short: the largest shifts are 0.0048 *over the grid of window edges and* 0.0126 *over the grid of the two PID cuts, whose samples are smaller (mean uncertainty* 0.0112*); the signal region moves* A_{raw} *by* 6×10^{-5}.

Numbers. Three results are used here, from the same notebook as the nominal fit; they do not enter the systematic uncertainty, whose variants and combination rule are those above.

- *Fit window*: the maximum shift with respect to the nominal fit is 0.0048 over the grid of lower and upper edges (standard deviation 0.0018, mean statistical uncertainty 0.0085); the extreme corners are the smallest and most asymmetric windows, where the background shape is least constrained.
- *Signal region* (used only to start the sideband fit): the shift is at most 6×10^{-5}, so the result does not depend on it.
- *Particle-identification requirements*: on the grid of the two cuts the maximum shift is 0.0126 (standard deviation 0.0046). This is larger than the statistical uncertainty of the nominal fit, 0.0085, but the grid contains much smaller samples, with a mean statistical uncertainty of 0.0112; the relevant comparison for nested samples is with σ_B, as in the one-cut-at-a-time scan above, where no shift exceeds $1.3\,\sigma_B$ on the tight side.

The window and signal-region scans are therefore consistent with a fit that does not depend on these choices, and the preselection remains the dominant contribution, as quantified by the systematic study.




FIG. XXI. Shift of $\hat{A}$ between the nominal sample and a tighter selection, divided by the median σ_B, in 300 bootstrap replicas of the data: (a) `ProbK` > 0.6, (b) `ProbPi` < 0.4, and standard normal distribution.

Appendix F: Validation of the Estimator and of the Intervals

In short: how the toy-MC experiments are generated, the bias, efficiency and coverage of the estimator, its asymptotic behavior, the size and power of the test of $A = 0$ and the goodness of fit as a function of the binning.

1. Generation of the Toy-MC Experiments

In short: toy experiments are generated from the fitted model (Poisson yields, masses by the transformation method) and fitted with the nominal procedure: 1000 *at the fitted value and* 300 *at each of four other values of A.* The frequentist results of Sec. IV rest on asymptotic properties of the maximum-likelihood estimator and of the profile-likelihood ratio, and Sec. IV D verified the coverage by resampling the data. Here the same questions are asked of toy-MC experiments generated from the fitted model, as was done for the Bayesian interval in Sec. D.16, which allows the true value of A to be chosen. The fitted model of Sec. III defines the toy-MC datasets: for each charge the yields of signal and background are Poisson-fluctuated, the signal masses are drawn from the truncated Gaussian and the background masses from the two truncated exponentials by the transformation method. Each dataset is fitted with the nominal frequentist procedure; the likelihood ratio $t(A_{\text{true}}) = -2\ln[L(A_{\text{true}})/L(\hat{A})]$ and the statistic of the test of $A = 0$, $q(0)$, are computed with the other eleven parameters profiled. One thousand toy-MC experiments were generated at the fitted value, $A_{\text{true}} = -0.0441$, and 300 at each of $A_{\text{true}} = -0.1$, -0.02, 0 and $+0.05$; between 1% and 3% of the fits are not flagged as valid by `MIGRAD` and are excluded.

2. Bias, Efficiency and Coverage

In short: the bias is at most 0.0008*, the spread equals the Hesse uncertainty within* 6%*, and the coverage at the fitted value is* 0.696 ± 0.015 *and* 0.954 ± 0.007. Table III summarizes the estimator over the five injected values. The bias of $\hat{A}$ is at most 0.0008 (1.6 standard errors), the spread of the estimates equals the mean Hesse uncertainty within 6% (ratio 0.94–1.05), and the pulls $(\hat{A} - A_{\text{true}})/\sigma(\hat{A})$ have width 0.94–1.05: the estimator is unbiased and reaches the minimum variance bound for this sample size, and the Hesse uncertainty is the correct statistical uncertainty. At the fitted value the coverage of the interval $t(A_{\text{true}}) \leq 1$ is 0.696 ± 0.015 for the 68.3% level and 0.954 ± 0.007 for the 95.4% level (Fig. XXII of Appendix F, panel b, shows the p-value of $t(A_{\text{true}})$, uniform as expected, Kolmogorov-Smirnov $p = 0.96$ for all values pooled). The 68% coverage at $A_{\text{true}} = +0.05$, 0.614 ± 0.028, is 2.5 standard deviations below nominal; the 95% coverage at the same value is 0.956, and with five injected values a deviation of this size is not unexpected.

3. Asymptotic Behavior of the Estimator

In short: with the number of candidates the bias tends to zero, the spread scales as $N^{-1/2}$ and the distribution of $\hat{A}$ becomes Gaussian, as the central limit theorem and the asymptotic properties of the maximum-likelihood estimate predict. The 300 toy-MC datasets for each size of the low-statistics study (Sec. H.11) show the approach to the asymptotic limit (Table XIII): the bias decreases from -0.0085 at $N = 50$ to $+0.0009$ at $N = 2000$, the product of the spread and $\sqrt{N}$ is constant, and the distribution of $\hat{A}$ is compatible with a Gaussian from $N = 150$ (Shapiro-Wilk $p > 0.3$), with a mild excess kurtosis at $N = 50$ ($p = 0.03$). Linear propagation of the Poisson uncertainty of the signal yield, $\sigma(A) = \sqrt{(1 - A^2)/N_s}$ with $N_s = 0.86\,N$, gives 0.153 at $N = 50$ and 0.024 at $N = 2000$, against spreads of 0.161 and 0.023: it is correct at the few-per-cent level, because the fit also has to determine the background, and the first-order expansion is not the limiting approximation. At the full sample the spread of the estimator equals the Hesse uncertainty (Sec. VII A).

N	bias	sd($\hat{A}$)	sd·$\sqrt{N}$	skewness	ex. kurtosis	Shapiro p
50	-0.0085	0.1611	1.14	-0.15	$+0.64$	0.03
150	-0.0051	0.0920	1.13	$+0.16$	$+0.31$	0.59
500	-0.0022	0.0501	1.12	-0.08	$+0.38$	0.52
2000	$+0.0009$	0.0225	1.01	-0.20	-0.20	0.32

TABLE XIII. Distribution of the maximum-likelihood estimate of A in 300 toy-MC datasets of N candidates (six-parameter model of Sec. B.4).

4. Size and Power of the Test of $A = 0$

In short: the test has the nominal size, the statistic $q(0)$ follows a χ^2 distribution with one degree of freedom, and an asymmetry of -0.02 would be rejected at 3σ in one toy-MC experiment out of four. Under $A_{\text{true}} = 0$ the fraction of toy-MC experiments with $q(0) > 3.84$ is 0.051 ± 0.013, against the nominal size $\alpha = 0.05$, and 2 of 293 exceed $q = 9$ (3σ), against the 0.8 expected; the distribution of $q(0)$ is compatible with χ^2_1 (Kolmogorov-Smirnov $p = 0.79$, Fig. XXIIa), and the tail used for the significance ($q = 27$) is an extrapolation of it. The power, the probability of rejecting $A = 0$ when the asymmetry is A_{true} (Fig. XXIIIc), is 1.00 at the fitted value and at $\pm$ the larger injected values, and 0.68 (0.27 at 3σ) at $A_{\text{true}} = -0.02$: with 19,731 candidates an asymmetry of 0.02 is detected in about two experiments out of three at the 5% size, and an asymmetry below that is not resolved.




FIG. XXII. Toy-MC experiments, validity of Wilks' theorem. (a) Fraction of experiments with $q(0)$ above the observed value, for $A_{\rm true} = 0$, compared with the χ^2_1 distribution. (b) p-value of $t(A_{\rm true})$ against a uniform distribution, at the fitted value and pooled over the injected values. (c) Distribution of $t(A_{\rm true})$ at the fitted value and χ^2_1.




FIG. XXIII. Toy-MC experiments at five injected asymmetries. (a) Bias of $\hat{A}$, with its standard error. (b) Spread of the estimates divided by the mean Hesse uncertainty, and width of the pulls. (c) Probability to reject $A = 0$ in the test with $q(0) > 3.84$ (size 5%) and $q(0) > 9$ (3σ), as a function of the injected value: size at $A_{\rm true} = 0$, power elsewhere.

5. Goodness of Fit and the Binning

In short: the binned χ^2 of toy-MC experiments follows its asymptotic distribution, so its p-values are accurate; the misfit of the data is 5.4 *standard deviations at* 10 *MeV bins and* 2.0 *at* 2.5 *MeV, because it lives on the scale of the flanks of the peak.* The p-values of the binned χ^2 rely on the asymptotic distribution with ndof degrees of freedom, which can fail for bins with few entries. The χ^2 of 1000 toy-MC experiments from the fitted model has mean 87.6 and standard deviation 13.3 for 2×50 bins (88 degrees of freedom, expected $\sqrt{2 \cdot 88} = 13.3$), and mean 387.6, standard deviation 27.1 for 2×200 bins (388 degrees of freedom): the asymptotic χ^2 distribution describes the toys (Fig. XXIV). The data give 159.5 for 2×50 bins of 10 MeV, 5.4 standard deviations above the toys (asymptotic $p = 4.8 \times 10^{-6}$, fewer than 1/1000 toys above), 253.1/188 for 5 MeV bins ($p = 1.1 \times 10^{-3}$) and 441.7/388 for 2.5 MeV bins (2.0 standard deviations, $p = 0.031$, in toys 0.032). The same fit therefore has $p = 0.03$ or $p = 5 \times 10^{-6}$ depending on the binning: the discrepancy between model and data is a feature of ~ 10 MeV width, the flanks of the peak (Fig. III), diluted in narrower bins, and the binned χ^2 is a test whose power depends on the binning, in the same way as the Bayes factor depends on the prior width.




FIG. XXIV. Binned χ^2 of 1000 toy-MC experiments from the fitted model, χ^2 distribution with the corresponding degrees of freedom, and value of the data: (a) 2×50 bins of 10 MeV, (b) 2×200 bins of 2.5 MeV.

Appendix G: Local Asymmetry in the Dalitz Plane

In short: the local deviations and their robustness, the null tests of the procedure, the region of the published analysis and the split by magnet polarity.

1. Local Deviations and Their Robustness

In short: two bins have $|S_i| > 3$ (-3.7 and -3.0) and the global homogeneity test gives $\chi^2 = 80.8$ for 31 degrees of freedom ($p = 3 \times 10^{-6}$), also in the signal window alone. Two of the 32 bins have $|S_i| > 3$, adjacent to each other at $m^2_{\rm lo} < 4$ GeV^2 and $7.5 < m^2_{\rm hi} < 11.25$ GeV^2 ($S_i = -3.7$ and -3.0), with A_i more negative than the global value; the bin below the first one ($m^2_{\rm lo} < 2$, $3.75 < m^2_{\rm hi} < 7.5$ GeV^2) has $S_i = -2.98$, just under the threshold (Fig. VII); about 0.09 would be expected if the asymmetry were uniform, with a probability of 3×10^{-3} for two or more. Because the bins are examined together this is a look-elsewhere situation: the number refers to any two of the 32 bins, not to the two found, and the global homogeneity test that follows does not depend on which bins are singled out. The largest positive deviations, close to $+3$, lie at high $m^2_{\rm hi}$. A global test of homogeneity, $\chi^2 = \sum_i S_i^2 = 80.8$ for 31 degrees of freedom ($p = 3 \times 10^{-6}$), confirms that the local asymmetry is not uniform. Restricting the sample to the signal window ($|m - \mu| < 2\sigma$, $n = 15{,}130$, where the fitted model gives a signal purity of ≈96%) gives the same picture – three bins with $|S_i| > 3$, $\chi^2 = 95.6/28$ – so the structure is not produced by the combinatorial background. Nor is it produced by the charm contributions: with the D^0 veto two bins still exceed 3σ ($\chi^2 = 79.2/31$), and with all three vetoes one remains ($\chi^2 = 74.4/31$, $p = 2 \times 10^{-5}$). It does not depend on the binning either: 6×6, 10×10 and 12×12 grids give $\chi^2 = 81.0/19$, $97.1/43$ and $120.0/60$ ($p \leq 7 \times 10^{-6}$).

2. Homogeneity Test: Signal Window, Vetoes, Grids and Magnet Split

In short: the homogeneity test holds in the signal window, with the charm vetoes and for other grids, and the structure is the same in the two magnet polarities ($\chi^2 = 23.0$ for 29 bins). Two of the 32 bins have $|S_i| > 3$, adjacent at $m^2_{\rm lo} < 4$ GeV^2 and $7.5 < m^2_{\rm hi} < 11.25$ GeV^2 ($S_i = -3.7$ and -3.0, Fig. VII); about 0.09 would be expected if the asymmetry were uniform (3×10^{-3} for two or more), a look-elsewhere number that refers to any two of the 32 bins. The global homogeneity test, $\chi^2 = \sum_i S_i^2 = 80.8$ for 31 degrees of freedom ($p = 3 \times 10^{-6}$), confirms that the local asymmetry is not uniform; it holds in the signal window alone ($\chi^2 = 95.6/28$, signal purity ≈96%), with the D^0 veto (79.2/31) and with all three charm vetoes (74.4/31), and for 6×6, 10×10 and 12×12 grids (81.0/19, 97.1/43, 120.0/60).

The structure does not depend on the magnet polarity either. The left-right part of the detection asymmetry changes sign when the polarity is reversed, so a local asymmetry produced by it would appear with opposite deviations in the two samples. In the region of the two significant bins ($m^2_{\rm lo} < 4$ GeV^2, $7.5 < m^2_{\rm hi} < 11.25$ GeV^2) the counting asymmetry is -0.129 ± 0.023 with the magnet down and -0.126 ± 0.028 with the magnet up, although the global asymmetries of the two samples differ (-0.051 and -0.032), and the two maps agree bin by bin ($\chi^2 = 23.0$ for 29 bins, $p = 0.78$). This split does not test the part of the detection asymmetry that does not change sign with the polarity, the different interaction of K^+ and K^- with the detector material.

3. Null Tests of the Local-Asymmetry Procedure

In short: permuting the charges gives $\chi^2 = 31.5 \pm 8.4$, as expected, and never reaches the observed 80.8; the phase-space simulation is consistent with no asymmetry,




FIG. XXV. Homogeneity statistic $\chi^2 = \sum_i S_i^2$ of the 8×8 Dalitz grid for 2000 random permutations of the charges (histogram), χ^2 distribution with 31 degrees of freedom (dashed) and observed value (line).

with slightly optimistic χ^2 p-values. Two null tests check the procedure itself. Shuffling the charges among the candidates removes any local asymmetry while keeping the occupancy of the bins and the global asymmetry of the data: over 2000 permutations the 8×8 grid gives $\chi^2 = 31.5 \pm 8.4$, as expected for 31 degrees of freedom, and no permutation reaches the observed value (permutation $p < 5 \times 10^{-4}$; $p = 1 \times 10^{-3}$ for the 12×12 grid). The phase-space simulation of the record, without asymmetry by construction, gives $\chi^2 = 40.0/30$ ($p = 0.11$) with the same procedure on the whole sample; on 500 subsamples of the size of the data the mean χ^2/ndof is 1.16 and 13% have $p < 0.05$, so the χ^2 p-values are slightly optimistic (the binomial uncertainty is evaluated at the observed A_i). The permutation test, whose null distribution is exact, gives the same conclusion: the non-uniformity is a property of the data.

4. Distribution of the Homogeneity Statistic under Charge Permutation

In short: shuffling the charges reproduces the χ^2 distribution of a uniform asymmetry, and the observed value lies far beyond all 2000 *permutations.* For each of 2000 random permutations of the charges among the candidates, which removes any local asymmetry while keeping the occupancy of the bins and the global asymmetry, the statistic $\chi^2 = \sum_i S_i^2$ was computed on the 8×8 grid (Fig. XXV). The distribution follows the χ^2 with 31 degrees of freedom, which confirms that the statistic is correctly normalized, and no permutation reaches the observed value (80.8).

5. Asymmetry in the $\phi(1020)$ Band and in the Significant Bins

In short: the band of the $\phi(1020)$ has no asymmetry; the negative asymmetry is outside it, and larger when the ϕ candidates are removed. The asymmetry is not produced by the $\phi(1020)$ (Fig. XXVI). The band $|m_{\rm lo} - m_\phi| < 10$ MeV around the mass of the ϕ (panel a) contains 1964 candidates and has a counting asymmetry compatible with zero, -0.004 ± 0.023, while the region of the two significant bins has -0.128 ± 0.018 and, without the candidates of the ϕ band, -0.145 ± 0.019; the 353 candidates of the ϕ band that lie in that region have $+0.003 \pm 0.053$. The region just below the ϕ ($m^2_{\rm lo} < 2$ GeV^2, outside the band) has -0.120 ± 0.014 (panel b).

6. Region of the Published Analysis

In short: the $\phi(1020)$ band has no asymmetry (-0.004 ± 0.023); the fit of the 2146 *candidates of the published region gives -0.234 ± 0.025 (Bayesian) and -0.237 ± 0.025 (frequentist), against -0.239 ± 0.020.* The ordered Dalitz plot (Fig. VIIIa) shows the two-body resonances of the decay: the narrow $\phi(1020)$ band at the kinematic edge, $m^2_{\rm lo} \approx 1.04$ GeV^2, and the charm bands of the D^0, J/ψ and χ_{c0}. The negative local asymmetry is not produced by the $\phi(1020)$: its band ($|m_{\rm lo} - m_\phi| < 10$ MeV, 1964 candidates) has $A_{\rm count} = -0.004 \pm 0.023$, while the significant bins without it have -0.145 ± 0.019. The asymmetry sits in the low-mass K^+K^- region just above the ϕ, which is where the published analysis of the same data located it, defining the region $1.2 < m^2_{\rm lo} < 2.0$ GeV^2, $m^2_{\rm hi} < 15$ GeV^2 and measuring there $A_{\rm raw} = -0.239 \pm 0.020$ [5]. Since that region is fixed by the paper and not chosen on our data, it allows an unbiased comparison: fitting the 2146 candidates in it with the nominal model gives $A_{\rm raw} = -0.234 \pm 0.025$ (Bayesian; the chain mixes slowly in the poorly constrained second background exponential, $N/\tau = 15$) and -0.237 ± 0.025 (frequentist, 9.4σ from zero), reproducing the published value; the mass spectra of the two charges in the region (Fig. VIIIb) show the difference directly. Three caveats apply: the binning is coarse and was not optimized; the asymmetries are computed on all candidates, without the sWeight-based background subtraction that a full analysis would use (Appendix J.5); and, since $A_{\rm raw}$ is not corrected for detection asymmetries, local variations can still be partly instrumental – the K^+/K^- interaction asymmetry depends on the kaon momenta, which are correlated with the position in the Dalitz plane, and is not removed by the polarity split. Appendix I.8 revisits the same question with an adaptive, data-driven partition (K-means).




FIG. XXVI. (a) Distribution of $m_{\rm lo}(K^+K^-)$ around the $\phi(1020)$ (5 MeV bins) and band used (shaded). (b) Counting asymmetry, with binomial uncertainty and number of candidates, of all candidates (dotted line), of the ϕ band, of the region $m^2_{\rm lo} < 2$ GeV^2 outside the band, and of the two significant bins of Fig. VII, with and without the candidates of the ϕ band, and of those candidates alone.

Appendix H: Comparison of the Two Approaches

In short: the numerical agreement of the two fits, the meaning of the intervals, the hypothesis tests, the profile likelihood against the posterior, and the two methods at low statistics.

1. Numerical Agreement

In short: the two fits agree to the quoted digits (estimates differ by 0.0002*, both uncertainties are* 0.0085*, systematic uncertainties* 0.0070 *and* 0.0072*); sideband subtraction gives* -0.0503 ± 0.0083. The two fits of the same model agree to the precision of the quoted digits: the estimates differ by 0.0002 (0.02 standard deviations), the uncertainties are both 0.0085, the 68% and 95% intervals differ by at most 0.0002, and the shifts of the thirteen systematic variants agree within 0.0005, with the same group uncertainties for the window (0.0014) and the preselection (0.0058) and slightly different ones for the background (0.0037 against 0.0040) and the signal model (0.0002 against 0.0006), so that the total systematic uncertainties are 0.0070 and 0.0072 (Sec. VI). A third estimate, by sideband subtraction (counting in the signal region after removing the background expected from the sideband fit), gives -0.0503 ± 0.0083, which differs from the nominal value by 0.006, less than one standard deviation, on the same candidates. This is expected, and it is not by itself a validation of the model. With $\sim 2 \times 10^4$ candidates and flat priors the posterior is proportional to the likelihood, the likelihood is close to a Gaussian in A, and the posterior mode coincides with the maximum-likelihood estimate; the posterior median, the posterior mean and the mode differ by less than 2×10^{-4}. What the agreement does show is that neither the sampling (MCMC) nor the asymptotic approximations (Hesse, Wilks) introduce a bias: the Bayesian intervals have the coverage of the frequentist ones (Sec. D.16) and the profile-likelihood interval has the coverage assumed by Wilks' theorem (Sec. IV D).

The agreement does not extend to the validity of the model itself. Both methods describe the same spectrum with the same Gaussian signal and double-exponential background, and neither reaches $\chi^2/{\rm ndof} \approx 1$ (Sec. D.13); what protects $A_{\rm raw}$ is that its value does not depend on the signal shape (shift -0.0002 with a Crystal Ball) and that the background treatment, the largest model dependence, is included in the systematic uncertainty.

2. Meaning of the Interval

In short: the credible interval is a probability on A given the data and the prior, the confidence interval a property of the procedure, which the bootstrap and the toy experiments verified. Meaning of the interval. The credible interval $[-0.0528, -0.0358]$ states that, given the data and the prior, A lies in it with 68% probability; the confidence interval $[-0.0526, -0.0356]$ is the set of values that the data do not reject at the 32% level, and the 68% is the fraction of repeated experiments in which such an interval contains the true value. The numbers are the same here; the statements are not, and only the second is a property of the procedure that can be checked by repeating the experiment, which is what the bootstrap and toy checks did.

3. Profile Likelihood, Posterior and the Two Polarities

In short: profiling and marginalizing the nuisance parameters give the same distribution of A; the two magnet polarities agree with each other, and combining their likelihoods reproduces the joint fit. The fre-

quentist way of eliminating the nuisance parameters is to profile them, $q(A) = -2\Delta \ln L$ with the other parameters re-maximized at each A, the Bayesian one is to marginalize them; Fig. IXa shows that $\exp[-q(A)/2]$, normalized, and the marginal posterior coincide, as expected for a near-Gaussian likelihood and flat priors. The two polarities were fitted separately and their profile curves added, which is the combination of likelihoods of two independent measurements of the same parameter (Fig. IXb): magnet down gives -0.0492 ± 0.0111, magnet up -0.0374 ± 0.0131, their sum -0.0443 ± 0.0084, against -0.0441 ± 0.0085 of the joint fit (the nuisance parameters are shared in the joint fit, separate in the sum). The Bayesian counterpart is sequential: the posterior of magnet down used as the prior of magnet up gives -0.0446 ± 0.0086 (Table XIV).

	Frequentist	Bayesian
Magnet down	-0.0492 ± 0.0111	-0.0497 ± 0.0112
Magnet up	-0.0374 ± 0.0131	-0.0378 ± 0.0129
Combination of the two	-0.0443 ± 0.0084	-0.0446 ± 0.0086
Joint fit	-0.0441 ± 0.0085	-0.0443 ± 0.0085

TABLE XIV. A_{raw} for the two magnet polarities and their combination, by the two methods; the frequentist combination is the sum of the profile likelihoods of the two polarities, the Bayesian one the sequential update of the posterior.




FIG. XXVII. 68% (solid) and 95% (dashed) regions in (A, A_b): frequentist, from the profile likelihood with χ^2_2 thresholds, and Bayesian, highest-posterior-density regions of the chain. Dots: best-fit and posterior median.

4. Confidence and Credible Regions in (A, A_b)

In short: the 68% *and* 95% *regions of the frequentist profile (*ΔNLL *at the* χ^2_2 *values* 2.30 *and* 5.99*) and of the Bayesian highest posterior density coincide, and* $A = 0$ *lies outside both.* For two parameters of interest the Wilks statistic follows a χ^2 with two degrees of freedom (Fig. XXVII). The background asymmetry is $A_b = -0.0402 \pm 0.0176$ (Hesse; Minos $[-0.0578, -0.0226]$), correlated with A by $\rho = -0.16$ in the likelihood, -0.17 in the posterior, and -0.17 in the bootstrap. The point $A_b = 0$ is at the edge of the 95% region.

5. Hypothesis Test

In short: the exact binomial test on all candidates gives 6.1σ*, the test of the fit* $p = 2 \times 10^{-7}$ (5.2σ) *and the Bayes factor* $BF_{10} \approx 9 \times 10^3$*; they answer different questions. Hypothesis test.* The simplest test, the exact binomial test of $N^- = N^+$ on all candidates of the window, gives 6.1σ ($p = 1.3 \times 10^{-9}$, $A_{\text{count}} = -0.0432 \pm 0.0071$); it sees the asymmetry of signal and background together and does not use the mass. The frequentist test of the fit gives a p-value, 2×10^{-7} ($Z = \sqrt{q(0)} = 5.2\sigma$ in the two-sided convention; the one-sided $Z = \Phi^{-1}(1-p)$ would give 5.06σ, Appendix C.1), which does not depend on the prior and is not the probability of $A = 0$; its size and power were verified with toy-MC experiments (Sec. F.4). In particle physics a significance of 5σ is the conventional threshold for a discovery claim, chosen to limit the effect of multiple testing and of unforeseen systematic effects; the 5.2σ found here crosses it, but it refers to a raw asymmetry that includes production and detection effects (Sec. IX B), and is not a discovery of CP violation. The Bayes factor, $BF_{10} \approx 9 \times 10^3$, compares the evidence of A free and $A = 0$ and depends on the prior width: with $A \in (-0.3, 0.3)$ instead of $(-1, 1)$ it decreases by a factor 3.3. For A the two give the same conclusion. For the background asymmetry, $A_b = -0.040 \pm 0.018$, they would lead to different statements: a 2.3σ deviation from zero, which a frequentist test would call a hint, and $BF_{01} \approx 3.5$, a mild preference for $A_b = 0$ because a free parameter with a prior of width two is penalized (Sec. V D). Neither is wrong; they answer different questions.

6. Goodness of Fit

In short: the same fit has $p = 0.03$ *in* 2.5 *MeV bins and* $p \approx 5 \times 10^{-6}$ *in* 10 *MeV bins: the p-value of the binned* χ^2 *is a property of the binning as much as of the model. Goodness of fit.* The same fit of the same data gives $\chi^2/\text{ndof} = 441.7/388$ ($p = 0.031$) in 2×200 bins of 2.5 MeV and $\approx 160/88$ ($p \approx 5 \times 10^{-6}$) in 2×50 bins of 10 MeV (the Bayesian fit, 164.1/88, $p = 1.6 \times 10^{-6}$, uses the second binning). The misfit is at the scale of ~ 10 MeV, the flanks of the peak, and is diluted in narrow bins where Poisson fluctuations dominate; the p-value of a binned χ^2 is a property of the binning as much as of the model. The Bayesian analysis adds the posterior predictive check, which accounts for the posterior uncertainty

of the parameters and shows that the misfit is concentrated in the B^+ sample (Sec. D.13).

7. Nuisance Parameters and Validation

In short: the frequentist analysis profiles the eleven nuisance parameters and the Bayesian one marginalizes them; the results coincide because A is almost uncorrelated with them. Nuisance parameters. The frequentist analysis profiles the eleven nuisance parameters, the Bayesian one marginalizes them. The two differ when the likelihood is far from Gaussian or the prior is informative; here A is almost uncorrelated with the others (largest correlation -0.17, with A_b), and the results coincide.

Validation. The Bayesian toys are generated from the fitted model, which makes them a closure test; the bootstrap replicas of the frequentist check are resampled from the data and carry the features that the model does not describe. That both give the nominal coverage supports the interval, but they test different things (Sec. IV D).

8. Published Analysis: Samples and Models

In short: the published analysis has $22{,}119 \pm 164$ *signal candidates, a Cruijff signal and misidentified and partially reconstructed backgrounds; with the same D^0 veto the Bayesian fit gives* -0.0496 ± 0.0088. The published analysis of the same data [5] finds $A_{\rm raw} = -0.060 \pm 0.007$ with $22{,}119 \pm 164$ signal candidates, against -0.0441 ± 0.0085 and $15{,}328 \pm 133$ here. The two differ in the sample (the Open Data record is a preselected subset, reduced to 23,895 candidates by the identification requirement alone) and in the model, which includes a Cruijff signal, partially reconstructed four-body decays and a misidentified peaking background from simulation. With the same D^0 veto as the published selection the Bayesian fit gives -0.0496 ± 0.0088, within one standard deviation of the quadrature sum of the two uncertainties (0.93σ), although the samples overlap and the selections differ.

9. Published Analysis: Raw versus CP Asymmetry

In short: $A_{CP} \approx A_{\rm raw} - A_P - A_D$; the corrections move the published result from -0.060 *to* $-0.043 \pm 0.009 \pm 0.003 \pm 0.007$ *and cannot be assumed for a different selection.* The raw asymmetry is not the CP asymmetry. The two are related by $A_{CP} \approx A_{\rm raw} - A_P - A_D$ (Eq. (2)) and, in the published analysis, the correction obtained from $B^\pm \to J/\psi K^\pm$ decays and from the acceptance moves the result from -0.060 to $-0.043 \pm 0.009 \pm 0.003 \pm 0.007$ (3.7σ) [5], i.e. by $+0.017$, of the order of two statistical uncertainties of this work. A correction of that size cannot be assumed for a different selection, and the 5.2σ of this work is the significance of a nonzero raw asymmetry, which includes the production asymmetry of $B^\pm$ mesons and the different interaction of K^+ and K^- with the detector, not a significance of CP violation. A later measurement on a larger data set gives $A_{CP} = -0.037 \pm 0.002 \pm 0.002 \pm 0.003$ (8.5σ) [1, 2]. Our result is consistent in sign and size with these values, which is what can be said without the control channel, absent from this record.

10. Machine-Learning Cross-Check

In short: a cut at 0.5 *on the classifier score keeps* 92% *of the signal and* 36% *of the background and gives* -0.0451 ± 0.0087 *(Bayesian) and* -0.0447 ± 0.0086 *(frequentist).* A classifier trained on weak labels from the mass spectrum (test AUC 0.86) was used as an independent selection: a cut at 0.5 keeps 92% of the fitted signal and 36% of the background and leaves $A_{\rm raw}$ within its expected statistical spread (-0.0451 ± 0.0087 Bayesian, -0.0447 ± 0.0086 frequentist), without improving the precision; it cannot remove the misidentified decays that peak under the signal. Details, including the K-means partition of the Dalitz plane, are in Appendix I.

11. Bayesian and Frequentist Intervals at Low Statistics

In short: credible and confidence intervals coincide for $\sim 2 \times 10^4$ *candidates with flat priors, and stay equal, in value and coverage, down to* 50 *candidates.* With $\sim 2 \times 10^4$ candidates the posterior is Gaussian and, with flat priors, the Bayesian and frequentist results coincide by construction (Sec. V C). To find where they could differ, the same likelihood was analysed with both methods on toy-MC datasets of $N = 50$, 150, 500 and 2000 candidates, 300 for each size. The model is the six-parameter one of Sec. B.4 (Gaussian signal, one exponential background shared by the charges), which stays identifiable down to a few tens of events; the parameters are those of its fit to the data, with the yields rescaled. Each replica is analysed with the MCMC (median and equal-tailed credible intervals) and by maximum likelihood with `iminuit` (Hesse intervals, and Minos intervals from the profile likelihood and Wilks' theorem).

The two methods agree at every size (Fig. XXVIII). The coverages of the three intervals are compatible with nominal and with each other, the posterior median and the maximum-likelihood estimate differ on average by 5% of the statistical uncertainty at $N = 50$ and by less above, and the credible intervals stay symmetric (at $N = 50$ the shortest chain is only ≈ 10 autocorrelation times long, so the Monte Carlo error on the interval endpoints is larger there). The asymmetry is a ratio of Poisson yields, whose likelihood is close to Gaussian already with a few tens of signal events: in this problem flat-prior credible intervals and profile-likelihood confidence intervals give the same




FIG. XXVIII. Coverage of the (a) 68% and (b) 95% intervals on A vs the number of candidates, for the Bayesian credible interval, the profile-likelihood (Minos) interval and the Hesse interval of the same likelihood; 300 toy-MC datasets per point, binomial uncertainties. The dashed lines mark the nominal coverage (1σ, 2σ; the Bayesian 95% interval is the 2.5–97.5% one).

answer, and Wilks' theorem holds, down to $N = 50$. The two approaches differ in what the interval means and in hypothesis testing (Sec. V D), not in the numbers. At $N = 2000$ all three intervals over-cover slightly (0.98–0.99 for the 95% interval); since the effect is the same for the three methods, it is a feature of this set of replicas rather than of either inference.

Appendix I: Machine-Learning Analysis

In short: classifiers trained on weak labels from the mass spectrum (there are no true labels) reach a test AUC of 0.86*; a cut on their score leaves $A_{\rm raw}$ unchanged, and an adaptive partition of the Dalitz plane shows the same pattern as the fixed binning.* As a complementary, data-driven approach to the signal/background separation of the mass fits (Secs. IV–V), supervised classifiers were trained on weak labels derived from the mass spectrum, and used to study whether an ML-assisted event selection changes $A_{\rm raw}$ or the mass spectrum, and whether an adaptive partition of the Dalitz plane resolves local structure.

1. Background: Regularization, Bias-Variance and Loss Functions

In short: cross-entropy is a Bernoulli negative log-likelihood, $L1$ and $L2$ penalties shrink the coefficients, the test error splits into bias, variance and noise, and class weights rescale the score by a known factor. For a classifier with output probability $p(x)$ the binary cross-entropy $-\sum_i [y_i \ln p_i + (1-y_i)\ln(1-p_i)]$ is the negative log-likelihood of a Bernoulli distribution, the same principle (minimization of $-\ln L$) as in the mass fits; the Brier score $\frac{1}{N}\sum_i (p_i - y_i)^2$ [19] is the mean squared error of the predicted probability. The logistic regression is regularized by adding a penalty to the negative log-likelihood, $\alpha \sum_j \beta_j^2$ ($L2$, ridge: shrinks all the coefficients) or $\alpha \sum_j |\beta_j|$ ($L1$, lasso [20]: sets some of them exactly to zero), with α chosen by cross-validation (in `scikit-learn`, $C = 1/\alpha$ multiplies the likelihood term). For squared error the expected test error of an estimator $\hat{f}$ decomposes as $E[(y-\hat{f})^2] = \mathrm{Bias}^2 + \mathrm{Var} + \sigma^2_{\rm irr}$: flexible models lower the bias and raise the variance, and the optimal complexity is where the sum is minimal, which the train-test gap of gradient boosting against the number of stages makes visible.

When the classes are weighted, with weight w_c for class c, the minimized loss is $\sum_i w_{c_i} \ell_i$ and the classifier estimates $\tilde{p} = w_s p/(w_s p + w_b(1-p))$ instead of p. A monotonic recalibration (isotonic regression [21]) restores the probability scale without changing the ranking, hence the AUC.

2. Background: Classification Without Labels

In short: the classifier that is optimal between two mixtures of the same signal and background is monotonic in the signal-to-background ratio, so labels derived from the mass are sufficient. Let two samples be mixtures of the same signal and background densities $s(x)$ and $b(x)$, with signal fractions $f_1 \neq f_2$. The ratio of their feature densities, $p_1(x)/p_2(x) = [f_1\, s + (1-f_1)\, b]/[f_2\, s + (1-f_2)\, b]$, is a monotonic function of $s(x)/b(x)$, so that the classifier that is optimal between the two mixtures is, up to a monotonic transformation, the one that is optimal between signal and background [22]. The proof requires that s and b be the same in the two samples, which is the condition discussed in Appendix I.4.

3. Background: Clustering and Uncertainty Estimators

In short: K-means converges to a local minimum of the inertia, the adjusted Rand index measures the stability of a partition, and the spread of an ensemble estimates the epistemic uncertainty. K-means [23] alternates the assignment of each point to the nearest centroid and the update of the centroids to the cluster means; each step lowers the inertia W, so the algorithm converges to a local minimum that depends on the initialization (hence several random starts). The adjusted Rand index [24] between two partitions is $(\mathrm{RI} - E[\mathrm{RI}])/(\max \mathrm{RI} - E[\mathrm{RI}])$: it is 1 for identical partitions and 0 on average for random ones, and it is used to quantify the stability of the partition under a change of the initialization. The spread of the predictions of an ensemble, $\sigma^2(x) = \frac{1}{M}\sum_m [p_m(x) - \bar{p}(x)]^2$, estimates the epistemic part of the uncertainty when the members differ in the training data (bootstrap, as in the Random Forest) or in the initialization (deep ensemble [25]), and MC dropout [26] obtains M such members from a single network by keeping the dropout active at inference.

4. Weak Labels and Feature Set

*In short: a classifier separates a signal-enriched window from the mass sidebands (*15,130 *signal-like and* 1940 *background-like candidates,* 20 *features,* 60/20/20 *split); by the Neyman-Pearson argument of the CWoLa approach it is also the signal/background classifier, if the features carry no information on the mass.* No candidate carries a true signal/background label, so the classifiers are trained with weak labels, following the classification-without-labels (CWoLa) approach [22]: a classifier is trained to separate two *mixed* samples with different signal fractions, here a signal-enriched window around the B peak and the mass sidebands. If, within each class, the feature distributions are the same in the two samples, the optimal classifier between the two mixtures is a monotonic function of the signal-to-background likelihood ratio $p_s(x)/p_b(x)$, which by the Neyman-Pearson lemma is the most powerful test statistic for signal against background: the classifier trained on the mixtures is also the optimal signal/background classifier [22]. Two conditions follow. The background in the sidebands must be representative of the background under the peak; and the features must not be correlated with the B mass. The second condition cannot hold exactly: the reconstructed mass is not an input feature, but the nine momentum components of the three kaons determine it, so a flexible model can partly learn the mass, separate the two mass regions rather than signal from background, and *sculpt* the mass distribution of the candidates that pass a cut on its score. Appendix I.9 tests this directly.

The labels use the nominal Bayesian fit (Sec. V C, $\mu = 5284.39$, $\sigma = 18.00$ MeV). Candidates with $|m - \mu| < 2\sigma$, $m \in [5248.4, 5320.4]$ MeV, are labeled signal-like (15,130); candidates in $[5150, \mu - 4\sigma] \cup [\mu + 4\sigma, 5600] = [5150, 5212.4] \cup [5356.4, 5600]$ MeV are labeled background-like (1940); the 2–4σ gap is left unlabeled. The region 5100–5150 MeV, inside the fit window, is excluded from the background-like class to satisfy the first condition: there the background rises towards partially reconstructed B decays (Sec. II A), whose kinematics differ from those of the combinatorial background under the peak. The classes are imbalanced (1940/15,130 = 0.128). Every model is trained with the same balanced class weights, $w_c = N/(2N_c)$, i.e. 4.40 for background-like and 0.56 for signal-like candidates (through `class_weight` for the scikit-learn models and the network, and through sample weights for gradient boosting), so that the two classes contribute equally to the loss. Figure XXIX shows the three regions on the mass spectrum. (Down-sampling the signal-like class and up-weighting it afterwards would have the same effect with less training data; the weights keep all candidates.)




FIG. XXIX. Weak labels on the mass spectrum of the fit window: signal-like candidates ($|m - \mu| < 2\sigma$, 15,130) and background-like candidates (sidebands beyond 4σ and above 5150 MeV, 1940); the 2–4σ gap is left unlabeled and the region below 5150 MeV, where the background rises towards partially reconstructed decays, is excluded.

The feature set consists of the 6 per-track kinematic/PID variables (`PX`, `PY`, `PZ`, `ProbK`, `ProbPi`, `IPChi2`) for each of the 3 final-state tracks (18 features), plus two B-candidate variables not used in the preselection: `B_FlightDistance` and `B_VertexChi2`, standard discriminants against combinatorial background (a genuine B flies further before decaying and has a better-reconstructed vertex than a random track combination). Both separate the weak-label classes (Kolmogorov-Smirnov $p < 10^{-60}$) and are not redundant with the per-track features (Spearman $|\rho| \leq 0.53$ with each of the 18), giving 20 features. A principal-component analysis of the standardized features needs 15 of 20 components to reach 90% of the variance, so no dimensionality reduction is applied; with 20 features and $\sim 10^4$ training

candidates the sample is not in the regime where the distance-based methods suffer from the curse of dimensionality. The labeled sample was split 60/20/20 into training, validation and test sets (10,242/3414/3414 candidates, 388 background-like in the test set), stratified on the label, with the feature scaler fit on the training set only.

5. Model Comparison

In short: six classifiers; the tree ensembles and the neural network reach test AUC 0.857–0.864*, about* 0.09 *above the linear models and indistinguishable from each other within the test-set uncertainty (*0.012*); gradient boosting, best in cross-validation, is the reference.*

Model	CV AUC	Test AUC	Brier
Logistic Regression	0.769 ± 0.016	0.771 ± 0.013	0.197
Logistic Regression (L1)	–	0.771 ± 0.013	0.198
Decision Tree	0.816 ± 0.011	0.824 ± 0.013	0.133
Random Forest	0.863 ± 0.017	0.860 ± 0.012	0.122
Gradient Boosting	$\mathbf{0.866 \pm 0.016}$	0.864 ± 0.012	0.116
FFNN	–	0.857 ± 0.012	0.134

TABLE XV. Cross-validated AUC (mean and standard deviation over the folds), test-set AUC (with bootstrap uncertainty) and test-set Brier score of the raw scores. Gradient boosting, best in cross-validation, is the reference model. The FFNN is validated on the validation set instead of by cross-validation.

Six classifiers were trained: logistic regression, without and with an $L1$ penalty (linear baselines), a decision tree (depth 4, fixed in advance as a shallow, interpretable baseline and not pruned), a Random Forest [27] (300 trees of depth 8), gradient boosting [28] (200 stages of depth-3 trees, learning rate 0.05), and a feed-forward neural network (FFNN: two hidden layers of 32 and 16 units with ReLU activation, dropout 0.2, binary cross-entropy, the Adam optimizer [29] with learning rate 10^{-3}, early stopping on the validation AUC, sigmoid output). The linear models and the network are parametric, with a fixed number of parameters, and the trees and ensembles non-parametric, whose complexity grows with the data; the more flexible models are less interpretable, which is why the linear and single-tree baselines are kept. Both the logistic-regression log-loss and the network's binary cross-entropy are the negative log-likelihood of a Bernoulli distribution with probability $p(x)$, the same quantity minimized by the mass fits. The models are compared through the area under the receiver operating characteristic (ROC) curve (AUC), which equals the probability that a randomly chosen signal-like candidate receives a higher score than a randomly chosen background-like one (0.5 for a random classifier, 1 for a perfect one). Table XV gives the 5-fold cross-validated AUC on the training set and the AUC on the untouched test set, with its bootstrap uncertainty. The reference model for the rest of the section was chosen on the cross-validated AUC, so that the test set is not used for any choice: gradient boosting (0.866), essentially tied with the Random Forest (0.863).




FIG. XXX. ROC curves on the test set, with the test AUC of each model in the legend (Table XV); the reference model is drawn thicker, and the two logistic regressions overlap. The circle marks the cut on the reference score used in Appendix I.9.

The three non-linear ensembles and the FFNN improve the AUC by ∼0.09 over the linear models and by ∼0.04 over the single tree (Fig. XXX), while their mutual differences (≤ 0.007) are smaller than the test-set uncertainty (0.012). The FFNN training is reproducible (fixed seeds, deterministic operations); its dependence on the random initialization, measured by retraining it with ten seeds, is 0.854 ± 0.005 in test AUC.

The two measures of the relevance of a feature available here – the Random Forest importance (the decrease in Gini impurity $G = 1 - \sum_c p_c^2$, p_c the class fractions in a node, produced by the splits on that feature) and the magnitude of the $L1$-regularized (Lasso [20]) logistic-regression coefficient of the standardized feature – agree only partly (Fig. XXXI; Spearman correlation of the two rankings 0.42, $p = 0.06$). The impact-parameter variables `IPChi2` dominate the Random Forest, while the Lasso coefficients are largest for `B_FlightDistance`, `H3_IPChi2`, `H2_PZ` and `B_VertexChi2`. The difference is expected: the Gini importance reflects the use of a feature in non-linear splits and interactions, and is shared among correlated features, while a linear coefficient measures the effect of a feature with the others held fixed. Both measures rank the two B-candidate variables among the top five (Random Forest 4th and 5th, Lasso 1st and 4th): they summarize in one number the displaced-vertex topology that the per-track variables encode only jointly.




FIG. XXXI. Random Forest feature importance (Gini, a) and magnitude of the Lasso coefficient of the standardized feature (b) for the 20 features, both ordered by Random Forest importance; the two B-candidate variables are highlighted.

6. Regularization, Bias-Variance, and Calibration

In short: the linear model does not overfit, gradient boosting overfits beyond ∼150 stages without losing test AUC, and class weights bias the probabilities, which an isotonic recalibration corrects (Brier score $0.122 \to 0.062$*) without changing the ranking.* A scan of the regularization strength of the logistic regression (penalties and loss functions are defined in Appendix I.1) (30 log-spaced values of $C = 1/\alpha$, cross-validated) gives the same optimal AUC for the $L1$ and $L2$ penalties (0.7686, at $C_{L1} = 0.12$ and $C_{L2} = 9.2$), and the optimal $L1$ penalty sets no coefficient to zero: the linear model does not overfit, and its limitation is the linear decision boundary, not the variance (Fig. XXXII). The two curves differ only at strong regularization: the L_2 model, whose coefficients are shrunk but not removed, is flat over the whole range, while the L_1 model loses ~ 0.2 of AUC for $C \lesssim 3 \times 10^{-3}$, where its coefficients are set to zero.




FIG. XXXII. Cross-validated AUC of the logistic regression against the inverse regularization strength $C = 1/\alpha$ (30 log-spaced values), for the L_1 (Lasso) and L_2 (Ridge) penalties; the dotted lines mark the optimum of each.

The bias-variance trade-off was studied with gradient boosting, whose complexity grows with the number of boosting stages (Fig. XXXIII). The test AUC rises up to ∼150 stages and then stays at 0.865, while the training AUC keeps rising: the train-test gap grows from 0.018 at 20 stages to 0.084 at 300. The additional stages fit fluctuations of the training sample without improving the ranking of new candidates – overfitting, which the default 200 stages of Table XV do not reach in the test AUC. The test accuracy at the 0.5 threshold stays below that of always predicting the majority class (0.886): with balanced class weights the classifier is trained to minimize the weighted error, in which a misclassified background-like candidate counts eight times more than a signal-like one, and not the unweighted error rate.

The AUC depends only on the ranking of the scores; whether a score can be read as a probability is measured by the Brier score [19], $\mathrm{BS} = \frac{1}{N}\sum_i (p_i - y_i)^2$, the mean squared difference between predicted probability p_i and label $y_i \in \{0, 1\}$ (lower is better). Class weights change the probabilities a model learns: a classifier trained with weights w_s, w_b on the two classes converges to

$$\tilde{p} = \frac{w_s p}{w_s p + w_b (1 - p)}$$

instead of the true p, i.e. it describes a world in which




FIG. XXXIII. Gradient boosting: (a) accuracy at the 0.5 threshold and (b) AUC on the training and test sets as a function of the number of boosting stages. The dotted line is the accuracy of always predicting the signal-like class.




FIG. XXXIV. Reliability diagram of the Random Forest on the test set: observed fraction of signal-like candidates vs. predicted probability, for the raw score (with binomial uncertainties) and for the isotonic-recalibrated one. Bottom: distribution of the raw score for the two weak-label classes; the low-probability bins contain few candidates.

the classes are re-balanced. With $w_b > w_s$ the raw scores therefore *underestimate* the probability of the signal-like class, which is what the reliability diagram of the Random Forest shows (Fig. XXXIV): at a raw score of 0.25 the observed signal-like fraction is already 0.58. An isotonic recalibration [21], fitted on held-out folds of the unweighted training sample, follows the diagonal and halves the Brier score, from 0.122 to 0.062 (the Brier score is the mean squared error of the predicted probability against the 0/1 label). Most of the gain undoes the re-weighting rather than correcting the model; the ranking, and hence the AUC and any selection based on a score threshold, is unchanged by a monotonic recalibration.

7. Uncertainty Quantification

In short: the three estimators give mean per-candidate uncertainties of 0.178 *(Random Forest),* 0.095 *(MC dropout) and* 0.050 *(deep ensemble); they measure different sources of variability, and only the Random Forest spread visibly decreases with the training-set size.*

The uncertainty of a prediction has two components: *aleatoric*, from the intrinsic overlap of the classes in feature space, which more training data cannot reduce, and *epistemic*, from the limited knowledge of the model parameters, which more training data reduces. Three estimators were compared, each giving an ensemble of M predictions $p_m(x)$ for each candidate and quoting their spread, $\sigma(x) = \left[\frac{1}{M}\sum_m (p_m(x) - \bar{p}(x))^2\right]^{1/2}$: the $M = 300$ trees of the Random Forest, each trained on a bootstrap sample; $M = 300$ forward passes of the FFNN with dropout kept active at inference (MC dropout), which approximates sampling from a Bayesian posterior over the network weights [26] (a full Bayesian network trained by variational inference was not used: it needs a dedicated implementation and MC dropout gives the same kind of ensemble at no extra cost); and a deep ensemble [25] of $M = 10$ FFNNs trained from different random initializations. The mean per-candidate uncertainty is 0.178 for the Random Forest, 0.095 for MC dropout (0.099 ± 0.006 over the ten initializations) and 0.050 for the deep ensemble (Fig. XXXV), with the same values within 10% for background-like and signal-like candidates.

The three estimators answer different questions, which explains the factor ∼3.5 between them. The deep ensemble only varies the initialization, on the same training data; MC dropout also perturbs the network structure; the trees of the Random Forest differ in their training data (bootstrap) and in the features they use, and each




FIG. XXXV. (a) Spread across the Random Forest trees vs. mean tree score, median and 16–84% band of the test candidates of each weak-label class. (b) Distributions of the per-candidate uncertainty from the Random Forest trees, MC dropout and the deep ensemble. (c) Mean Random Forest spread vs. training-set size, for all, background-like and signal-like test candidates.

tree, of limited depth, is individually a poor and discrete estimator of p. Their per-candidate values are correlated but not interchangeable (Spearman correlations 0.41–0.57 between pairs). The Random Forest spread contains both components. It is largest for scores around 0.5 (Fig. XXXVa), where the classes overlap and the trees disagree most – an aleatoric contribution; and it decreases with the size of the training set, from 0.220 with a quarter of it to 0.204 with half and 0.179 with the full set, while the test AUC stays at 0.860–0.862 – the epistemic contribution, which more data reduces.

8. K-means on the Dalitz Plot

In short: for $K = 8$ two clusters deviate by more than 3σ ($\chi^2 = 46.4$ for 7 degrees of freedom): a negative asymmetry at low $m^2_{\rm lo}$ and positive ones at high $m^2_{\rm hi}$, as in the fixed binning. K-means is the unsupervised part of this analysis: it uses no labels, and the partition of the Dalitz plane is learned from the density of the candidates rather than fixed by hand. As an adaptive alternative to the fixed Dalitz-plot binning of Sec. VIII, the candidates of the fit window were partitioned with K-means [23] on the two standardized Dalitz variables $(m^2_{\rm lo}, m^2_{\rm hi})$. K-means partitions the points x_i into K clusters C_k minimizing the within-cluster sum of squares (inertia) $W = \sum_k \sum_{i\in C_k} \|x_i - \bar{x}_k\|^2$, with $\bar{x}_k$ the cluster centroids; K has to be fixed in advance.

The quality of a partition is measured by the silhouette [30]: for each point, $s_i = (b_i - a_i)/\max(a_i, b_i)$ compares its mean distance a_i to the other points of its cluster with the mean distance b_i to the points of the nearest other cluster, and the average over all points ($-1 \le s \le 1$, higher is better) measures how well separated the clusters are. $K = 8$ was fixed in advance, as a compromise between resolution and the number of candidates per cluster (at least 1233), and $K = 3$, the silhouette optimum over $K = 2$–12 (0.46, against 0.41 for $K = 8$, a local maximum), was added. Both partitions are stable: repeating the clustering with a different random initialization gives an adjusted Rand index [24] above 0.998. Figure XXXVI shows the inertia and the silhouette against K: the inertia decreases monotonically and has no marked elbow, the silhouette has a global maximum at $K = 3$ and a local one at $K = 8$.

The asymmetry of each cluster was compared with the global one as in Sec. VIII, $S_k = (A_k - A_{\rm glob})/\sigma(A_k)$. For $K = 8$ (Fig. XXXVII), two clusters deviate by more than 3σ (probability 2×10^{-4} for two or more if the asymmetry were uniform), and the homogeneity test gives $\chi^2 = 46.4/7$ ($p = 7 \times 10^{-8}$). The most significant one lies at low $m^2_{\rm lo}$ (1–2.5 GeV^2) and intermediate $m^2_{\rm hi}$, with $A_k = -0.127 \pm 0.017$ ($n = 3601$, $S_k = -5.1$); two clusters at high $m^2_{\rm hi}$ have positive asymmetries, $A_k = +0.020 \pm 0.020$ ($S_k = +3.2$) and $+0.040 \pm 0.028$ (+2.9). In the signal window ($|m - \mu| < 2\sigma$) the same pattern is sharper – three clusters beyond 3σ, with $S_k = -5.6$, +4.4 and +3.0, and $\chi^2 = 62.3/7$ – so it is not produced by the background. The coarser $K = 3$ partition gives $\chi^2 = 15.2/2$ ($p = 5\times10^{-4}$). The adaptive partition therefore confirms the picture of the fixed binning, a negative local asymmetry at low $m^2_{\rm lo}$ and positive ones at high $m^2_{\rm hi}$, with the same caveats (no background subtraction, no correction for detection asymmetries).

9. Stability of $A_{\rm raw}$ under ML Selection

In short: a cut on the score at 0.5 keeps 92% of the signal and 36% of the background and leaves $A_{\rm raw}$ within its expected statistical spread with both fits and with counting; it does not improve the precision and cannot remove the misidentified background.

The reference (gradient-boosting) score was applied to all candidates of the fit window (on the test set, a cut at 0.5 keeps 88% of the signal-like and 28% of the background-like candidates, Fig. XXX), not only to the




FIG. XXXVI. (a) Inertia and (b) mean silhouette of the K-means partition of the Dalitz plane for $K = 2$–12; the dashed line marks the $K = 8$ fixed in advance and the dash-dotted line the maximum of the silhouette, $K = 3$.




FIG. XXXVII. K-means partition ($K = 8$) of the Dalitz plane, all candidates of the fit window, same axes as Fig. VII. Color and labels: local significance of each cluster with respect to the global asymmetry, $S_k = (A_k - A_{\rm glob})/\sigma(A_k)$, on the same scale as panel (c) of Fig. VII.




FIG. XXXVIII. Counting asymmetry after a cut on the gradient-boosting score, vs. threshold, with its binomial uncertainty. Dotted line: counting asymmetry of the full fit window; band: expected statistical spread of a nested subsample around it, $\pm\sqrt{\sigma^2_{\rm sub} - \sigma^2_{\rm full}}$.

labeled ones used in training, and the counting asymmetry was recomputed after cuts on the score from 0.1 to 0.9 (Fig. XXXVIII). Since every selected sample is contained in the full one, the expected statistical spread of the difference is $\sqrt{\sigma^2_{\rm sub} - \sigma^2_{\rm full}}$, not the uncertainty of either estimate. Over all thresholds, which keep from 18,443 to 2621 candidates, the shift with respect to the full window ($A_{\rm count} = -0.0432$) stays within 1.4 times this spread: the selection does not introduce a detectable bias in the counting asymmetry. Split by magnet polarity at a cut of 0.5, the counting asymmetries are -0.060 ± 0.010 (down) and -0.024 ± 0.012 (up), 2.2σ apart; the frequentist fit of the same samples, which separates the background, gives -0.054 ± 0.011 and -0.032 ± 0.013, 1.3σ apart (against 0.7σ for the full sample, Sec. D.15), so most of the larger difference of the counting estimates comes from the background.

Counting cannot reveal whether the selection deforms the mass spectrum. To test this, the selected sample (score > 0.5, 15,709 candidates) was refitted with the nominal mass model by both inference methods of this work, the Bayesian fit of Sec. V and the frequentist maximum-likelihood fit of Sec. IV, and the three estimates were compared with those of the full sample (Table XVI, Fig. XXXIX). The asymmetry is stable: all three shifts are within their expected nested-sample spread, and the significance of $A \neq 0$ from the profile likelihood (Wilks' theorem [8]) is 5.2σ in both samples. The selection keeps 92% of the fitted signal and 36% of the background ($N_s = 14{,}121$, $N_b = 1590$), but this does not improve the precision on $A_{\rm raw}$ (0.0087 against 0.0085): the background under the peak is already small (≈4% in the signal window) and the fit separates it from the signal anyway.

The mass spectrum, instead, changes. The goodness

Method	Full sample	Score > 0.5	Shift (expected)
Bayesian fit	-0.0443 ± 0.0085	-0.0451 ± 0.0087	$-0.0008\,(0.0018)$
Frequentist fit	-0.0441 ± 0.0085	-0.0447 ± 0.0086	$-0.0006\,(0.0012)$
Counting	-0.0432 ± 0.0071	-0.0446 ± 0.0080	$-0.0014\,(0.0036)$

TABLE XVI. A_{raw} on the full fit window (19,731 candidates) and after the cut on the gradient-boosting score (15,709), by the three methods. The expected shift is the statistical spread of a nested subsample, $\sqrt{\sigma^2_{\text{sub}} - \sigma^2_{\text{full}}}$. Counting includes the background.




FIG. XXXIX. A_{raw} from the Bayesian fit, the frequentist fit and counting, for the full sample and the ML-selected one (Table XVI).

of fit of the nominal model degrades from $\chi^2 = 164/88$ to 253/88 in the Bayesian fit and from 442/388 to 557/388 in the frequentist one (with a different binning), and the background shape changes: the slope of the slow exponential grows from 0.0028 to 0.0071 MeV^{-1} (B^+) and from 0.0016 to 0.0057 MeV^{-1} (B^-), and the fraction of the steep one from 0.11 to 0.28 and from 0.21 to 0.50.

Two mechanisms can produce this: the classifier could reconstruct the mass from the kaon momenta and sculpt the background (Appendix I.4), or the selection could change the composition of the background. The first is excluded: the same classifier trained without the nine momentum components (test AUC 0.860) gives the same degradation ($\chi^2 = 533/388$).

The efficiency of the cut as a function of the mass shows the second (Fig. XL): it is 0.89–0.90 at the peak, ≈ 0.70 below 5150 MeV and 0.06–0.13 above 5375 MeV. The classifier does what it is trained for – it removes the combinatorial background and keeps genuine B decays, including the partially reconstructed ones and the other non-combinatorial components that the weak labels cannot separate from the signal. In the selected sample these components weigh relatively more, and the nominal model, which has no dedicated component for them, describes it worse, mostly under the peak (the χ^2 of the $\pm 4\sigma$ region grows from 75 to 134 for 30 bins) and below it. A Crystal Ball signal gives the same degradation (407 → 522 for 386 degrees of freedom), so it is not due to the signal tails.

A_{raw} is unaffected because it is determined by the peak and the background shape is free in the fit, but a measurement that relied on the background shape, or on a background model taken from the sidebands of the full sample, would be biased by the selection.

In the nominal sample the selection cannot improve the precision because there is little background to remove. To test it where the background is larger, the same study was repeated on a sample with a looser kaon identification, `ProbK` > 0.3 (33,328 candidates, signal purity 94% within $\pm 2\sigma$ of the peak), with the reference classifier retrained on the weak labels of this sample (test AUC 0.895). Cleaning it with a cut on the classifier score is compared with tightening the identification cut, at equal signal efficiency (Fig. XLI). The classifier removes the combinatorial background with little signal loss – a cut at 0.5 keeps 93% of the signal and raises the purity to 99% – and at any efficiency gives a slightly smaller $\sigma(A_{\text{raw}})$ than the identification cut. The gain is small (0.0072 → 0.0070 at best): loosening the identification adds mostly signal (+47% with respect to the nominal selection), and the background under the peak stays a small fraction.

The classifier also cannot remove the background that matters for A_{raw}. With the looser identification the asymmetry moves to -0.055, as expected from misidentified $B^\pm \to \pi^\pm K^+ K^-$ decays (Sec. VI); tightening the identification cut to `ProbK` > 0.4 moves it back by $+0.008$ (3.3 times the nested-sample spread), while the score cut that keeps 93% of the signal moves it by -0.001. Misidentified B decays peak close to the B mass, so the mass-based weak labels mark them as signal-like: the CWoLa classifier, by construction, cannot learn to reject a background that sits under the peak, and particle identification remains necessary.

10. Hyperparameter Optimization

In short: the best configurations improve the test AUC by at most 0.005*, less than its uncertainty, and the cross-validated maximum overestimates the test AUC (winner's curse).* The hyperparameters of the Random Forest (120 configurations), gradient boosting (100) and the FFNN (168: seven architectures, four dropout rates, six learning rates) were optimized by repeated cross-validation on the training set (Tables XVII–XVIII). The best con-




FIG. XL. (a) Mass spectrum of all candidates and of those with gradient-boosting score > 0.5, for the 20-feature classifier and for one trained without the kaon momentum components (25 MeV bins). (b) Fraction of candidates passing the cut. Hatched: the region below 5150 MeV excluded from the background-like labels; grey: the signal-like window.




FIG. XLI. High-background sample (`ProbK` > 0.3): (a) statistical uncertainty and (b) value of A_{raw} from the frequentist fit vs signal efficiency, for tighter identification cuts (`ProbK` $> 0.3, \ldots, 0.8$) and for cuts on the score of the classifier retrained on this sample $(0, \ldots, 0.9)$. The dotted line marks the nominal selection (`ProbK` > 0.5). The samples are nested, so their uncertainties are correlated.

figurations improve the test AUC over the defaults of Table XV by 0.005 (Random Forest), 0.001 (gradient boosting) and 0.004 (FFNN), all well below the test-set uncertainty (0.012) and, for the FFNN, comparable with its spread over initializations (0.005). For the two tree ensembles the best cross-validated AUC is higher than the test AUC of the same configuration (0.871 vs 0.866 and 0.868 vs 0.865), as expected when the maximum is taken over many noisy cross-validation estimates (the "winner's curse"): the selection favors configurations with a favorable fluctuation, and the cross-validated score of the selected configuration is an optimistic estimate of its performance. Figure XLII shows the cross-validated AUC of the 168 network configurations by architecture: the best value rises from 0.839 (one layer of 8 units) to 0.858 (two or three layers) and then saturates (0.857 and 0.854 for four and five layers), while the spread within each architecture is dominated by the learning rate and the dropout. The searches show that the defaults are already close to optimal for this problem, so the defaults are used throughout.

11. Machine Learning and the Fits

In short: the machine-learning selection is a cross-check: a cut at 0.5 *leaves* A_{raw} *within its expected spread with both methods, does not improve the precision, and cannot remove the misidentified decays.* The machine-learning selection was used as a cross-check of the fits, not as a replacement. A cut on the gradient-boosting score at 0.5 keeps 92% of the fitted signal and 36% of the background and does not change A_{raw} beyond its expected statistical spread, with both inference methods (-0.0451 ± 0.0087 Bayesian, -0.0447 ± 0.0086 frequentist, against -0.0443 and -0.0441 for the full window). It does not improve the precision, 0.0087 against 0.0085, because the nominal sample contains little background.

Model	Grid	Best configuration	CV AUC / Test AUC (default)
Random Forest	`n_estimators` 200–1200, `max_depth` 6–12 or none, `min_samples_leaf` 5–40	400, 12, 5	0.8706 / 0.8656 (0.8604)
Gradient Boosting	`n_estimators` 100–500, `max_depth` 1–5, `learning_rate` 0.01–0.1	300, 2, 0.1	0.8684 / 0.8650 (0.8643)

TABLE XVII. Hyperparameter search for the tree ensembles (120 and 100 configurations, 5-fold cross-validation repeated 10 times, balanced class weights). The test AUC of the best configuration is compared with that of the default of Table XV (in parentheses); the bootstrap uncertainty of each test AUC is 0.012.

Architecture	Layers	Dropout	Learning rate	CV AUC
`deep_3layer` (best)	[64, 32, 16]	0.2	0.003	0.8585 ± 0.0174
`wide_2layer`	[64, 32]	0.3	0.003	0.8582 ± 0.0175
`deep_3layer`	[64, 32, 16]	0.3	0.003	0.8580 ± 0.0187
`wide_2layer`	[64, 32]	0.3	0.005	0.8579 ± 0.0184
`wide_2layer`	[64, 32]	0.4	0.005	0.8579 ± 0.0187

TABLE XVIII. Top five of the 168 FFNN configurations (7 architectures from one hidden layer of 8 units to five layers [128, 64, 32, 16, 8], dropout 0.1–0.4, learning rates 0.0002–0.005; 5-fold cross-validation repeated twice). The five are within 0.0007 of each other, far below the fold-to-fold spread: the optimum is a plateau. The best configuration has test AUC 0.8606, against 0.8568 for the default architecture (±0.005 over initializations).




FIG. XLII. Cross-validated AUC of the 168 FFNN configurations (dots) for each of the seven architectures, ordered by number of units; the black segment marks the best configuration of each architecture.

In a sample with a looser kaon identification (33,328 candidates) the classifier raises the purity from 94% to 99% and reduces $\sigma(A_{\text{raw}})$ from 0.0072 to 0.0070 at best, and it cannot remove the misidentified decays that move A_{raw} to −0.055: they peak under the signal, and weak labels derived from the mass spectrum mark them as signal-like (Appendix I.9). The useful role of the machine-learning analysis is therefore that of a consistency test of the asymmetry under a different event selection, and the partition of the Dalitz plane that it provides (below). Gaining precision or controlling the peaking background would require labels that carry information beyond the mass, such as simulation.

12. Local Asymmetries: Fixed Binning and K-Means

In short: the fixed grid and the K-means partition show the same pattern, a negative asymmetry at low m^2_{lo} and positive ones at high m^2_{hi} ($\chi^2 = 80.8/31$ and $46.4/7$). Two partitions of the Dalitz plane were used, a fixed 8×8 grid (Sec. VIII) and an adaptive K-means partition with $K = 8$ (Appendix I.8). Both find that the asymmetry is not uniform: $\chi^2 = 80.8$ for 31 degrees of freedom for the grid and 46.4 for 7 for K-means ($p = 3 \times 10^{-6}$ and 7×10^{-8}). Both place the negative local asymmetry at low m^2_{lo} and intermediate m^2_{hi} ($A = -0.128 \pm 0.018$ in the region of the two significant bins, -0.127 ± 0.017 in the most significant cluster) and positive deviations at high m^2_{hi}. The two partitions have different bins, so the numbers are not independent, and the K-means p-value benefits from the larger bins; the agreement is in the pattern. In the region defined by the published analysis, fixed before looking at the data, both fits give a compatible value (-0.234 ± 0.025 Bayesian, -0.237 ± 0.025 frequentist, 9.4σ from zero) with -0.239 ± 0.020. All local numbers are counting asymmetries or fits of the integrated mass spectrum in a region, with no background subtraction by event weights and no correction for detection asymmetries (Appendix J.5).

Appendix J: Conclusions, Scope and Limitations

In short: the scope of the study and the channels of the record that are not analysed, the complete text of the conclusions, and the limitations of the result.

1. Scope of the Study and Channels of the Record

In short: the aim is a methodological comparison of the two frameworks on the same data, not a measurement of A_{CP}, which this record cannot give for lack of the control channel; the other three channels are not analysed. In this work we use the LHCb Open Data record for $B^{\pm} \to K^{\pm}K^{+}K^{-}$ [3, 4], a subset of the 2011 data on which the first evidence of an inclusive CP asymmetry in this decay was published [5], to address the question: *to what extent do frequentist likelihood fits and Bayesian inference produce consistent, well-controlled estimates of the raw asymmetry $A_{\rm raw}$ in this decay?* The aim is not a competitive measurement of A_{CP}, but a methodological comparison of the two inference frameworks on the same data and the same physical question. Of the four channels in the record, $B^{\pm} \to K^{\pm}K^{+}K^{-}$ has the lowest background; the others ($\pi^{\pm}\pi^{+}\pi^{-}$, $K^{\pm}\pi^{+}\pi^{-}$, $\pi^{\pm}K^{+}K^{-}$) need a different particle-identification selection and a model of larger and peaking (misidentified) backgrounds, and are not analysed, to keep the focus on the comparison of the methods. The published A_{CP} cannot be reproduced from this record, which does not contain the control channel used to correct $A_{\rm raw}$ for the production and detection asymmetries (Appendix J.5); what we extend is the statistical treatment: a common model fitted with two inference methods, a common systematic study, a validation of the intervals, and a machine-learning (ML) selection, described in Appendix I, whose effect on the measurement is tested. Section II describes the dataset and the preselection common to the analyses and Sec. III the mass model and likelihood common to the two mass fits; Secs. IV and V present the frequentist and Bayesian fits, Sec. VI their common systematic study, Sec. VII the validation of the estimator and of the intervals with toy-MC experiments, Sec. VIII the local asymmetry in the Dalitz plane, and Sec. IX compares the two approaches; the machine-learning selection is in Appendix I. The main results are collected in Table I.

2. Headline Result

In short: $A_{\rm raw} = -0.0441 \pm 0.0085 \pm 0.0072$ (frequentist) and $-0.0443 \pm 0.0085 \pm 0.0070$ (Bayesian), 5.2 standard deviations from zero, stable under the thirteen variants; it is a raw asymmetry, not a significance of CP violation. We have measured the raw charge asymmetry of $B^{\pm} \to K^{\pm}K^{+}K^{-}$ decays in the 2011 LHCb Open Data record with a common selection and a common mass model, and compared how two inference frameworks treat the same question. The frequentist profile-likelihood fit and the Bayesian posterior give $A_{\rm raw} = -0.0441 \pm 0.0085 \pm 0.0072$ and $-0.0443 \pm 0.0085 \pm 0.0070$ (statistical, systematic), a nonzero asymmetry at 5.2 standard deviations that is stable under the thirteen variants of the fit and selection. The result is compatible in sign and size with the published analysis of the same data and with its later, more precise measurements, and an asymmetry concentrated in a region of the Dalitz plane, outside the $\phi(1020)$, is reproduced a priori in the region defined by the published analysis. It is a raw asymmetry: without the control channel that corrects for the production of $B^{\pm}$ mesons and for the detection of $K^{\pm}$, it is not a measurement of A_{CP}, and its significance is not that of CP violation (Sec. IX B).

3. Lessons of Each Method

In short: the frequentist fit gives intervals with nominal coverage and an unbiased estimator; the Bayesian one the same interval with flat priors plus diagnostics to check, a Bayes factor that depends on the prior and a predictive check; the ML selection leaves $A_{\rm raw}$ unchanged. Each method taught something specific.

- *Frequentist.* With the other eleven parameters profiled, Wilks' theorem gives a 68% interval with the nominal coverage ($68.4 \pm 1.5\%$ in 1000 bootstrap replicas of the data, 0.70 ± 0.01 and 0.95 ± 0.01 in toy-MC experiments from the fitted model, where the estimator is also unbiased and reaches the minimum variance bound), and the Hesse and Minos uncertainties agree, so the likelihood is parabolic in A at this sample size.
- *Bayesian.* With $\sim 2 \times 10^4$ candidates and flat priors the posterior is the likelihood, and the credible interval coincides with the confidence interval and has the same coverage in toy experiments. The Bayesian analysis adds what the frequentist one lacks: sampling diagnostics that must be checked rather than assumed (autocorrelation time, dispersed starts, label switching of the two background slopes), a Bayes factor that depends on the prior width, and a posterior predictive check whose charge statistics, being fitted, have no power.
- *Machine learning (Appendix I).* Weak labels from the mass spectrum are enough to train classifiers of test AUC $\simeq 0.86$, but the selection does not improve the precision on $A_{\rm raw}$ in a sample with little background, and it cannot remove the misidentified decays that peak under the signal. Its value is as an independent test of the stability of $A_{\rm raw}$ under a different selection and, with K-means, as an adaptive description of the Dalitz plane that agrees with the fixed binning.

4. Systematic Study and Dependence on Choices

In short: the particle-identification requirement dominates, and several statistics depend on choices: the binned χ^2 on the binning, the Bayes factor on the prior, the coverage on how the toy experiments are generated. On the systematic study, the dominant contribution is the particle-identification requirement, which a finer scan

shows to be consistent with a statistical fluctuation of a nested sample; the background model is second, and the signal shape does not matter. The same variants and the same combination rule give the same uncertainty in the two methods. More generally, several statistics that look informative depend on choices: the p-value of a binned χ^2 on the binning, the Bayes factor on the prior, the coverage on whether the toy-MC experiments come from the model or from the data.

5. Limitations and Future Work

In short: no signal model reaches $\chi^2/\mathrm{ndof} \approx 1$ *although* A_{raw} *does not depend on it, the machine-learning labels are weak and imbalanced,* A_{raw} *is not corrected for production and detection asymmetries, and the natural next step is a control-channel correction.* Several modeling choices and known weaknesses limit how far the results of Secs. V–VI and Appendix I can be pushed, and are collected here rather than scattered through the results sections.

On the mass fits (Secs. V–VI):

- no tested signal model reaches $\chi^2/\mathrm{ndof} \approx 1$ (the best, a Crystal Ball, gives 131.0/86, Sec. D.13; the p-value of the binned χ^2 is 5×10^{-6} at 10 MeV and 0.03 at 2.5 MeV, Sec. F.5), although A_{raw} does not depend on the signal shape;
- the systematic uncertainty, 0.007, is almost as large as the statistical one and is dominated by a PID variant that a finer scan identifies as a statistical fluctuation (Sec. VI); it is therefore conservative, but the combination rule cannot separate the two;
- the Savage-Dickey Bayes factor of Sec. V D relies on a Gaussian approximation of the posterior far in its tail and depends on the prior width, although the conclusion (decisive evidence against $A = 0$) is robust;
- the toy-MC coverage check of Sec. D.16 is a closure test and cannot detect model misspecification;
- the local Dalitz asymmetries (Secs. VIII and I.8) use a coarse partition, a mass-window cut rather than a proper background subtraction, and no correction for momentum-dependent detection asymmetries.

On the machine-learning side (Appendix I): the weak labels are imbalanced (1940 background-like vs. 15,130 signal-like candidates, Appendix I.4), and the test set contains only 388 background-like candidates, so differences in test AUC below $\sim$0.01 cannot be resolved; the three uncertainty estimators differ by a factor $\sim$3.5 in scale and measure different sources of variability (Appendix I.7); and the score-based selection changes the composition of the background, keeping the partially reconstructed and other non-combinatorial B decays that the nominal model does not describe (Appendix I.9); removing the momentum features does not help, since the weak labels cannot separate these decays from the signal.

More generally, no detector-level systematic correction (selection efficiency, production asymmetry) is applied anywhere in this work, so A_{raw} as reported throughout is explicitly *not* a measurement of A_{CP}; in the language of unfolding, the measured asymmetry is the folded one, and mapping it back to the true asymmetry would require the response of the detector (acceptance, charge-dependent efficiency, interaction of kaons with matter), which is not available in this record (Sec. IX). The published analysis corrects for $A_P + A_D$ with $B^{\pm} \to J/\psi K^{\pm}$, $J/\psi \to \mu^+\mu^-$ [1], which is not part of this record; the $J/\psi \to K^+K^-$ decays inside our sample cannot replace it, since this mode is rare and the 568 candidates under the B peak in the J/ψ band are dominated by charmless decays ($A_{\mathrm{count}} = -0.165 \pm 0.041$, a local Dalitz-plot asymmetry rather than a control measurement).

The most natural next steps, in rough priority order, are: correcting A_{raw} for the production and detection asymmetries with a control channel such as $B^{\pm} \to J/\psi K^{\pm}$, $J/\psi \to \mu^+\mu^-$; replacing the fixed-cut local Dalitz asymmetries (both binned, Sec. VIII, and K-means, Appendix I.8) with a proper sWeight-based background subtraction; and an explicit model of the partially reconstructed and other non-combinatorial B backgrounds, which would also make an ML-selected sample usable for a fit.

Appendix K: Code and Data Availability

In short: Python code with the package versions listed here; frequentist, Bayesian and machine-learning code and notebooks are in the accompanying repository. All code is written in Python $\geq$3.12, with `scikit-learn` [31] for the classical machine-learning models and `TensorFlow/Keras` [32] for the neural networks. Key package versions: `uproot 5.7`, `iminuit 2.32.0`, `emcee 3.1.6`, `corner 2.3.0`, `scikit-learn 1.9`, `tensorflow 2.21.0`, `keras 3.15.1`, `scipy 1.18`, `numpy 2.4`, `pandas 3.0`.

The dataset is the CERN Open Data record 4900 ("Matter Antimatter Differences – B meson decays to three hadrons"). The analysis code is available in the accompanying repository, in `Code_general/Frequentist/` (frequentist fits, the common systematic study, and `freq_validation.py` with the toy-MC experiments, profile likelihoods, confidence belt and bootstrap studies of Sec. VII and Appendix F), `Code_general/Bayesian/` (notebooks `01_bayes_mass_fit_and_asymmetry`: nominal fit, Bayes factors, systematic study, toy-MC closure test, magnet polarities and Dalitz plane; `02_bayes_model_checks_and_priors`: posterior predictive checks, convergence diagnostics, sensitivity to the priors and comparison with the profile likelihood; `03_bayes_vs_frequentist_low_statistics`: the two approaches at low statistics; the mass model, the sampler and the toy generator are in `bayes_model.py`), and `Code_general/Machine_Learning/` (notebooks

`04_ml_signal_background_classifiers`: weak labels, classifiers, calibration, uncertainty and hyperparameters; `05_ml_selection_and_local_asymmetry`: effect of the selection on $A_{\rm raw}$, K-means and null tests on the Dalitz plane, comparison with the published analysis; helper functions in `ml_tools.py`). Each of these notebooks opens with a table of the sections and figures of this paper that it produces and generates those figures from its saved results, so that they can be rebuilt without repeating the long computations; the common style is in `Code_general/plot_style.py`. The figures of the frequentist analysis are produced from its saved results by `regen_figures_validation.py`, `regen_figures_systematics.py` and `regen_figures_extras.py` in the figures directory of the report. All random numbers are generated with fixed seeds, so the toy-MC experiments can be reproduced.

Appendix L: Notation

Symbol	Meaning
$A_{\rm raw}$	raw charge asymmetry $(N^- - N^+)/(N^- + N^+)$ of the signal, parameter A of the fit
A_{CP}, A_P, A_D	CP asymmetry; production asymmetry of $B^\pm$; detection asymmetry
$A_{\rm count}$	counting asymmetry of all candidates of a sample
A_b	asymmetry of the background yields
N_s, N_b	signal and background yields (both charges)
μ, σ	mean and width of the Gaussian signal
$\lambda^\pm_{1,2}$, $f_\pm$	slopes and fraction of the double-exponential background, per charge
θ	vector of the twelve parameters
$\mathcal{L}$, π	likelihood and prior
$q(A) = 2\Delta\text{NLL}$	profile likelihood ratio of A; $t(A_{\rm true})$ at the injected value
Z, p	significance and p-value of the test of $A = 0$
BF_{01}, BF_{10}	Bayes factor of $A = 0$ against A free, and its inverse
p_B	posterior predictive p-value
σ_B	expected spread of the shift between nested samples, $\sqrt{\lvert\sigma^2_{\rm var} - \sigma^2_{\rm nom}\rvert}$
N/τ	chain length in integrated autocorrelation times
S_i, S_k	local significance of a Dalitz bin or cluster with respect to the global asymmetry

TABLE XIX. Main symbols used in the text.

[1] R. Aaij *et al.* (LHCb), Direct CP violation in charmless three-body decays of $B^\pm$ mesons, Phys. Rev. D **108**, 012008 (2023), arXiv:2206.07622 [hep-ex].

[2] L. Falcão, CP violation measurements in three-body charmless B decays, on behalf of the LHCb Collaboration, 12th Workshop on the CKM Unitarity Triangle (2024), arXiv:2403.19921 [hep-ex].

[3] LHCb Collaboration, Matter Antimatter Differences – B meson decays to three hadrons (2017), record 4900.

[4] LHCb Collaboration, Matter Antimatter Differences – Project Notebook (2017), record 4902.

[5] LHCb Collaboration, R. Aaij, *et al.*, Measurement of CP violation in the phase space of $B^\pm \to K^\pm\pi^+\pi^-$ and $B^\pm \to K^\pm K^+ K^-$ decays, Physical Review Letters **111**, 101801 (2013), arXiv:1306.1246.

[6] H. Dembinski, P. Ongmongkolkul, *et al.*, scikit-hep/iminuit, Zenodo (2020).

[7] F. James and M. Roos, Minuit – a system for function minimization and analysis of the parameter errors and correlations, Computer Physics Communications **10**, 343 (1975).

[8] S. S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses, Annals of Mathematical Statistics **9**, 60 (1938).

[9] J. Goodman and J. Weare, Ensemble samplers with affine invariance, Commun. Appl. Math. Comput. Sci. **5**, 65 (2010).

[10] D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman, emcee: The MCMC Hammer, Publ. Astron. Soc. Pac. **125**, 306 (2013), arXiv:1202.3665.

[11] J. M. Dickey, The weighted likelihood ratio, linear hypotheses on normal location parameters, Ann. Math. Statist. **42**, 204 (1971).

[12] H. Jeffreys, *Theory of Probability*, 3rd ed. (Oxford University Press, Oxford, 1961).

[13] R. Barlow, Systematic errors: facts and fictions, in *Conference on Advanced Statistical Techniques in Particle Physics* (Durham, 2002) arXiv:hep-ex/0207026.

[14] G. Cowan, *Statistical Data Analysis* (Oxford University Press, 1998).

[15] J. Neyman and E. S. Pearson, On the problem of the most efficient tests of statistical hypotheses, Philosophical Transactions of the Royal Society of London A **231**, 289 (1933).

[16] R. E. Kass and A. E. Raftery, Bayes Factors, J. Am. Stat. Assoc. **90**, 773 (1995).

[17] A. Gelman, X.-L. Meng, and H. Stern, Posterior predictive assessment of model fitness via realized discrepancies, Statistica Sinica **6**, 733 (1996).

[18] T. Skwarnicki, *A study of the radiative cascade transitions between the* Υ' *and* Υ *resonances*, Ph.D. thesis, Institute of Nuclear Physics, Krakow (1986), dESY-F31-86-02.

[19] G. W. Brier, Verification of forecasts expressed in terms of probability, Mon. Weather Rev. **78**, 1 (1950).

[20] R. Tibshirani, Regression shrinkage and selection via the lasso, Journal of the Royal Statistical Society B **58**, 267 (1996).

[21] B. Zadrozny and C. Elkan, Transforming classifier scores into accurate multiclass probability estimates, in *Proceedings of the 8th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (2002) pp. 694–699.

[22] E. M. Metodiev, B. Nachman, and J. Thaler, Classification without labels: Learning from mixed samples in high energy physics, JHEP **10**, 174, arXiv:1708.02949.

[23] S. P. Lloyd, Least squares quantization in PCM, IEEE Transactions on Information Theory **28**, 129 (1982).

[24] L. Hubert and P. Arabie, Comparing partitions, Journal of Classification **2**, 193 (1985).

[25] B. Lakshminarayanan, A. Pritzel, and C. Blundell, Simple and scalable predictive uncertainty estimation using deep ensembles, in *Advances in Neural Information Processing Systems 30* (2017) arXiv:1612.01474.

[26] Y. Gal and Z. Ghahramani, Dropout as a Bayesian Approximation: Representing Model Uncertainty in Deep Learning, in *Proceedings of the 33rd International Conference on Machine Learning (ICML)* (2016) arXiv:1506.02142.

[27] L. Breiman, Random forests, Machine Learning **45**, 5 (2001).

[28] J. H. Friedman, Greedy function approximation: a gradient boosting machine, Annals of Statistics **29**, 1189 (2001).

[29] D. P. Kingma and J. Ba, Adam: A method for stochastic optimization, in *3rd International Conference on Learning Representations (ICLR)* (2015) arXiv:1412.6980.

[30] P. J. Rousseeuw, Silhouettes: A graphical aid to the interpretation and validation of cluster analysis, J. Comput. Appl. Math. **20**, 53 (1987).

[31] F. Pedregosa *et al.*, Scikit-learn: Machine learning in Python, Journal of Machine Learning Research **12**, 2825 (2011).

[32] M. Abadi *et al.*, TensorFlow: Large-scale machine learning on heterogeneous distributed systems (2016), arXiv:1603.04467.