

Vectors practice

Perform the calculations below.

EXAMPLE:

Question: Vector \vec{a} has magnitude (length) of 5 and direction of 45° , vector \vec{b} has magnitude of 10 and direction of 30° . Find components of vectors \vec{a} and \vec{b} , calculate $\vec{c} = \vec{a} + \vec{b}$, $\vec{a} \cdot \vec{b}$, the length $c = |\vec{c}|$, and the direction ϕ_c of vector \vec{c} . (Use calculator.)

Answer:

1. Components of vector \vec{a} are $\vec{a} = (a \cos 45^\circ, a \sin 45^\circ) = (5 \cos 45^\circ, 5 \sin 45^\circ) = (3.5, 3.5)$.
2. $\vec{b} = (b \cos 30^\circ, b \sin 30^\circ) = (10 \cos 30^\circ, 10 \sin 30^\circ) = (8.6, 5)$.
3. $\vec{c} = (3.5, 3.5) + (8.6, 5) = (13.1, 8.5)$.
4. $\vec{a} \cdot \vec{b} = (3.5, 3.5) \cdot (8.6, 5) = 8.6 * 3.5 + 5 * 3.5 = 46.6$
5. $c = |\vec{c}| = \sqrt{13.1^2 + 8.5^2} = 15.62$
6. $\tan \phi_c = c_y/c_x = 8.5/13.1 = 0.65$ (finding $\tan \phi_c$ is enough!).

ASSIGNMENT:

A. Find components of vectors $\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}$.

1. Vector \vec{a} has magnitude of 5 and direction of 60° .
2. Vector \vec{b} has magnitude of 2.5 and direction of 135° .
3. Vector \vec{c} has magnitude of 15 and direction of -45° .
4. Vector \vec{d} has magnitude of 1 and direction of 120° .
5. Vector \vec{e} has magnitude of 10 and direction of -30° .

B. Using above $\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}$, find

1. $\vec{a} + \vec{b}, |\vec{a} + \vec{b}|$ (length of $\vec{a} + \vec{b}$), and direction of $\vec{a} + \vec{b}$, ϕ_{a+b} .
2. $\vec{a} + 2\vec{b}, |\vec{a} + 2\vec{b}|$, and direction.
3. $\vec{b} + \vec{c}, |\vec{b} + \vec{c}|$, and direction.
4. $3\vec{a} - 4\vec{e}, |3\vec{a} - 4\vec{e}|$, and direction.
5. $\vec{a} + 2\vec{c} - \vec{e}, |\vec{a} + 2\vec{c} - \vec{e}|$, and direction.
6. $1.5\vec{a} + \vec{c} - \vec{d} - \vec{e}, |1.5\vec{a} + \vec{c} - \vec{d} - \vec{e}|$, and direction.
7. $\vec{c} + \vec{d} + 0.5\vec{a} + 1.5\vec{b}, |\vec{c} + \vec{d} + 0.5\vec{a} + 1.5\vec{b}|$, and direction.
8. $\vec{c} + \vec{d} - (\vec{a} + \vec{b}) + \vec{e}, |\vec{c} + \vec{d} - (\vec{a} + \vec{b}) + \vec{e}|$, and direction.
9. $\vec{c} + 2(\vec{d} - \vec{b}) + (\vec{a} + \vec{e}), |\vec{c} + 2(\vec{d} - \vec{b}) + (\vec{a} + \vec{e})|$, and direction.

C. Using above $\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}$ find $\vec{a} \cdot \vec{b}, \vec{a} \cdot \vec{c}, \vec{b} \cdot \vec{d}, \vec{a} \cdot \vec{e}, \vec{a} \cdot (\vec{b} + \vec{e}), \vec{a} \cdot (2\vec{c} + 3\vec{e}), \vec{a} \cdot (2\vec{e}), (2\vec{a} + \vec{d}) \cdot \vec{b}$.