## MATH 308 D200, Fall 2019

# 21. Examples of matrix games (based on notes from Dr. J. Hales, Dr. L. Stacho, and Dr. L. Godyyn)

asca on notes from Dr. 5. Hales, Dr. E. Stacho, and Dr. E. Godyyn

Dr. Masood Masjoody

SFU Burnaby

Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1,2,3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

### Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1, 2, 3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

#### Six step solution:

### Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1,2,3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

#### Six step solution:

1. Set up the payoff matrix in terms of Ron's winnings.

			oleen even)		
		1	2	3	
	1	2	-3	2	
Ron (odd)	2	-3	4	-3	
	3	2	-3	6	

### Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1,2,3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

#### Six step solution:

- 1. Set up the payoff matrix in terms of Ron's winnings.
- 2. Domination reduce the payoff matrix (optional).

		_	oleen even)		
		1	2	3	
	1	2	-3	2	
Ron (odd)	2	-3	4	-3	
	3	2	-3	6	

### Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1,2,3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

#### Six step solution:

- 1. Set up the payoff matrix in terms of Ron's winnings.
- 2. Domination reduce the payoff matrix (optional).

			oleen even)					Cole (eve	
		1	2	3				1	2
	1	2	-3	2	$\mapsto$		1	2	-3
Ron (odd)	2	-3	4	-3		Ron (odd)	2	-3	4
, ,	3	2	-3	6		, ,	3	2	-3

2/15

### Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1,2,3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

#### Six step solution:

- 1. Set up the payoff matrix in terms of Ron's winnings.
- 2. Domination reduce the payoff matrix (optional).

			oleen even)					Cole (eve	
		1	2	3				1	2
	1	2	-3	2	$\mapsto$		1	2	-3
Ron (odd)	2	-3	4	-3		Ron (odd)	2	-3	4
, ,	3	2	-3	6		, ,	3	2	-3

2/15

### Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1,2,3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins \$x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

#### Six step solution:

- 1. Set up the payoff matrix in terms of Ron's winnings.
- 2. Domination reduce the payoff matrix (optional).

	Coleen (even)				Coleen (even)						Coleen (even)				
		1	2	3				1	2				(606	211)	
	1	2	-3	2	$\mapsto$		1	2	-3	$\mapsto$			1		
Ron	_					Ron	_					2	-3	-4	l
(odd)	2	-3	4	-3		(odd)	2	-3	4		Ron	3	2	-3	]
( /	3	2	-3	6		,	3	2	-3		(odd)	Ū	_		J

SFU department of mathematics

2/15

### Game 1: Modified Rock, Paper, Scissors (Texbook Example 1, p. 117)

#### Find the optimal mixed strategies for the following matrix game

Two players, say Ron ("odd" player) and Coleen ("even" player), each secretly choose an integer from  $\{1,2,3\}$ . Both players reveal their numbers simultaneously.

- o If the sum of the numbers is odd, Coleen wins \$3 from Ron.
- o If the sum of the numbers is even, Ron wins x from Coleen where:
  - x equals to the difference of the numbers, provided they are different numbers.
  - x equals the sum of the numbers, provided they are the same number

#### Six step solution:

- 1. Set up the payoff matrix in terms of Ron's winnings.
- 2. Domination reduce the payoff matrix (optional).

Coleen (even)					Coleen (even)						Coleen (even)				
		1	2	3				1	2				(eve	:11)	
	1	2	-3	2	$\mapsto$		1	2	-3	$\mapsto$			1		7
Ron	_	_		_		Ron	^	_	_		_	2	-3	-4	
Ron (odd)	2	-3	4	-3		(odd)	2	-3	4		Ron	3	2	-3	
` /	3	2	-3	6		,	3	2	-3	1	(odd)	-		<u> </u>	J

3. Set up the game tableau:

SFU department of mathematics

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1	-1	-1	= -0
<b>p</b> 2	-1	-3	4	0	$=-t_{2}$
<b>p</b> 3	-1	2	-3	0	$=-t_{3}$
-1	-1	0	0	0	= f
	= 0	$= s_2$	$= s_3$	= g	,

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1	-1	-1	= -0
<b>p</b> 2	-1	-3	4	0	$=-t_{2}$
<b>p</b> 3	-1	2°	-3	0	$=-t_{3}$
-1	-1	0	0	0	= f
	= 0	$= s_2$	= <b>s</b> <sub>3</sub>	= g	,

The 2 and 4 are largest in their column.

The  $2^{\circ}$  is the smaller of these.

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1*	-1	-1	= -0
<b>p</b> 2	-1	-3	4	0	$=-t_{2}$
<b>p</b> <sub>3</sub>	-1**	2°	-3	0	$=-t_{3}$
-1	-1	0	0	0	= f
	= 0	$= s_2$	= <b>s</b> <sub>3</sub>	= g	•

The 2 and 4 are largest in their column.

The  $2^{\circ}$  is the smaller of these.

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1*	-1	-1	= -0
<b>p</b> 2	-1	-3	4	0	$=-t_{2}$
<b>p</b> 3	-1**	2°	-3	0	$=-t_{3}$
-1	-1	0	0	0	= f
	= 0	$= s_2$	= <b>s</b> <sub>3</sub>	= g	•

	V	0	<b>q</b> 2	-1	
<b>s</b> 2	0	-1	1	1	$=-q_1$
<b>p</b> 2	-1	-3	7	3	$=-t_2$
<b>p</b> 3	-1**	2°	-5	-2	$=-t_3$
-1	-1	0	0	0	= f
	= 0	<u>u</u>	$= s_3$	=g	,

The 2 and 4 are largest in their column.

The  $2^{\circ}$  is the smaller of these.

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1*	-1	-1	=-0
<b>p</b> <sub>2</sub>	-1	-3	4	0	=-t
<b>p</b> 3	-1**	2°	-3	0	=-t
-1	-1	0	0	0	= f
	= 0	$= s_2$	= <b>s</b> <sub>3</sub>	= g	,

	V	0	<b>q</b> 2	-1	
<b>s</b> 2	0	-1	1	1	$=-q_1$
<b>p</b> <sub>2</sub>	-1	-3	7	3	$=-t_2$
<i>p</i> <sub>3</sub>	-1**	2°	-5	-2	$=-t_3$
-1	-1	0	0	0	= f
	= 0	(u)	$= s_3$	= g	•

The 2 and 4 are largest in their column.

The  $2^{\circ}$  is the smaller of these.

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1*	-1	-1	= -0
<b>p</b> <sub>2</sub>	-1	-3	4	0	$=-t_{2}$
<b>p</b> <sub>3</sub>	-1**	2°	-3	0	$=-t_{3}$
-1	-1	0	0	0	= f
	= 0	$= s_2$	= <b>s</b> <sub>3</sub>	= g	-

The 2 and 4 are largest in their column.

	V	0	<b>q</b> 2	-1	
<b>s</b> 2	0	-1	1	1	$=-q_1$
<b>p</b> <sub>2</sub>	-1	-3	7	3	$=-t_2$
<b>p</b> 3	-1**	2°	-5	-2	$=-t_3$
-1	-1	0	0	0	= f
	= 0	<u>u</u>	$= s_3$	= g	,

		V	<b>q</b> 2	-1	
	<b>s</b> <sub>2</sub>	0	1	1	$=-q_1$
	<b>p</b> <sub>2</sub>	-1	7	3	$=-t_2$
•	<b>p</b> 3	-1**	-5	-2	$=-t_3$
	-1	-1	0	0	= f
		= 0	$= s_3$	=g	•

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1*	-1	-1	= -0
<b>p</b> 2	-1	-3	4	0	$=-t_{2}$
<b>p</b> 3	-1**	2°	-3	0	$=-t_{3}$
-1	-1	0	0	0	= f
	= 0	$= s_2$	$= s_3$	= g	,

The 2 and 4 are largest in their column.

	V	0	<b>q</b> 2	-1				V	<b>q</b> 2	-1	
<b>s</b> 2	0	-1	1	1	$= -q_1$		<b>s</b> 2	0	1	1	$=-q_1$
<b>p</b> <sub>2</sub>	-1	-3	7	3	$=-t_2$		<b>p</b> <sub>2</sub>	-1	7	3	$=-t_2$
<b>p</b> <sub>3</sub>	-1**	2°	-5	-2	$=-t_3$	$\rightarrow$	<b>p</b> <sub>3</sub>	-1**	-5	-2	$=-t_3$
-1	-1	0	0	0	= f		-1	-1	0	0	= f
·	= 0	<u>u</u>	$= s_3$	= g	•			= 0	$= s_3$	= g	

	V	$q_1$	<b>q</b> 2	-1	
<u>u</u>	0	-1*	-1	-1	= -0
<b>p</b> <sub>2</sub>	-1	-3	4	0	$=-t_2$
<b>p</b> <sub>3</sub>	-1**	2°	-3	0	$=-t_3$
-1	-1	0	0	0	= f
	= 0	$= s_2$	$= s_3$	= g	,

The 2 and 4 are largest in their column.

	V	0	<b>q</b> 2	-1				V	<b>q</b> 2	-1	
					$=-q_1$						$=-q_1$
<b>p</b> 2	-1	-3	7	3	$=-t_{2}$		<b>p</b> 2	-1	7	3	$\begin{vmatrix} =-t_2\\ =-t_3 \end{vmatrix}$
<b>p</b> 3	-1**	2°	-5	-2	$=-t_3$	$\rightarrow$	<b>p</b> 3	-1**	-5	-2	$=-t_3$
-1	-1	0	0	0	= f		-1	-1	0	0	= f
	= 0	<u>u</u>	$= s_3$	= g	•			= 0	$= s_3$	= g	•

	<i>t</i> <sub>3</sub>	$q_2$	-1	
<b>s</b> <sub>2</sub>	0	1	1	$=-q_1$
<b>p</b> <sub>2</sub>	-1	12	5	$=-t_{2}$
0	-1	5	2	= -0
-1	-1	5	2	= f
	$= p_3$	$= s_3$	= g	•

The 2 and 4 are largest in their column.

	<b>v</b>	0	<b>q</b> 2	-1			V	<b>q</b> 2	-1		
<b>s</b> <sub>2</sub>	0	-1	1	1	$=-q_1$	<b>s</b> <sub>2</sub>	0	1	1	$=-q_1$	
$p_2$	-1	-3	7	3	$=-t_2$	<b>p</b> <sub>2</sub>	-1	7	3	$=-t_{2}$	
<b>p</b> <sub>3</sub>	-1**	2°	-5	-2	$=-t_3$	<b>p</b> <sub>3</sub>	-1**	-5	-2	$=-t_{3}$	
-1	-1	0	0	0	= f	-1	-1	0	0	= f	
	= 0	<u>u</u>	= <b>s</b> <sub>3</sub>	=g	-		= 0	$= s_3$	=g	-	

-1(V)  $q_2$  $q_1$ 0 -1\* -1 <u>u</u> -1 = -0-1 -3 4 0  $p_2$ -1\*\* 2° -3 0 **p**<sub>3</sub>  $= -t_3$ -1-1 0 0 0 = f= 0 $= s_2$  $= s_3$ = g

4. Find first two pivots:

The 2 and 4 are largest in their column. The  $2^{\circ}$  is the smaller of these.

Pivot and delete twice.

5. Apply SA Phase 2 to the feasible tableau.

-1(V)  $q_2$  $q_1$ 0 -1\* -1 <u>u</u> -1 = -0-1 -3 4 0  $p_2$ -1\*\* 2° -3 0 **p**<sub>3</sub>  $= -t_3$ -1-1 0 0 0 = f= 0 $= s_2$  $= s_3$ = g

4. Find first two pivots:

The 2 and 4 are largest in their column. The  $2^{\circ}$  is the smaller of these.

Pivot and delete twice.

**5**. Apply SA Phase 2 to the feasible tableau.

6. Summarize the optimal strategies and the Von Neumann value:

Ron chooses 2 with probability 5/12, chooses 3 with probability 7/12, and never chooses 3. Coleen chooses 1 with probability 7/12, chooses 2 with probability 5/12, and never chooses 3. Coleen expects an average payoff of 1/12 per game.

### Game 2: Guess the Number:

Find the optimal mixed strategies for the following matrix game

Ron and Coleen simultaneously writes either the number 2 or 3. Each player also writes a number which they believe the opponents has written down.

- If neither player guesses right, Ron wins \$3 from Coleen.
- If both players guess right, Coleen wins \$3 from Ron.
- Otherwise, the player who guesses correctly wins (from the other player) the sum (in \$) of the (first) numbers written by two players.

SFU department of mathematics

#### Game 2: Guess the Number:

Find the optimal mixed strategies for the following matrix game

Ron and Coleen simultaneously writes either the number 2 or 3. Each player also writes a number which they believe the opponents has written down.

- If neither player guesses right, Ron wins \$3 from Coleen.
- If both players guess right, Coleen wins \$3 from Ron.
- Otherwise, the player who guesses correctly wins (from the other player) the sum (in \$) of the (first) numbers written by two players.

Notation: The pair (a, b) denotes a player's choice of writing a and guessing b, for  $a, b \in \{2, 3\}$ .

SFU department of mathematics

#### Coleen

	(2, 2)	(2,3)	(3, 2)	(3, 3)
(2,2)	-3	4	-5	3
(2,3)	-4	3	-3	5
(3, 2)	5	-3	3	-6
(3,3)	3	-5	6	-3

(Already domination reduced)

Ron

#### Coleen

	(2,2)	(2,3)	(3, 2)	(3,3)
(2,2)	-3	4	-5	3
(2,3)	-4	3	-3	5
(3, 2)	5	-3	3	-6
(3,3)	3	-5	6	-3

(Already domination reduced)

	(3, 2)	5	-3	3	-6			
	(3,3)	3	-5	6	-3			
	V	<b>q</b> 22	<b>q</b> 23	<b>q</b> 32	<b>q</b> 33	-1		
<u>u</u>	0	-1	-1	-1	-1	-1	= -0	
<b>p</b> 22	-1	-3	4°	-5	3	0	$=-t_{22}$	
<b>p</b> 23	-1	-4	3	-3	5	0	$=-t_{23}$	(Using natural variable names.)
<b>p</b> <sub>32</sub>	-1	5	-3	3	-6	0	$=-t_{32}$	(Osing natural variable names.)
<b>p</b> 33	-1	3	-5	6	-3	0	$=-t_{33}$	
-1	-1	0	0	0	0	0	= f	
	=0	$= s_{22}$	$= s_{23}$	$=s_{32}$	$= s_{33}$	=g	-	

Ron

```
Coleen
                         (2,3)
               (2,2)
                                   (3, 2)
                                            (3,3)
     (2, 2)
                 -3
                           4
                                    -5
                                               3
                                                      (Already domination reduced)
                                               5
Ron (2, 3)
                 -4
                           3
                                    -3
     (3, 2)
                 5
                           -3
                                     3
                                              -6
     (3, 3)
                 3
                           -5
                                     6
                                              -3
        (V)
                                                     -1
                q22
                         q23
                                   q32
                                            q33
        0
                -1
                          -1
                                   -1
                                             -1
                                                      -1
                                                             = -0
        -1
                -3
                          4°
                                   -5
                                             3
                                                      0
                                                             =-t_{22}
        -1
                          3
                                   -3
                                             5
                -4
                                                      0
                                                             =-t_{23}
                                                                        (Using natural variable names.)
        -1
                 5
                          -3
                                    3
                                             -6
                                                      0
                                                             =-t_{32}
        -1
                 3
                          -5
                                    6
                                             -3
                                                      0
                                                             =-t_{33}
       -1
                 0
                          0
                                    0
                                             0
                                                      0
                                                             = f
       =0
               = s_{22}
                        = s_{23}
                                  = s_{32}
                                           = 533
                                                     =g
                   t<sub>32</sub>
                              t<sub>23</sub>
                                                    -1
         t33
                                         q33
       -1/25
                 -2/75
                             1/15
                                       17/15
                                                  13/25
                                                             = -q_{23}
        1/5
                               0
                                                   2/5
                  -1/5
                                          1
                                                             = -q_{32}
       -4/25
                 17/75
                            -1/15
                                      -17/15
                                                   2/25
                                                                        (I used computer to solve this)
        3/5
                  -3/5
                              -1
                                          0
                                                   1/5
                                                             =-t_{22}
       -2/25
                 -29/75
                            -8/15
                                       -1/15
                                                   1/25
                                                             = f
       = p_{33}
                  = p_{32}
                             = p_{23}
                                        = s_{33}
                                                   =g
```

**(u)** 

**p**22

**p**23

p<sub>32</sub>

**p**33 -1

**5**23

**5**32

**s**22

**p**22

-1

		(2,2)	(2,3)	(3, 2)	(3,	3)						
(2,2)		-3	4	-5	3		( )	adu damir	amination raduced)			
Ron (2, 3)		-4	3	-3	5		(Alle	(Already domination reduced)				
	(3, 2)	5	-3	3	-6	5						
	(3, 3)	3	-5	6	-3	3						
	V	<b>q</b> 22	<b>q</b> 23	<b>q</b> 32	<b>q</b> 33		-1					
<u>u</u>	0	-1	-1	-1	-1		-1	= -0				
<b>p</b> 22	-1	-3	4°	-5	3		0	$=-t_{22}$				
<b>p</b> 23	-1	-4	3	-3	5		0	$=-t_{23}$	(Haine natural variable names)			
<b>p</b> <sub>32</sub>	-1	5	-3	3	-6		0	$=-t_{32}$	(Using natural variable names.)			
<b>p</b> 33	-1	3	-5	6	-3		0	$=-t_{33}$				
-1	-1	0	0	0	0		0	= f				
$=0$ $=s_{22}$ $=s_{23}$ $=s_{3}$				$= s_{32}$	$s_{32} = s_{33} = g$			,				
	t <sub>33</sub>	t <sub>32</sub>	t <sub>23</sub>	3 9	33		-1					
<i>s</i> <sub>23</sub>	-1/25	-2/7	5   1/1	15 17	/15		3/25	$=-q_{23}$				
<b>s</b> 32	1/5	-1/5	5 0		1		2/5	$=-q_{32}$				
<b>s</b> <sub>22</sub>	-4/25	17/7	5 -1/2	15 -17	-17/15		2/25	$=-q_{22}$	(I used computer to solve this			
<b>p</b> 22	3/5	-3/5	5 -1		0	:	1/5	$=-t_{22}$				
-1	-2/25	-29/7	'5 -8/	15 -1	/15	1	./25	= f				
	$= p_{33}$	$= p_3$	$_2 = p$	23 =	<b>s</b> 33		=g					
Ron wins 4 cents per game on average. Optimal strategies:												
For Ron $(p_{22}, p_{23}, p_{32}, p_{33}) = (0, \frac{8}{15}, \frac{29}{75}, \frac{2}{25})$ . For Coleen $(q_{22}, q_{23}, q_{32}, q_{33}) = (\frac{2}{25}, \frac{13}{25}, \frac{2}{5}, 0)$ .												

Coleen

SFU department of mathematics

Game 3. Guess an Adjacent Number:

1 A 2 B 3 C 4 D 5

Player  $P_1$  secretly chooses one of four letters A–D, Player  $P_2$  secretly chooses one of the five boxes 1–5. They reveal their choices

- If  $P_1$  chooses a letter that is adjacent to the  $P_2$ 's box,  $P_1$  wins \$1 from  $P_2$ .
- Otherwise,  $P_2$  wins \$1 from  $P_1$ .

Find

- a) Payoff matrix for  $P_1$ .
- b) Simplify it by dominance.
- c) Suppose  $P_2$  plays mixed strategy (1/2,0,1/4,0,1/4). With respect to this strategy find a mixed strategy for  $P_1$  that will guarantee a fair game.

a) b)

c)

SFU faculty of science department of mathematics

### Game 4: Little Casino

Ron antes \$5 and Coleen antes \$3 to the pot. Ron then secretly tosses a coin and checks the result. Ron now has two options:

FOLD—Coleen wins the pot

BET—Ron adds \$4 to the pot

If Ron bets, Coleen has two options

FOLD—Ron wins the pot



SEE—Coleen adds \$8 to the pot If Coleen sees, the coin is revealed. If the coin is Heads, Ron wins the pot. Otherwise, Coleen wins the pot. Find the von Neumann value and the optimal strategies for this game.

- Compute payoff matrix on cell at a time. For example, suppose Ron selects BF and Coleen selects S. If the coin is Heads, then Ron will Bet and add \$4 to the pot, Coleen will See and add \$8. When Ron reveals that the coin was Heads, he collects the 3+8 dollars that Coleen had put in the pot. If the coin is Tails, then Ron will Fold and lose \$5.

- Compute payoff matrix on cell at a time. For example, suppose Ron selects BF and Coleen selects S. If the coin is Heads, then Ron will Bet and add \$4 to the pot, Coleen will See and add \$8. When Ron reveals that the coin was Heads, he collects the 3+8 dollars that Coleen had put in the pot. If the coin is Tails, then Ron will Fold and lose \$5. Each of these events has probability  $\frac{1}{2}$  so we record the game matrix entry

$$a_{BF,S} = \frac{1}{2} \cdot (3+8) + \frac{1}{2} \cdot (-5) = 3.$$

- Compute payoff matrix on cell at a time. For example, suppose Ron selects BF and Coleen selects S. If the coin is Heads, then Ron will Bet and add \$4 to the pot, Coleen will See and add \$8. When Ron reveals that the coin was Heads, he collects the 3+8 dollars that Coleen had put in the pot. If the coin is Tails, then Ron will Fold and lose \$5. Each of these events has probability  $\frac{1}{2}$  so we record the game matrix entry

$$a_{BF,S} = \frac{1}{2} \cdot (3+8) + \frac{1}{2} \cdot (-5) = 3.$$

If Ron selects BB and Coleen selects S, then Ron wins 3+8 dollars with Heads, but loses 5+4 dollars with Tails, so  $a_{BB,S}=\frac{(3+8)-(5+4)}{2}=1$ . We obtain the following game matrix.

- Compute payoff matrix on cell at a time. For example, suppose Ron selects BF and Coleen selects S. If the coin is Heads, then Ron will Bet and add \$4 to the pot, Coleen will See and add \$8. When Ron reveals that the coin was Heads, he collects the 3+8 dollars that Coleen had put in the pot. If the coin is Tails, then Ron will Fold and lose \$5. Each of these events has probability  $\frac{1}{2}$  so we record the game matrix entry

$$a_{BF,S} = \frac{1}{2} \cdot (3+8) + \frac{1}{2} \cdot (-5) = 3.$$

If Ron selects BB and Coleen selects S, then Ron wins 3+8 dollars with Heads, but loses 5+4 dollars with Tails, so  $a_{BB,S}=\frac{(3+8)-(5+4)}{2}=1$ . We obtain the following game matrix.

# Game 5: Little Poker (Textbook Example 5, p. 127)

Ron and Coleen play with three cards, J, Q, and K which are face-down. They each ante 25¢ each and select a card. Ron looks at his card, then chooses to either:

Fold (F) — here Coleen wins the pot, or

Bet (B) — here Ron adds 10¢ to the pot

If Ron bets, then Coleen looks at her card, a chooses to either:

Fold (F) — here Ron wins the pot, or

See — here Coleen also adds 10¢ to the pot

If Coleen sees, then both cards are revealed and the player with the higher card (where J < Q < K) wins the pot.

Find the von Neumann value and the optimal strategies for this game.



	A.A.										
	FFF	FFS	FSF	SFF	FSS	SFS	SSF	SSS			
FFF	-25	-25	-25	-25	-25	-25	-25	-25	7		
FFB	-25/3	-25/3	-20/3	-20/3	-20/3	-20/3	-5	-5			
<b>FBF</b>	-25/3	-55/3	-25/3	-20/3	-55/3	-50/3	-20/3	-50/3			
BFF	-25/3	-55/3	-55/3	-25/3	-85/3	-55/3	-55/3	-85/3			
<b>FBB</b>	25/3	-5/3	10	35/3	0	5/3	40/3	10/3			
BFB	25/3	-5/3	0	10	-10	0	5/3	-25/3			
BBF	25/3	-35/3	-5/3	10	-65/3	-10	0	-20			
BBB	_ 25	5	50/3	85/3	-10/3	25/3	20	0			
					11	7100E00E0000					