

MATH 308 D200, Fall 2019

20. Mixed strategies

(based on notes from Dr. J. Hales, Dr. L. Stacho, and Dr. L. Goddyn)

Dr. Masood Masjoody

SFU Burnaby

Mixed (Probabilistic) Strategy

Many games are not strictly determined and hence lack a best pure strategy according to the Fundamental Principle of game theory. It is dangerous for either player to stick to a pure strategy. So we let players to choose their moves at random in each round of the game.

Definition (Mixed (Probabilistic) Strategy)

Let $A = [a_{ij}]$ be an $m \times n$ matrix game. A *mixed (probabilistic) strategy* for the row player is a vector $\mathbf{p} = (p_1, p_2, \dots, p_m)$ such that

$$\sum_{i=1}^m p_i = 1, \quad p_i \geq 0 \text{ for all } i.$$

A *mixed (probabilistic) strategy* for the column player is a vector $\mathbf{q} = (q_1, q_2, \dots, q_n)$ such that

$$\sum_{j=1}^n q_j = 1, \quad q_j \geq 0 \text{ for all } j.$$

Any mixed strategy containing an entry of 1 is a pure strategy. e.g. $\mathbf{p} = \mathbf{e}_3 = (0, 0, 1, 0, 0, 0)$.

So in our earlier modified Rock-Paper-Scissors game, Ron can choose mixed strategy $\mathbf{p} = (1/6, 1/2, 1/3)$ and Coleen can choose $\mathbf{q} = (1/3, 1/2, 1/6)$.

Question: If Ron and Coleen use these mixed strategies, how much, on average, can Ron expect to win (or lose) on each round of the game?

Mixed (Probabilistic) Strategy

Example (Original Rock-Paper-Scissors)

Suppose that Ron plays *paper* half the time, and each of the other two moves a quarter of the time, and Coleen always plays *rock*. This gives $\mathbf{p} = (1/4, 1/2, 1/4)$ and $\mathbf{q} = (1, 0, 0)$. Then Ron's expected payoff (per game) is

Mixed (Probabilistic) Strategy

Example (Original Rock-Paper-Scissors)

Suppose that Ron plays *paper* half the time, and each of the other two moves a quarter of the time, and Coleen always plays *rock*. This gives $\mathbf{p} = (1/4, 1/2, 1/4)$ and $\mathbf{q} = (1, 0, 0)$. Then Ron's expected payoff (per game) is

$$1/4 \cdot 0 + 1/2 \cdot 1 + 1/4 \cdot (-1) = 1/4$$

Mixed (Probabilistic) Strategy

Example (Original Rock-Paper-Scissors)

Suppose that Ron plays *paper* half the time, and each of the other two moves a quarter of the time, and Coleen always plays *rock*. This gives $\mathbf{p} = (1/4, 1/2, 1/4)$ and $\mathbf{q} = (1, 0, 0)$. Then Ron's expected payoff (per game) is

$$\begin{aligned} & 1/4 \cdot 0 + 1/2 \cdot 1 + 1/4 \cdot (-1) = 1/4 \\ & = \begin{bmatrix} 1/4 & 1/2 & 1/4 \end{bmatrix} \begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \end{aligned}$$

Mixed (Probabilistic) Strategy

Example (Original Rock-Paper-Scissors)

Suppose that Ron plays *paper* half the time, and each of the other two moves a quarter of the time, and Coleen always plays *rock*. This gives $\mathbf{p} = (1/4, 1/2, 1/4)$ and $\mathbf{q} = (1, 0, 0)$. Then Ron's expected payoff (per game) is

$$1/4 \cdot 0 + 1/2 \cdot 1 + 1/4 \cdot (-1) = 1/4$$

$$= \begin{bmatrix} 1/4 & 1/2 & 1/4 \end{bmatrix} \begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$= \mathbf{p}^T A \mathbf{q}.$$

Mixed (Probabilistic) Strategy

Example (Original Rock-Paper-Scissors)

Suppose that Ron plays *paper* half the time, and each of the other two moves a quarter of the time, and Coleen always plays *rock*. This gives $\mathbf{p} = (1/4, 1/2, 1/4)$ and $\mathbf{q} = (1, 0, 0)$. Then Ron's expected payoff (per game) is

$$\begin{aligned} & 1/4 \cdot 0 + 1/2 \cdot 1 + 1/4 \cdot (-1) = 1/4 \\ &= \begin{bmatrix} 1/4 & 1/2 & 1/4 \end{bmatrix} \begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \\ &= \mathbf{p}^T \mathbf{A} \mathbf{q}. \end{aligned}$$

Theorem

For any payoff matrix \mathbf{A} if Ron adopts mixed strategy \mathbf{p} and Coleen adopts mixed strategy \mathbf{q} (\mathbf{p} and \mathbf{q} are column vectors), then Ron's expected payoff (per round) is $\mathbb{E}(\mathbf{p}, \mathbf{q}) = \mathbf{p}^T \mathbf{A} \mathbf{q}$.

Mixed (Probabilistic) Strategy

Example (Original Rock-Paper-Scissors)

Suppose that Ron plays *paper* half the time, and each of the other two moves a quarter of the time, and Coleen always plays *rock*. This gives $\mathbf{p} = (1/4, 1/2, 1/4)$ and $\mathbf{q} = (1, 0, 0)$. Then Ron's expected payoff (per game) is

$$\begin{aligned} & 1/4 \cdot 0 + 1/2 \cdot 1 + 1/4 \cdot (-1) = 1/4 \\ &= \begin{bmatrix} 1/4 & 1/2 & 1/4 \end{bmatrix} \begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \\ &= \mathbf{p}^T \mathbf{A} \mathbf{q}. \end{aligned}$$

Theorem

For any payoff matrix \mathbf{A} if Ron adopts mixed strategy \mathbf{p} and Coleen adopts mixed strategy \mathbf{q} (\mathbf{p} and \mathbf{q} are column vectors), then Ron's expected payoff (per round) is $\mathbb{E}(\mathbf{p}, \mathbf{q}) = \mathbf{p}^T \mathbf{A} \mathbf{q}$.

Definition (Optimal mixed strategy)

An **optimal mixed strategy** is a mixed strategy that minimizes the potential loss against the opponent's best counter-strategy. For example, strategy \mathbf{p}^* is optimal for Ron, if

$$\min_{\mathbf{q}} \mathbb{E}(\mathbf{p}^*, \mathbf{q}) \geq \min_{\mathbf{q}} \mathbb{E}(\mathbf{p}, \mathbf{q})$$

for every mixed strategy \mathbf{p} for Ron. Finding optimal mixed strategies for both players in a matrix game is called **solving the game**.

Optimal Mixed Strategy

Natural Protocol:

Ron can estimate \mathbf{q} by keeping statistics over many rounds of play. Assume that Ron knows \mathbf{q} .

Ron selects \mathbf{p} to maximize $\mathbb{E}(\mathbf{p}, \mathbf{q})$. Then Coleen examines \mathbf{p} and selects \mathbf{q} to minimize $\mathbb{E}(\mathbf{p}, \mathbf{q})$.

This process repeats/converges to optimal strategies $(\mathbf{p}^*, \mathbf{q}^*)$ where

$$\text{val}(A) = \max_{\mathbf{p}} \mathbb{E}(\mathbf{p}, \mathbf{q}^*) = \max_{\mathbf{p}} \min_{\mathbf{q}} \mathbb{E}(\mathbf{p}, \mathbf{q}) \quad \text{and}$$

$$\text{val}(A) = \min_{\mathbf{q}} \mathbb{E}(\mathbf{p}^*, \mathbf{q}) = \min_{\mathbf{q}} \max_{\mathbf{p}} \mathbb{E}(\mathbf{p}, \mathbf{q}).$$

Optimal Mixed Strategy

Natural Protocol:

Ron can estimate \mathbf{q} by keeping statistics over many rounds of play. Assume that Ron knows \mathbf{q} .

Ron selects \mathbf{p} to maximize $\mathbb{E}(\mathbf{p}, \mathbf{q})$. Then Coleen examines \mathbf{p} and selects \mathbf{q} to minimize $\mathbb{E}(\mathbf{p}, \mathbf{q})$.

This process repeats/converges to optimal strategies $(\mathbf{p}^*, \mathbf{q}^*)$ where

$$\text{val}(A) = \max_{\mathbf{p}} \mathbb{E}(\mathbf{p}, \mathbf{q}^*) = \max_{\mathbf{p}} \min_{\mathbf{q}} \mathbb{E}(\mathbf{p}, \mathbf{q}) \quad \text{and}$$

$$\text{val}(A) = \min_{\mathbf{q}} \mathbb{E}(\mathbf{p}^*, \mathbf{q}) = \min_{\mathbf{q}} \max_{\mathbf{p}} \mathbb{E}(\mathbf{p}, \mathbf{q}).$$

Coleen's Modified Protocol:

Coleen selects \mathbf{q} and shows it to Ron. Then Ron examines \mathbf{q} and chooses a **pure strategy** $\mathbf{p} = \mathbf{e}_i$. The expected payoff is

$$F_i(\mathbf{q}) := \mathbb{E}(\mathbf{e}_i, \mathbf{q}).$$

Ron selects i to maximize this quantity. Coleen selects \mathbf{q} to minimize the expected payoff

$$\max_i F_i(\mathbf{q}).$$

Coleen optimal strategy, call it \mathbf{q}' , will result in an expected loss

$$v' = \max_i F_i(\mathbf{q}') = \min_{\mathbf{q}} \max_i F_i(\mathbf{q}).$$

Optimal Mixed Strategy

Natural Protocol:

Ron can estimate \mathbf{q} by keeping statistics over many rounds of play. Assume that Ron knows \mathbf{q} .

Ron selects \mathbf{p} to maximize $\mathbb{E}(\mathbf{p}, \mathbf{q})$. Then Coleen examines \mathbf{p} and selects \mathbf{q} to minimize $\mathbb{E}(\mathbf{p}, \mathbf{q})$.

This process repeats/converges to optimal strategies $(\mathbf{p}^*, \mathbf{q}^*)$ where

$$\text{val}(A) = \max_{\mathbf{p}} \mathbb{E}(\mathbf{p}, \mathbf{q}^*) = \max_{\mathbf{p}} \min_{\mathbf{q}} \mathbb{E}(\mathbf{p}, \mathbf{q}) \quad \text{and}$$

$$\text{val}(A) = \min_{\mathbf{q}} \mathbb{E}(\mathbf{p}^*, \mathbf{q}) = \min_{\mathbf{q}} \max_{\mathbf{p}} \mathbb{E}(\mathbf{p}, \mathbf{q}).$$

Coleen's Modified Protocol:

Coleen selects \mathbf{q} and shows it to Ron. Then Ron examines \mathbf{q} and chooses a **pure strategy** $\mathbf{p} = \mathbf{e}_i$. The expected payoff is

$$F_i(\mathbf{q}) := \mathbb{E}(\mathbf{e}_i, \mathbf{q}).$$

Ron selects i to maximize this quantity. Coleen selects \mathbf{q} to minimize the expected payoff

$$\max_i F_i(\mathbf{q}).$$

Coleen optimal strategy, call it \mathbf{q}' , will result in an expected loss

$$v' = \max_i F_i(\mathbf{q}') = \min_{\mathbf{q}} \max_i F_i(\mathbf{q}).$$

Ron's Modified Protocol:

Arguing similarly, if Ron selects strategy \mathbf{p} , and Coleen selects pure strategy \mathbf{e}_j which minimizes

$$E_j(\mathbf{p}) := \mathbb{E}(\mathbf{p}, \mathbf{e}_j),$$

then Ron's optimal strategy \mathbf{p}' is a vector which maximizes $\min_j E_j(\mathbf{p})$. His expected payoff is

$$u' = \max_{\mathbf{p}} E_j(\mathbf{p}') = \max_{\mathbf{p}} \min_j E_j(\mathbf{p}).$$

$$A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mn} \end{bmatrix}$$

Coleen's modification

If Ron plays row i , the expected payoff is

$$F_i(\mathbf{q}) = \mathbf{e}_i^\top A \mathbf{q} = q_1 a_{i1} + \dots + q_n a_{in}$$

Coleen chooses $\mathbf{q} = \mathbf{q}'$ in order to minimize

$$\max_{1 \leq i \leq m} F_i(\mathbf{q})$$

Expected payoff of Coleen's modification is

$$v' = \max_{1 \leq i \leq m} F_i(\mathbf{q}') = \min_{\mathbf{q}} \max_{1 \leq i \leq m} F_i(\mathbf{q})$$

Ron's modification

If Coleen plays column j , the expected payoff is

$$E_j(\mathbf{p}) = \mathbf{p}^\top A \mathbf{e}_j = p_1 a_{1j} + \dots + p_m a_{mj}$$

Ron chooses $\mathbf{p} = \mathbf{p}'$ in order to maximize

$$\min_{1 \leq j \leq n} E_j(\mathbf{p})$$

Expected payoff of Ron's modification is

$$u' = \min_{1 \leq j \leq n} E_j(\mathbf{p}') = \mathbf{p} \min_{1 \leq j \leq n} E_j(\mathbf{p})$$

$$A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mn} \end{bmatrix}$$

Coleen's modification

If Ron plays row i , the expected payoff is

$$F_i(\mathbf{q}) = \mathbf{e}_i^\top A \mathbf{q} = q_1 a_{i1} + \dots + q_n a_{in}$$

Coleen chooses $\mathbf{q} = \mathbf{q}'$ in order to minimize

$$\max_{1 \leq i \leq m} F_i(\mathbf{q})$$

Expected payoff of Coleen's modification is

$$v' = \max_{1 \leq i \leq m} F_i(\mathbf{q}') = \min_{\mathbf{q}} \max_{1 \leq i \leq m} F_i(\mathbf{q})$$

Since \mathbf{e}_i might not be optimal for Ron,

$$v' \leq \text{val}(A).$$

Ron's modification

If Coleen plays column j , the expected payoff is

$$E_j(\mathbf{p}) = \mathbf{p}^\top A \mathbf{e}_j = p_1 a_{1j} + \dots + p_m a_{mj}$$

Ron chooses $\mathbf{p} = \mathbf{p}'$ in order to maximize

$$\min_{1 \leq j \leq n} E_j(\mathbf{p})$$

Expected payoff of Ron's modification is

$$u' = \min_{1 \leq j \leq n} E_j(\mathbf{p}') = \mathbf{p} \min_{1 \leq j \leq n} E_j(\mathbf{p})$$

Since \mathbf{e}_j might not be optimal for Coleen,

$$\text{val}(A) \leq u'.$$

Optimal Mixed Strategy

$$A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mn} \end{bmatrix}$$

Coleen's modification

If Ron plays row i , the expected payoff is

$$F_i(\mathbf{q}) = \mathbf{e}_i^\top A \mathbf{q} = q_1 a_{i1} + \dots + q_n a_{in}$$

Coleen chooses $\mathbf{q} = \mathbf{q}'$ in order to minimize

$$\max_{1 \leq i \leq m} F_i(\mathbf{q})$$

Expected payoff of Coleen's modification is

$$v' = \max_{1 \leq i \leq m} F_i(\mathbf{q}') = \min_{\mathbf{q}} \max_{1 \leq i \leq m} F_i(\mathbf{q})$$

Since \mathbf{e}_i might not be optimal for Ron,

$$v' \leq \text{val}(A).$$

Ron's modification

If Coleen plays column j , the expected payoff is

$$E_j(\mathbf{p}) = \mathbf{p}^\top A \mathbf{e}_j = p_1 a_{1j} + \dots + p_m a_{mj}$$

Ron chooses $\mathbf{p} = \mathbf{p}'$ in order to maximize

$$\min_{1 \leq j \leq n} E_j(\mathbf{p})$$

Expected payoff of Ron's modification is

$$u' = \min_{1 \leq j \leq n} E_j(\mathbf{p}') = \mathbf{p} \min_{1 \leq j \leq n} E_j(\mathbf{p})$$

Since \mathbf{e}_j might not be optimal for Coleen,

$$\text{val}(A) \leq u'.$$

Surprise! We have $v' = u'$ so $\text{val}(A) = v' = u'$ and $(\mathbf{p}', \mathbf{q}')$ are optimal for the natural game!

Theorem

Let $A = [a_{ij}]$ be an $m \times n$ matrix game. Then the mixed strategies \mathbf{p} and \mathbf{q} obtained by solving the dual non-canonical LP problems with tableau

	Ⓥ	q_1	...	q_n	-1	
Ⓢ	0	-1	...	-1	-1	$= -0$
p_1	-1				0	$= -t_1$
\vdots	\vdots				\vdots	\vdots
p_m	-1				0	$= -t_m$
-1	-1	0	...	0	0	$= f$
	$= 0$	$= s_1$...	$= s_n$	$= g$	

are optimal for the Ron and Coleen player respectively. Both linear programs always have an optimal solution. The dual non-canonical tableau above is called the **game tableau** for A .

The dual non-canonical tableau above is called the **game tableau** for A .

Proof.

For any \mathbf{q} , we have

$$\max_{1 \leq i \leq m} F_i(\mathbf{q}) = \min\{v \mid v \geq F_i(\mathbf{q}), \text{ for } i = 1, 2, \dots, m \}.$$

Proof.

For any \mathbf{q} , we have

$$\max_{1 \leq i \leq m} F_i(\mathbf{q}) = \min\{v \mid v \geq F_i(\mathbf{q}), \text{ for } i = 1, 2, \dots, m\}.$$

Coleen's optimal vector $\mathbf{q}' = (q_1, q_2, \dots, q_n)$ minimizes this among all probability vectors \mathbf{q} .

$$v' = \min\{v \mid v \geq F_i(\mathbf{q}) \text{ (} i = 1, 2, \dots, m \text{) and } q_1 + q_2 + \dots + q_n = 1 \text{ and } q_1, q_2, \dots, q_n \geq 0\}.$$

Thus $v', q'_1, q'_2, \dots, q'_n$ is the optimal solution of a linear program with variables v, q_1, q_2, \dots, q_n .

Proof.

For any \mathbf{q} , we have

$$\max_{1 \leq i \leq m} F_i(\mathbf{q}) = \min\{v \mid v \geq F_i(\mathbf{q}), \text{ for } i = 1, 2, \dots, m\}.$$

Coleen's optimal vector $\mathbf{q}' = (q_1, q_2, \dots, q_n)$ minimizes this among all probability vectors \mathbf{q} .

$$v' = \min\{v \mid v \geq F_i(\mathbf{q}) \text{ (} i = 1, 2, \dots, m \text{) and } q_1 + q_2 + \dots + q_n = 1 \text{ and } q_1, q_2, \dots, q_n \geq 0\}.$$

Thus $v', q'_1, q'_2, \dots, q'_n$ is the optimal solution of a linear program with variables v, q_1, q_2, \dots, q_n .

$$v' = \text{minimize } v$$

$$v \geq F_1(\mathbf{q})$$

$$v \geq F_2(\mathbf{q})$$

$$\vdots$$

$$v \geq F_m(\mathbf{q})$$

$$q_1 + q_2 + \dots + q_n = 1$$

$$q_1, q_2, \dots, q_n \geq 0$$

$$v \text{ unconstrained}$$



Proof.

For any \mathbf{q} , we have

$$\max_{1 \leq i \leq m} F_i(\mathbf{q}) = \min\{v \mid v \geq F_i(\mathbf{q}), \text{ for } i = 1, 2, \dots, m\}.$$

Coleen's optimal vector $\mathbf{q}' = (q_1, q_2, \dots, q_n)$ minimizes this among all probability vectors \mathbf{q} .

$$v' = \min\{v \mid v \geq F_i(\mathbf{q}) \text{ (} i = 1, 2, \dots, m \text{) and } q_1 + q_2 + \dots + q_n = 1 \text{ and } q_1, q_2, \dots, q_n \geq 0\}.$$

Thus $v', q'_1, q'_2, \dots, q'_n$ is the optimal solution of a linear program with variables v, q_1, q_2, \dots, q_n .

$$v' = \text{minimize } v$$

$$v \geq F_1(\mathbf{q})$$

$$v \geq F_2(\mathbf{q})$$

$$\vdots$$

$$v \geq F_m(\mathbf{q})$$

$$q_1 + q_2 + \dots + q_n = 1$$

$$q_1, q_2, \dots, q_n \geq 0$$

$$v \text{ unconstrained}$$

$$\text{maximize } -v = f(v, q_1, q_2, \dots, q_n)$$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + F_1(\mathbf{q}) \leq 0$$

$$-v + F_2(\mathbf{q}) \leq 0$$

$$\vdots$$

$$-v + F_m(\mathbf{q}) \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

$$v \text{ unconstrained}$$



Proof.

For any \mathbf{q} , we have

$$\max_{1 \leq i \leq m} F_i(\mathbf{q}) = \min\{v \mid v \geq F_i(\mathbf{q}), \text{ for } i = 1, 2, \dots, m\}.$$

Coleen's optimal vector $\mathbf{q}' = (q_1, q_2, \dots, q_n)$ minimizes this among all probability vectors \mathbf{q} .

$$v' = \min\{v \mid v \geq F_i(\mathbf{q}) \text{ (} i = 1, 2, \dots, m \text{) and } q_1 + q_2 + \dots + q_n = 1 \text{ and } q_1, q_2, \dots, q_n \geq 0\}.$$

Thus $v', q'_1, q'_2, \dots, q'_n$ is the optimal solution of a linear program with variables v, q_1, q_2, \dots, q_n .

(P) maximize $-v = f$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + a_{11}q_1 + a_{12}q_2 + \dots + a_{1n}q_n \leq 0$$

$$-v + a_{21}q_1 + a_{22}q_2 + \dots + a_{2n}q_n \leq 0$$

$$\vdots$$

$$-v + a_{m1}q_1 + a_{m2}q_2 + \dots + a_{mn}q_n \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

v unconstrained

maximize $-v = f(v, q_1, q_2, \dots, q_n)$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + F_1(\mathbf{q}) \leq 0$$

$$-v + F_2(\mathbf{q}) \leq 0$$

$$\vdots$$

$$-v + F_m(\mathbf{q}) \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

v unconstrained



Proof.

Similarly Ron's optimal vector $\mathbf{p}' = (p_1, p_2, \dots, p_m)$ is found by a linear program with variables u, p_1, p_2, \dots, p_m .

$$u' = \max_{\mathbf{p}} \min_j E_j(\mathbf{p})$$

$$= \max\{u \mid u \leq E_j(\mathbf{p}), (j = 1, 2, \dots, n) \text{ and } p_1 + p_2 + \dots + p_m = 1 \text{ and } p_1, p_2, \dots, p_m \geq 0\}$$

(P) maximize $-v = f$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + a_{11}q_1 + a_{12}q_2 + \dots + a_{1n}q_n \leq 0$$

$$-v + a_{21}q_1 + a_{22}q_2 + \dots + a_{2n}q_n \leq 0$$

$$\vdots$$

$$-v + a_{m1}q_1 + a_{m2}q_2 + \dots + a_{mn}q_n \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

v unconstrained

Proof.

Similarly Ron's optimal vector $\mathbf{p}' = (p_1, p_2, \dots, p_m)$ is found by a linear program with variables u, p_1, p_2, \dots, p_m .

$$u' = \max_{\mathbf{p}} \min_j E_j(\mathbf{p})$$

$$= \max\{u \mid u \leq E_j(\mathbf{p}), (j = 1, 2, \dots, n) \text{ and } p_1 + p_2 + \dots + p_m = 1 \text{ and } p_1, p_2, \dots, p_m \geq 0\}$$

(P) maximize $-v = f$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + a_{11}q_1 + a_{12}q_2 + \dots + a_{1n}q_n \leq 0$$

$$-v + a_{21}q_1 + a_{22}q_2 + \dots + a_{2n}q_n \leq 0$$

$$\vdots$$

$$-v + a_{m1}q_1 + a_{m2}q_2 + \dots + a_{mn}q_n \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

$$v \text{ unconstrained}$$

$u' = \text{maximize } u$

$$u \leq F_1(\mathbf{p})$$

$$u \leq F_2(\mathbf{p})$$

$$\vdots$$

$$u \leq F_m(\mathbf{p})$$

$$p_1 + p_2 + \dots + p_m = 1$$

$$p_1, p_2, \dots, p_m \geq 0$$

$$u \text{ unconstrained}$$

Proof.

Similarly Ron's optimal vector $\mathbf{p}' = (p_1, p_2, \dots, p_m)$ is found by a linear program with variables u, p_1, p_2, \dots, p_m .

$$u' = \max_{\mathbf{p}} \min_j E_j(\mathbf{p})$$

$$= \max\{u \mid u \leq E_j(\mathbf{p}), (j = 1, 2, \dots, n) \text{ and } p_1 + p_2 + \dots + p_m = 1 \text{ and } p_1, p_2, \dots, p_m \geq 0\}$$

(P) maximize $-v = f$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + a_{11}q_1 + a_{12}q_2 + \dots + a_{1n}q_n \leq 0$$

$$-v + a_{21}q_1 + a_{22}q_2 + \dots + a_{2n}q_n \leq 0$$

$$\vdots$$

$$-v + a_{m1}q_1 + a_{m2}q_2 + \dots + a_{mn}q_n \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

v unconstrained

minimize $-u = g(u, p_1, p_2, \dots, p_m)$

$$-p_1 - p_2 - \dots - p_m = -1$$

$$-u + F_1(\mathbf{p}) \geq 0$$

$$-u + F_2(\mathbf{p}) \geq 0$$

$$\vdots$$

$$-u + F_n(\mathbf{p}) \geq 0$$

$$p_1, p_2, \dots, p_m \geq 0$$

u unconstrained

Proof.

Similarly Ron's optimal vector $\mathbf{p}' = (p_1, p_2, \dots, p_m)$ is found by a linear program with variables u, p_1, p_2, \dots, p_m .

$$u' = \max_{\mathbf{p}} \min_j E_j(\mathbf{p})$$

$$= \max\{u \mid u \leq E_j(\mathbf{p}), (j = 1, 2, \dots, n) \text{ and } p_1 + p_2 + \dots + p_m = 1 \text{ and } p_1, p_2, \dots, p_m \geq 0\}$$

(P) maximize $-v = f$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + a_{11}q_1 + a_{12}q_2 + \dots + a_{1n}q_n \leq 0$$

$$-v + a_{21}q_1 + a_{22}q_2 + \dots + a_{2n}q_n \leq 0$$

$$\vdots$$

$$-v + a_{m1}q_1 + a_{m2}q_2 + \dots + a_{mn}q_n \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

v unconstrained

(D) minimize $-u = g$

$$-p_1 - p_2 - \dots - p_m = -1$$

$$-u + a_{11}p_1 + a_{21}p_2 + \dots + a_{m1}p_m \geq 0$$

$$-u + a_{12}p_1 + a_{23}p_2 + \dots + a_{m2}p_m \geq 0$$

$$\vdots$$

$$-u + a_{1m}p_1 + a_{2m}p_2 + \dots + a_{mm}p_m \geq 0$$

$$p_1, p_2, \dots, p_m \geq 0$$

u unconstrained

Proof.

Similarly Ron's optimal vector $\mathbf{p}' = (p_1, p_2, \dots, p_m)$ is found by a linear program with variables u, p_1, p_2, \dots, p_m .

$$u' = \max_{\mathbf{p}} \min_j E_j(\mathbf{p})$$

$$= \max\{u \mid u \leq E_j(\mathbf{p}), (j = 1, 2, \dots, n) \text{ and } p_1 + p_2 + \dots + p_m = 1 \text{ and } p_1, p_2, \dots, p_m \geq 0\}$$

(P) maximize $-v = f$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + a_{11}q_1 + a_{12}q_2 + \dots + a_{1n}q_n \leq 0$$

$$-v + a_{21}q_1 + a_{22}q_2 + \dots + a_{2n}q_n \leq 0$$

$$\vdots$$

$$-v + a_{m1}q_1 + a_{m2}q_2 + \dots + a_{mn}q_n \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

v unconstrained

(D) minimize $-u = g$

$$-p_1 - p_2 - \dots - p_m = -1$$

$$-u + a_{11}p_1 + a_{21}p_2 + \dots + a_{m1}p_m \geq 0$$

$$-u + a_{12}p_1 + a_{23}p_2 + \dots + a_{m2}p_m \geq 0$$

$$\vdots$$

$$-u + a_{1m}p_1 + a_{2m}p_2 + \dots + a_{mm}p_m \geq 0$$

$$p_1, p_2, \dots, p_m \geq 0$$

u unconstrained

These are the dual non-canonical LPs for the game tableau.

The optimal value is $f = g = -u = -v = -\text{val}(A)$.

Proof.

Similarly Ron's optimal vector $\mathbf{p}' = (p_1, p_2, \dots, p_m)$ is found by a linear program with variables u, p_1, p_2, \dots, p_m .

$$\begin{aligned} u' &= \max_{\mathbf{p}} \min_j E_j(\mathbf{p}) \\ &= \max\{u \mid u \leq E_j(\mathbf{p}), (j = 1, 2, \dots, n) \text{ and } p_1 + p_2 + \dots + p_m = 1 \text{ and } p_1, p_2, \dots, p_m \geq 0\} \end{aligned}$$

(P) maximize $-v = f$

$$-q_1 - q_2 - \dots - q_n = -1$$

$$-v + a_{11}q_1 + a_{12}q_2 + \dots + a_{1n}q_n \leq 0$$

$$-v + a_{21}q_1 + a_{22}q_2 + \dots + a_{2n}q_n \leq 0$$

$$\vdots$$

$$-v + a_{m1}q_1 + a_{m2}q_2 + \dots + a_{mn}q_n \leq 0$$

$$q_1, q_2, \dots, q_n \geq 0$$

v unconstrained

(D) minimize $-u = g$

$$-p_1 - p_2 - \dots - p_m = -1$$

$$-u + a_{11}p_1 + a_{21}p_2 + \dots + a_{m1}p_m \geq 0$$

$$-u + a_{12}p_1 + a_{23}p_2 + \dots + a_{m2}p_m \geq 0$$

$$\vdots$$

$$-u + a_{1m}p_1 + a_{2m}p_2 + \dots + a_{mm}p_m \geq 0$$

$$p_1, p_2, \dots, p_m \geq 0$$

u unconstrained

These are the dual non-canonical LPs for the game tableau.

The optimal value is $f = g = -u = -v = -\text{val}(A)$.

Both optima exist, since making v and $-u$ very large makes both LP's feasible.



Von Neumann Minimax Theorem

Theorem (Von Neumann Minimax Theorem)

Let $A = [a_{ij}]$ be an $m \times n$ matrix game. Then there exist optimal mixed strategies \mathbf{p}^* and \mathbf{q}^* for the row player and the column player respectively. Furthermore,

$$\min_{1 \leq j \leq n} E_j(\mathbf{p}^*) = \max_{\mathbf{p}} \min_{1 \leq j \leq n} E_j(\mathbf{p}) = \min_{\mathbf{q}} \max_{1 \leq i \leq m} F_i(\mathbf{q}) = \min_{1 \leq i \leq m} F_i(\mathbf{q}^*)$$

and this common value is said to be the **von Neumann value** of the game.

The von Neumann value is the amount of expected winnings of the row player and the expected losses of the column player.

A positive von Neumann value indicates that the game favours the row player, a negative von Neumann value indicates that the game favours the column player, and a von Neumann value 0 indicates that the game is fair.

Saddle Point and von Neumann Value of a Game

Recall the definition of a saddle point of a matrix game:

Definition (Saddle Point)

An entry a_{ij} of a matrix game $A \in \mathbb{R}^{m \times n}$ is called a *saddle point* if it is a minimum entry in the row $\#i$ and the maximum entry in the column $\#j$.

Theorem

If a payoff matrix A has a saddle point a_{ij} , then the von Neumann value of the game is a_{ij} .

Proof.



Transforming Maximum Game Tableau to a Basic Feasible Game Tableau

We need at least two pivot transformations to get a Basic Feasible Tableau.

And we do not need more! We can skip Phase 1 of SA by selecting the first two pivots carefully.

Algorithm (Transforming the original game tableau into the Feasible Tableau)

1. Find the maximum entry in each column of the matrix game A .
2. Choose the minimum of these maximum entries, say it is the entry a_{ij} of A .
3. Pivot on the -1 in the top of column $\#j$ and the -1 in the left of row $\#i$ (in either order).
Remove 0 -column and 0 -row after pivoting.
We do not need to file equations for u and v as they can be obtained from objective functions.
4. The resulting tableau is Feasible Tableau.

We do not prove that this procedure works as claimed.