

SUBJECT INFORMATION SHEET

Study Title	A Randomized, Double-masked, Active Controlled, Parallel Group, Multicentre Bioequivalence Study with Clinical Endpoint Comparing Brinzolamide 1% Ophthalmic Suspension (Sponsor), to the reference product (Brinzolamide) 1% Ophthalmic Suspension of (the originator manufacturer) in the Treatment of Chronic Open Angle Glaucoma or Ocular Hypertension in Both Eyes.
Protocol Number	Sample/2026/01
Protocol Version Number and Date	[Version, Date]
Subject Information Sheet and Informed Consent Form – Version and Date	[Version, Date]
Sponsor	Sponsor details
Site ID	
Subject Name	
Principal Investigator Name	
Site Address	
Principal Investigator/ Designee Contact Details	
Subject Number	
Subject Initials	

Dear Prospective Subject,

You are invited to participate in this study, the title of which is given above. This is a research study and if you do not wish to participate, your medical care will not be affected in any way. The purpose of this informed consent document is to clearly state the procedures of the study and to ensure that you participate in this study only after knowing all the relevant facts. An oral presentation of this document will be made after you go through this Subject Information Sheet and Informed Consent Form. Please read the information carefully and ask the study doctor or the study staff to explain anything you do not understand and clarify any doubts if any.

You can take home a copy of this document to think about or talk with family or friends before making your decision. It is important that you must be aware of the risks and benefits of the study to make an informed decision. Only if you have understood all the salient aspects of the study procedures and decide to take part in this study, you will be asked to sign this Informed Consent Form (ICF) to confirm that you understood the information and that you have given your voluntary consent to participate in the research study.

A photocopy of the signed informed consent form will be provided to you for your reference and records.

Nature and Purpose of the Study

This study shall be conducted in subjects with Chronic Open Angle Glaucoma or Ocular Hypertension. We are asking you to take part in this study because you have Chronic Open Angle Glaucoma/Ocular Hypertension in both eyes.

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

Eye pressure (called intraocular pressure or IOP) is measured in millimeters of mercury (mm Hg). Normal eye pressure ranges from 12-21 mm Hg, and eye pressure greater than 21 mm Hg is considered higher than normal. When the IOP is higher than normal is referred to as Ocular Hypertension. It is caused by the trabecular blockage. Because the microscopic passageways are blocked, the pressure builds up in the eye and causes imperceptible or very gradual vision loss. Peripheral vision is affected first, but eventually, the entire vision will be lost if not treated.

Open-angle glaucoma is when optic nerve damage results in a progressive loss of the visual field. This is associated with increased pressure in the eye. Not all people with primary open-angle glaucoma have eye pressure that is elevated beyond normal, but decreasing the eye pressure further has been shown to stop progression even in these cases.

In this study, the drug tested is Brinzolamide 1% ophthalmic suspension which is an approved drug, available in the market as a safe and effective treatment for Chronic Open Angle Glaucoma or Ocular Hypertension. There are similarly marketed products of the same medications also available on the market.

In this study, there are two study medications i.e., Test product, and Reference product will be tested. The test product Brinzolamide 1% ophthalmic suspension is an experimental product developed by (Sponsor) and has not been approved for use. It is being tested as an experimental product in comparison with the reference product (Brinzolamide) 1% ophthalmic suspension of (the originator manufacturer) to find out whether it has a similar effect in the treatment of chronic open angle glaucoma or ocular hypertension. During the study, you will receive any one of the study medications.

Neither you nor your study doctor (double-masked) will know which of the study medications you are receiving. The decision as to which drug you will receive is determined using a computerized process called randomization (like tossing a coin) and it cannot be changed by anyone. Every eligible subject who participates in the study will receive randomly any one of the two study medications. This is to prevent you and your study doctor from preferring one treatment to another. You will have a 50:50 chance of receiving either the test or reference product.

There is also a possibility that the Study Medication fails to provide the intended therapeutic/drug effect. Even though the study is double-masked, the mask can be broken in case of an emergency so that the study doctor can quickly find out to which study group you are assigned.

This research study procedure has been thoroughly reviewed by the concerned doctors, medical experts, ethics committee, and applicable regulatory body.

Number of Subjects in the Study

Approximately 670 male or non-pregnant females aged at least 18 years shall be enrolled/randomized into the study.

Study Procedures

If you decided to participate in the study, you will have to come to the hospital for a check-up. The study duration will be approximately a maximum of 13 to 17 weeks (including 6 weeks screening period).

You should complete four scheduled clinic visits as follows

- Visit 1 - Screening Visit (Day -42 to Day -1)

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

- Visit 2 - Baseline/Randomization Visit (Day 0)
- Visit 3 - Interim Visit (Day 14 $\pm$ 2 days)
- Visit 4 - End of Treatment/Early Termination Visit (Day 42 $\pm$ 2 days)

A telephonic call shall be done from the study site personnel to you after 7 days $\pm$ 2 days of the end of treatment visit 4. This is to ensure your overall well-being.

However, you are free to visit the study doctor at any time if you experience any discomfort related to the use of the study medication. You must bring your study medication and duly filled subject diary during such visits.

The following activities will be performed during specific visits. You are requested to reach the investigator site well ahead of time to facilitate protocol specific assessments.

Visit 1: Screening Visit (Day -42 to Day -1)

- Informed consent Process: The study doctor or a member of the study staff will explain to you the details of the study. You will be provided sufficient time to read and understand the study details and any doubts that you may have will be clarified by delegated study person or study doctor. If you decide to participate you will be asked to sign and date this informed consent form
- Demographic data (Height, weight, age, sex, ethnicity, race, irides colour, etc.) will be noted
- Medical/surgical/medication/ocular history: Information regarding the subject's medical, and surgical history, and any underlying condition that began within the last 30 days information regarding your current and past medical conditions will be taken including surgical procedures. Tell your doctor that you are breastfeeding or plan to breastfeed, you are pregnant or plan to become pregnant. Tell your doctor about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements etc. and also have taken but discontinued within the 30 days prior to screening Also, tell your doctor about the contraceptive which you are adopting currently (if any)
- Vital signs such as heart rate and blood pressure will be measured and recorded after you have rested for at least 5 minutes
- Assessment of COVID-19 - After obtaining the ICF, you shall be assessed on general health status related to COVID-19 and documented.
- Urine pregnancy test will be performed (if you are female with childbearing potential) and negative result of a urine pregnancy test should be obtained for the participation
- You will be evaluated for Best-spectacle corrected visual acuity, Corneal pachymetry, Slit lamp biomicroscopy, IOP measurement in both eyes by Goldmann Tonometer, Gonioscopy, Pupil dilation, and Fundus ophthalmoscopy
 - Automated perimetry will be done only if there is no data within the past 6 months
 - Best-spectacle corrected visual acuity: Subjects must have a score of 0.8 logMAR or better in each eye in order to qualify
 - Slit lamp biomicroscopy: A slit lamp exam shall be performed to exclude subjects with disallowed ocular conditions
 - Intraocular pressure (Goldmann applanation tonometry): IOP measurements shall be taken using a Goldmann applanation tonometer. IOP shall be measured at least twice in each eye with the right eye measured first. If the difference of the first 2 measurements is ≤ 1 mm Hg, the average of these 2 measurements shall be used as the IOP data for this eye. If the difference is > 1 mm Hg, a third measurement shall be taken, and the median of these 3 measurements shall be used as the IOP data for this eye
 - Corneal pachymetry for central corneal thickness: (required only if there is no data in within past 30 days). Corneal thickness shall be measured using an ultrasound pachymeter. Subjects shall be excluded in corneal thickness is > 600 μ m.

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

- Gonioscopy: To examine iridocorneal angles (open, closed, or partially closed). The procedure shall be conducted per the investigator's standard of practice with grading based on the Shaffer scale
- Dilated ophthalmoscopy including C/D ratio: A dilated ophthalmoscopy shall be performed by the investigator to evaluate the presence or absence of clinically significant fundus abnormalities, vitreous pathology.
- Discontinuation of current IOP lowering medications, if any
- Evaluation of inclusion and exclusion criteria - to check whether you are eligible to participate in the study. When you have completed all screening procedures, compliance with the inclusion and exclusion criteria will be reviewed. After the inclusion and exclusion criteria have been confirmed, you will enter the washout period.
- Washout period- A minimum washout period of 4 days to 4 weeks has to be followed if you selected during the screening visit. To minimize potential risk to you, due to intraocular pressure elevations during the washout period, your study doctor may substitute medicine in place of your current medication if required. After the appropriate washout period, a baseline visit (visit II) will be performed
- Inform your doctor if you experience any adverse events and taken any medication upon signing the informed consent form
- Date of next visit to the hospital and actions will be informed to you after the appropriate washout period.

Visit 2: Baseline/Randomization Visit (Day 0)

- Assessment of general health status related to COVID-19 shall be performed
- You will be randomized in the study.
- The use of any medications since the previous study visit shall be assessed and recorded
- The occurrence of all adverse events at each visit will be assessed and recorded
- Vital signs such as heart rate and blood pressure will be measured and recorded after you have rested for at least 5 minutes
- Urine pregnancy test will be performed (Only for women of childbearing potential) Best-spectacle corrected visual acuity, Slit-lamp biomicroscopy, Intraocular pressure 8:00 AM $\pm$ 30 minutes: by Goldmann applanation tonometer (this shall be the qualifying IOP for enrolment) Intraocular pressure at 10:00 AM $\pm$ 30 minutes: by Goldmann applanation tonometer
- Inclusion and exclusion criteria will be reviewed and study medication will be provided to you if found eligible for the study
- Study Medication shall be dispensed to you by the site staff. Instructions will be provided to you on how to use the study medication and completion of subject diaries. You should start using the Study Medication at home starting the next day of visit 2. The study medication should not be opened at the site. You should come for the next visit without instilling the morning dose on the visit date. You have to shake the bottle well and then instill Investigational Product into the inferior cul-de-sac of both eyes at 8.00 AM. ($\pm$ 60 minutes), 4.00 PM. ($\pm$ 60 minutes) and 10.00 PM. ($\pm$ 60 minutes)
- A subject diary, containing instructions for subjects, shall be dispensed to you. Apart from the instill dose details, you will be instructed to document any adverse event, concomitant medication taken and any additional application made
- At the end of each visit, your next visit will be scheduled by the investigator or his team member. You are required to bring the used study medication along with its outer box at the next study visit.

Visit 3: Interim Visit (Day 14 ± 2 days)

- Assessment of general health status related to COVID-19 shall be performed
- The use of any medications since the previous study visit shall be assessed and recorded
- The occurrence of all adverse events at each visit will be assessed and recorded
- Vital signs such as heart rate and blood pressure will be measured and recorded after you have rested for at least 5 minutes
- Best-spectacle corrected visual acuity, Slit-lamp biomicroscopy, Intraocular pressure in both eyes (at 8:00 AM ± 30 minutes and at 10:00 AM ± 30 minutes) by Goldmann applanation tonometer
- In house Study Medication instillation (8:00 AM ± 30 minutes dose) shall be done after the IOP reading. The dose shall be self-administered by you in a secluded room without any investigational site staff present.
- The quantity of study medication remaining in the container and the number of doses mentioned in the subject diary is verified to ensure compliance. Any discrepancy found with the number of doses mentioned in the diary and the study medication quantity used by you will be discussed by your study doctor, to verify whether you are following the correct quantity and dosage instructions
- Study Medication shall be dispensed by the site staff to you. You should come for the next visit without instilling the morning dose on the visit date
- Review of subject diary – You should be instructed to continue the assigned medication as per the dosage
- Schedule next visit - You are required to bring the used study medication along with its outer box at the next study visit.

Visit 4: End of Treatment/Early Termination Visit (Day 42 ± 2 days)

Visit 4 is to be scheduled for the day after the last day of dosing.

- Assessment of general health status related to COVID-19 shall be performed
- The use of any medications since the previous study visit shall be assessed and recorded
- The occurrence of all adverse events at each visit will be assessed and recorded
- Vital signs such as heart rate and blood pressure will be measured and recorded after you have rested for at least 5 minutes
- Urine pregnancy test will be performed (Only for women of childbearing potential)
- Best-spectacle corrected visual acuity, Slit-lamp biomicroscopy, Intraocular pressure in both eyes (at 08:00 AM ± 30 minutes and at 10:00 AM ± 30 minutes) by Goldmann applanation tonometer and Dilated ophthalmoscopy
- In house study medication instillation (8:00 AM ± 30 minutes dose) shall be done after the IOP reading. The dose shall be self-administered by you in a secluded room without any investigational site staff present
- Study Medication with the remaining study medication quantity will be collected back from you. The quantity of study medication remaining in the container and the number of doses mentioned in the subject diary is verified to ensure compliance. Any discrepancy found with the number of doses mentioned in the diary and the study medication quantity used by you will be discussed by your study doctor, to verify whether you are following the correct quantity and dosage instructions
- The completed subject diary will be reviewed and collected at this visit
- You will be instructed to contact the investigator in case of any adverse event that he considers as significant, up to 7 days after the visit.

In case of early discontinuation after randomization, following assessments to be performed:

- Assessment of general health status related to COVID-19 shall be performed
- The use of any medications since the previous study visit shall be assessed and recorded
- The occurrence of all adverse events at each visit will be assessed and recorded
- Vital signs such as heart rate and blood pressure will be measured and recorded after you have rested for at least 5 minutes
- Urine pregnancy test will be performed (Only for women of childbearing potential)
- Best-spectacle corrected visual acuity, Slit-lamp biomicroscopy, intraocular pressure in both eyes (at 08:00 AM $\pm$ 30 minutes and at 10:00 AM $\pm$ 30 minutes) by Goldmann applanation tonometer and dilated ophthalmoscopy
- In house Study Medication instillation (8:00 AM $\pm$ 30 minutes dose) shall be done after the IOP reading. The dose shall be self-administered by you in a secluded room without any investigational site staff present
- Study Medication with the remaining study medication quantity will be collected back from you. The quantity of study medication remaining in the container and the number of doses mentioned in the subject diary is verified to ensure compliance. Any discrepancy found with the number of doses mentioned in the diary and the study medication quantity used by you will be discussed by your study doctor, to verify whether you are following the correct quantity and dosage instructions
- The completed subject diary will be reviewed and collected at this visit
- You will be instructed to contact the investigator in case of any adverse event that he considers as significant, up to 7 days after the visit.

End of Study/Safety follow-up 7 days ($\pm$ 2 days) after end of treatment visit

Study personnel will contact you telephonically after 7 days $\pm$ 2 days of visit 4. This is to ensure your overall wellbeing of you.

Early termination Visit

Even after you discontinue from the trial, you will be asked to come in for one more visit in order to return the unused medication and completed/blank subject diary. Also, to complete the end of treatment assessments.

Unscheduled Visit

You can also visit the Investigator's study site anytime, should you experience severe/intolerable Adverse Events or feel that your condition is deteriorating, for assessment.

How to Use the Study Medication

- To apply eye drops, wash your hands first
- Shake the bottle well before using
- To avoid contamination, do not touch the dropper tip or let it touch your eye or surrounding structures, or other surfaces. The product can become contaminated by common bacteria known to cause eye infections. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions
- Tilt your head back, look upward, and pull down the lower eyelid to make a pouch. Hold the dropper directly over your eye and place one drop into the pouch. Look downward and gently close your eyes for 1 to 2 minutes. Place one finger at the corner of your eye (near the nose) and apply gentle pressure. This will prevent the medication from draining out. Try not to blink and do not rub your eye. Repeat these steps for your other eye if so directed

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

- Do not rinse the dropper. Replace the dropper cap after each use
- If you are using another kind of eye medication (such as drops or ointments), wait at least 10 minutes before applying other medications
- Apply this medication to each eye
- Instill one drop 3 times daily at 8.00 AM. (± 30 minutes), 4.00 PM. (± 30 minutes) and 10.00 PM. (± 30 minutes). Remember to use it at the same times each day
- Do not dose at home before your study visits to the hospital on Week 2 and Week 6 morning
- Vision may be temporarily blurred following dosing with study medication. You have to be careful in operating machinery or driving a motor vehicle
- If serious or unusual ocular or systemic reactions or signs of hypersensitivity occur, you should discontinue the use of the study product and contact your doctor immediately
- If you see a physician for another medical problem while you are participating in this study, please contact your study doctor.
- You should not stop the application of study medication during the treatment period unless instructed by the doctor/site staff.
- You should not take any other medicine without asking your doctor/site staff
- You should bring your study medication with you at each visit in order to determine if you are using the study medication properly. You will be given an additional bottle as needed. All used and unused study medication must be returned to your doctor's clinic during the last visit
- Store the medication at room temperature below 25°C, Protect it from light, Do not freeze, Keep away from heat, Keep out of the reach of small children. Put the top back on the bottle right away after use to avoid contaminating the eye drops. Do not leave the top off the bottle for any length of time.

Other Medications while you are participating in the Study

You should not take any other medicine without asking your doctor/site staff. You must mention all your current medications and any medications taken before the start of the study. Your doctor will discuss with you the medications prohibited during the study. If you take such medications during the study then you will be discontinued from the study. In case of any ailment or sickness, avoid taking any medication or home remedies without consulting your doctor. Inform the doctor and the doctor will decide to add/remove any medication to your current Study Medication.

Please find below the drugs/procedures that are prohibited while using the Study Medication

- Ocular hypotensive drug products other than a study treatment, e.g., acetazolamide (Diamox[®]), betaxolol solution (Betoptic[®]), betaxolol and pilocarpine (Betoptic[®] Pilo), bimatoprost (Lumigan[®]), brimonidine tartrate (Alphagan[®], Alphagan[®] P), brimonidine tartrate and brinzolamide (Simbrinza[®]), brimonidine tartrate and timolol maleate (Combigan[®]), brinzolamide (Azopt[®]), carbachol (Miostat[®]), carteolol (Ocupress[®]), dorzolamide hydrochloride (Trusopt[®]), dorzolamide hydrochloride and timolol maleate (Cosopt[®]), epinephrine (Epifrin[®]), latanoprost (Xalatan[®]), levobetaxolol (Betaxon[®]), levobunolol (Akbeta[®], Betagan[®]), mannitol (Osmitol[®]), metipranolol (OptiPranolol[®]), pilocarpine (Isopto[®] Carpine, Pilopine HS[®]), tafluprost (Zioptan[™]), timolol (Betimol[®], Istalol[®], Timoptic[®], Timoptic XE[®]), travoprost (Travatan[®], Travatan Z[®])
- Ophthalmic over-the-counter or prescription products, other than study treatment and the occasional use of artificial tears
- Oral carbonic anhydrase inhibitor

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

- High-dose salicylate therapy
- Topical or systemic corticosteroid
- Topical ophthalmic corticosteroid
- Intraocular corticosteroid implant
- Intravitreal or subtenon injection of ophthalmic corticosteroid
- Systemic beta-adrenergic blocking drug product
- Change in concurrent treatment or initiation of treatment with agents potentially affecting IOP, e.g., antihypertensive medication
- Contact lenses
- Ocular surgery

Subject's Responsibilities

You are requested to cooperate with site staff and maintain discipline during the course of the study. Please do not hide anything. This is for your own safety. You have to attend all the scheduled visits as explained in this document and by your doctor. You must follow all the instructions given by your doctor

- You are expected to cooperate with the attending doctor and you are expected to administer the medication as prescribed by the doctor
- You are also expected to follow the advice given to you regarding open-angle glaucoma or ocular hypertension by the doctor
- You have to tell the study doctor truthfully about your complete medical history
- You should report to the study site at the specified time given by the doctor/staff
- Take the study medication at about the same time every day
- You should administer the ophthalmic suspension as per the instruction given
- You should be able to comply with all study requirements
- You must follow the precautions related to the use of other medications.
- Do not use any other medication without the permission of the study doctor
- You have to report any new problems, illnesses, or changes in medication during the study
- You have to consult your physician before taking any other medications

Significance of Restrictions/Requirements in the Study

The above-mentioned restrictions must be followed to avoid the chances of getting false results in the study. Now as you are aware of the need for such restrictions/requirements, we request you comply with all these restrictions.

Risks or Discomforts

The study has been designed to be conducted on subjects who would otherwise be prescribed this drug for their illness in the dose prescribed. The study is planned to be conducted at the site under the supervision of a qualified investigator experienced in the treatment and management of such subjects. The study procedures described in the protocol are keeping in mind the subject's safety and wellbeing.

Any increase in risk to the study subject as a result of participation in the study is thus minimal. The study protocol would be approved by the ethics committee at each site ensuring the subject's safety, rights, and well-being is reviewed and approved by the ethics committee.

Since the active ingredient in the test product is similar to that of the reference product, it is expected that the test product, will have the same side effects as the originator product. The most common adverse reactions are blurred vision (care should be exercised in operating heavy machinery or driving a motor vehicle), bitter sour or unusual taste, blepharitis (inflammation of the eyelid), dermatitis (inflammation of the skin), dry eye foreign body sensation, headache, hyperemia (increase of blood flow to different tissues in the body), ocular discharge, ocular

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

discomfort, ocular keratitis, ocular pain, ocular pruritus (itch- is a sensation that causes the desire to scratch) and rhinitis (irritation and inflammation of the soft skin inside the nose), allergic reactions, hair loss, chest pain, conjunctivitis (redness, watering and pain in eye), diarrhea (loose and increased frequency of stools), diplopia (double vision), dizziness (spinning sensation in head), dry mouth, dyspnea (difficulty in breathing), dyspepsia (impaired digestion, recurrent pain in the upper abdomen, upper abdominal fullness), eye fatigue, hypertonia (increased muscle tone), kerato-conjunctivitis (inflammation of cornea and conjunctiva leading to redness, watering and pain in eye), keratopathy (any corneal disease, damage, dysfunction or abnormality), kidney pain (abdominal pain in kidney region), lid margin crusting or sticky sensation, nausea, pharyngitis, tearing and urticarial (itching of skin).

Risks during Pregnancy

The Study Medication may cause harm to an unborn child when administered to pregnant women. It is not known whether the ingredients of this cream can pass into breast milk.

Pregnant or nursing women cannot participate in this study. If you are a female of childbearing potential, you will be asked to have a pregnancy test, which must be negative to participate in this study. You should not be pregnant, planning pregnancy, or breastfeeding to participate in this study. You should discuss with your doctor the methods of effective contraception to avoid pregnancy. If you are planning to become pregnant during the period of this study, you should not agree to participate.

If you suspect pregnancy during the study, you must notify your study doctor immediately. You may be withdrawn from the study. You must not continue in the study if you become pregnant. Your study doctor will have additional instructions and information for you.

You must follow any of the methods of birth control mentioned below:

- Sexual abstinence: If you practice abstinence, then you have to practice a second method of birth control, as mentioned below if you become sexually active during the study participation.
- Oral contraceptives for at least 3 months before the start of the study. and remaining in place during the study period
- Intrauterine device (IUD) for at least 3 months before the start of the study and remaining in place during the study period
- Double-barrier method
- Transdermal, injection, or implants
- Non-hormonal or hormonal method
- Condom plus spermicide
- Diaphragm plus spermicide, or vaginal spermicidal suppository

The method should be followed for the duration of the study as judged by the doctor and agree to follow the same should be used during treatment. Female subjects must not change the method and/or medications of birth control during the course of the study.

For Male subjects: You should immediately inform your study doctor if your partner become pregnant during the study or after 7 days from the end of the treatment visit. Your partner will be asked to provide separate informed consent before collecting any information about pregnancy or a child.

Unforeseeable Risks

An allergic reaction is always a possibility associated with the usage of any drug and if any such reaction happens, we are well prepared to deal with it. As with any drug, unexpected side effects could occur and there is also the possibility of unexpected risks and serious adverse events, which could be severe and life-threatening.

Benefits to Subjects/Society/Community

You may or may not receive direct medical benefits from participating in this study. It is hoped that this study treatment may improve the treatment of chronic open-angle glaucoma or ocular hypertension; however, such a benefit cannot be guaranteed.

But society, in general, will benefit from this study because these studies help in introducing many formulations of the same drug on the market.

It is also possible that this study may contribute to the development of Brinzolamide 1% ophthalmic suspension for open-angle glaucoma or ocular hypertension and the drug might be available with lower costs than the currently available treatments.

The information collected from this study may help other people who suffer from open-angle glaucoma or ocular hypertension in the future.

Alternative Procedures or Any Other Treatments

If you do not wish to participate in this study, your doctor will prescribe other medications available in India for the treatment of your disease. If you do not participate in this study, it will not affect your relationship with your doctor or any of the staff; and it will not affect the care or concern you are currently receiving from them.

Your doctor will discuss other treatment options with you including the potential benefits and risks.

Confidentiality of Records

Your records of being in this study will be kept private except when ordered by law. The following people will have access to your study records:

- Study personnel
- Sponsor company or research institution or authorized designee
- Drugs Controller General of India
- Other country, state, or federal regulatory agencies
- Ethics Committee.

The Ethics Committee (EC), Regulatory Authorities, and accrediting agencies may inspect and copy your records, which may have your name on them. Therefore, your total privacy cannot be guaranteed. In case of publication of the study results, your identity will, in no case be made public.

Insurance and Compensation to the Subject

For participants in this study, there is appropriate insurance coverage. This insurance covers damages or injury or death resulting from participation in a study and which are mentioned during the participation in the study. Any injury or damages or death will be informed to the insurer.

You should report any discomforts, problems, or research-related injuries immediately to your study doctor or study staff.

In the case of an injury occurring to you during the clinical study, you will be given free medical management as long as required or till such time it is established that the injury is not related to the clinical study, whichever is earlier.

In the case of study-related injury or death, the sponsor or sponsor's representative will provide you with complete medical care along with compensation for the injury or death. In case there is no permanent injury, the quantum of compensation shall be commensurate with the nature of the non-permanent injury and loss of wages. In case of death, your nominee will be provided with appropriate compensation. The financial compensation will be over and above any expenses incurred on your medical management.

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

The amount of compensation will be decided by the regulatory authority. The independent expert committee (constituted by the regulatory authority in case of death) and the ethics committee will provide their opinion on the compensation to be provided.

You or your nominee or legal representative can also contact the sponsor to make claims in the case of study-related injury or death.

Sponsor's Name	:	
Sponsor's Address	:	
Sponsor Representative's Name	:	
Sponsor Representative's Address	:	

You are not waiving off your legal rights by signing this form.

Voluntary Nature of Participation

Please remember that your participation in this study is entirely voluntary and this document is presented to you to help you in making an informed decision. If you have any doubts regarding the study, you can ask questions of any of our study staff and clarify them to your utmost satisfaction.

Refusal to participate in this study will involve no penalty or loss of benefits, to which you might otherwise be entitled.

Your Participation in the Study and Your Rights

Your participation in this study is voluntary and you can withdraw from the study at any time. In that case, you will be given proper treatment for your condition. If at any time you feel worse or suffer from any other illness, then please inform your attending doctor. If the treatment appears to be unsuitable for you, it will be stopped. It is possible that the study could be stopped without your consent. We will inform you if any new information becomes known during the study, which may affect your willingness to continue the study.

Subject Withdrawal:

Subjects will be advised that they are free to withdraw from the study at any time for any reason or, if necessary, the Investigator may withdraw a subject from the study to protect the health of that subject. A subject may also be withdrawn for not complying with study procedures. The clinical report will include all reasons for early withdrawals.

All subjects who are randomized and received Study Medication will be included in the safety analysis comparing the Test/Reference. If a randomized subject terminates from the study early, all efforts will be made to complete the end-of-study procedures. In case of early termination, the Investigator shall fully document the reason for early termination. Reasons for the early termination may include the following:

- If the subject withdraws his or her consent for any reason
- The subject's condition has worsened to the degree that the investigator feels it is unsafe for the subject to continue in the study
- The subject's condition has worsened to the degree that the investigator feels it is unsafe for the subject to continue in the study and requires alternate or supplemental therapy (e.g., IOP $\geq$ 36 mm Hg in either eye)
- If the subject misses more than 03 consecutive days of Investigational Product instillation
- If the subject's treatment is un-masked
- If an adverse event occurs for which the subject desires to discontinue treatment or the investigator determines that it is in the subject's best interest to be discontinued

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

- If there is a significant major protocol deviation (such as the use of prohibited medication during the study conduct)
- If concomitant therapy is reported or required which is liable to interfere with the results of the study
- If the subject is lost to follow-up. (The investigator shall document attempts to reach the subject by telephone in source documents before considering that the subject lost to follow-up)
- If the female subject becomes pregnant.

You will be requested to return the used and unused medication along with your diary. Even when you are terminated from participating in the study, if you require any medical care it will be provided. If any time, the doctor identifies that the infection has become worse, is not responding to treatment, or that the study treatment is not sufficient to treat the present condition, your doctor removes you from the study and prescribes appropriate treatment or refers you to another physician.

Termination of the Study

Under certain circumstances, it is possible that your study doctor or sponsor of this study will decide that you should be withdrawn prematurely from the clinical study without your prior consent. This may happen if your participation in the clinical study is no longer acceptable for medical reasons; or if it is decided to discontinue the whole clinical study. But the same will be intimated to you for information.

The study can be terminated for the following reasons

1. The doctor and/or sponsor feel that the number and/or the severity of the adverse events justify the discontinuation of the study.
2. Data not known before become available and raised a concern about the safety of the Study Medication so that continuation would pose potential risks to the subjects.
3. The Ethics Committee and/or regulatory authority may terminate the study if the continuation of the study compromises subject safety.

When the study is terminated prematurely if the study subjects need any medical care it will be provided.

Financial Reimbursement

You will not be compensated for any loss of wages for the days of your participation.

All study products, site visits, and procedures related to this study will be provided to you at no cost. You will not be paid for your participation in this study. However, during each visit to the site, the traveling charges if any, incurred to you will be reimbursed on an actual basis and it will be a maximum of up to Rs.500/- per visit.

New Findings

During the course of this study, new information regarding the Study Medications or the risks and benefits of the study may become known to the study doctors. If this occurs, the doctor will inform you about this new information in a timely manner. Your willingness to participate in this research study is voluntary and you may withdraw from the study at any time.

The new information may mean that you can no longer participate in the research, or that you may not want to continue in the research study. If this occurs, the study doctor will stop your participation in the study and you will be offered all available care to suit your needs and medical conditions.

Additional Information during the COVID-19

This is to inform you that the novel coronavirus, COVID-19, has been declared a worldwide pandemic by the World Health Organization. COVID-19 is extremely contagious and is

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

believed to spread primarily through droplets of saliva or discharge from the nose when an infected person coughs or sneezes and also by person-to-person contact with a COVID subject. COVID-19 affects different people in different ways. Most infected people will develop mild to moderate illness and recover without hospitalization.

On average it takes 5-6 days from when someone is infected with the virus for symptoms to show, however, it can take up to 14 days.

You may be exposed to an infection that causes COVID-19 during your visits to the clinical site when coming into contact with the infected person. All procedures and precautions will be taken by the study staff to ensure that the risk of this exposure is minimized during your visits to the clinical site. In accordance with local Government guidelines, the clinical site has established additional rules and precautionary measures to prevent the spread of the infection and limit physical contact at the hospital.

Subject Responsibilities:

As a subject in this study, you have certain responsibilities listed below to ensure your safety and the safety of others during your visit to the clinical site.

- Maintain at least two meters distance between you and people coughing or sneezing
- Avoid social gatherings and unnecessary movement outside
- Wash your hands regularly with soap and water at least for 20 seconds or clean your hand with alcohol-based hand rubs or sanitiser
- Wear a face mask always
- Cover your mouth and nose when coughing or sneezing
- Avoid touching your face
- Do not share items, such as food, pens, etc.,
- Follow the instruction of the study coordinator for specific study visit date and timing
- Report any symptoms of being sick immediately to the study staff
- Do not attend any visits if you are sick, living with someone who is sick, have been potentially exposed to COVID-19, or living with anyone who has travelled outside the state/country
- Follow the Government rules and hospital regulations specifically for COVID-19.

Others:

In case of a serious adverse event, the physician has the authority to provide emergency medical care to ensure your well-being without regard to the provisions of this document.

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

Key Study Contacts:

You may ask questions that you have about the study at any time. If you later have a new question or in the case of research-related injury, you can always call your study doctor. (For Study-related injury and Information)

Investigator (Name and Address):

Contact Number: _____

If you have any questions, comments, or concerns about the research, you can talk to the sponsor

Ethics committee:

If you have questions about your rights as a research subject, or you have concerns or suggestions and you want to talk to someone other than the researchers, you may contact;

Chairman/Chairperson (Name and Address):

Contact Number: _____

A description of this clinical study will be available on www.ctri.nic.in/ www.clinicaltrials.gov

This website will not include information that can identify you. At most, the website will include a summary of the results. You can search this website at any time and the information will be in English. You can request information from the study staff at any time and have access to data that is publicly available.

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

INFORMED CONSENT FORM

Study Title A Randomized, Double-masked, Active Controlled, Parallel Group, Multicentre Bioequivalence Study with Clinical Endpoint Comparing Brinzolamide 1% Ophthalmic Suspension (Sponsor), to the reference product (Brinzolamide) 1% Ophthalmic Suspension of (the originator manufacturer) in the Treatment of Chronic Open Angle Glaucoma or Ocular Hypertension in Both Eyes.

Protocol Number Sample/2026/01

Subject Number:

Subject Initials:

Name of the Subject:

Date of Birth:

Age of the Subject:

Address of the Subject:

Occupation of the Subject: Student/Self-employed/Service/Housewife/Others (Please tick as appropriate)

Qualification of the Subject:

Annual Income of the Subject:

Name of the Nominee:

(For the purpose of compensation in case of study related Death)

Address of the Nominee:

Relationship of the Nominee with the Subject:

SUBJECT INFORMATION SHEET AND INFORMED CONSENT FORM

Declaration by the Subject	Initial in the box
i. I confirm that I have read and understood the Informed Consent Document (Subject information sheet and Informed consent form) Version (version number), dated (date) for the above study, and have had the opportunity to ask questions.	
ii. I understand that my participation in the study is voluntary and that I am free to withdraw at any time, without giving any reason, and without my medical care or legal rights being affected.	
iii. I understand that the Sponsor of the clinical study, others working on the Sponsor's behalf, the Ethics Committee, and the regulatory authorities will not need my permission to look at my health records both in respect of the current study and any further research that may be conducted in relation to it, even if I withdraw from the study. I agree to this access. However, I understand that my identity will not be revealed in any information released to third parties or published.	
iv. I agree not to restrict the use of any data or results that arise from this study provided such a use is only for the scientific purpose(s).	
v. I agree to take part in the above study.	
vi. I understand that there is a risk of COVID-19 virus infection during the study even with the additional precautions in place. I also understand my responsibilities as a participant to help minimize the risk of infection	

Name of the Subject	Signature of Subject	Date

Name of the Impartial Witness	Signature of Impartial Witness	Date

Principal Investigator/ Designee's Name	Principal Investigator/ Designee's Signature	Date

(Copy of the Subject Information Sheet and duly filled Informed Consent Form shall be handed over to the Subject)