

Composition Lemma for Lean4

Edinah Gnang

Parikshit Chalise

Contents

6	Chapter 1. Composition Lemma	5
7	1.1. Overview	5
8	1.2. Functional Directed Graphs	5
9	1.3. Quotient-Remainder Theorem and Lagrange Interpolation	6
10	1.4. The Composition Lemma	14
11	Bibliography	23

Composition Lemma

1.1. Overview

The *Composition Lemma* was developed and refined over 6 years, beginning in 2018, as a novel approach to settle in the affirmative the *Graceful Tree Conjecture*. The first of such papers was posted in [3] by Gngang. A further developed series of papers resolving the same conjecture again appeared in [4] and [5]. Recently, the same method has been applied to settle other longstanding conjectures in [1] and [2]. We comment that the series of papers shared on the open-source platform arXiv reflect the evolving landscape of Gngang's thought process, and the frequent re-uploads were driven by the natural progression and refinement of ideas. However, we recognize that these numerous edits may have unintentionally caused confusion and raised questions regarding the success of the method. In the current work, we aim to address these concerns by presenting a detailed blueprint of the proof, with the goal of formalizing it in Lean4.

1.2. Functional Directed Graphs

For notational convenience, let \mathbb{Z}_n denote the set whose members are the first n natural numbers, i.e.,

$$(1.2.1) \quad \mathbb{Z}_n := \{0, \dots, n-1\}.$$

For a function $f : \mathbb{Z}_m \rightarrow \mathbb{Z}_n$, we write $f \in \mathbb{Z}_n^{\mathbb{Z}_m}$. For $X \subseteq \mathbb{Z}_m$, $f(X)$ denotes the image of X under f , i.e.,

$$(1.2.2) \quad f(X) = \{f(i) : i \in X\},$$

and $|f(X)|$ denotes its cardinality. For $Y \subseteq \mathbb{Z}_n$, $f^{-1}(Y)$ denotes the pre-image of Y under f i.e.

$$(1.2.3) \quad f^{-1}(Y) = \{j \in \mathbb{Z}_m : f(j) \in Y\}$$

DEFINITION 1.2.4 (Functional digraphs). For an arbitrary $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$, the *functional directed graph* prescribed by f , denoted G_f , is such that the vertex set $V(G_f)$ and the directed edge set $E(G_f)$ are respectively as follows:

$$V(G_f) = \mathbb{Z}_n, \quad E(G_f) = \{(v, f(v)) : v \in \mathbb{Z}_n\}.$$

DEFINITION 1.2.5 (Graceful functional digraphs). The functional directed graph prescribed by $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ is graceful if there exist a bijection $\sigma \in S_n \subset \mathbb{Z}_n^{\mathbb{Z}_n}$ such that

$$(1.2.6) \quad \{|\sigma f \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\} = \mathbb{Z}_n.$$

If $\sigma = \text{id}$ (the identity function), then G_f — the functional directed graph prescribed by f — is gracefully labeled.

DEFINITION 1.2.7 (Automorphism group). For a functional directed graph G_f , its automorphism group, denoted $\text{Aut}(G_f)$, is defined as follows:

$$\text{Aut}(G_f) = \{\sigma \in S_n : \{(i, f(i)) : i \in \mathbb{Z}_n\} = \{(j, \sigma f \sigma^{-1}(j)) : j \in \mathbb{Z}_n\}\}.$$

For a polynomial $P \in \mathbb{C}[x_0, \dots, x_{n-1}]$, its automorphism group, denoted $\text{Aut}(P)$, is defined as follows:

$$\text{Aut}(P) = \{\sigma \in S_n : P(x_0, \dots, x_i, \dots, x_{n-1}) = P(x_{\sigma(0)}, \dots, x_{\sigma(i)}, \dots, x_{\sigma(n-1)})\}.$$

DEFINITION 1.2.8 (Graceful re-labelings). The set of distinct gracefully labeled functional directed graphs isomorphic to G_f is

$$\text{GrL}(G_f) := \left\{ G_{\sigma f \sigma^{-1}} : \begin{array}{l} \sigma \text{ is a representative of a coset in } S_n / \text{Aut}(G_f) \text{ and} \\ \mathbb{Z}_n = \{|\sigma f \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\} \end{array} \right\}$$

DEFINITION 1.2.9 (Complementary labeling involution). If $\varphi = n - 1 - \text{id}$, i.e. $\varphi \in \mathbb{Z}_n^{\mathbb{Z}_n}$ such that

$$\varphi(i) = n - 1 - i, \forall i \in \mathbb{Z}_n,$$

then for an arbitrary $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ the complementary labeling involution is defined as the map

$$f \mapsto \varphi f \varphi^{-1}$$

Observe that for all $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ the complementary labeling involution fixes the induced edge label of each edge as seen from the equality

$$(1.2.10) \quad |f(i) - i| = |\varphi f(i) - \varphi(i)|, \quad \forall i \in \mathbb{Z}_n.$$

In other words, induced edge labels are fixed by the vertex relabeling effected by φ . We call this induced edge label symmetry the *complementary labeling symmetry* of the functional directed graph G_f .

1.3. Quotient-Remainder Theorem and Lagrange Interpolation

PROPOSITION 1.3.1 (Multivariate Quotient-Remainder). Let $d(x) \in \mathbb{C}[x]$ be a degree n monic polynomial with simple roots, i.e.,

$$(1.3.2) \quad d(x) = \prod_{i \in \mathbb{Z}_n} (x - \alpha_i) \quad \text{and} \quad 0 \neq \prod_{0 \leq u < v < n} (\alpha_v - \alpha_u),$$

where $\{\alpha_u : u \in \mathbb{Z}_n\} \subset \mathbb{C}$. For all $P \in \mathbb{C}[x_0, \dots, x_{m-1}]$, there exists a unique remainder $r(x_0, \dots, x_{m-1}) \in \mathbb{C}[x_0, \dots, x_{m-1}]$ of degree at most $n - 1$ in each variable such that for quotients: $\{q_k(x_0, \dots, x_{n-1}) : k \in \mathbb{Z}_n\} \subset \mathbb{C}[x_0, \dots, x_{n-1}]$, we have

$$(1.3.3) \quad P(x_0, \dots, x_{m-1}) = r(x_0, \dots, x_{m-1}) + \sum_{u \in \mathbb{Z}_m} q_u(x_0, \dots, x_{m-1}) d(x_u).$$

PROOF. We prove by induction on the number of variables that the remainder admits the expansion

$$(1.3.4) \quad r(x_0, \dots, x_{m-1}) = \sum_{g \in \mathbb{Z}_n^m} P(\alpha_g) \prod_{i \in \mathbb{Z}_m} \left(\prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right) \right),$$

where for notational convenience $P(\alpha_g) := P(\alpha_{g(0)}, \dots, \alpha_{g(m-1)})$. The base case stems from the univariate quotient-remainder theorem over the field \mathbb{C} . The univariate-quotient remainder theorem over the field \mathbb{C} asserts that there exist a unique quotient-remainder pair $(q(x_0), r(x_0)) \in \mathbb{C}[x_0] \times \mathbb{C}[x_0]$ subject to

$$(1.3.5) \quad H(x_0) = q(x_0) d(x_0) + r(x_0),$$

where $r(x_0) \in \mathbb{C}[x_0]$ is of degree at most $n - 1$. It is completely determined by its evaluation over $\{\alpha_i : i \in \mathbb{Z}_n\}$, and by Lagrange interpolation we have

$$(1.3.6) \quad r(x_0) = \sum_{g \in \mathbb{Z}_n^1} H(\alpha_{g(0)}) \prod_{j_0 \in \mathbb{Z}_n \setminus \{g(0)\}} \left(\frac{x_0 - \alpha_{j_0}}{\alpha_{g(0)} - \alpha_{j_0}} \right),$$

thus establishing the claim in the base case. For the induction step, assume as our induction hypothesis that for all $F \in \mathbb{C}[x_0, \dots, x_{m-1}]$, we have

$$(1.3.7) \quad F = \sum_{k \in \mathbb{Z}_m} q_k(x_0, \dots, x_{m-1}) d(x_k) + \sum_{g \in \mathbb{Z}_n^m} F(\alpha_g) \prod_{i \in \mathbb{Z}_m} \left(\prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right) \right).$$

We proceed to show that the hypothesis implies that every polynomial in $m + 1$ variables also admits a similar expansion, thus establishing the desired claim. Consider a polynomial $H \in \mathbb{C}[x_0, \dots, x_m]$. We view H as a univariate polynomial in the variable x_m whose coefficients lie in the field of fraction $\mathbb{C}(x_0, \dots, x_{m-1})$. The univariate quotient-remainder theorem over the field of fractions $\mathbb{C}(x_0, \dots, x_{m-1})$ asserts that there exist a unique quotient-remainder pair

$$(q(x_m), r(x_m)) \in (\mathbb{C}(x_0, \dots, x_{m-1}))[x_m] \times (\mathbb{C}(x_0, \dots, x_{m-1}))[x_m]$$

subject to

$$(1.3.8) \quad H(x_0, \dots, x_m) = q(x_0, \dots, x_m) d(x_m) + r(x_0, \dots, x_m),$$

where $r(x_0, \dots, x_m) \in (\mathbb{C}(x_0, \dots, x_{m-1}))[x_m]$ is of degree at most $n-1$ in the variable x_m . We write

$$(1.3.9) \quad r(x_0, \dots, x_m) = \sum_{k \in \mathbb{Z}_n} a_k(x_0, \dots, x_{m-1}) (x_m)^k.$$

We now show that coefficients $\{a_k(x_0, \dots, x_{m-1}) : k \in \mathbb{Z}_n\}$ all lie in the polynomial ring $\mathbb{C}[x_0, \dots, x_{m-1}]$ via the equality

$$(1.3.10) \quad \left(\text{Vander} \begin{pmatrix} \alpha_0 \\ \vdots \\ \alpha_u \\ \vdots \\ \alpha_{n-1} \end{pmatrix} \right) \cdot \begin{pmatrix} a_0(x_0, \dots, x_{m-1}) \\ \vdots \\ a_u(x_0, \dots, x_{m-1}) \\ \vdots \\ a_{n-1}(x_0, \dots, x_{m-1}) \end{pmatrix} = \begin{pmatrix} H(x_0, \dots, x_{m-1}, \alpha_0) \\ \vdots \\ H(x_0, \dots, x_{m-1}, \alpha_u) \\ \vdots \\ H(x_0, \dots, x_{m-1}, \alpha_{n-1}) \end{pmatrix},$$

where

$$(1.3.11) \quad \left(\text{Vander} \begin{pmatrix} \alpha_0 \\ \vdots \\ \alpha_u \\ \vdots \\ \alpha_u \end{pmatrix} \right) [i, j] = (\alpha_i)^j, \quad \forall 0 \leq i, j < n.$$

Since the Vandermonde matrix is invertible by the fact

$$(1.3.12) \quad 0 \neq \det \left(\text{Vander} \begin{pmatrix} \alpha_0 \\ \vdots \\ \alpha_u \\ \vdots \\ \alpha_u \end{pmatrix} \right) = \prod_{0 \leq u < v < n} (\alpha_v - \alpha_u),$$

we indeed have

$$(1.3.13) \quad \begin{pmatrix} a_0(x_0, \dots, x_{m-1}) \\ \vdots \\ a_u(x_0, \dots, x_{m-1}) \\ \vdots \\ a_{n-1}(x_0, \dots, x_{m-1}) \end{pmatrix} = \left(\text{Vander} \begin{pmatrix} \alpha_0 \\ \vdots \\ \alpha_u \\ \vdots \\ \alpha_u \end{pmatrix} \right)^{-1} \cdot \begin{pmatrix} H(x_0, \dots, x_{m-1}, \alpha_0) \\ \vdots \\ H(x_0, \dots, x_{m-1}, \alpha_u) \\ \vdots \\ H(x_0, \dots, x_{m-1}, \alpha_{n-1}) \end{pmatrix}.$$

Therefore, we have

$$(1.3.14) \quad H(x_0, \dots, x_m) = q_m(x_0, \dots, x_m) d(x_m) + \sum_{g(m) \in \mathbb{Z}_n} H(x_0, \dots, x_{m-1}, \alpha_{g(m)}) \prod_{j \in \mathbb{Z}_n \setminus \{g(m)\}} \left(\frac{x_m - \alpha_{j_m}}{\alpha_{g(m)} - \alpha_{j_m}} \right).$$

Applying the induction hypothesis to coefficients

$$\{H(x_0, \dots, x_{m-1}, \alpha_{g(m)}) : \alpha_{g(m)} \in \mathbb{C}\} \subset \mathbb{C}[x_0, \dots, x_{m-1}]$$

yields the desired expansion. Finally, quotients $\{q_k(x_0, \dots, x_{m-1}) : k \in \mathbb{Z}_m\}$ lie in the polynomial ring $\mathbb{C}[x_0, \dots, x_{m-1}]$ since the polynomial $H(x_0, \dots, x_{m-1}) - r(x_0, \dots, x_{m-1})$ lies in the ideal generated by members of the set $\{d(x_u) : u \in \mathbb{Z}_m\}$. \square

PROPOSITION 1.3.15 (Ring Homomorphism). *For an arbitrary $H \in \mathbb{C}[x_0, \dots, x_{n-1}]$, let \overline{H} denote the remainder of the congruence class*

$$H \mod \{d(x_i) : i \in \mathbb{Z}_n\},$$

where

$$d(x) = \prod_{i \in \mathbb{Z}_n} (x - \alpha_i) \quad \text{and} \quad 0 \neq \prod_{0 \leq u < v < n} (\alpha_v - \alpha_u),$$

Then the following hold:

- (i) For all $g \in \mathbb{Z}_n^{\mathbb{Z}_n}$, we have $\overline{H}(\alpha_g) = H(\alpha_g)$.
- (ii) If $H = H_0 + H_1$, where $H_0, H_1 \in \mathbb{C}[x_0, \dots, x_{n-1}]$, then $\overline{H_0} + \overline{H_1} = \overline{H}$.
- (iii) If $H = H_0 \cdot H_1$, where $H_0, H_1 \in \mathbb{C}[x_0, \dots, x_{n-1}]$, then $\overline{H} \equiv \overline{H_0} \cdot \overline{H_1}$.

PROOF. The first claim follows from Proposition 1.3.1 for we see that the divisor vanishes over the lattice. To prove the second claim we recall that

$$\begin{aligned} \overline{H} &= \sum_{g \in \mathbb{Z}_n^{\mathbb{Z}_n}} H(\alpha_g) \prod_{i \in \mathbb{Z}_n} \left(\prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right) \right), \\ \Rightarrow \overline{H} &= \sum_{g \in \mathbb{Z}_n^{\mathbb{Z}_n}} (H_0(\alpha_g) + H_1(\alpha_g)) \prod_{i \in \mathbb{Z}_n} \left(\prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right) \right), \\ \Rightarrow \overline{H} &= \sum_{k \in \mathbb{Z}_2} \sum_{g \in \mathbb{Z}_n^{\mathbb{Z}_n}} H_k(\alpha_g) \prod_{i \in \mathbb{Z}_n} \left(\prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right) \right). \end{aligned}$$

Thus $\overline{H_0} + \overline{H_1} = \overline{H}$ as claimed. Finally the fact (iii) is a straightforward consequence of Proposition 1.3.16, which is proved next. \square

For notational convenience, we denote by $L_g(\mathbf{x})$ the Lagrange basis polynomial associated with $g \in \mathbb{Z}_n^{\mathbb{Z}_n}$

$$L_g(\mathbf{x}) := \prod_{i \in \mathbb{Z}_n} \left(\prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right) \right)$$

and for any $S \subseteq \mathbb{Z}_n$ we denote by $L_g(\mathbf{x}|_S)$ the factor of $L_g(\mathbf{x})$ which only features variables indexed by S as follows

$$L_g(\mathbf{x}|_S) := \prod_{i \in S} \left(\prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right) \right)$$

PROPOSITION 1.3.16. *Let $f, g \in \mathbb{Z}_n^{\mathbb{Z}_n}$. For congruence classes modulo the ideal generated by $\{d(x_i) : i \in \mathbb{Z}_n\}$, we have*

$$L_f(\mathbf{x}) \cdot L_g(\mathbf{x}) \equiv \begin{cases} L_f(\mathbf{x}) & \text{if } f = g \\ 0 & \text{otherwise,} \end{cases}$$

PROOF. Observe that

$$L_f(\mathbf{x}) \cdot L_g(\mathbf{x}) = \prod_{i \in \mathbb{Z}_n} \left(\left(c_{i,f} \frac{d(x_i)}{x_i - \alpha_{f(i)}} \right) \left(c_{i,g} \frac{d(x_i)}{x_i - \alpha_{g(i)}} \right) \right),$$

where

$$c_{i,f} = \prod_{j_i \in \mathbb{Z}_n \setminus \{f(i)\}} (\alpha_{f(i)} - \alpha_{j_i})^{-1} \quad \text{and} \quad c_{i,g} = \prod_{j_i \in \mathbb{Z}_n \setminus \{g(i)\}} (\alpha_{g(i)} - \alpha_{j_i})^{-1}.$$

If $f \neq g$, then there exists $j \in \mathbb{Z}_n$ such that $f(j) \neq g(j)$ and $L_f(\mathbf{x}) \cdot L_g(\mathbf{x})$ is a multiple of $(x_j)^n$, as a result of which we obtain $L_f(\mathbf{x}) \cdot L_g(\mathbf{x}) \equiv 0$. Alternatively if $f = g$, then

$$L_f(\mathbf{x}) \cdot L_g(\mathbf{x}) = (L_f(\mathbf{x}))^2 = L_f(\mathbf{x}) + \left((L_f(\mathbf{x}))^2 - L_f(\mathbf{x}) \right).$$

We now show that $(L_f(\mathbf{x}))^2 - L_f(\mathbf{x}) \equiv 0$

$$\begin{aligned} (L_f(\mathbf{x}))^2 - L_f(\mathbf{x}) &= L_f(\mathbf{x}) (L_f(\mathbf{x}) - 1) \\ &= L_f(\mathbf{x}) \left(L_f(\mathbf{x}) - \sum_{g \in \mathbb{Z}_n^{\mathbb{Z}_n}} L_g(\mathbf{x}) \right) \\ &= -L_f(\mathbf{x}) \left(\sum_{g \in \mathbb{Z}_n^{\mathbb{Z}_n} \setminus \{f\}} L_g(\mathbf{x}) \right) \\ &\equiv 0, \end{aligned}$$

where the latter congruence identity stems from the prior setting where $f \neq g$. \square

LEMMA 1.3.17 (Symmetrization Lemma). Let $P, F \in \mathbb{Q}[x_0, \dots, x_{n-1}]$ and $\mathcal{S}_u \subseteq \mathbb{Z}_n$ for all $u \in \mathbb{Z}_m$. For positive integers $\{D_u : u \in \mathbb{Z}_m\}$ let

$$(1.3.18) \quad P(\mathbf{x}) = F(\mathbf{x}) \sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (x_i)^{g(i)} \right),$$

such that for some integer κ we have $F(\alpha_{f\tau^{-1}}) = (-1)^\kappa F(\alpha_f)$ for all $f \in \mathbb{Z}_n^{\mathbb{Z}_m}$, where τ denotes a transposition which exchanges a fixed pair of variables. If $F(\alpha_f)$ vanishes for all $f \in \mathbb{Z}_n^{\mathbb{Z}_m} \setminus S_n$, then

$$\tau \in \text{Aut}(\overline{P}(\mathbf{x})) \implies \tau \in \text{Aut} \left(\sum_{\sigma \in S_n} F(\alpha_\sigma) \sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) \right)$$

PROOF. Observe that

$$\sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (x_i)^{g(i)} \right) \equiv \sum_{j \in \mathbb{Z}_m} \sum_{h_j \in \mathbb{Z}_n^{s_j}} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{h_j(i)})^{g(i)} \right) L_{h_j}(\mathbf{x}|_{\mathcal{S}_j}).$$

By our premise $\overline{F}(\alpha_{f\tau^{-1}}) = -\overline{F}(\alpha_f)$ for all $f \in \mathbb{Z}_n^{\mathbb{Z}_m}$ or alternatively $\overline{F}(\alpha_{f\tau^{-1}}) = \overline{F}(\alpha_f)$ for all $f \in \mathbb{Z}_n^{\mathbb{Z}_m}$. Furthermore by the premise that $F(\alpha_f)$ vanishes for all $f \in \mathbb{Z}_n^{\mathbb{Z}_m} \setminus S_n$,

$$\begin{aligned} \overline{P} &\equiv \overline{F} \sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (x_i)^{g(i)} \right) \\ &\implies \overline{P} \equiv \left(\sum_{\sigma \in S_n} F(\alpha_\sigma) L_\sigma(\mathbf{x}) \right) \sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (x_i)^{g(i)} \right) \\ &\equiv \left(\sum_{\sigma \in S_n} F(\alpha_\sigma) L_\sigma(\mathbf{x}) \right) \sum_{j \in \mathbb{Z}_m} \sum_{h_j \in \mathbb{Z}_n^{s_j}} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{h_j(i)})^{g(i)} \right) L_{h_j}(\mathbf{x}|_{\mathcal{S}_j}) \\ &\implies \left(\sum_{\sigma \in S_n} F(\alpha_\sigma) L_\sigma(\mathbf{x}) \right) \sum_{j \in \mathbb{Z}_m} \sum_{h_j \in \mathbb{Z}_n^{s_j}} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{h_j(i)})^{g(i)} \right) L_{h_j}(\mathbf{x}|_{\mathcal{S}_j}) = \\ &\sum_{j \in \mathbb{Z}_m} \sum_{(\sigma, h_j) \in S_n \times \mathbb{Z}_n^{s_j}} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{h_j(i)})^{g(i)} \right) F(\alpha_\sigma) L_{h_j}(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\overline{\mathcal{S}_j}}), \end{aligned}$$

where $\overline{\mathcal{S}_j} := \mathbb{Z}_n \setminus \mathcal{S}_j$. Recall that $L_\sigma(\mathbf{x})$ has a trivial automorphism group for all $\sigma \in S_n$. Each summand in the sum above is either such that

$$\left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) F(\alpha_\sigma) L_{h_j}(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\overline{\mathcal{S}_j}}) \equiv 0$$

in which case the said summand imposes no constraints at all on the membership of any permutation into the automorphism group of \overline{P} . Thus such summands play no role in ascertaining $\text{Aut}(\overline{P})$. Alternatively a summand

$$\left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) F(\alpha_\sigma) L_{h_j}(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\overline{\mathcal{S}_j}}) \neq 0$$

is a member of an orbit

$$\left\{ \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{s_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) F(\alpha_\sigma) \underbrace{L_{h_j}(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|_{\overline{\mathcal{S}_j}})}_{\equiv L_\sigma(\mathbf{x})} \neq 0 : \begin{array}{l} \sigma \in S_n \\ \text{s.t.} \\ \sigma|_{\mathcal{S}_j} = h_j \end{array} \right\}$$

which impose constraints on membership of permutations into the automorphism group of \overline{P} . Therefore the remainder of such summand must be accounted for in order to ascertain $\text{Aut}(\overline{P})$. If τ lies in the automorphism group of \overline{P} then there are

two cases to consider for each variable indexing set \mathcal{S}_j . In the first case both variables interchanged by the transposition τ either lie in \mathcal{S}_j or both lie in its complement $\overline{\mathcal{S}_j}$. In this setting the orbit

$$\left\{ \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{\mathcal{S}_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) F(\alpha_\sigma) L_\sigma(\mathbf{x}) : \begin{array}{l} \sigma \in S_n \\ \text{s.t.} \\ \sigma|_{\mathcal{S}_j} = h_j \end{array} \right\}$$

is up to signed fixed by the transposition τ . Otherwise in the setting where only one of the variables exchanged by τ lies in \mathcal{S}_j , τ induces a transposition of the orbit of summands

$$\left\{ \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{\mathcal{S}_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) F(\alpha_\sigma) L_\sigma(\mathbf{x}) : \begin{array}{l} \sigma \in S_n \\ \text{s.t.} \\ \sigma|_{\mathcal{S}_j} = h_j \end{array} \right\}$$

with some other orbit of summands

$$\left\{ \left(\sum_{g \in (1+\mathbb{Z}_{D_k})^{\mathcal{S}_k}} c_{k,g} \prod_{i \in \mathcal{S}_k} (\alpha_{\gamma(i)})^{g(i)} \right) F(\alpha_\gamma) L_\gamma(\mathbf{x}) : \begin{array}{l} \gamma \in S_n \\ \text{s.t.} \\ \gamma|_{\mathcal{S}_k} = h_k \end{array} \right\}$$

for we see that if that $F(\alpha_{f\tau^{-1}}) = -F(\alpha_f)$ for all $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ or alternatively $F(\alpha_{f\tau^{-1}}) = F(\alpha_f)$ for all $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$, then the orbit summands

$$\left\{ \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{\mathcal{S}_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) F(\alpha_\sigma) L_\sigma(\mathbf{x}) : \begin{array}{l} \sigma \in S_n \\ \text{s.t.} \\ \sigma|_{\mathcal{S}_j} = h_j \end{array} \right\}$$

is not up to sign fixed by τ . Thus pairing by this interchange \mathcal{S}_j with \mathcal{S}_k and vice versa.

I do not see what was missing and is fixed here in the argument by requiring each \mathcal{S}_j to be distinct as we have done, by requiring degree to be greater than 0 for each variable.

In other words $\tau(\mathcal{S}_j) = \mathcal{S}_k$. Observe that

$$(1.3.19) \quad \overline{P}(\mathbf{x}) = \sum_{\sigma \in S_n} F(\alpha_\sigma) \sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{\mathcal{S}_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|\overline{\mathcal{S}_j})$$

and the associated symmetrized polynomial is

$$\sum_{\sigma \in S_n} F(\alpha_\sigma) \sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{\mathcal{S}_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) \sum_{h \in \mathbb{Z}_n^{\overline{\mathcal{S}_j}}} L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_h(\mathbf{x}|\overline{\mathcal{S}_j}).$$

Note that to obtain the symmetrized polynomial, for each $\sigma \in S_n$, in equation (1.3.19) within the j -th summand for each $j \in \mathbb{Z}_m$, we have replaced the Lagrange basis

$$L_\sigma(\mathbf{x}) = L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_\sigma(\mathbf{x}|\overline{\mathcal{S}_j})$$

with the sum over Lagrange bases polynomials

$$\sum_{f \in \mathbb{Z}_n^{\overline{\mathcal{S}_j}}} L_\sigma(\mathbf{x}|_{\mathcal{S}_j}) L_f(\mathbf{x}|\overline{\mathcal{S}_j}).$$

These sums over Lagrange bases are as block subject to the same pairing argument described above. Thus τ also fixes the symmetrized polynomial. By properties of Lagrange basis polynomials it follows that the symmetrized polynomial equals

$$\sum_{\sigma \in S_n} F(\alpha_\sigma) \sum_{j \in \mathbb{Z}_m} \left(\sum_{g \in (1+\mathbb{Z}_{D_j})^{\mathcal{S}_j}} c_{j,g} \prod_{i \in \mathcal{S}_j} (\alpha_{\sigma(i)})^{g(i)} \right) L_\sigma(\mathbf{x}|_{\mathcal{S}_j}).$$

By which we conclude that the desired claim holds. □

DEFINITION 1.3.20 (Polynomial of Grace). We define $P_f \in \mathbb{C}[x_0, \dots, x_{n-1}]$ for all $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ as follows:

$$(1.3.21) \quad P_f(\mathbf{x}) := \underbrace{\prod_{0 \leq u < v < n} (x_v - x_u)}_{V(x_0, \dots, x_{n-1})} \underbrace{\prod_{0 \leq u < v < n} ((x_{f(v)} - x_v)^2 - (x_{f(u)} - x_u)^2)}_{E_f(x_0, \dots, x_{n-1})}.$$

DEFINITION 1.3.22 (Congruence class). For polynomials $P, Q \in \mathbb{C}[x_0, \dots, x_{n-1}]$, if

$$(1.3.23) \quad P(\mathbf{x}) \equiv Q(\mathbf{x}) \pmod{\left\{ \prod_{j \in \mathbb{Z}_n} (x_i - j) : i \in \mathbb{Z}_n \right\}},$$

we simply write $P \equiv Q$.

Unless otherwise stated, all congruence relations in this paper are prescribed modulo the ideal of polynomials generated by members of the set

$$\left\{ \prod_{j \in \mathbb{Z}_n} (x_i - j) : i \in \mathbb{Z}_n \right\}$$

PROPOSITION 1.3.24 (Certificate of Grace). Let $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$. The functional directed graph G_f prescribed by f is graceful if and only if $P_f(\mathbf{x}) \not\equiv 0$.

PROOF. Observe that the vertex Vandermonde factor $V(\mathbf{x})$ is of degree exactly $n-1$ in each variable and therefore equal to its remainder, i.e.,

$$(1.3.25) \quad V(\mathbf{x}) = \sum_{\theta \in S_n} \text{sgn}(\theta) \prod_{i \in \mathbb{Z}_n} (x_i)^{\theta(i)} = \prod_{v \in \mathbb{Z}_n} (v!) \sum_{\theta \in S_n} \text{sgn}(\theta) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\theta(i)\}}} \left(\frac{x_i - j_i}{\theta(i) - j_i} \right),$$

where

$$(1.3.26) \quad \text{sgn}(\theta) := \prod_{0 \leq u < v < n} \left(\frac{\theta(v) - \theta(u)}{v - u} \right), \quad \forall \theta \in S_n.$$

When $n > 2$, for every $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$, the induced edge label Vandermonde factor $E_f(\mathbf{x})$ is of degree $> (n-1)$ in some of its variables. Therefore, by Proposition 1.3.1, we have

$$(1.3.27) \quad E_f(\mathbf{x}) = \sum_{l \in \mathbb{Z}_n} q_l(\mathbf{x}) \prod_{k \in \mathbb{Z}_n} (x_l - k) + \prod_{v \in \mathbb{Z}_n} (v!) \frac{(n-1+v)!}{(2v)!} \sum_{\substack{g \in \mathbb{Z}_n^{\mathbb{Z}_n} \\ |gf - g| \in S_n}} \text{sgn}(|gf - g|) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{g(i)\}}} \left(\frac{x_i - j_i}{g(i) - j_i} \right).$$

Observe that by the expansions in 1.3.25 and 1.3.27,

$$(1.3.28) \quad P_f(\mathbf{x}) = \sum_{l \in \mathbb{Z}_n} q_l(\mathbf{x}) V(\mathbf{x}) \prod_{k \in \mathbb{Z}_n} (x_l - k) + \left(\prod_{v \in \mathbb{Z}_n} v! \sum_{\theta \in S_n} \text{sgn}(\theta) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\theta(i)\}}} \left(\frac{x_i - j_i}{\theta(i) - j_i} \right) \right) \left(\prod_{v \in \mathbb{Z}_n} (v!) \frac{(n-1+v)!}{(2v)!} \sum_{\substack{g \in \mathbb{Z}_n^{\mathbb{Z}_n} \\ |gf - g| \in S_n}} \text{sgn}(|gf - g|) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{g(i)\}}} \left(\frac{x_i - j_i}{g(i) - j_i} \right) \right).$$

is congruent to


$$(1.3.28) \quad \prod_{v \in \mathbb{Z}_n} (v!)^2 \frac{(n-1+v)!}{(2v)!} \sum_{\substack{\sigma \in S_n \\ \text{s.t.} \\ |\sigma f - \sigma| \in S_n}} \text{sgn}(\sigma |\sigma f - \sigma|) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right),$$

where the permutation $|\sigma f - \sigma|$ denotes the induced edge label permutation associated with a graceful relabeling $G_{\sigma f \sigma^{-1}}$ of G_f . The congruence above stems from Prop. 1.3.16. A graceful labeling necessitates the integer coefficient

$$\prod_{0 \leq i < j < n} (j-i)(j^2-i^2) = \prod_{0 \leq i < j < n} (j-i)^2(j+i) = \prod_{v \in \mathbb{Z}_n} (v!)^2 \frac{(n-1+v)!}{(2v)!} \neq 0,$$

thus establishing the desired claim. \square

EXAMPLE 1.3.29. We present an example of a path on 5 vertices. This is known to be graceful, so we expect a non-zero remainder.



221 Run the SageMath script `ex1325.sage` to verify.

222 PROPOSITION 1.3.30 (Complementary Labeling Symmetry). *Let $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ and the remainder of P_f be*

$$223 \quad (1.3.31) \quad \bar{P}_f(\mathbf{x}) := \prod_{v \in \mathbb{Z}_n} (v!)^2 \frac{(n-1+v)!}{(2v)!} \sum_{\substack{\sigma \in S_n \\ \text{s.t.} \\ |\sigma f - \sigma| \in S_n}} \text{sgn}(\sigma |\sigma f - \sigma|) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right).$$

224 *The complementary labeling map $x_i \mapsto x_{n-1-i}$, for all $i \in \mathbb{Z}_n$, fixes \bar{P}_f up to sign.*

225 PROOF. For notational convenience, let $\mathbf{x}_\varphi := (x_{\varphi(0)}, \dots, x_{\varphi(i)}, \dots, x_{\varphi(n-1)})$. Observe that for any permutation $\varphi \in S_n$,
 226 the action of φ on P_f yields equalities

$$\begin{aligned} P_f(\mathbf{x}_\varphi) &= \prod_{0 \leq u < v < n} (x_{\varphi(v)} - x_{\varphi(u)}) ((x_{\varphi f(v)} - x_{\varphi(v)})^2 - (x_{\varphi f(u)} - x_{\varphi(u)})^2), \\ &= \prod_{0 \leq \varphi^{-1}(i) < \varphi^{-1}(j) < n} (x_j - x_i) ((x_{\varphi f \varphi^{-1}(j)} - x_j)^2 - (x_{\varphi f \varphi^{-1}(i)} - x_i)^2). \end{aligned}$$

228 The last equality above features the indexing change of variable $u = \varphi^{-1}(i)$ and $v = \varphi^{-1}(j)$. If $\varphi \in \text{Aut}(G_f)$ then
 229 $P_f(x_{\varphi(0)}, \dots, x_{\varphi(n-1)})$ is up to sign equal to $P_{\varphi f \varphi^{-1}}$, in accordance with Definition 1.3.20. Furthermore, by the proof of
 230 Proposition 1.3.24, the action of φ on P_f yields the congruence identity

$$231 \quad P_f(\mathbf{x}_\varphi) \equiv \bar{P}_f(\mathbf{x}_\varphi).$$

232 Hence,

$$\begin{aligned} 233 \quad \bar{P}_f(\mathbf{x}_\varphi) &= \prod_{v \in \mathbb{Z}_n} ((v!)^2 \frac{(n-1+v)!}{(2v)!}) \sum_{\substack{\sigma \in S_n \\ \text{s.t.} \\ |\sigma f - \sigma| \in S_n}} \text{sgn}(\sigma |\sigma f - \sigma|) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_{\varphi(i)} - j_i}{\sigma(i) - j_i} \right), \\ 234 \quad &= \text{sgn}(\varphi) \prod_{v \in \mathbb{Z}_n} \left((v!)^2 \frac{(n-1+v)!}{(2v)!} \right) \sum_{\substack{\sigma \in S_n \\ \text{s.t.} \\ |\sigma f - \sigma| \in S_n}} \text{sgn}(\sigma |\sigma f - \sigma| \varphi^{-1}) \prod_{\substack{u \in \mathbb{Z}_n \\ v_u \in \mathbb{Z}_n \setminus \{\sigma \varphi^{-1}(u)\}}} \left(\frac{x_u - v_u}{\sigma \varphi^{-1}(u) - v_u} \right). \end{aligned}$$

235 If $\varphi = n-1 - \text{id}$, then, by the complementary labeling symmetry, we have

$$236 \quad G_{\sigma f \sigma^{-1}} \in \text{GrL}(G_f) \iff G_{\sigma \varphi^{-1} f (\sigma \varphi^{-1})^{-1}} \in \text{GrL}(G_f)$$

237 Let \mathfrak{G} denote the subgroup of S_n whose members are $\{\text{id}, \varphi\}$. We write

$$\begin{aligned} 238 \quad \bar{P}_f(\mathbf{x}_\varphi) &= \\ 239 \quad &\prod_{v \in \mathbb{Z}_n} ((v!)^2 \frac{(n-1+v)!}{(2v)!}) \sum_{\substack{\sigma \in S_n / \mathfrak{G} \\ \gamma = |\sigma f - \sigma| \in S_n}} \text{sgn}(\sigma \gamma) \left(\text{sgn}(\varphi^{-1}) \prod_{\substack{u \in \mathbb{Z}_n \\ v_u \in \mathbb{Z}_n \setminus \{\sigma(u)\}}} \left(\frac{x_u - v_u}{\sigma(u) - v_u} \right) + \prod_{\substack{u \in \mathbb{Z}_n \\ v_u \in \mathbb{Z}_n \setminus \{\sigma \varphi^{-1}(u)\}}} \left(\frac{x_u - v_u}{\sigma \varphi^{-1}(u) - v_u} \right) \right). \end{aligned}$$

241 Similarly,

$$\begin{aligned} 242 \quad \bar{P}_f(\mathbf{x}) &= \\ 243 \quad &\prod_{v \in \mathbb{Z}_n} ((v!)^2 \frac{(n-1+v)!}{(2v)!}) \sum_{\substack{\sigma \in S_n / \mathfrak{G} \\ \gamma = |\sigma f - \sigma| \in S_n}} \text{sgn}(\sigma \gamma) \left(\prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right) + \text{sgn}(\varphi^{-1}) \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\sigma \varphi^{-1}(i)\}}} \left(\frac{x_i - j_i}{\sigma \varphi^{-1}(i) - j_i} \right) \right). \end{aligned}$$


245 We conclude that the complementary labeling symmetry yields the equality

$$246 \quad \bar{P}_f(\mathbf{x}) = \text{sgn}(\varphi) \bar{P}_f(\mathbf{x}_\varphi) = \bar{P}_{\varphi f \varphi^{-1}}(\mathbf{x}),$$

247 thus establishing the desired claim. □

248 EXAMPLE 1.3.32. We present an example of a path on 5 vertices.

249



251 Run the SageMath script `ex1328.sage` to verify.

LEMMA 1.3.33 (Variable Dependency). *Let $P \in \mathbb{Q}[x_0, \dots, x_{n-1}]$ and $S \subsetneq \mathbb{Z}_n$. If*

$$(1.3.34) \quad P(\mathbf{x}) = \sum_{g \in \mathbb{Z}_n^S} c_g \prod_{i \in S} (x_i)^{g(i)},$$

where $c_g \in \mathbb{C}$ for all $g \in \mathbb{Z}_n^S$, then for any positive integer m , the polynomial $(P(\mathbf{x}))^m$ admits a quotient-remainder expansion of the form

$$(1.3.35) \quad (P(x_0, \dots, x_{n-1}))^m = \sum_{j \in S} q_j(x_0, \dots, x_{n-1}) \prod_{k \in \mathbb{Z}_n} (x_j - \alpha_k) + \sum_{g \in \mathbb{Z}_n^S} a_g \prod_{i \in S} (x_i)^{g(i)}$$

where $\alpha_k, a_g \in \mathbb{C}$ for all $k \in \mathbb{Z}_n$ such that $n = |\{\alpha_k : k \in \mathbb{Z}_n\}|$ and $g \in \mathbb{Z}_n^S$.

PROOF. By the premise, the polynomial $P(\mathbf{x})$ is of degree at most $n-1$ in its variables. Thus by Proposition 1.3.1, the polynomial $P(\mathbf{x})$ is equal to its remainder, i.e.,

$$(1.3.36) \quad P(\mathbf{x}) = \sum_{g \in \mathbb{Z}_n^S} c_g \prod_{i \in S} (x_i)^{g(i)} = \sum_{g \in \mathbb{Z}_n^S} P(g) \prod_{\substack{i \in S \\ j_i \in \mathbb{Z}_n \setminus \{g(i)\}}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right).$$

The remainder of $(P(\mathbf{x}))^m$ is obtained by repeatedly replacing each occurrence of $(x_i)^n$ with $(x_i)^n - \prod_{k \in \mathbb{Z}_n} (x_i - \alpha_k)$, followed by expanding the resulting polynomials, starting from the expanded form of

$$(1.3.37) \quad \left(\sum_{g \in \mathbb{Z}_n^S} c_g \prod_{i \in S} (x_i)^{g(i)} \right)^m,$$

until we obtain a polynomial of degree at most $n-1$ in each variable. The transformation never introduces a variable indexed by a member of the complement of S . We obtain that

$$\left(\sum_{g \in \mathbb{Z}_n^S} c_g \prod_{i \in S} (x_i)^{g(i)} \right)^m = \sum_{j \in \mathbb{Z}_m} q_j(\mathbf{x}) \prod_{k \in \mathbb{Z}_n} (x_j - \alpha_k) + \sum_{g \in \mathbb{Z}_n^S} (P(g))^m \prod_{\substack{i \in S \\ j_i \in \mathbb{Z}_n \setminus \{g(i)\}}} \left(\frac{x_i - \alpha_{j_i}}{\alpha_{g(i)} - \alpha_{j_i}} \right)$$

by which it follows that

$$(1.3.38) \quad \left(\sum_{g \in \mathbb{Z}_n^S} c_g \prod_{i \in S} (x_i)^{g(i)} \right)^m = \sum_{j \in \mathbb{Z}_m} q_j(\mathbf{x}) \prod_{k \in \mathbb{Z}_n} (x_j - \alpha_k) + \sum_{g \in \mathbb{Z}_n^S} a_g \prod_{i \in S} (x_i)^{g(i)},$$

where $\alpha_k, a_g \in \mathbb{C}$ for all $k \in \mathbb{Z}_n$ and $n = |\{\alpha_k : k \in \mathbb{Z}_n\}|$ as claimed. \square

LEMMA 1.3.39 (Monomial support). *Let $P \in \mathbb{Q}[x_0, \dots, x_{n-1}]$ be such that it is not identically constant. If*

$$(1.3.40) \quad P(\mathbf{x}) = \sum_{\sigma \in S_n} a_\sigma \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right),$$

then there exist a minimal non-empty set $\mathcal{M}_P \subset \mathbb{Z}_n^{\mathbb{Z}_n}$ subject to $|f^{-1}(\{0\})| \leq 1$ for all $f \in \mathcal{M}_P$ such that

$$(1.3.41) \quad P(\mathbf{x}) = \sum_{f \in \mathcal{M}_P} c_f \prod_{i \in \mathbb{Z}_n} x_i^{f(i)},$$

where $c_f \in \mathbb{Q} \setminus \{0\}$.

PROOF. Stated otherwise, every term in the expanded form of P is a multiple of at least $n-1$ distinct variables. Consider a Lagrange basis polynomial associated with an arbitrary $\sigma \in S_n$:

$$L_\sigma(\mathbf{x}) = \prod_{\substack{i \in \mathbb{Z}_n \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right) = \prod_{\substack{i \in \mathbb{Z}_n \setminus \{\sigma^{-1}(0)\} \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right) \prod_{j_{\sigma^{-1}(0)} \in \mathbb{Z}_n \setminus \{0\}} \left(\frac{x_{\sigma^{-1}(0)} - j_{\sigma^{-1}(0)}}{0 - j_{\sigma^{-1}(0)}} \right).$$

On the right-hand side of the second equal sign immediately above, the univariate polynomial in $x_{\sigma^{-1}(0)}$ encompassed within the scope of the second Π indexed by $j_{\sigma^{-1}(0)} \in \mathbb{Z}_n \setminus \{0\}$ has (in its expanded form) a non-vanishing constant term equal to one. However, the constant term vanishes within the expanded form of each univariate factor

$$\prod_{j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right)$$

encompassed within the scope of the first Π indexed by $i \in \mathbb{Z}_n \setminus \{\sigma^{-1}(0)\}$. Indeed, we have

$$L_\sigma(\mathbf{x}) = \underbrace{\prod_{\substack{i \in \mathbb{Z}_n \setminus \{\sigma^{-1}(0)\} \\ j_i \in \mathbb{Z}_n \setminus \{\sigma(i)\}}} \left(\frac{x_i - j_i}{\sigma(i) - j_i} \right)}_{\text{does not feature the variable } x_{\sigma^{-1}(0)}} \left(\frac{(x_{\sigma^{-1}(0)})^{n-1} + \dots + (-1)^{n-1}(n-1)!}{(-1)^{n-1}(n-1)!} \right).$$

Observe that each summand term in the expanded form of the Lagrange basis polynomial $L_\sigma(\mathbf{x})$ above which is a non-vanishing monomial multiple of $x_{\sigma^{-1}(0)}$ is a multiple of every variable in $\{x_0, \dots, x_{n-1}\}$. By contrast, every non-vanishing monomial summand term which is not a multiple of $x_{\sigma^{-1}(0)}$ is a multiple of every other variables, i.e., variables in the set $\{x_0, \dots, x_{n-1}\} \setminus \{x_{\sigma^{-1}(0)}\}$. Applying the same argument to each $\sigma \in S_n$ yields the desired claim. \square

1.4. The Composition Lemma

LEMMA 1.4.1 (Transposition Invariance). *Let $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ be such that its functional directed graph G_f has at least two sibling leaf nodes, i.e., G_f has vertices $u, v \in \mathbb{Z}_n$ such that $f^{-1}(\{u, v\}) = \emptyset$ and $f(u) = f(v)$. If the transposition $\tau \in S_n$ exchanges u and v , i.e.,*

$$\tau(i) = \begin{cases} v & \text{if } i = u \\ u & \text{if } i = v \\ i & \text{otherwise} \end{cases} \quad \forall i \in \mathbb{Z}_n.$$

Then

$$(1.4.2) \quad \tau \in \text{Aut}(P_f(\mathbf{x})),$$

where P_f is the polynomial certificate of grace as defined in 1.3.20.

PROOF. Stated otherwise, the claim asserts that the polynomial P_f is fixed by a transposition of any pair of variables associated with sibling leaf vertices. By construction of $P_f(\mathbf{x})$, the changes in its Vandermonde factors induced by the action of τ are as follows:

$$(1.4.3) \quad P_f(x_{\tau(0)}, \dots, x_{\tau(i)}, \dots, x_{\tau(n-1)}) = \prod_{0 \leq i < j < n} (x_{\tau(j)} - x_{\tau(i)}) \prod_{0 \leq i < j < n} ((x_{\tau f(j)} - x_{\tau(j)})^2 - (x_{\tau f(i)} - x_{\tau(i)})^2).$$

Note that there is a bijection

$$(1.4.4) \quad x_i \mapsto (x_{f(i)} - x_i)^2, \quad \forall i \in \mathbb{Z}_n.$$

Hence, the transposition $\tau \in \text{Aut}(G_f)$ of the leaf nodes induces a transposition τ of the corresponding leaf edges outgoing from the said leaf nodes. More precisely, the maps

$$\begin{pmatrix} x_0 & \dots & x_i & \dots & x_{n-1} \\ \downarrow & & \downarrow & & \downarrow \\ x_{\tau(0)} & \dots & x_{\tau(i)} & \dots & x_{\tau(n-1)} \end{pmatrix}$$

and

$$\begin{pmatrix} (x_{f(0)} - x_0)^2 & \dots & (x_{f(i)} - x_i)^2 & \dots & (x_{f(n-1)} - x_{n-1})^2 \\ \downarrow & & \downarrow & & \downarrow \\ (x_{\tau f(0)} - x_{\tau(0)})^2 & \dots & (x_{\tau f(i)} - x_{\tau(i)})^2 & \dots & (x_{\tau f(n-1)} - x_{\tau(n-1)})^2 \end{pmatrix}$$

prescribe the same permutation τ of the vertex variables and induced edges label binomials respectively. Observe that

$$P_f(x_{\tau(0)}, \dots, x_{\tau(i)}, \dots, x_{\tau(n-1)}) = \left(\prod_{0 \leq i < j < n} \frac{x_{\tau(j)} - x_{\tau(i)}}{x_j - x_i} \prod_{0 \leq i < j < n} (x_j - x_i) \right) \left(\prod_{0 \leq i < j < n} \frac{(x_{\tau f(j)} - x_{\tau(j)})^2 - (x_{\tau f(i)} - x_{\tau(i)})^2}{(x_{f(j)} - x_j)^2 - (x_{f(i)} - x_i)^2} \prod_{0 \leq i < j < n} ((x_{f(j)} - x_j)^2 - (x_{f(i)} - x_i)^2) \right)$$

$$\begin{aligned}
&= \left(\operatorname{sgn}(\tau) \prod_{0 \leq i < j < n} (x_j - x_i) \right) \left(\operatorname{sgn}(\tau) \prod_{0 \leq i < j < n} ((x_{f(j)} - x_j)^2 - (x_{f(i)} - x_i)^2) \right) \\
&= \left((-1) \prod_{0 \leq i < j < n} (x_j - x_i) \right) \left((-1) \prod_{0 \leq i < j < n} ((x_{f(j)} - x_j)^2 - (x_{f(i)} - x_i)^2) \right) \\
&\implies P_f(x_{\tau(0)}, \dots, x_{\tau(n-1)}) = P_f(x_0, \dots, x_{n-1}),
\end{aligned}$$

thus establishing the desired claim. \square

PROPOSITION 1.4.6 (Composition Inequality). *Consider an arbitrary $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ subject to the fixed point condition $|f^{(n-1)}(\mathbb{Z}_n)| = 1$. The following statements are equivalent:*

(i)

$$\max_{\sigma \in S_n} |\{|\sigma f^{(2)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}| \leq \max_{\sigma \in S_n} |\{|\sigma f \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}|.$$

(ii)

$$P_{f^{(2)}}(\mathbf{x}) \neq 0 \implies P_f(\mathbf{x}) \neq 0.$$

(iii)

$$\operatorname{GrL}(G_f) \neq \emptyset$$

PROOF. If $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ is identically constant, then G_f is graceful. We see this from the fact that the functional digraph of the identically zero function is gracefully labeled and the fact that functional digraphs of identically constant functions are all isomorphic. It follows that all functional directed graphs having diameter less than 3 are graceful. Consequently, all claims hold for all functional digraphs of diameter less than 3. We now turn our attention to functional trees of diameter greater or equal to 3. It follows by definition

$$(1.4.7) \quad n = \max_{\sigma \in S_n} |\{|\sigma f \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}| \iff P_f(\mathbf{x}) \neq 0 \iff \operatorname{GrL}(G_f) \neq \emptyset.$$

We now proceed to show (i) \iff (iii). The backward claim is the simplest of the two claims. We see that if f is contractive, so too is $f^{(2)}$. Then the assertions

$$(1.4.8) \quad n = \max_{\sigma \in S_n} |\{|\sigma f^{(2)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}| \text{ and } n = \max_{\sigma \in S_n} |\{|\sigma f \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}|$$

indeed implies the inequality

$$(1.4.9) \quad \max_{\sigma \in S_n} |\{|\sigma f^{(2)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}| \leq \max_{\sigma \in S_n} |\{|\sigma f \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}|.$$

We now establish the forward claim by contradiction. Assume for the sake of establishing a contradiction that for some contractive map $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$ we have

$$(1.4.10) \quad n > \max_{\sigma \in S_n} |\{|\sigma f^{(2)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}|,$$

for we know by the number of edges being equal to n that it is impossible that

$$(1.4.11) \quad n < \max_{\sigma \in S_n} |\{|\sigma f^{(2)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}|.$$

Note that the range of f is a proper subset of \mathbb{Z}_n . By the premise that f is contractive, it follows that $f^{(\lceil 2^{\lg(n-1)} \rceil)}$ is identically constant and thus

$$(1.4.12) \quad n = \max_{\sigma \in S_n} |\{|\sigma f^{(\lceil 2^{\lg(n-1)} \rceil)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}|,$$

where \lg denotes the logarithm base 2. Consequently there must be some integer $0 \leq \kappa < \lg(n-1)$ such that

$$(1.4.13) \quad \max_{\sigma \in S_n} |\{|\sigma f^{(\lceil 2^\kappa \rceil)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}| > \max_{\sigma \in S_n} |\{|\sigma f^{(\lceil 2^{\kappa-1} \rceil)} \sigma^{-1}(i) - i| : i \in \mathbb{Z}_n\}|.$$

This contradicts the assertion of statement (i), thereby establishing the backward claim. The exact same reasoning as above establishes (ii) \iff (iii), for we have

$$(1.4.14) \quad P_{f^{(\lceil 2^{\lg(n-1)} \rceil)}}(\mathbf{x}) \neq 0.$$

□

Having assembled together the pieces required to prove our main result, we proceed to fit the pieces together to state and prove the *Composition Lemma*.

LEMMA 1.4.15 (Composition Lemma). *For all contractive $f \in \mathbb{Z}_n^{\mathbb{Z}_n}$, i.e., subject to $|f^{(n-1)}(\mathbb{Z}_n)| = 1$, we have*

$$(1.4.16) \quad \max_{\sigma \in S_n} |\{\sigma f^{(2)} \sigma^{-1}(i) - i : i \in \mathbb{Z}_n\}| \leq \max_{\sigma \in S_n} |\{\sigma f \sigma^{-1}(i) - i : i \in \mathbb{Z}_n\}|.$$

PROOF. Owing to Proposition 1.4.6, we prove the statement by establishing

$$P_{f^{(2)}}(\mathbf{x}) \not\equiv 0 \implies P_f(\mathbf{x}) \not\equiv 0.$$

For simplicity, we prove a generalization of the desired claim. Given that the diameter of G_f is greater than 2, we may assume without loss of generality that $f^{-1}(\{n-1\}) = \emptyset$ and $f^{(2)}(n-1) \neq f(n-1)$. Let the contractive map $g \in \mathbb{Z}_n^{\mathbb{Z}_n}$ be devised from f such that

$$(1.4.17) \quad g(i) = \begin{cases} f^{(2)}(i) & \text{if } i \in f^{-1}(\{f(n-1)\}) \\ f(i) & \text{otherwise} \end{cases}, \quad \forall i \in \mathbb{Z}_n.$$

We show that

$$(1.4.18) \quad P_g(\mathbf{x}) \not\equiv 0 \implies P_f(\mathbf{x}) \not\equiv 0.$$

Note that the assertion immediately above generalizes the composition lemma since, the function f is only partially iterated. More precisely, f is iterated only on the restriction $f^{-1}(\{f(n-1)\}) \subset \mathbb{Z}_n$. Iterating this slight generalization of the composition lemma yields that all functional trees are graceful, which in turn implies that the *Composition Lemma* as stated in Lemma 1.4.16 holds. For notational convenience, we assume without loss of generality that

$$(1.4.19) \quad f(n-1) = n - |f^{-1}(\{f(n-1)\})| - 1 \quad \text{and} \quad f^{-1}(\{f(n-1)\}) = \mathbb{Z}_n \setminus \mathbb{Z}_{1+f(n-1)}.$$

If the conditions stated above are not met, we relabel the vertices of G_f to ensure that such is indeed the case. Note that such a relabeling does not affect the property we seek to prove. We prove the contrapositive claim

$$(1.4.20) \quad P_f(\mathbf{x}) \equiv 0 \implies P_g(\mathbf{x}) \equiv 0.$$

By construction, the polynomial

$$(1.4.21) \quad P_f(\mathbf{x}) = \prod_{0 \leq i < j < n} (x_j - x_i) \times \prod_{\substack{0 \leq u < v \leq f(n-1) \\ t \in \{0,1\}}} \left(x_{f(v)} - x_v + (-1)^t (x_{f(u)} - x_u) \right) \times \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} \left(x_{f(v)} - x_v + (-1)^t (x_{f(u)} - x_u) \right) \times \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v \\ t \in \{0,1\}}} \left(x_{f(v)} - x_v + (-1)^t (x_{f(u)} - x_u) \right),$$

differs only slightly from

$$\begin{aligned}
 P_g(\mathbf{x}) = & \prod_{0 \leq i < j < n} (x_j - x_i) \times \\
 & \prod_{\substack{0 \leq u < v \leq f(n-1) \\ t \in \{0,1\}}} \left(x_{f(v)} - x_v + (-1)^t (x_{f(u)} - x_u) \right) \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} \left(x_{f^{(2)}(v)} - x_v + (-1)^t (x_{f(u)} - x_u) \right) \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v \\ t \in \{0,1\}}} \left(x_{f^{(2)}(v)} - x_v + (-1)^t (x_{f^{(2)}(u)} - x_u) \right).
 \end{aligned}
 \tag{1.4.22}$$

We setup a variable *telescoping* within each induced edge label binomial $(x_{f^{(2)}(v)} - x_v)$ where $v \in f^{-1}(\{f(n-1)\})$ (i.e. induced edge binomials of edges outgoing from the subset of vertices where f is iterated) as follows:

$$\underbrace{(x_{f^{(2)}(v)} - x_v)}_{x_v \longrightarrow x_{f^{(2)}(v)}} = \underbrace{(\textcolor{red}{x}_{f(v)} - x_v)}_{x_v \longrightarrow \textcolor{red}{x}_{f(v)}} + \underbrace{(x_{f^{(2)}(v)} - \textcolor{blue}{x}_{f(v)})}_{\textcolor{blue}{x}_{f(v)} \longrightarrow x_{f^{(2)}(v)}}.$$

The telescoping enables us to express induced edge binomials of edges outgoing from the subset of vertices where f is iterated in terms of induced edge binomials which feature in E_f . Note that

$$(\textcolor{blue}{x}_{f^{(2)}(v)} - x_v) = \left((x_{f^{(2)}(v)} - \textcolor{blue}{x}_{f(v)}) + (\textcolor{red}{x}_{f(v)} - x_v) \right) = (\textcolor{blue}{x}_{f^{(2)}(n-1)} - \textcolor{blue}{x}_{f(n-1)}) + (\textcolor{red}{x}_{f(v)} - x_v),$$

given that $f(v) = f(n-1)$ for all $v \in f^{-1}(\{f(n-1)\})$. Thus

$$\begin{aligned}
 P_g = & \prod_{0 \leq i < j < n} (x_j - x_i) \times \\
 & \prod_{\substack{0 \leq u < v \leq f(n-1) \\ t \in \{0,1\}}} \left(x_{f(v)} - x_v + (-1)^t (x_{f(u)} - x_u) \right) \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} \left((\textcolor{blue}{x}_{f^{(2)}(n-1)} - \textcolor{blue}{x}_{f(n-1)}) + (\textcolor{red}{x}_{f(v)} - x_v) + (-1)^t (x_{f(u)} - x_u) \right) \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v \\ t \in \{0,1\}}} \left((\textcolor{blue}{x}_{f^{(2)}(n-1)} - \textcolor{blue}{x}_{f(n-1)}) + (\textcolor{red}{x}_{f(v)} - x_v) + (-1)^t ((\textcolor{blue}{x}_{f^{(2)}(n-1)} - \textcolor{blue}{x}_{f(n-1)}) + (\textcolor{red}{x}_{f(u)} - x_u)) \right).
 \end{aligned}$$

For notational convenience, we set induced edge binomials to be $b_u := (x_{f(u)} - x_u)$ for all $u \in \mathbb{Z}_n$ we write

$$\begin{aligned}
 P_g = & \prod_{0 \leq i < j < n} (x_j - x_i) \times \\
 & \prod_{0 \leq u < v \leq f(n-1)} ((b_v)^2 - (b_u)^2) \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (\textcolor{blue}{b}_{u,v,t} + \textcolor{blue}{b}_{f(n-1)}) \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{blue}{b}_v - \textcolor{blue}{b}_u + 0 \textcolor{blue}{b}_{f(n-1)}) (\textcolor{blue}{b}_v + \textcolor{blue}{b}_u + 2 \textcolor{blue}{b}_{f(n-1)})
 \end{aligned}
 \tag{1.4.23}$$

where chromatic induced edge label binomial expressions are

$$\begin{aligned}
 \textcolor{blue}{b}_{f(n-1)} &:= (x_{f(2)(n-1)} - \textcolor{blue}{x}_{f(n-1)}), \\
 \textcolor{red}{b}_i &:= (\textcolor{red}{x}_{f(i)} - x_i), \quad \forall i \in f^{-1}(\{f(n-1)\}),
 \end{aligned}$$

and

$$\textcolor{red}{b}_{u,v,t} := \textcolor{red}{b}_v + (-1)^t b_u = ((\textcolor{red}{x}_{f(v)} - \textcolor{red}{x}_v) + (-1)^t (x_{f(u)} - x_u)), \quad \forall \substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}.$$

Note that in equation 1.4.23, we have successfully re-expressed the edge label Vandermonde factor E_g exclusively in terms of induced edge label binomials from G_f . We re-write the expression of P_f in equation (1.4.21) using the shorthand notation for induced edge binomials as follows:

$$\begin{aligned}
 P_f = & \prod_{0 \leq i < j < n} (x_j - x_i) \times \\
 & \prod_{0 \leq u < v \leq f(n-1)} ((b_v)^2 - (b_u)^2) \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} \textcolor{red}{b}_{u,v,t} \times \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v - \textcolor{red}{b}_u) (\textcolor{red}{b}_v + \textcolor{red}{b}_u).
 \end{aligned}
 \tag{1.4.24}$$

Invoking the multi-binomial identity on the two bichromatic factors of P_g in equation (1.4.23) yields equalities

$$\begin{aligned}
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v + \textcolor{red}{b}_u + 2 \textcolor{blue}{b}_{f(n-1)}) = \\
 & \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v + \textcolor{red}{b}_u) + \sum_{\substack{r_{u,v} \in \{0,1\} \\ 0 = \prod r_{u,v}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v + \textcolor{red}{b}_u)^{r_{u,v}} (2 \textcolor{blue}{b}_{f(n-1)})^{1-r_{u,v}}.
 \end{aligned}$$

and

$$\prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (\textcolor{red}{b}_{u,v,t} + \textcolor{blue}{b}_{f(n-1)}) =$$

$$\prod_{\substack{t \in \{0,1\} \\ v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1)}} \textcolor{red}{b}_{u,v,t} + \sum_{\substack{s_{u,v,t} \in \{0,1\} \\ 0 = \prod s_{u,v,t}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (\textcolor{red}{b}_{u,v,t})^{s_{u,v,t}} (\textcolor{blue}{b}_{f(n-1)})^{1-s_{u,v,t}}.$$

Substituting equalities immediately above into equation (1.4.23) yields an expression of P_g of the form

$$(1.4.25) \quad P_g = \textcolor{red}{P}_f + R_{f,g}.$$

The monochromatic red expressions in the multi-binomial expansion collect to result in $\textcolor{red}{P}_f$ as written in equation (1.4.24).

The second part denoted $R_{f,g}$ simply collects the remaining bichromatic summands and is given by

$$\begin{aligned} R_{f,g} = & \prod_{0 \leq i < j < n} (x_j - x_i) \prod_{0 \leq u < v \leq f(n-1)} ((b_v)^2 - (b_u)^2) \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v - \textcolor{red}{b}_u) \times \\ & \left[\left(\prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v + \textcolor{red}{b}_u) \right) \left(\sum_{\substack{s_{u,v,t} \in \{0,1\} \\ 0 = \prod s_{u,v,t}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (\textcolor{red}{b}_{u,v,t})^{s_{u,v,t}} (\textcolor{blue}{b}_{f(n-1)})^{1-s_{u,v,t}} \right) + \right. \\ & \left(\prod_{\substack{t \in \{0,1\} \\ v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1)}} \textcolor{red}{b}_{u,v,t} \right) \left(\sum_{\substack{r_{u,v} \in \{0,1\} \\ 0 = \prod r_{u,v}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v + \textcolor{red}{b}_u)^{r_{u,v}} (2\textcolor{blue}{b}_{f(n-1)})^{1-r_{u,v}} \right) + \\ & \left. \left(\sum_{\substack{s_{u,v,t} \in \{0,1\} \\ 0 = \prod s_{u,v,t}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (\textcolor{red}{b}_{u,v,t})^{s_{u,v,t}} (\textcolor{blue}{b}_{f(n-1)})^{1-s_{u,v,t}} \right) \left(\sum_{\substack{r_{u,v} \in \{0,1\} \\ 0 = \prod r_{u,v}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v + \textcolor{red}{b}_u)^{r_{u,v}} (2\textcolor{blue}{b}_{f(n-1)})^{1-r_{u,v}} \right) \right]. \end{aligned}$$

The color scheme above is meant to help track the location of telescoping variables. We now proceed with the main contradiction argument. Assume for the sake of establishing a contradiction that the claim (1.4.20) is false, i.e., for some f subject to conditions described in our premise, we have

$$(1.4.26) \quad 0 \equiv \textcolor{red}{P}_f \text{ and } 0 \not\equiv P_g.$$

Then by equation (1.4.25), we obtain

$$(1.4.27) \quad P_g \equiv R_{f,g} \not\equiv 0 \iff \overline{P}_g = \overline{R}_{f,g} \not\equiv 0.$$

Observe that every summand in $R_{f,g}$ is a multiple of a positive power of the induced edge label binomial

$$\textcolor{blue}{b}_{f(n-1)} = (x_{f(2)(n-1)} - x_{f(n-1)})$$

We focus in particular on the summand within $R_{f,g}$ which is a multiple of the largest possible power of the blue induced edge label binomial $\textcolor{blue}{b}_{f(n-1)}$, namely the summand associated with binary exponent assignments

$$(1.4.28) \quad s_{u,v,t} = 0, \text{ for all } \begin{matrix} v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\} \end{matrix} \quad \text{as well as } r_{u,v} = 0, \text{ for all } \begin{matrix} v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \end{matrix}.$$

The said summand is

$$c \prod_{0 \leq i < j < n} (x_j - x_i) \left(\prod_{0 \leq u < v \leq f(n-1)} ((b_v)^2 - (b_u)^2) \right) \left(\prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (\textcolor{red}{b}_v - \textcolor{red}{b}_u) \right) (\textcolor{blue}{b}_{f(n-1)})^m,$$

where

$$(1.4.29) \quad m = \left| \left\{ \begin{matrix} v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v \end{matrix} \right\} \right| + \left| \left\{ \begin{matrix} v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\} \end{matrix} \right\} \right| \text{ and } c = 2^{\left| \left\{ \begin{matrix} v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v \end{matrix} \right\} \right|}.$$

It follows from the premise $0 \neq P_g$ that the remainder of the chosen summand is non-vanishing. Observe that the factor

$$(1.4.30) \quad \prod_{0 \leq i < j < n} (x_j - x_i) \left(\prod_{0 \leq u < v \leq f(n-1)} ((b_v)^2 - (b_u)^2) \right) \left(\prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (b_v - b_u) \right)$$

is a common factor to every summand in $R_{f,g}$. We factor out from the said common factor a smaller factor up to sign invariant to the transposition τ which exchanges the variable x_{n-1} with $x_{f(n-1)}$. We write

$$\begin{aligned} R_{f,g} = & \prod_{0 \leq i < j < n} (x_j - x_i) \prod_{0 \leq u < v < f(n-1)} ((b_v)^2 - (b_u)^2) \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \setminus \{n-1\} \\ f(n-1) < u < v}} (b_v - b_u) \times \\ & \prod_{0 \leq u < f(n-1)} ((b_{f(n-1)})^2 - (b_u)^2) \prod_{f(n-1) < u < n-1} (b_{n-1} - b_u) \times \\ & \left[\left(\prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (b_v + b_u) \right) \left(\sum_{\substack{s_{u,v,t} \in \{0,1\} \\ 0 = \prod s_{u,v,t}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (b_{u,v,t})^{s_{u,v,t}} (b_{f(n-1)})^{1-s_{u,v,t}} \right) + \right. \\ & \left(\prod_{\substack{t \in \{0,1\} \\ v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1)}} b_{u,v,t} \right) \left(\sum_{\substack{r_{u,v} \in \{0,1\} \\ 0 = \prod r_{u,v}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (b_v + b_u)^{r_{u,v}} (2b_{f(n-1)})^{1-r_{u,v}} \right) + \\ & \left. \left(\sum_{\substack{s_{u,v,t} \in \{0,1\} \\ 0 = \prod s_{u,v,t}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (b_{u,v,t})^{s_{u,v,t}} (b_{f(n-1)})^{1-s_{u,v,t}} \right) \left(\sum_{\substack{r_{u,v} \in \{0,1\} \\ 0 = \prod r_{u,v}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (b_v + b_u)^{r_{u,v}} (2b_{f(n-1)})^{1-r_{u,v}} \right) \right]. \end{aligned}$$

Thus

$$R_{f,g}(\mathbf{x}) = \left(\prod_{0 \leq i < j < n} (x_j - x_i) \prod_{0 \leq u < v < f(n-1)} ((b_v)^2 - (b_u)^2) \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \setminus \{n-1\} \\ f(n-1) < u < v}} (b_v - b_u) \right) Q_{f,g}(\mathbf{x}),$$

where

$$\begin{aligned} Q_{f,g}(\mathbf{x}) = & \prod_{0 \leq u < f(n-1)} ((b_{f(n-1)})^2 - (b_u)^2) \prod_{f(n-1) < u < n-1} (b_{n-1} - b_u) \times \\ & \left[\left(\prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (b_v + b_u) \right) \left(\sum_{\substack{s_{u,v,t} \in \{0,1\} \\ 0 = \prod s_{u,v,t}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (b_{u,v,t})^{s_{u,v,t}} (b_{f(n-1)})^{1-s_{u,v,t}} \right) + \right. \\ & \left(\prod_{\substack{t \in \{0,1\} \\ v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1)}} b_{u,v,t} \right) \left(\sum_{\substack{r_{u,v} \in \{0,1\} \\ 0 = \prod r_{u,v}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (b_v + b_u)^{r_{u,v}} (2b_{f(n-1)})^{1-r_{u,v}} \right) + \\ & \left. \left(\sum_{\substack{s_{u,v,t} \in \{0,1\} \\ 0 = \prod s_{u,v,t}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0,1\}}} (b_{u,v,t})^{s_{u,v,t}} (b_{f(n-1)})^{1-s_{u,v,t}} \right) \left(\sum_{\substack{r_{u,v} \in \{0,1\} \\ 0 = \prod r_{u,v}}} \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v}} (b_v + b_u)^{r_{u,v}} (2b_{f(n-1)})^{1-r_{u,v}} \right) \right]. \end{aligned}$$

Let

$$\Phi(g) := \{ \theta \in S_n : G_{\theta g \theta^{-1}} \in \text{GrL}(G_g) \}.$$

By Proposition 1.3.1, the premise $\overline{P}_g = \overline{R}_{f,g}$ implies

$$(1.4.31) \quad P_g(h) = R_{f,g}(h) = v_h Q_{f,g}(h) = \begin{cases} \text{sgn}(h |hg - h|) \prod_{v \in \mathbb{Z}_n} (v!)^2 \frac{(n-1+v)!}{(2v)!} & \text{if } h \in \Phi(g) \\ 0 & \text{otherwise} \end{cases}, \forall h \in \mathbb{Z}_n^{\mathbb{Z}_n}.$$

Recall that $Q_{f,g}$ is the remaining factor of $R_{f,g}$ after we exclude the invariant common factor (up to sign invariant to the transposition τ) equal to

$$\prod_{0 \leq i < j < n} (x_j - x_i) \prod_{0 \leq u < v < f(n-1)} ((b_v)^2 - (b_u)^2) \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \setminus \{n-1\} \\ f(n-1) < u < v}} (b_v - b_u)$$

Specifically, $Q_{f,g}$ is a polynomial resulting from the sum over chromatic summands resulting from the multibinomial expansions. Let us view $Q_{f,g}$ as a sum of $|\Sigma|$ summands and denote by $Q_{f,g}^{[s]}$ the summands $1 \leq s \leq |\Sigma|$ of $Q_{f,g}$. By Proposition 1.3.15, we can write the remainder of $R_{f,g}$ as follows:

$$\bar{R}_{f,g} = \sum_{1 \leq s \leq |\Sigma|} \left(\sum_{\sigma \in \Phi(g)} v_\sigma \cdot Q_{f,g}^{[s]}(\sigma) \cdot L_\sigma(\mathbf{x}) \right),$$

Let us denote by I_s the set of indices associated with variables featured in $Q_{f,g}^{[s]}$. Recall that $L_\sigma(\mathbf{x}|_{I_s})$ denotes the factor of $L_\sigma(\mathbf{x})$ which only features variables indexed by I_s . Note that if any permutation τ lies in the automorphism group of the polynomial

$$\bar{R}_{f,g} = \sum_{1 \leq s \leq |\Sigma|} \left(\sum_{\sigma \in \Phi(g)} v_\sigma \cdot Q_{f,g}^{[s]}(\sigma) \cdot L_\sigma(\mathbf{x}|_{I_s}) \cdot L_\sigma(\mathbf{x}|_{\mathbb{Z}_n \setminus I_s}) \right),$$

then, by Lemma 1.3.17, τ also lies in the automorphism group of the polynomial

$$\sum_{1 \leq s \leq |\Sigma|} \left(\sum_{\sigma \in \Phi(g)} v_\sigma \cdot Q_{f,g}^{[s]}(\sigma) \cdot L_\sigma(\mathbf{x}|_{I_s}) \right).$$

By Lemma 1.4.1, the premise $\bar{P}_g = \bar{R}_{f,g}$ implies that the transposition τ fixes $\bar{R}_{f,g}$. That is to say, we obtain

$$\tau \in \text{Aut} \left(\sum_{1 \leq s \leq |\Sigma|} \left(\sum_{\sigma \in \Phi(g)} v_\sigma \cdot Q_{f,g}^{[s]}(\sigma) \cdot L_\sigma(\mathbf{x}|_{I_s}) \right) \right)$$

As a consequence of the multi-binomial expansion, the sum expressing $Q_{f,g}$ features as one of its summand a unique monochromatic blue binomial summand, say $Q_{f,g}^{[1]}$, given by

$$Q_{f,g}^{[1]} = c (\textcolor{red}{b}_{n-1})^{n-1-f(n-1)-2} (\textcolor{blue}{b}_{f(n-1)})^\ell.$$

where

$$\ell = \left| \left\{ \begin{array}{l} v \in f^{-1}(\{f(n-1)\}) \\ f(n-1) < u < v \end{array} \right\} \right| + \left| \left\{ \begin{array}{l} v \in f^{-1}(\{f(n-1)\}) \\ 0 \leq u \leq f(n-1) \\ t \in \{0, 1\} \end{array} \right\} \right| + 2f(n-1).$$

So that we are interested in

$$\left(\prod_{0 \leq i < j < n} (x_j - x_i) \prod_{0 \leq u < v < f(n-1)} ((b_v)^2 - (b_u)^2) \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \setminus \{n-1\} \\ f(n-1) < u < v}} (b_v - b_u) \right) Q_{f,g}^{[1]} =$$

$$\left(\prod_{0 \leq i < j < n} (x_j - x_i) \prod_{0 \leq u < v < f(n-1)} ((b_v)^2 - (b_u)^2) \prod_{\substack{v \in f^{-1}(\{f(n-1)\}) \setminus \{n-1\} \\ f(n-1) < u < v}} (b_v - b_u) \right) c (\textcolor{red}{b}_{n-1})^{n-1-f(n-1)-2} (\textcolor{blue}{b}_{f(n-1)})^\ell$$

Note that $\textcolor{red}{b}_{n-1}$ is up to sign invariant to transposition τ . By Lemmas 1.3.33 and 1.3.17, the invariance of $Q_{f,g}^{[1]}$ to the transposition τ is predicated upon the τ -invariance up to sign of the polynomial

$$c \sum_{\sigma \in \Phi(g)} (\sigma f^{(2)}(n-1) - \sigma f(n-1))^\ell \times$$

$$\prod_{j_{f(2)(n-1)} \in \mathbb{Z}_n \setminus \{\sigma f^{(2)}(n-1)\}} \left(\frac{x_{f(2)(n-1)} - j_{f(2)(n-1)}}{\sigma f^{(2)}(n-1) - j_{f(2)(n-1)}} \right) \prod_{j_{f(n-1)} \in \mathbb{Z}_n \setminus \{\sigma f(n-1)\}} \left(\frac{\textcolor{blue}{x}_{f(n-1)} - j_{f(n-1)}}{\sigma f(n-1) - j_{f(n-1)}} \right),$$

which is seen to be up to sign asymmetric to the transposition τ .

Let us now focus on the action of the transposition τ on individual summands of some polynomial resulting from an arbitrary but fixed partition of its non-vanishing monomial terms. There are exactly three distinct ways that a candidate

transposition of a pair of variables can lie in the automorphism group of a given polynomial. Assume that we reason about a particular summand denoted as S .

- (i) Option 1: The candidate transposition of a pair of variables fixes the chosen summand S . This occurs when S is symmetric in the chosen pair of variables being transposed.
- (ii) Option 2: The candidate transposition of the chosen pair of variables does not fix S (i.e., Option 1 does not apply) but induces in turn a transposition which exchanges the chosen summand S with some other summand from the partition say, S' . This occurs, for instance, if we consider the sum $S + S'$ where $S = (x_0)^2 x_1$ and $S' = x_0 (x_1)^2$. In this example, we see that transposition which exchanges variables x_0 with x_1 does not fix S , but it induces a transposition which exchanges the summand S with the summand S' .
- (iii) Option 3: The candidate transposition of a pair of variables neither fixes S nor does it induce a transposition which exchanges S with some other summand (i.e., neither Option 1 nor Option 2 applies). Instead, S is such that a symmetry broadening cancellation occurs. Such a cancellation must involve interactions between the non-vanishing monomials within the monomial support of S with the non-vanishing monomials within the support of other summands. Option 3 occurs, for instance, if we take $S = -x_1$ and $S' = x_0 + 2x_1$. We see that in this example neither Option 1 nor Option 2 applies when the candidate transposition is the transposition which exchanges variables x_0 with x_1 . However $S + S' = x_0 + x_1$ is symmetric and thus admits the said transposition in its automorphism group. This fact is due to the symmetry broadening cancellation of like terms : $-x_1 + 2x_1$.

We have already established that $Q_{f,g}^{[1]}$ is not up to sign fixed by the transposition $\tau \in S_n$. This first observation accounts for Option 1. Also note that the remainder of the chosen summand $Q_{f,g}^{[1]}$ does not exchange with the remainder of any other summands when we exchange $x_{f(n-1)}$ with x_{n-1} since by Lemma 1.3.39, when discounting the τ -invariant (up to sign) factor edge binomial b_{n-1} , the non-vanishing remainders of other bi-chromatic summand in $Q_{f,g}$ depends on 3 or more variables. This second observation accounts for Option 2. We now account for Option 3 and show that there are no symmetry-broadening cancellations which adjoin τ to the automorphism group. Again by Lemma 1.3.39, such a symmetry broadening cancellation can occur only for Lagrange bases

$$\prod_{j_{f(2)(n-1)} \in \mathbb{Z}_n \setminus \{\sigma f(2)(n-1)\}} \left(\frac{x_{f(2)(n-1)} - j_{f(2)(n-1)}}{\sigma f(2)(n-1) - j_{f(2)(n-1)}} \right) \prod_{j_{f(n-1)} \in \mathbb{Z}_n \setminus \{\sigma f(n-1)\}} \left(\frac{x_{f(n-1)} - j_{f(n-1)}}{\sigma f(n-1) - j_{f(n-1)}} \right).$$

where $\sigma \in \Phi(g)$ is subject to $\sigma(n-1) = 0$ and G_f is such that $1 = |f^{-1}(\{f(n-1)\})|$. In that setting, non-vanishing monomials occurring in the expanded form of said Lagrange bases summands possibly cancel out non-vanishing monomials occurring in the expanded form of Lagrange bases expressing remainders of bi-chromatic summands in $Q_{f,g}$ of the form



$$\left(b_{f(n-1), n-1, t} \right)^r \left(b_{f(n-1)} \right)^s = \left(b_{n-1} + (-1)^t b_{f(n-1)} \right)^r \left(b_{f(n-1)} \right)^s$$

However, the restriction imposed by $\sigma \in \Phi(g)$ where $\sigma(n-1) = 0$ breaks the complementary-labeling symmetry. Indeed by Proposition 1.3.30, the remainder is up to sign invariant to the involution prescribed by the map: $x_i \mapsto x_{n-1-i}$ for all $i \in \mathbb{Z}_n$. But the complementary labeling involution maps any Lagrange basis associated with $\sigma \in \Phi(g)$ such that $\sigma(n-1) = 0$ to different Lagrange basis associated $\sigma' \in \Phi(g)$ such that $\sigma'(n-1) = n-1$ and thus negates the symmetry broadening cancellations. We see that a symmetry broadening cancellation which adjoins τ to the automorphism group of the remainder of $R_{f,g}$ would break the complementary labeling symmetry, thereby resulting in the contradiction

$$\tau = (f(n-1), n-1) \notin \text{Aut}(\overline{R}_{f,g}).$$

We conclude that the desired claim $P_g(\mathbf{x}) \not\equiv 0 \implies P_f(\mathbf{x}) \not\equiv 0$. holds. □

EXAMPLE 1.4.32. We present a verification of Lemma 1.4.15 with an example of a path on 5 vertices.



Run the SageMath script `ex1434.sage` to verify.

Bibliography

- 511 [1] Parikshit Chalise, Antwan Clark, and Edinah K. Gnan. Every tree on n edges decomposes $k_{nx,nx}$ and k_{2nx+1} , 2024.
- 512 [2] Parikshit Chalise, Antwan Clark, and Edinah K. Gnan. A proof of the tree packing conjecture, 2024.
- 513 [3] Edinah K. Gnan. On graceful labelings of trees, 2020.
- 514 [4] Edinah K. Gnan. On the composition lemma, 2022.
- 515 [5] Edinah K. Gnan. A proof of the kotzig-ringel-rosa conjecture, 2023.