

Midterm 2, MA 265 Section: Spring 2015

Name:

PUID:

Exam B:

No calculators may be used on this exam.

Problem Number	Possible Points	Points Earned
I	80pts	
II	20pts	
Total Points	100pts	

I (80pts)

Answer the following questions by clearly circling your answers. No partial credit is awarded for the questions in Part I

(1) Given the matrices

$$\mathbf{A} = \begin{pmatrix} 0 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 5 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 1 & 3 & 7 \\ 0 & 0 & 5 \\ 0 & 0 & 4 \end{pmatrix} \quad (1)$$

which one of the matrix operations below yields an invertible matrix (circle all that apply)

- (a) $\mathbf{A} \cdot \mathbf{B}$
- (b) $\mathbf{A} \cdot \mathbf{B}^T$
- (c) $\mathbf{A}^T \cdot \mathbf{B}$
- (d) $\mathbf{A} + \mathbf{B}$
- (e) None of the above

(2) Let

$$\mathbf{v}_1 = \begin{bmatrix} 1 & 1 & 2 & 3 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 2 & 3 & 4 & 5 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}, \mathbf{v}_4 = \begin{bmatrix} 3 & 4 & 4 & 4 \end{bmatrix}.$$

be vectors in $\mathbb{R}^{1 \times 4}$ spanning the subspace W . Which of the following subset forms a basis for W ? (circle all that apply)

- (a) $\{\mathbf{v}_1, \mathbf{v}_2\}$.
- (b) $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_4\}$.
- (c) $\left\{ \begin{bmatrix} 1 & 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 1 & 1 \end{bmatrix} \right\}$.
- (d) $\left\{ \begin{bmatrix} 1 & 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 & -1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 2 \end{bmatrix} \right\}$.
- (e) $\left\{ \begin{bmatrix} 1 & 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 & 2 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & -1 \end{bmatrix} \right\}$.

(3) Let V denote the vector space of 3×3 matrices endowed with the usual matrix addition and the usual product of matrices with numbers. Which ones of the subsets of V below are not subspaces of V ? (circle all that apply)

- (a) The set of diagonal 3×3 matrices.
- (b) The set of 3×3 matrices for which $\mathbf{A} \cdot \mathbf{A}^T \neq 0$.

- (c) The set of 3×3 matrices for which $\mathbf{A}^{-1} = \mathbf{A}^T$.
- (d) The set of 3×3 matrices for which $\mathbf{A} + 2\mathbf{A}^T = \mathbf{0}_{3 \times 3}$.
- (e) The set of 3×3 matrices for which $2\mathbf{A} + \mathbf{A}^T = \mathbf{I}$.

(4) Which statements below are true (circle all that apply)

- (a) Every vector in the null space of a matrix \mathbf{A} is necessarily orthogonal to every vector in the row space of \mathbf{A} .
- (b) There are vector spaces of dimension 0.
- (c) The row rank can sometimes differ from the column rank.
- (d) Orthonormal set of vectors in a basis can sometimes be linearly dependent

(5) What is the determinant of the matrix

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 0 & 3 \\ 2 & 3 & 0 \end{pmatrix}$$

- (a) 0
- (b) 3
- (c) 4
- (d) 12

(6) What is the dimension of the null space of the matrix below

$$\mathbf{A} = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 0 & 6 \\ 0 & 6 & 1 \end{pmatrix}$$

- (a) 0
- (b) 1
- (c) 2
- (d) 3

(7) The row echelon form of a matrix \mathbf{A} yields the following matrix

$$\begin{pmatrix} 1 & 0 & 3 & 1 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

then we can conclude that the row space of \mathbf{A} is of dimension

- (b) 0
- (c) 1
- (d) 2
- (e) 3
- (f) None of the above

(8) Which among the statements below are true about $n \times n$ matrix \mathbf{Q} with orthonormal columns? (circle all that apply)

- (a) $\mathbf{Q} \cdot \mathbf{Q}^T = \mathbf{I}$.
- (b) $\mathbf{Q}^{-1} = \mathbf{Q}^T$.
- (c) $\mathbf{Q} \cdot \mathbf{Q} = \mathbf{I}$.
- (c) $\mathbf{Q} + \mathbf{Q}^T = 2\mathbf{I}$.
- (d) None of the above.

II (20pts)

Answer the following questions by showing details of your work.

Let

$$\mathbf{A} = \begin{pmatrix} 1 & 0 & -1 & 2 \\ 1 & -1 & 0 & 0 \\ 4 & -5 & 1 & 2 \end{pmatrix}$$

(a) Find a basis for the null space of \mathbf{A} and the rank of \mathbf{A} .

Answer:

(b) Let $\mathbf{q}_1 = \frac{\sqrt{2}}{2} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$, $\mathbf{q}_2 = \begin{pmatrix} x_1 \\ 0 \\ x_2 \end{pmatrix}$ and $\mathbf{q}_3 = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$. Solve for x_1, x_2, y_1, y_2, y_3 so as to ensure that set of vectors below

$$\{\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3\}$$

form an orthonormal basis for $\mathbb{R}^{3 \times 1}$.

Answer:

(c) Let \mathbf{A} denote the matrix which collects the column vectors \mathbf{A} . Find a basis for a basis for the column space of \mathbf{A} formed by a subset of columns of \mathbf{A} .

Answer:

