

MACHINE LEARNING-BASED TRANSIENT BROKERS FOR REAL-TIME CLASSIFICATION OF THE LSST ALERT STREAM

GAUTHAM NARAYAN
STSCI
LASKER DATA SCIENCE FELLOW

 @gsnarayan



NOAO: Monika Soraisam, Abhijit Saha, Tom Matheson

UA CS: Rick Snodgrass, John Kececioglu, Carlos Scheidegger

UA Math/Statistics: Robert Maier

UA Astro: Tim Axelrod

CSS: Rob Seaman

LSST: Tim Jenness

Grads: Zhe Wang, Clark Taylor, Eric Welch,
Zhenge Zhao, Eric Evans,
Shuo Yang, Navdeep Singh

Former students: Tayeb Zaidi, Jackson Toeniskotter

B
I

NING-B
EAL-TIM
SST ALE

NT
ON


GA
STSCI

LASKER DATA SCIENCE FELLOW

 @gsnarayan



ANTARES



NOAO: [Monika Soraisam](#), Abhijit Saha, Tom Matheson

UA CS: Rick Snodgrass, John Kececioglu, Carlos Scheidegger

UA Math/Statistics: Robert Maier

UA Astro: Tim Axelrod

CSS: Rob Seaman

LSST: Tim Jenness

Grads: Zhe Wang, Clark Taylor, Eric Welch,
Zhenge Zhao, Eric Evans,
Shuo Yang, Navdeep Singh

Former students: [Tayeb Zaidi](#), Jackson Toeniskotter

Motivation




The LSST era is now! From this morning on Cerro Pachon.

We need useful alert brokers before first light.



Example broker with real stages and real data in Narayan+ (2018, submitted)

PS1 produced $O(10^2)$ new alerts per night.
We sorted them visually, and could follow ~5/night spectroscopically.


LSST will produce $O(10^7)$ per night.
There's no hope to sort this visually.
We can still follow ~5.



**Really big,
imbalanced, multi
class data set of time-
series goes in to
pipeline.**






Narayan et al, (2018, submitted) pipeline implements core processing of ANTARES



Use known Galactic stellar variability distribution from Kepler to characterize common variables vs explosive transients even with only a few observations (Ridgway et al, 2014)


Can be extended to multi-phase, multi-band (Soraisam et al, in prep)






Account for correlations in features.

Intermediate Phase Classification




● Double Periodic Variables ● Ellipsoidal Binaries ● δ Scuti ● Type II Cepheids ● Long Period Variables
● Double-mode Cepheids ● Supernovae ● Classical Cepheids ● Eclipsing Binaries ● RR Lyrae

t-SNE with Imbalanced Dataset



t-SNE with Rebalanced Dataset




t-SNE to visualize this
feature set

**SMOTE to deal with class imbalance
dramatically helps with distinguishing classes**

Intermediate-phase classification


Point estimates + PCA + imbalanced learning are pretty good at getting most of the large classes right, even with sparse light curves (~10 observations).

Probably useful for most science cases during LSST operations





Use Gaussian Processes to interpolate and model time-series observations




B1-392 / OGLE-BLG-RRLYR-11078




DES SN193775



Use wavelet transformation to describe shape



True Positive Rate



ROC Curves

Very comparable to Lochner '16 for
SNPhotCC. Real heterogenous data is
not like homogenous sims.

Take aways

- If you are visually inspecting survey data to find transients and variables, we can help.
- Narayan, Zaidi, Soraisam et al (2018, ApJS, submitted)
- Next: Apply to PS1 MDS light curves + contextual information, and ongoing Kepler K2 C16.
- Apply to PLAsTiCC simulated data, ZTF public alerts (~April)
- Postage stamps as features with CNNs, unsupervised learning techniques
- **WE ARE HIRING!**
Associate or Assistant Scientist at NOAO
(<https://goo.gl/bceKu7>)

