



Equipped for success: Genomes and metabolomes of the European *Amanita muscaria* are conserved in its novel South African range

Journal:	<i>New Phytologist</i>
Manuscript ID	Draft
Manuscript Type:	Full Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Nickles, Grant; University of Wisconsin-Madison, Medical Microbiology and Immunology Stokes, Cecelia; University of Wisconsin-Madison, Botany Narh, Deborah; University of Pretoria Forestry and Agricultural Biotechnology Institute, Departments of Biochemistry, Genetics and Microbiology; Stanford University, Department of Biology Lynn, Kira; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology Fuqua, Savannah; University of Arizona, Department of Ecology and Evolutionary Biology Bryan, Corbin; University of Wisconsin-Madison, Departments of Botany and Bacteriology Allen, Brooke; University of Mississippi, Department of Biology, College of Liberal Arts Bivins, Christopher; University of California Merced, School of Natural Sciences Woo Bok, Jin; University of Wisconsin-Madison, Department of Medical Microbiology and Immunology Brewer, Stephen; University of Mississippi, Biology; Buthlezi, Sikelela; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology Clark, Jahiya; University of California Santa Cruz, Ecology and Evolutionary Biology Department Coon, Keri; University of Wisconsin-Madison, Department of Bacteriology Corby, Lauren; Jackson State University, Department of Chemistry and Biochemistry Coetzee, Martin; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology Dewing, Claudette; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology Duong, Tuan; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology</p>

	<p>Harris, Mathew; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Plant and Soil Sciences</p> <p>Keller, Nancy; University of Wisconsin, Medical Microbiology and Immunology</p> <p>Kopotsa , Katlego; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology</p> <p>Lane, Frances; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology</p> <p>Nichols, Holly; University of Wisconsin-Madison, Department of Bacteriology; University of Wisconsin-Madison, Microbiology Doctoral Training Program</p> <p>Nieuwoudt, Alandie; University of Pretoria Forestry and Agricultural Biotechnology Institute, Departments of Biochemistry, Genetics and Microbiology</p> <p>Nunez, Martin; INIBIOMA, Grupo de Ecología de Invasiones</p> <p>Medina Munoz, Miguel; University of Wisconsin-Madison, Department of Bacteriology</p> <p>Park, Sung Chul; University of Wisconsin-Madison, Department of Medical Microbiology and Immunology</p> <p>Pham, Nam; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology</p> <p>Ryan, Kaetlyn; University of Wisconsin-Madison, Department of Pathobiological Sciences</p> <p>Solís, Myriam; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology</p> <p>Vilgalys, Rytas; Duke University, Biology Department</p> <p>Wallace, Jasmine; Mississippi Valley State University, Department of Natural Sciences and Environmental Health</p> <p>Wang, Yen-Wen; Yale University, Yale School of Public Health</p> <p>Wingfield, Brenda; University of Pretoria Forestry and Agricultural Biotechnology Institute, Department of Biochemistry, Genetics and Microbiology</p> <p>Wingfield, Mike; University of Pretoria Forestry and Agricultural Biotechnology Institute, FABI</p> <p>Worley, Travis; University of Wisconsin-Madison, Department of Bacteriology; University of Wisconsin-Madison, Microbiology Doctoral Training Program</p> <p>Zallek, Taylor; University of Pittsburgh, Department of Biological Sciences</p> <p>Zamanian, Mostafa; University of Wisconsin-Madison, Department of Pathobiological Sciences</p> <p>Hoeksema, Jason; University of Mississippi, Department of Biology</p> <p>Drott, Milton; USDA-ARS Cereal Disease Laboratory, Plant Pathology</p> <p>Pringle, Anne; University of Wisconsin-Madison, Botany and Bacteriology</p>
Key Words:	Biosynthetic gene clusters, bioactivity, invasive mushrooms, Global Natural Product Social Molecular Networking (GNPS) analysis, nematocidal, primary or specialized metabolism, South Africa
Suggested Handling Editor:	

Data Availability Statement

Our [Mandates Data Policy](#) requires data to be shared and a Data Availability Statement, so please enter one in the space below. Sample statements can be found [here](#). Please note that this statement will be published alongside your manuscript, if it is accepted for publication.

All genomic data analyzed in this study were generated by the authors. Raw reads and assemblies are publicly available through NCBI BioProject [###]. Other data and methods enabling replication are provided with this manuscript and in the Supporting Information (which contains Supporting Methods, Tables, and Figures).

For Peer Review

Equipped for success: Genomes and metabolomes of the European *Amanita muscaria* are conserved in its novel South African range

AUTHORS & AFFILIATIONS

First: Grant R. Nickles¹

Second:

Cecelia K. Stokes^{2,3}, Deborah L. Narh^{4,5}, Kira M.T. Lynn⁴, Savannah R. Fuqua⁶, Corbin Bryan²

Middle:

Brooke M. Allen⁷, Christopher P. Bivins⁸, Jin Woo Bok¹, J. Stephen Brewer⁷, Sikelela T. Buthelezi⁹, Jahiya P.R.M. Clark¹⁰, Kerri L. Coon¹¹, Lauren R. Corby¹², Martin P.A. Coetzee⁴, Claudette Dewing⁴, Tuan A. Duong⁴, Mathew A. Harris^{9,13}, Nancy P. Keller^{1,14}, Katlego Kopotsa^{9,15}, Frances A. Lane⁴, Holly L. Nichols^{11,16}, Alandie Nieuwoudt⁹, Martin A. Nuñez¹⁷, Miguel E. Medina Munoz¹¹, Sung Chul Park¹, Nam Q. Pham⁴, Kaetlyn T. Ryan¹⁸, Myriam Solís⁴, Rytas Vilgalys¹⁹, Jasmine Marea Wallace²⁰, Yen-Wen Wang²¹, Brenda D. Wingfield⁴, Michael J. Wingfield⁹, Travis K. Worley^{11,16}, Taylor A. Zallek²², Mostafa Zamanian¹⁸

Last: Jason D. Hoeksema⁷, Milton T. Drott²³, Anne Pringle^{2,11}

ORCID:

GRN 0000-0002-0234-8049, CKS (0009-0006-7257-1947), DLN 0000-0003-2000-2321, KMTL 0000-0003-0533-0458, SRF 0000-0001-5139-9603, CTB 0009-0006-8075-9018, BMA 0000-0002-6000-9061, CPB 0000-0001-6228-8212, JWB 0000-0002-4394-2749, JSB 0000-0002-6720-242X, STB (none), JPRMC 0009-0007-3563-6796, KLC 0000-0002-2701-5195, LRC 0000-0002-2928-8154, MPAC 0000-0001-7848-4111, CD 0000-0001-6208-1721, TAD 0000-0001-5110-1854, MAH 0000-0003-4310-1146, NPK 0000-0002-4386-9473, KK 0000-0003-4062-9762, FAL 0000-0002-9103-7938, HLN 0000-0003-2158-1767, AN (none), MAN 0000-0003-0324-5479, MEMM 0000-0001-6932-3105, SCP 0009-0008-9081-7992, NQP 0000-0002-4938-9067, KTR 0000-0002-2051-7969, MS (none), RV 0000-0001-8299-3605, JMW (none), YWW 0000-0002-6851-1646, BDW 0000-0002-6189-1519, MJW 0000-0001-9346-2009, TKW 0009-0007-7065-0032, TAZ 0009-0006-8137-6449, MZ 0000-0001-9233-1760, JDH 0000-0003-2338-8442, MTD 0000-0001-9715-2200, AP 0000-0002-1526-6739

KEYWORDS

Biosynthetic gene clusters, bioactivity, invasive mushrooms, invasive ectomycorrhizal fungi, Global Natural Product Social Molecular Networking (GNPS) analysis, fly agaric, nematicidal, primary or specialized metabolism, secondary metabolites, South Africa

SUMMARY

- Plants and soils have been moved around the world for centuries, but invasive mushrooms receive scant attention. The *Amanita muscaria* species complex was introduced to South

Africa in the context of forestry, but its origins, ecology, and recent evolution are unstudied.

- We sequenced the genomes of 24 Northern and Southern Hemisphere *A. muscaria*, built phylogenies, and reconstructed its South African history. We identified the biosynthetic gene clusters (BGCs) encoding specialized metabolites (SMs). We subsequently extracted mushrooms' metabolites and used mass spectrometry data to group SMs into unique molecular families (MFs). We tested metabolites for bioactivity against diverse microbes and animals.
- We identify Europe as the origin of South African *A. muscaria*. A highly conserved suite of BGCs are found in nearly all European and African genomes, and only 13 of 273 MFs are unique to South Africa. Metabolites extracted from all mushrooms kill nematodes, while microbes and flies appear unaffected.
- The nearly global distribution of the fly agaric results from multiple introductions of a single European clade to the Southern Hemisphere. Despite its long history in South Africa, the fungus has not lost any of its BGCs, suggesting a conservation of function(s) across multiple continents.

WORD COUNTS

Introduction: 1,483 → 1,057

Box 1: 516 → 384

Materials and Methods: 2,612

Results: 3,166 → 1,586 (figure captions 792 → 712).

Discussion: 1,418

No. of Tables: 2

No. of Figures: 5

Supporting Information: 2,977

INTRODUCTION

Humans have moved plants and soils among continents for centuries, resulting in the co-introduction of fungal pathogens and symbionts (Anagnostakis, 1987; Dickie *et al.*, 2010; Cleary *et al.*, 2016). Fungi mediate basic ecological processes (Treseder & Lennon, 2015) and when introduced fungi become invasive, they can dramatically alter local ecosystems (Anagnostakis, 1987; Desprez-Loustau *et al.*, 2007; Rossman, 2009; Boyd *et al.*, 2013). Interest in invasive nonpathogenic fungi is growing (Pringle *et al.*, 2009; Dickie *et al.*, 2010; Vargas *et al.*, 2019; Milani *et al.*, 2022), but most research continues to focus on the biology and impacts of introduced and invasive pathogens. As mutualistic fungi spread across the globe (Vellinga *et al.*, 2009; Berch *et al.*, 2017; Vargas *et al.*, 2019; Pildain *et al.*, 2021), discovering the mechanisms enabling their successful invasions and documenting impacts on native ecosystems emerge as key research priorities.

The most intensively studied plant-fungal co-invasions involve fungi and plant species of *Pinus* (family Pinaceae). Pines have been introduced across the Southern Hemisphere for commercial forestry, reforestation, and afforestation. These non-native trees now cover more than 6 million hectares in countries including South Africa, Argentina, Brazil, and New Zealand (Rouget *et al.*, 2001; Simberloff *et al.*, 2010; Dickie *et al.*, 2014; Brancatelli *et al.*, 2020). Species like *Pinus elliottii*, *P. patula*, and *P. radiata* have spread beyond forestry plantations and into natural forest and grassland biomes, including into biodiversity hotspots, for example, South Africa’s fynbos (Richardson & Higgins, 1998; van Wilgen & Richardson, 2012). Invading pines can reduce water availability, suppress native plant and fungal species, and disrupt ecosystem services (Le Maitre *et al.*, 2002; Dickie *et al.*, 2014; Nuñez *et al.*, 2017; Brewer *et al.*, 2018; Sapsford *et al.*, 2022).

Pines are obligately associated with ectomycorrhizal (ECM) fungi, a symbiosis enabling the trees to obtain essential nutrients and water (Smith & Read, 2008; Nuñez *et al.*, 2009; Dickie *et al.*, 2010). Historically, plantings of pines in the Southern Hemisphere failed until soils were imported to supply their symbionts (Mikola, 1970), and introduced ECM fungi have subsequently facilitated invasions of pine trees (Hayward *et al.*, 2015; Policelli *et al.*, 2019; Policelli *et al.*, 2023). Despite the critical role of ECM fungi in pine invasions and the potential for ECM fungi to invade and harm native forests on their own (Dickie *et al.*, 2017), the ecological and evolutionary dynamics of invading ECM fungi remain poorly understood, perhaps because invading mutualists are perceived as unequivocally beneficial (but see Schwartz *et al.*, 2006). In fact, ECM fungi may function across a spectrum of mutualism and parasitism (Karst *et al.*, 2008). Efforts to elucidate the ecology of ECM fungi are further exacerbated by their ephemeral and often cryptic biology (Wang *et al.*, 2023), leaving us with incomplete, often inaccurate historical records of species’ native ranges (Pringle & Vellinga, 2006). Strategies combining historical research with modern ‘omics methods offer a novel approach to identify the drivers of ECM fungal invasions.

Invasion biology often invokes interactions among species as a mechanism driving spread, and fungal interactions are often hypothesized to be mediated by specialized metabolites (SMs) (Drott *et al.*, 2017; Keller, 2019; Tannous *et al.*, 2022). Although ECM fungi produce a plethora of SMs (Schöffler, 2018; Walton, 2018; Obermaier & Müller, 2020; Mudbhari *et al.*, 2024) and must routinely interact with pathogens, predators, and competitors, interactions between

introduced ECM fungi and local organisms are rarely considered. However, the toxins of at least one invasive ECM fungus, *Amanita phalloides*, appear to be evolving dynamically in its invasive range (Drott *et al.*, 2023). This finding and the ubiquity of SMs raise questions about the translation of the “enemy release” and “novel weapons” hypotheses, often invoked to explain the success of non-native plants and animals, to invasive fungi (Blossey & Nötzold, 1995; Torchin *et al.*, 2003; Callaway & Ridenour, 2004; Liu & Stiling, 2006). The “enemy release” hypothesis proposes introduced species succeed by escaping natural enemies, while the “novel weapons” hypothesis suggests they spread by producing compounds against which native species have no resistance (Gillett, 1962; Callaway & Ridenour, 2004).

The ecological roles and evolution of SMs remain poorly understood, even in iconic ECM species like *Amanita muscaria* sensu lato (s.l.) (Geml *et al.*, 2006), the famous red-and-white-spotted “fly agaric” known for its psychoactive properties (Camazine, 1983; Michelot & Melendez-Howell, 2003) and for the folklore suggesting it can repel or kill flies (Lumpert & Kreft, 2016). *A. muscaria* produces ibotenic acid and muscimol using a single biosynthetic pathway (Størmer *et al.*, 2004; Su *et al.*, 2023), and muscimol is a potent GABA_A receptor agonist (Johnston, 2014), a feature which contributes to its distinctive biological effects. Native to nearly all temperate and boreal biomes of the Northern Hemisphere, the *A. muscaria* species complex includes multiple cryptic lineages associated with diverse plants, including pines. While some of these cryptic species are now formally recognized (for example, *A. persicina* (Tulloss *et al.*, 2015)), the true number of cryptic species within the *A. muscaria* species complex remains unknown. For simplicity, we refer to all as-yet unnamed lineages as *A. muscaria*. Introduced to South America, Australia, and Africa, *A. muscaria* forms novel ECM partnerships with native trees in Columbia (*Quercus humboldtii*; (Vargas *et al.*, 2019), Chile (*Nothofagus*; (Márquez Parraguez, 2024) and Australia (*Nothofagus* and *Allocasuarina*; (Fuhrer, 1992; Dunk *et al.*, 2012; Lebel *et al.*, 2024). Relatively little is known about its introductions across Africa (Vellinga *et al.*, 2009), but its South African history is particularly well documented (Box 1), offering a rare opportunity to study the impact of *A. muscaria*’s introduction on the fungus’s evolution and production of SMs.

We sequenced the genomes of 24 *A. muscaria* s.l. mushrooms (9 South African, 11 European, 2 Australian, and 2 Californian) and 1 outgroup (*Amanita pantherina*) to address three hypotheses: First, South African *A. muscaria* would be most closely related to a European lineage of the fungus. Our hypothesis is based on the history of *A. muscaria* in South Africa (Box 1). Second, a release from antagonistic interactions found in Europe but absent in South Africa would relax selection on specialized metabolites, enabling the fungus to invest more energy into growth and reproduction and potentially causing the loss of SM genes or gene function. Third, adaptation in the invasive range would result in shifts in metabolic profiles and impact the bioactivity of extracts against laboratory model bacteria, fungi, nematodes, mosquitoes, and flies.

BOX 1: The history of *Amanita muscaria* in South Africa

Doidge (Doidge, 1950) lists a number of early collections of the fungus; by 1874 *A. muscaria* was growing with pines in and around Cape Town (Fig. 1). By 1945 it was growing in what are now named as

the provinces of Mpumulanga to the north and Free State in the east. She identifies *A. muscaria* as growing with “shrubs”, pine trees, and “near living roots of *Quercus* sp.” (Doidge 1950 p. 553).

The history of *A. muscaria* is linked to the history of Europeans on the continent. Early European colonists did not find the timber species native to South Africa suitable for their uses, and they introduced fast-growing pines. Any plant moved to South Africa by European colonists would have been accompanied by fungi in its tissues, and plants may also have been moved as seedlings growing in soil, enabling the introduction of soil fungi. Doidge (1950) succinctly describes the dynamic of plants arriving to South Africa with Europeans,

“The history of plant introduction into South Africa begins in 1651, with the arrival of Johan van Riebeeck to establish a settlement at the Cape; van Riebeeck was accompanied by a gardener... To provide timber, sacks of acorns were sent from Holland and kernels of stone and cluster pines from Italy; these were planted in quantity and flourished amazingly... the majority being species of *Pinus*, *Eucalypts* and *poplars*...” Doidge (1950 pp. 15-16)

Mikola (1970) goes on to describe the subsequent deliberate export of soils from South Africa to enable plantation forestry elsewhere, including in Zimbabwe and Kenya (Mikola 1970; Zimbabwe discussed as Rhodesia). For example, three tons of soil from the Transvaal were imported to establish pine plantations in Eswatini (formerly Swaziland), and apparently *A. muscaria* was moved as well, “The commonest mushrooms in Usutu Forest of Swaziland [now Eswatini] in March 1967 were... *Laccaria laccata*, *Amanita muscaria*, *Boletus edulis*, *Rhizopogon roseolus*.” (Mikola, 1970).

The fungus is acknowledged as a European introduction in the most recent scientific literature on the *Amanita* of South Africa (Reid & Eicker, 1991) and in a current field guide (Goldman & Gryzenhout, 2019). While *A. muscaria* may have been introduced with European “stone and cluster pines” (Doidge, 1950), in modern South African forestry plantations it associates with pines from other continents, for example Mexican *Pinus patula* and Californian *P. radiata*.

.....

MATERIALS AND METHODS

Historical Records of South African *A. muscaria*

Historical records of *A. muscaria* from South Africa (from before 1945) were obtained from Doidge (1950). To compare the historical records with the current distribution of the fungus, we downloaded all publicly available South African records from the Global Biodiversity Information Facility (GBIF) on 24 July 2023 (GBIF, 2023). We looked at the photographs linked to each record to identify surrounding tree species. Because every record suggested an association of *A. muscaria* with *Pinus* spp., we extracted information about plantation forestry land cover within South Africa from the 2022 South African National Land Cover dataset (Department of Forestry, Fisheries and Environment). We mapped the historical and current records of *A. muscaria*. We note historical records often use older place names for both the country and its provinces, for example Transvaal for Gauteng or Mpumalanga.

Mushroom Collections

We sequenced genomes from a total of 24 *A. muscaria* and 1 *A. pantherina* mushrooms collected from South Africa, Australia, Europe and North America (Table 1). South African ($n = 9$) and Australian ($n = 2$) specimens were collected from pine plantations while European *A. muscaria* ($n = 11$), *A. pantherina* ($n = 1$) and Californian ($n = 2$) mushrooms were collected from native

forests (Table 1; Fig. S1). Specimens were dried within 1-2 days using conventional food driers at low heat (less than 35 °C). Our sampling scheme was designed to contextualize South African *A. muscaria* by integrating new data within published phylogenies, primarily Geml et al. (2008); a comprehensive survey of the global species complex was beyond our scope. We targeted as many South African specimens as possible, European specimens from three countries (based on what we found and could collect), and other specimens from additional clades identified by Geml et al. (2008), for example Clade I, represented by our two Californian mushrooms. Out of curiosity, we also sequenced two Australian specimens. Prior to sequencing, all specimens were stored as dried mushrooms at room temperature in either the Pringle or Vilgalys Laboratory fungaria at the University of Wisconsin-Madison and Duke University, respectively.

Table 1: Collection metadata for the specimens collected and analyzed.

Specimen ID	Collection Date	Location	Region	Country	Latitude	Longitude	Species	Habitat	Host	Collector
11662	27-Feb-19	Bulwer	KwaZulu-Natal	South Africa	29.8077 S	29.76 E	<i>A. muscaria</i>	plantation	Pinus patula	J. Roux
11663	27-Feb-19	Bulwer	KwaZulu-Natal	South Africa	29.8077 S	29.76 E	<i>A. muscaria</i>	plantation	Pinus patula	J. Roux
11664	27-Feb-19	Bulwer	KwaZulu-Natal	South Africa	29.8077 S	29.76 E	<i>A. muscaria</i>	plantation	Pinus patula	J. Roux
11665	6-Mar-19	Bulwer	KwaZulu-Natal	South Africa	29.8077 S	29.76 E	<i>A. muscaria</i>	plantation	Pinus patula	J. Roux
11666	25-Apr-19	Kranskop / Greytown	KwaZulu-Natal	South Africa	28.9667 S	30.8642 E	<i>A. muscaria</i>	plantation	Pinus eliottii	I. Greyling
11667	25-Apr-19	Kranskop / Greytown	KwaZulu-Natal	South Africa	28.9667 S	30.8642 E	<i>A. muscaria</i>	plantation	Pinus eliottii	I. Greyling
11668	25-Apr-19	Kranskop / Greytown	KwaZulu-Natal	South Africa	28.9667 S	30.8642 E	<i>A. muscaria</i>	plantation	Pinus eliottii	I. Greyling
11669	25-Apr-19	Kranskop / Greytown	KwaZulu-Natal	South Africa	28.9667 S	30.8642 E	<i>A. muscaria</i>	plantation	Pinus eliottii	I. Greyling
11670	25-Apr-19	Kranskop / Greytown	KwaZulu-Natal	South Africa	28.9667 S	30.8642 E	<i>A. muscaria</i>	plantation	Pinus eliottii	I. Greyling
Road_3	12-Sep-19	Østfold	Østlandet	Norway	59.4237 N	10.5937 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Sogn_5	15-Sep-19	Oslo	Østlandet	Norway	59.9822 N	10.7415 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Skrup_3	17-Sep-19	Oslo	Østlandet	Norway	59.8657 N	10.8562 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Nes_1	17-Sep-19	Akershus	Østlandet	Norway	59.8666 N	10.5181 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Frag_1	18-Sep-19	Oslo	Østlandet	Norway	59.9820 N	10.6742 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Ring_1	6-Oct-19	Zurich	Zurich	Switzerland	47.3617 N	8.4853 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Wirz_3	7-Oct-19	Nidwalden	Nidwalden	Switzerland	46.9053 N	8.3696 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Gril_1	7-Oct-19	Nidwalden	Nidwalden	Switzerland	46.9010 N	8.3571 E	<i>A. muscaria</i>	native habitat	Not recorded	S.L. Harrow
Kara_3	7-Oct-19	Zurich	Zurich	Switzerland	47.3617 N	8.4853 E	<i>A. muscaria</i>	native habitat	Not recorded	K. O'Keefe
Nagy_Heves_A	12-Oct-13	Heves	Heves	Hungary	47.3182 N	13.8414 E	<i>A. muscaria</i>	native habitat	Picea abies or Betula pendula	L. Nagy
Nagy_Heves_B	12-Oct-13	Heves	Heves	Hungary	47.3173 N	13.3415 E	<i>A. muscaria</i>	native habitat	Picea abies or Fagus sylvatica	L. Nagy
20031	17-Dec-21	Point Reyes National Seashore	California	United States	38.0519 N	122.8310 W	<i>A. muscaria</i>	native habitat	Not recorded	Wang/Pringle
20045	19-Dec-21	Point Reyes National Seashore	California	United States	37.9702 N	122.7304 W	<i>A. muscaria</i>	native habitat	Not recorded	Wang/Pringle
NzAUS95	27-May-19	Railton	Tasmania	Australia	41.2231 S	146.236 E	<i>A. muscaria</i>	plantation	Pinus radiata	Vilgalys/Henderson/Uehling
Aus332	31-Mar-18	Penrose	NSW	Australia	34.3748 S	150.1241 E	<i>A. muscaria</i>	plantation	Pinus radiata	Vilgalys/Henderson/Uehling
Nes_pan3	17-Sep-19	Akershus	Østlandet	Norway	59.8666 N	10.5181 E	<i>A. pantherina</i>	native habitat	Not recorded	S.L. Harrow

Genome Extraction, Sequencing, Assembly, Annotation, and Quality Control

DNA extraction and sequencing

High molecular weight DNA (HMW-DNA) was extracted from samples of dried mushroom tissues using previously published protocols involving a solution of 25:24:1 phenol/chloroform/isoamyl alcohol (Bok et al., 2005; Nickles et al., 2023). Sequencing was performed with an Illumina NovaSeq 6000 platform with the aim of obtaining six million reads per sample, to achieve a coverage of 30x or higher. We estimated the genome size as 44 Mb.

Genome assembly, annotation, and quality analyses

Raw sequencing data were trimmed with Trimmomatic v0.36 (Bolger et al., 2014) with LEADING:20 TRAILING:20 SLIDINGWINDOW:4:25 MINLEN:50, and genomes were assembled with SPAdes v3.11.1 (Bankevich et al., 2012; Prjibelski et al., 2020) using default parameters. Assemblies were filtered to keep only the contigs/scaffolds larger than 500 bp. Assembled genomes were annotated with Augustus v3.4.0 using models pre-built for *Laccaria bicolor* (Stanke et al., 2008; Hoff & Stanke, 2013). Genome statistics of each genome were generated using seqtk (Li, 2013) and Benchmarking Universal Single-Copy Orthologs (BUSCO) (Simao et al., 2015) completeness was determined using BUSCO v3.0.2 with the agaricales_odb10 database (Manni et al., 2021).

Prior to examining the assembled genomes, basic assembly metrics were determined using seqtk. To screen for potential fungal contamination, duplicate BUSCO genes were identified with BUSCO (v3.0.2) run in genome mode using the Fungi Odb10 database with default settings. Only one genome showed a single duplicated BUSCO gene, suggesting all genomes were free from fungal contaminants. The first 200 nucleotides of each contig from each fully assembled genome was then used to query NCBI's prokaryote nucleotide database (accessed July 27, 2023) in a local BLASTn search with default settings using NCBI's BLAST+ software package (v2.6.0) (Altschul *et al.*, 1990; McGinnis & Madden, 2004). Contigs with matches showing an e-value less than 1e-5 were considered contaminated and removed from each genome. Genome quality metrics were then recomputed with seqtk. The decontaminated genomes were used for all subsequent analyses (Dataset S2).

Generating and filtering SNP predictions

Short reads were quality controlled with BMap v38.32 (Bushnell, 2015) and aligned to the NCBI reference genome (accession GCA_001691765.1) using BWA mem v0.7.17 (Li & Durbin, 2009). Variants were called using Genome Analysis Toolkit v4.0.12.0 (McKenna *et al.*, 2010) using parameters and hard filters identical to Drott *et al.* (2023). Single nucleotide polymorphism (SNP) data were filtered using VCFtools (v0.1.13) (Danecek *et al.*, 2011) such that only biallelic sites were retained. Split decomposition was employed to reconstruct the phylogenetic tree using the filtered SNP dataset to identify potential errors or artifacts in the dataset (Supplemental Methods).

Clone Correction and Population Genetics

Clone correction

To test if different specimens are clones of each other, i.e., mushrooms generated from a single mycelium, or are distinct genetic individuals, we estimated pairwise kinships using KING-robust (Manichaikul *et al.*, 2010) using the function snpgdsIBDKING implemented in the R library SNPRelate.

Genetic differentiation and estimation of gene flow

Pairwise F -statistics (F_{ST}) and Euclidian distances (D) between individuals as well as the identification of genetic variants and private alleles between and within population were calculated or undertaken using the dartR package in R (Gruber *et al.*, 2018). Tajima's D was estimated in 5,000 bp sliding windows using VCFtools (v0.1.13) (Danecek *et al.*, 2011) with a subset of Clade II genomes from Europe and South Africa. Nucleotide diversity (π) was calculated across the entire genome using pixy (Korunes & Samuk, 2021) and heterozygosity was calculated across all variant sites using VCFtools.

Phylogenomic Analyses

Nomenclature of the A. muscaria species complex and closely related species and downloading of publicly available A. muscaria gene markers

We contextualized newly sequenced genomes using sequences of the internal transcribed spacer (ITS), α -tubulin (β -tub), nuclear large ribosomal subunit (LSU), and translation elongation factor 1-alpha (TEF-1 α) genes from three key publications (Oda *et al.*, 2004; Geml *et al.*, 2008;

Vargas *et al.*, 2019) (Dataset S1). Specimen ID codes from Oda *et al.* (2004) were shortened to only include the numeric ID. The taxonomy and nomenclature of *A. muscaria* s.l. is complex, and we used published literature to develop naming protocols.

Following recent precedent (Vargas *et al.*, 2019), we used Geml *et al.*'s (2008) clade labels to identify our specimens. While taxonomists used to consider *A. muscaria* as a complex of subspecific varieties existing under a single species concept and identifiable on the basis of cap color (Miller & Jenkins, 1978), when specimens with different cap colors were integrated into single phylogenies using DNA sequence data, new species concepts emerged (Oda *et al.*, 2004; Geml *et al.*, 2006). Geml *et al.* (2008) represents the last major phylogenetic revision of the taxa, but instead of naming clades as species, Geml *et al.* (2008) chose to label distinct clades with roman numerals. We use the same numbering system for our specimens and for the specimens associated with already published data (Oda *et al.*, 2004; Geml *et al.*, 2008; Vargas *et al.*, 2019), with two exceptions: after Geml *et al.* (2008) was published, *A. persicina* was formally recognized as a species, and we use its name (Tulloss *et al.*, 2015). We also use the name *A. regalis* because it was formally published in a previous revision of the species complex (Neville & Poumarat, 2004) and is supported by the later phylogenetic evidence (Geml *et al.*, 2008).

We used the same logic to identify specimens associated with other public data. Any specimen originally classified as and submitted to GenBank as one of the widely recognized and currently accepted infraspecific taxa of *A. muscaria* was given the working designation *A. muscaria* s.l. and later identified using Geml *et al.*'s (2008) clade labels. For our purposes, *A. muscaria* s.l. includes *A. muscaria* var. *guessowii*, *A. muscaria* var. *muscaria*, and *A. muscaria* var. *flavivolvata*. However, if a specimen had a GenBank accession for ITS locus sequence data and was submitted as *Amanita regalis* (or *A. muscaria* var. *regalis*) or *Amanita persicina* (or *A. muscaria* var. *persicina*) we used the modern nomenclature (Tulloss & Yang, 2016), in other words the names *A. regalis* or *A. persicina*. These two taxa are easily differentiated from *A. muscaria* s.l. on the basis of gross morphology. Specimens classified as *A. pantherina* were kept as such. For the two specimens lacking an ITS accession, the metadata associated with the LSU accession was used instead. All collection location data were extracted from the publication corresponding to each given collection.

Species tree construction using multi-loci barcodes

We aligned ITS (n = 189), β -tubulin (n = 102), LSU (n = 98), and TEF-1 α (n = 52) sequences with MAFFT v7.511 (Katoh & Standley, 2013) with an automated model selection (L-INS-i, FFT-NS-i and FFT-NS-2) and adjustment of reverse complements. Alignments were trimmed with Trimal v1.4.rev15 (Capella-Gutierrez *et al.*, 2009) with the gappyout parameter. Once trimmed, alignments for all four regions were used to construct a maximum likelihood phylogeny using an edge-linked partition model with IQ-TREE multicore version 2.2.0 (Minh *et al.*, 2020), with the optimal substitution model determined by ModelFinder (Kalyaanamoorthy *et al.*, 2017) and 1,000 rapid bootstraps.

Specialized Metabolite Biosynthetic Gene Cluster Prediction and Characterization

Genome mining for specialized metabolite BGCs

To identify canonical specialized metabolite BGCs (i.e., those with experimentally vetted class-defining core biosynthetic genes [termed backbone genes], e.g., polyketide BGCs, which are a

canonical class of SM defined by polyketide synthase backbone genes), Fungal antiSMASH (v5) (Blin *et al.*, 2019) was run on all of the genomes using the default settings. (However, specimen 11662 was erroneously omitted from genome mining analyses.) Closely related terpene or PKS BGCs were networked into gene cluster families (GCFs) using BiG-SCAPE v1.0.1 (Navarro-Munoz *et al.*, 2020). An optimal networking cutoff of 0.3 was determined across a testing range of 0.1 to 0.6 (Fig. S2 and S3).

We ran cblaster (v1.3.18) (Gilchrist *et al.*, 2021) using a previously characterized ibotenic acid BGC sequence as the query to identify the non-canonical ibotenic acid BGC (Obermaier & Müller, 2020), which is made up of the genes (from left to right) *iboA*, *iboF*, *iboD*, *iboC*, *iboG1*, *iboH*, and *iboG2*. We further investigated the distribution of genes in the ibotenic acid BGC using reciprocal best-hit blast analysis. Briefly: characterized protein sequences were used to replace corresponding protein predictions of IboA, IboC, IboD, IboF, IboG1, IboG2, and IboH (Kohler *et al.*, 2015) in the file containing the proteome of mushroom Gril_1. The resulting proteome was queried against a dataset of all annotated publicly available protein sequences downloaded from NCBI in the spring of 2023 with methods similar to those described previously (Drott *et al.*, 2020). Any two hits that were within 30 kb of each other were considered physically clustered. The presence of a putative ortholog was mapped to a modified version of the whole-kingdom phylogeny from Nickles *et al.* (2023) using ggtree (Yu *et al.*, 2017).

We searched for the presence of amatoxin-encoding “MSDIN” genes and the toxin-processing *popB* gene using methods and publicly available scripts from (Drott *et al.*, 2023), as detailed in the Supplemental Methods.

Phylogeny of Agaricomycete polyketide synthase (PKS) proteins

A representative query protein was selected from the two conserved PKSs found within the *A. muscaria* genomes. Each genome’s protein was BLASTp (Altschul *et al.*, 1990; Schaffer *et al.*, 2001) searched against every annotated Agaricomycetes genome publicly available on NCBI as of December 1, 2022 ($n = 292$ genomes) with an e-value cutoff of $1e^{-5}$. Only the top hit for each BLAST search was retained for further analysis. A gene tree of resulting sequences was constructed with the same methodology used to construct the species tree. To confirm if the observed PKS duplication was unique to the *A. muscaria* species complex, we also conducted a targeted analysis of the *Amanita* genus. Genes encoding all PKS enzymes were identified from the six available non-*muscaria* *Amanita* genomes. The resulting 18 PKS protein sequences were used to construct a gene tree with the same methodology described to construct the species tree. Domain analysis revealed a single hybrid NRPS-PKS protein in *A. inopinata* (KAF8636077.1), which was subsequently used to root the tree.

Metabolomics for Chemical Identification of Metabolites

Methanolic extraction of metabolites

We extracted metabolites from nearly every dried mushroom that had its genome sequenced ($n = 22$); the two Australian specimens (Aus332 and NzAUS95) were not included in metabolomics because there were insufficient tissues left. Two additional North American (20031-a and 20045-b) and one additional South African (11671) mushrooms whose genomes were not sequenced were assayed, bringing the total number of metabolite extracts to 26 (*A. muscaria*: 25 and *A. pantherina*: 1; Table S1).

All samples were imaged prior to extraction (see Fig. S1). Mushroom caps were pulverized and extracted overnight in 20-100 mL (depending on the volume of the mushroom tissue) of HPLC-grade methanol. Extracts were filtered with a 0.2 μ m syringe filter and evaporated to dryness on a rotary evaporator. Compounds were resuspended at 1 mg/mL in methanol or dimethylsulfoxide (DMSO) for chemical analysis or biological assays, respectively.

UHPLC–MS/MS analysis

Mushroom cap extracts were characterized on Ultra-high-performance liquid chromatography coupled with tandem mass spectrometry (UHPLC–MS/MS) using only spectroscopic grade solvents. Data was acquired using a Thermo Fisher Scientific Q Exactive Orbitrap mass spectrometer (Waltham, MA, USA) coupled to a Vanquish UHPLC (Waltham, MA, USA) operated in positive ionization mode. For all runs, we used a Waters XBridge BEH-C18 column (2.1 mm \times 100 mm, 1.7 μ m) and added 0.05% formic acid to our spectroscopic grade acetonitrile and water (flow rate 0.2 mL/min). The 37 min screening gradient method for the samples is as follows: Starting at 10% organic for 5 min, followed by a linear increase to 90% organic over 20 min, another linear increase to 98% organic for 2 min, holding at 98% organic for 5 min, decreasing back to 10% organic for 3 min, and holding at 10% organic for the final 2 min.

Global Natural Product Social Molecular Networking (GNPS) analysis

To construct a molecular network, the raw mass spectra were converted to the mzXML format using RawConverter software (ver.1.2.01., The Scripps Research Institute) (He *et al.*, 2015). We utilized GNPS analysis, the web-based server, with the default parameters provided by the platform: Min pairs cos = 0.7, network TopK = 10, max connected components size = 100, min matched fragment ions = 6, min cluster size = 2. A molecular network was created with a precursor ion mass tolerance of 2.0 Da and fragment ion mass tolerance of 0.5 Da. The resulting molecular network was visualized in Cytoscape (ver.3.10.2.) (Shannon *et al.*, 2003).

Preparing metabolites for bioactivity screening

Every crude extract was tested against a range of the laboratory model organisms available to us at the University of Wisconsin-Madison. Organisms included the pathogenic bacterium Methicillin-resistant *Staphylococcus aureus* (MRSA), *Pseudomonas aeruginosa*, and the pathogenic fungus *Candida auris*. Because experiments involving invertebrates are more complex, crude extracts from only a subset of the samples were used in tests against the nematodes *Caenorhabditis elegans* and *Brugia spp.*, and the dipterans *Aedes aegypti* and *Musca domestica*. For nematode screening, we used three extracts from each region (USA, South Africa, and Europe) in addition to extracts from the outgroup, *A. pantherina*. For insect screening, we used one sample from each of the three regions in addition to the outgroup (Table S1). Crude extracts were redissolved in methanol (MeOH) to test for antimicrobial activity or DMSO for invertebrate assays. Details of bioactivity assays are provided in the Supplemental Methods section. Our focus on model organisms was a practical choice and enables basic insights into bioactivity; future tests with species and populations of South African antagonists are needed.

RESULTS




Figure 1: Records of *Amanita muscaria* in South Africa. Occurrence data illustrate the spread of *A. muscaria* in South Africa over the last century. Panel (a) plots historical records up until 1945 as compiled by Doidge (1950). Insert i) is an enlargement of the Cape region. Panel (b) plots contemporary records as of 24 July 2023, also illustrating plantation forestry land use. Inserts are enlargements of ii) Cape Town and the southern Cape, iii) the Garden Route, and iv) the Eastern Cape. Numbered occurrences (1-3) are not associated with plantation forestry and instead record *A. muscaria* associated with pine trees in urban areas as in 1) Pretoria or 2) Clarens, or in rural areas as in 3), where pines are often used as windbreaks or to adorn the edges of farm roads. Panels (c) and (d) are images of *A. muscaria* specimens from Wynberg and Rondebosch in Cape Town collected in 1883 and 1874, respectively. Both records provide valuable information about associations with plants, e.g. “under Pine trees” in panel (d). Panel (e) is a photograph of *A. muscaria* collected near pine trees on a golf course (Simola Golf and Country Estate) in 2023 in the Western Cape province. Panel (f) is an image from a card bought at Kirstenbosch Botanical Gardens in Cape Town in 2023, reflecting widespread awareness of the fungus in South Africa.

South African *A. muscaria* Derive from a Single European Species

Amanita muscaria has long been associated with imported pines (Box 1, Fig. 1), and its history in South Africa is remarkably well recorded. The earliest herbarium specimen dates to 1874 (Box 1), but historical records suggest the fungus was already established by then. Today, it is common in planted forests, cities, and on rural farms (Fig. 1).

To investigate the origin of South African *A. muscaria*, we reconstructed phylogenies of *A. muscaria* using data from our own genomes and public repositories (Dataset S1); the topologies of created trees were largely in agreement with minor differences (Sup. Methods; Fig. S4, S5). The nine South African individuals we sequenced belong to Clade II of the *A. muscaria* complex (Fig. 2a), a lineage also including specimens from Australia and South America, where *A. muscaria* is also introduced and spreading (Vargas *et al.*, 2019). Clade II broadly corresponds to *A. muscaria* var. *muscaria*, widespread across Eurasia and into Alaska.

Site frequency spectrum analyses support a historical introduction of *A. muscaria* into South Africa. European Clade II individuals show a near-zero value (Fig. 2b), while South African individuals display a positive shift, consistent with the loss of rare alleles during a founder event. The European mushrooms consistently have higher nucleotide diversity (0.00166) and mean heterozygosity (0.0799 ± 0.0132 [Table S2]) as compared to the South African population (0.00118 and 0.0684 ± 0.0045 , respectively). Although sampling differences might influence results, the results align with historical records and phylogenetic inferences. Additionally, though the South African population shows signs of a bottleneck, the population remains diverse and is sexual: no clones were detected in either population and kinships are typically below 0.2 (Dataset S3). However, one pair of mushrooms from a single *P. elliottii* plantation in KwaZulu-Natal has a higher kinship (~ 0.33), suggesting some level of inbreeding, perhaps between multiple monokaryotic mycelia of a shared parent.

Roed_3 was the only European individual we sequenced falling into Clades I-I/A. Clades I-I/A represent a complex of two cryptic taxa widely distributed throughout North America. Interestingly, mitochondrial DNA sequences placed Roed_3 closer to Clade II than Clades I-I/A, a pattern we speculate as reflecting a hybridization event (Figs. S4 and S5).




Figure 2: A phylogenetic species reconstruction of 184 *A. muscaria* and 5 *A. pantherina* mushrooms. (a) Phylogeny constructed from an edge-linked partition model of ITS, LSU, β -tub, and TEF-1 α gene sequences. Clades are labeled with either the numbers used in Geml *et al.* (2008) or as *A.*

persicina or *A. regalis*; many of the labels from Geml *et al.* (2008) (e.g. clade VIII) represent undescribed taxa. Our phylogeny groups two specimens from Clade I in Geml *et al.* (2008) with Clade I/A. The tree is rooted with *A. pantherina*. Bootstrap support values are shown in red. Each mushroom's continent of origin is indicated by tip color, and available images of our sequenced mushrooms are displayed to the right of the phylogeny. Colors of the circles surrounding individual images mark the mushroom's continent of origin. (b) The distribution of Tajima's D was estimated from 5,000 bp windows across the reference genome using biallelic SNPs from the European (blue) and South African (red) Clade II genomes sequenced by us.

Specialized metabolite gene clusters are conserved across geographic regions and phylogenetic clades.

AntiSMASH predicted between 6–8 BGCs per genome, all of which were Type-1 polyketide synthases (PKSs) or isoprenoid(s) biosynthesis (terpene) backbone enzymes, with the exception of a single nonribosomal peptide synthetase (NRPS) found in the South African specimen 11667. All genomes, including the *A. pantherina* outgroup, contained the full ibotenic acid BGC (Fig. 3a, S6, S7) as defined previously (Obermaier & Müller, 2020). While the products of most of these *A. muscaria* BGCs are unknown, we confirmed the ibotenic acid BGC produced muscimol in each of the *A. muscaria* mushrooms we sequenced, regardless of phylogenetic grouping or geographic origin. Muscimol was also made by the *A. pantherina* outgroup (Nes_pan3 in Fig. S8).

BGCs were networked into the GCFs (Fig. 3b, Fig. S2, S3) likely to produce identical or closely related specialized metabolites (Navarro-Munoz *et al.*, 2020; Bağcı *et al.*, 2025). All *A. muscaria* genomes encoded two PKSs, with both copies (referred to as PKS1.1 and PKS1.2 [Fig. 3C]) grouping into the same GCF, termed polyketide (PK) GCF-1. The terpene BGCs, which showed more variability in copy number, were grouped into two major GCFs termed terpene GCF-1 and terpene GCF-2. Terpene GCF-1 was universally present as a single copy in every *A. muscaria* genome but was absent in the outgroup. By contrast, the copy number of terpene GCF-2 varied from 1–3 across samples, with no synteny observed among the neighboring BGC genes (Fig. 3c). The final group, termed terpene GCF-“other” (Fig. 3), represents all terpenes whose predicted BGC locus was found only in a single isolate. Variation in terpene gene copy number may reflect genome fragmentation or incomplete assemblies. However, expansions of terpenes are common across the Basidiomycota. A thorough analysis of these BGCs is beyond our scope but emerges as an interesting direction for future research. Only a small fraction of genes in antiSMASH-predicted BGCs housed domains associated with specialized metabolism in model Ascomycetes, suggesting a different biosynthetic logic in *A. muscaria* – either through stand-alone backbone enzymes or from the recruitment of novel tailoring genes.

Although *A. muscaria* has been reported to produce lethal amatoxins (Faulstich & Cochet-Meilhac, 1976), we found no genomic or chemical evidence—using bioinformatic searches, relaxed BLAST queries, and direct metabolite analysis—to support these claims (see Supplemental Methods).





Figure 3: Specialized metabolism (SM) in *Amanita muscaria* genomes. (a) Numbers of biosynthetic gene clusters (BGCs) per genome, sorted by type based on backbone enzymes. Gene copy numbers for each BGC locus also depicted. The maximum likelihood phylogeny of the isolates was generated using SNP data and rooted with *A. pantherina* (Nes_pan3). (b) Gene cluster family networks of the two major SM classes within the genomes: polyketides (PK) and terpenes. (c) Domain and genetic architecture of the antiSMASH-predicted BGC loci using Californian sample 20031-a (Clade I) as an example. Genes are depicted approximately to scale.

Conserved PKS enzymes in *A. muscaria* originated from duplication in a common ancestor of the species complex

The strong similarity between the two copies of PKS enzymes in all *A. muscaria* genomes led us to hypothesize the copies arose from a recent duplication event. In a phylogenetic analysis, all of *A. muscaria*'s PKS sequences form a monophyletic clade, while all other PKS sequences from other *Amanita* spp. share more ancient ancestry (Fig. 4b). A targeted follow-up analysis confirms the PKS duplication as unique to *A. muscaria* (Fig. S9). This analysis also revealed an interesting and distinct evolutionary history of the PKS enzyme in *A. inopinata*, a lineage which has independently duplicated the ancestral non-reducing PKS and acquired both a reducing-PKS and a hybrid NRPS-PKS (Fig. S9). The lack of trans-species representation in the duplicated *A. muscaria* PKS clade is strong evidence for the duplication having occurred relatively recently, likely in a common ancestor of the *A. muscaria* species complex.

To test whether the duplication of PKS-1.1 and PKS1.2 enabled functional divergence, we compared domain variation between the two copies and a sampling of highly similar PKS enzymes across diverse fungi. The PKS1.2 backbone in all *A. muscaria* genomes, regardless of geographic region or clade, has lost its C-terminal thioesterase (TE) domain (Fig. 4a). This result is not an annotation error; we used tBLASTn to verify that the coding sequence for this domain was only present in the genomic DNA associated with PKS1.1. The C-terminal domain is typically involved in the chain release and cyclization of polyketide products (Tang *et al.*, 2019). While most PKS hits from closely related *Amanita* species were found in single copies within their genomes and retained the C-terminal thioesterase, a distinct PKS clade - from *Amanita brunnescens* - showed a similar pattern of duplication and loss, suggesting the pattern has evolved multiple times in the genus (Fig. S9).

The domain structure of *A. muscaria*'s PKS enzymes closely mirrors the structure of known orsellinic acid-producing PKSs in other Agaricomycetes (Lackner *et al.*, 2013; Braesel *et al.*, 2017) and in the Ascomycete *Aspergillus nidulans* (Schroeckh *et al.*, 2009) (Fig. 4a). While these findings may suggest *A. muscaria* is producing an orsellinic-acid-like compound, a similarly strong hit to a PKS in the *Aspergillus fumigatus* genome is associated with production of a melanin intermediate (Fujii *et al.*, 2000; Brakhage & Liebmman, 2005), raising some doubt about the true end-product of the *A. muscaria* PKS enzymes.



muscaria genomes (highlighted in pink box). Left to right: Similarity scores were calculated by aligning each PKS sequence to the Sogn_5 PKS-1.1 backbone enzyme. The domain architecture of each PKS is color-coded. Known metabolites associated with specific PKSs are linked by dashed lines to the corresponding structures on the right. (b) Midpoint-rooted phylogeny of 244 Agaricomycete PKS proteins with additional hand-selected PKS proteins (e.g., reducing and non-reducing PKSs, *orsA* from *Aspergillus nidulans*). The tree was midpoint-rooted to facilitate clade visualization. Clades with an average branch length of less than 0.3 to their leaves were collapsed and displayed as triangles sized proportionally to the number of collapsed leaves. The monophyletic clade containing all *Amanita* PKS proteins is shaded in gray. A detailed analysis of proteins from the genus is provided in Fig. S9. The clades for *A. muscaria* PKS-1.1 and PKS-1.2 are highlighted in yellow and purple, respectively. Nodes with > 80% and > 90% support are indicated with orange and red stars, respectively. A subset of the proteins used to build the phylogeny are labeled for reference.

GNPS-based untargeted mass spectrometry analysis suggests the *A. muscaria* metabolome is highly conserved across native and novel ranges

GNPS analysis identified 273 unique molecular families (MFs) across all spectra (Fig. S10) including both primary and specialized metabolic features (Fig. 5a; see Methods). Of the 273 MFs, 215, including all MFs with at least 11 nodes, were detected in every geographic region. The largest MF (MF-21), comprising 100 unique nodes, was predicted to belong to a primary metabolism phospholipid family. We identified 25 MFs unique to *A. muscaria* Clade II samples (South Africa and Europe) and 13 MFs specific to South African *A. muscaria*. The largest South Africa-specific MF (MF-9) was made up of 10 unique nodes (Fig. 5b). These nodes had precursor m/z values between 900 and 1,200, with no matches to known library entries or previously characterized metabolites.












Figure 5: The metabolome of *A. muscaria* contextualized by geography. (a) Count of molecular families (MF) grouped by geographic region. (b) Network visualization of a subset of MFs. Node size reflects the total precursor intensity. Each node includes a pie chart marking the geographic regions where samples containing that MF were located. The full network of all detected MFs is in Fig. S10.

***A. muscaria* extracts reveal biological activity/ inactivity, with no population-specific patterns**

To understand the potential of *A. muscaria* metabolites to have broad-spectrum activity against bacteria, filamentous fungi, and yeasts, with potential relevance for ecological inferences and drug discovery, we performed disk-diffusion assays against Methicillin-resistant *Staphylococcus aureus* (MRSA; Gram-positive; Fig. S11), *Pseudomonas aeruginosa* (Gram-negative; Fig. S12), and *Candida auris* (a *Saccharomyces* yeast; Fig. S13). None of these organisms were inhibited by extracts (Table 2). In contrast to the microbial assays, extracts from nine geographically representative *A. muscaria* samples and one *A. pantherina* sample all strongly inhibited the

growth of the clade V nematode *Caenorhabditis elegans* (Table 2; Fig. S14). Extracts also immobilized the clade III parasitic nematodes *Brugia pahangi* and *Brugia malayi*, and assays with *B. malayi* reveal the impact of extracts on *B. malayi* was similar to heat-killed controls (Fig. S14). Insect assays using model organisms revealed a more selective pattern: while adult *Musca domestica* and *Aedes aegypti* were unaffected by the extracts, first-instar *A. aegypti* larvae were highly susceptible. In general, bioactivity declined as animals reached later developmental stages (Fig. S15). While the model organisms available to us might be unlikely to interact with *A. muscaria* in nature, our findings suggest there is variation in dipteran susceptibility, even despite the presence of GABA_A receptors in flies. Our data also suggest potential bioactivity may be limited to early developmental stages in certain species. We note we commonly find dipteran eggs laid among the gills of *A. muscaria* (Drott, Pringle and Stokes, pers. obs.). None of the bioactivities observed were different among the mushrooms collected from different continents. Our experiments targeted model systems, but important differences may exist between the models we used and the populations of species interacting with *A. muscaria* in South Africa and elsewhere.

Table 2: Summary of the bioactivity of *A. muscaria* extracts against Methicillin-resistant *Staphylococcus aureus* (MRSA), *Pseudomonas aeruginosa*, *Candida auris*, *Caenorhabditis elegans*, *Brugia* spp., *Musca domestica*, and *Aedes aegypti*. The table displays (from left to right) the number of extracts (mushrooms) tested in each experiment, the viability of the tested organism when confronted with extracts, the concentrations used, the method of inoculation, and a reference to the Sup. Figure(s) with full results.

Organism	Number of extracts tested (n)			Bioactivity <div><div>● Strong</div><div>● Mild</div><div>● None</div></div>	Lowest conc. with bioactivity	Highest conc. tested	Bioactivity assay ran	Sup. Figures with all results
	Europe	SA	USA					
 MRSA (Gram positive pathogen)	12	10	4	●	n/a	50 mg/ml	Disc diffusion assay	S10
 <i>Pseudomonas aeruginosa</i> (Gram negative pathogen)	12	10	4	●	n/a	50 mg/ml	Disc diffusion assay	S11
 <i>Candida auris</i> (Yeast pathogen)	12	10	4	●	n/a	50 mg/ml	Disc diffusion assay	S12
 <i>Caenorhabditis elegans</i> (nematode)	4	3	3	●	50 µg/ml	1 mg/ml	96-well liquid assays	S13
 <i>Brugia spp.</i> (nematode parasite)	4	3	3	●	10 µg/ml	1 mg/ml	96-well liquid assays	S13
 Adult <i>Musca domestica</i> (Housefly)	1	1	1	●	n/a	1 mg/ml	Administered by cotton wick	S14
 First/Late Instar <i>Aedes aegypti</i> (yellow fever mosquito)	1	1	1	● ● First Late	1 mg/ml	1 mg/ml	24-well liquid assays	S14
 Adult <i>Aedes aegypti</i> (yellow fever mosquito)	1	1	1	●	n/a	1 mg/ml	Administered by cotton wick	S14

DISCUSSION

The mushrooms of the fungus *A. muscaria* s.l. are famous, featured in settings ranging from Super Mario™ to treatises of Hindu texts. Despite its prominent cultural status, the human-mediated dispersal of this fungus is poorly documented. In this study, we used genomics and metabolomics to investigate the history and evolution of *A. muscaria* following its introduction to South Africa. Our phylogenetic analyses reveal South African *A. muscaria* populations belong to Clade II, a cryptic lineage of European origin. Historical records suggest the fungus was likely introduced more than 150 years ago, perhaps with pine seedlings brought by European settlers to Cape Town. Its subsequent spread was facilitated by the expansion of pine forestry in southern Africa, and South African *A. muscaria* now grow with pines imported from other continents, including the Mexican *P. patula*. The fungus is sexual, and its populations in South Africa are

genetically diverse, with multiple distinct individuals present within single forest stands. Despite its extended history in South Africa, the specialized metabolome of *A. muscaria* remains highly conserved, with gene clusters encoding muscimol, terpene, and polyketide products present in all genomes. Bioactivity assays with laboratory model organisms reveal the potent nematocidal effects of the fungus's metabolites, suggesting their potential (as yet untested) role in mediating ecological interactions.

The European Clade II lineage found in South Africa is also the lineage introduced to Colombia (Vargas *et al.*, 2019) and Australia (Lebel *et al.*, 2024) (Fig. 2a). Remarkably, the global dispersal of *A. muscaria* appears to involve just one of its many cryptic species. Europe has long been suspected as the source of Australian *A. muscaria*, but we are the first to identify Clade II as the specific lineage introduced there. More intensive sampling will be needed to pinpoint the exact European source(s) of South African *A. muscaria*, and additional sequencing may reveal whether other clades within the complex have been introduced elsewhere in the country. However, based on available data, Clade II appears to be the primary lineage driving invasions, suggesting underlying genetic, ecological, or historical factors contributing to its spread. The fungus may also have been introduced to South Africa from another invasive range, for example, Colombia or Australia. But historical records suggest the most parsimonious explanation for the *A. muscaria* in South Africa is a direct introduction(s) from Europe (Box 1, Fig. 1). Interestingly, several of our phylogenies (but not all) suggest South African *A. muscaria* share their most recent common ancestry with Australian populations, a pattern which may reflect a shared European source population (Fig. S4, S5).

Kinship analyses indicate no two South African mushrooms are clones, although many mushroom pairs are closely related; the data are consistent with ongoing sexual reproduction. While invasive fungal populations often exhibit shifts in mating systems or reproductive modes (Wang *et al.*, 2023), the conservation of sexual reproduction and mating type diversity in South Africa suggests either a weak genetic bottleneck or continued admixture from multiple introductions.

The introduction of *A. muscaria* into South Africa is also reflected in site-frequency spectra, where a positive shift in Tajima's *D* estimates among Clade II mushrooms (Fig. 2b) is consistent with a founder event followed by insufficient time and/or evolution to regenerate rare alleles. A similar pattern has been observed in *A. phalloides* populations introduced to California from Europe (Drott *et al.*, 2023). Other population genetic statistics, including *F_{st}* estimates, also suggest South African populations are differentiated from their European counterparts (Fig. S16-a). Loci with the highest *F_{st}* scores encode genes associated with oxidoreductase activity, metal binding, and cell membranes—functions previously linked to host and environmental stress responses in fungi (Fig. S16-b) (Staerck *et al.*, 2018; Feng *et al.*, 2023). Further research is needed to clarify the significance of the differences.

Despite the different ecological contexts of Europe, South Africa, and Australia, all sequenced *A. muscaria* genomes housed a similar complement of BGCs, including the ibotenic acid BGC (Fig. 3, S6, S7), an expanded set of terpenes, and two PKS gene clusters (Fig. 4). Research on the ecology of *A. muscaria* has focused primarily on its ecological niche; the ECM symbiosis has one of its origins in the *Amanita* genus (Wolfe *et al.*, 2012). Although little is known about the role of specialized metabolites in the ECM symbiosis, we identified an expansion of an orsellinic

acid-like PKS ortholog maintained across populations (Fig. 3, 4; how the duplicate genes are expressed remains a target for future research). One of these orthologs has lost a TE domain, suggesting the mature product is not cyclized or may depend on other enzymes for processing, similar to lovastatin biosynthesis (Xu *et al.*, 2013). Orsellinic acid, found in several basidiomycetes (Lackner *et al.*, 2012), has derivatives with potential herbicidal properties (Peres *et al.*, 2009), which we speculate may influence fungal-plant interactions. Orsellinic acid-sesterpene hybrids occur in some fungi, resulting in highly diverse bioactivities (Gao *et al.*, 2023). Trans-BGC interactions are increasingly being discovered (Won *et al.*, 2022) but are not well understood, particularly in Basidiomycete lineages; we speculate that such epistatic interactions modulate the chemical diversity associated with the conserved terpene BGCs in *A. muscaria*.

The conservation of primary and specialized metabolites across introduced and native populations provides no evidence for relaxed selection (Fig. S17) as might be expected if metabolic resources were being reappropriated to other pathways (Blossey & Nötzold, 1995). Instead, our findings raise the possibility of conserved functional interactions with communities in both native and introduced ranges. Perhaps imported European soils harbored antagonistic microbes and invertebrates now interacting with *A. muscaria* in South Africa, or perhaps South African *A. muscaria* have equivalent kinds of interactions with native African antagonists. It is also possible there has not been enough time for genomic signatures of relaxed selection to emerge, although SMs are often subject to strong selection and are readily lost, causing their highly patchy distribution across taxa (Robey *et al.*, 2021).

While *A. muscaria*'s core biosynthetic gene clusters (BGCs) are highly conserved, we also identified 13 molecular families unique to South African populations. Slight differences in mushroom handling after collection and differences among the microbial communities inhabiting mushrooms may have impacted identified metabolites, but we speculate these novel molecular families may represent metabolic innovations influencing fungal interactions in the novel range. The differentiation of metabolic space between fungal populations is often small but it can sometimes be defined by the production of a small subset of ecologically important metabolites (Drott *et al.*, 2021).

The association between *A. muscaria* and fly-killing dates back centuries, and both historical and modern accounts suggest insecticidal properties. The traditional recipe for using *A. muscaria* to kill flies involves soaking *A. muscaria* in milk or water (Lumpert & Kreft, 2016). As recently as 2021, Carboué and Lopez (2021) cited *A. muscaria* extracts as having fly-killing bioactivity. However, skepticism about its ability to kill insects has persisted for just as long as claims of its efficacy. As early as 1779, French botanist Jean Bulliard challenged claims of insecticidal activity, noting that flies appeared unaffected by the mushroom (Wasson, 1969). Some folklore suggests *A. muscaria* may act as an attractant rather than a direct toxin, luring flies to consume the mushroom before they ultimately drown in the *A. muscaria*-soaked milk. Our findings align with historical skepticism—while *A. muscaria* extracts strongly suppressed nematodes (*C. elegans* and *Brugia* spp.) and inhibited mosquito larvae, they had no detectable effect on any adult dipterans, bacteria, or yeast (Table 2). The strong nematicidal properties of *A. muscaria*, which our study has reproduced, have previously been attributed to its specialized metabolites, ibotenic acid and muscimol, which are known agonists to the GABA_A receptor complex (Johnston, 2014). Muscimol resistance has been documented among mycophagous dipterans but

not their frugivorous counterparts (Tuno *et al.*, 2007), emphasizing questions about the susceptibility of relevant antagonists in the introduced range.

In the aggregate, our data highlight the genetic and metabolic versatility of *A. muscaria*, a globally invasive ectomycorrhizal fungus. While our results offer insights into the origin of introduced populations, the evolution of the *A. muscaria* lineage, and the genomic signatures of these introductions (or their lack), these findings also emphasize many unanswered questions: how does *A. muscaria* interact with local antagonists in its novel ranges, and does enemy release or do novel weapons play a role in the spread of the fungus? Do population-specific molecular families contribute to its success? Our results offer insights into the global distribution of *A. muscaria*, creating a strong foundation for future research exploring how fungal invasions and co-invasions shape ecosystems across the globe.

For Peer Review

ACKNOWLEDGEMENTS:

We thank Jolanda Roux, Izette Greyling and Bernard Slippers for facilitating collections in South Africa; Jessie Uehling and Liz Henderson for Australian collections; and Inger Skrede and her laboratory, Lynn Harrow, László Nagy, and Kara O’Keefe for facilitating collections in Europe. We thank David W. Johnson and the staff of Future Africa and FABI for logistical and other support. We thank Irene Barnes for support during our initial workshop. We thank the Royal Botanic Gardens, Kew and especially Lee Davies for providing us with information about the South African *A. muscaria* specimens held in their fungarium. We thank Lyric Bartholomay and Kathy Vaccaro (UW-Madison Department of Pathobiological Sciences) for assistance with maintaining the *Ae. aegypti* colonies used in this study. Genomes were sequenced at the University of Wisconsin–Madison Biotechnology Center’s DNA Sequencing Facility (Research Resource Identifier (RRID) SCR_017759). We dedicate our work to Elizabeth N. Feliciano, a graduate student who died before she was able to join us in South Africa, and we honor her legacy of service to mycology.

COMPETING INTERESTS

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Our study stems from a unique collaboration between scientists working in South Africa and the United States, and together we span diverse countries and career stages. Our collaboration began at a U.S. National Science Foundation-funded workshop (Grant No. 1953299 awarded to JDH and JSB) held at Future Africa (www.futureafrica.science) in November 2022. The workshop initially focused on introducing students to invasion biology and providing bioinformatics training, using *Amanita muscaria* s.l. genomes as a teaching tool. AP and JDH conceived the study, and it was developed fully over the following 14 months. GRN designed and drafted the manuscript, leading the bioinformatic and natural product experiments and analyses. SRF, KMTL, DLN, and CKS (listed in reverse alphabetical order as equal co-second authors) contributed substantially to analyses and writing. CB also contributed significantly to analyses and writing. GRN, NPK, JWB, and AP sequenced genomes. Historical research was led by MAH, AN, and AP. Y-WW led kinship analyses. Metabolic extractions and mass spectrometry analyses were conducted by GRN, SCP, and NPK. Bioactivity work was designed, executed, and analyzed as follows: microbial assays by GRN, CKS, and NPK; insect assays by GRN, TKW, HLN, MEMM, and KLC; and nematode assays by GRN, KTR, and MZ. JDH, MTD, and AP contributed to data analysis, interpretation, and writing, with MTD and AP leading revisions to interpretations and the manuscript. All authors participated in workshop discussions, contributed to initial analyses, and helped edit the final manuscript.

DATA AVAILABILITY

All genomic data analyzed in this study were generated by the authors. Raw reads and assemblies are publicly available through NCBI BioProject [####]. Other data and methods enabling replication are provided with this manuscript and in the Supporting Information (which contains Supporting Methods, Tables, and Figures).

REFERENCE LIST

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990.** Basic local alignment search tool. *Journal of Molecular Biology* DOI: 10.1016/S0022-2836(05)80360-2
- Anagnostakis SL. 1987.** Chestnut blight - the classical problem of an introduced pathogen. *Mycologia* DOI: 10.2307/3807741
- Bağcı C, Nuhamunada M, Goyat H, Ladanyi C, Sehnal L, Blin K, Kautsar SA, Tagirdzhanov A, Gurevich A, Mantri S, et al. 2025.** BGC Atlas: a web resource for exploring the global chemical diversity encoded in bacterial genomes. *Nucleic Acids Research* DOI: 10.1093/nar/gkae953
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, et al. 2012.** SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* DOI: 10.1089/cmb.2012.0021
- Berch SM, Kroeger P, Finston T. 2017.** The death cap mushroom (*Amanita phalloides*) moves to a native tree in Victoria, British Columbia. *Botany* DOI: 10.1139/cjb-2016-0183
- Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, Medema MH, Weber T. 2019.** antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Research* DOI: 10.1093/nar/gkz310
- Blossey B, Nötzold R. 1995.** Evolution of increased competitive ability in invasive nonindigenous plants: A hypothesis. *Journal of Ecology* DOI: 10.2307/2261425
- Bok JW, Balajee SA, Marr KA, Andes D, Nielsen KF, Frisvad JC, Keller NP. 2005.** LaeA, a regulator of morphogenetic fungal virulence factors. *Eukaryotic Cell* DOI: 10.1128/EC.4.9.1574-1582.2005
- Bolger AM, Lohse M, Usadel B. 2014.** Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* DOI: 10.1093/bioinformatics/btu170
- Boyd IL, Freer-Smith PH, Gilligan CA, Godfray HC. 2013.** The consequence of tree pests and diseases for ecosystem services. *Science* DOI: 10.1126/science.1235773
- Braesel J, Fricke J, Schwenk D, Hoffmeister D. 2017.** Biochemical and genetic basis of orsellinic acid biosynthesis and prenylation in a stereaceous basidiomycete. *Fungal Genetics and Biology* DOI: 10.1016/j.fgb.2016.11.007
- Brakhage AA, Liebmann B. 2005.** *Aspergillus fumigatus* conidial pigment and cAMP signal transduction: significance for virulence. *Medical Mycology* DOI: 10.1080/13693780400028967
- Brancatelli GIE, Amodeo MR, Cuevas YA, Zalba SM. 2020.** Invasive pines in Argentinian grasslands: lessons from control operations. *Biological Invasions* DOI: 10.1007/s10530-019-02103-9
- Brewer JS, Souza FM, Callaway RM, Durigan G. 2018.** Impact of invasive slash pine (*Pinus elliottii*) on groundcover vegetation at home and abroad. *Biological Invasions* DOI: 10.1007/s10530-018-1734-z
- Bushnell B. 2015.** BMap short-read aligner, and other bioinformatics tools. *University of California, Berkeley, CA.*

- Callaway RM, Ridenour WM. 2004.** Novel weapons: invasive success and the evolution of increased competitive ability. *Frontiers in Ecology and the Environment* DOI: 10.1890/1540-9295(2004)002[0436:Nwisat]2.0.Co;2
- Camazine S. 1983.** Mushroom chemical defense: food aversion learning induced by hallucinogenic toxin, muscimol. *Journal of Chemical Ecology* DOI: 10.1007/BF00988513
- Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009.** trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* DOI: 10.1093/bioinformatics/btp348
- Carboué Q, Lopez M. 2021.** *Amanita muscaria*: ecology, chemistry, myths. *Encyclopedia* DOI: 10.3390/encyclopedia1030069
- Cleary M, Nguyen D, Marciulyniene D, Berlin A, Vasaitis R, Stenlid J. 2016.** Friend or foe? Biological and ecological traits of the European ash dieback pathogen *Hymenoscyphus fraxineus* in its native environment. *Scientific Reports* DOI: 10.1038/srep21895
- Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, et al. 2011.** The variant call format and VCFtools. *Bioinformatics* DOI: 10.1093/bioinformatics/btr330
- Desprez-Loustau ML, Robin C, Buee M, Courtecuisse R, Garbaye J, Suffert F, Sache I, Rizzo DM. 2007.** The fungal dimension of biological invasions. *Trends in Ecology & Evolution* DOI: 10.1016/j.tree.2007.04.005
- Dickie IA, Bolstridge N, Cooper JA, Peltzer DA. 2010.** Co-invasion by *Pinus* and its mycorrhizal fungi. *New Phytologist* DOI: 10.1111/j.1469-8137.2010.03277.x
- Dickie IA, Bufford JL, Cobb RC, Desprez-Loustau ML, Grelet G, Hulme PE, Klironomos J, Makiola A, Nuñez MA, Pringle A, et al. 2017.** The emerging science of linked plant-fungal invasions. *New Phytologist* DOI: 10.1111/nph.14657
- Dickie IA, St John MG, Yeates GW, Morse CW, Bonner KI, Orwin K, Peltzer DA. 2014.** Belowground legacies of *Pinus contorta* invasion and removal result in multiple mechanisms of invasional meltdown. *AoB Plants* DOI: 10.1093/aobpla/plu056
- Doidge EM. 1950.** The South African fungi and lichens. *Bothalia* 5: 1–1094.
- Drott MT, Bastos RW, Rokas A, Ries LNA, Gabaldon T, Goldman GH, Keller NP, Greco C. 2020.** Diversity of secondary metabolism in *Aspergillus nidulans* clinical isolates. *mSphere* DOI: 10.1128/mSphere.00156-20
- Drott MT, Lazzaro BP, Brown DL, Carbone I, Milgroom MG. 2017.** Balancing selection for aflatoxin in *Aspergillus flavus* is maintained through interference competition with, and fungivory by insects. *Proceedings of the Royal Society B Biological Sciences* DOI: 10.1098/rspb.2017.2408
- Drott MT, Park SC, Wang YW, Harrow L, Keller NP, Pringle A. 2023.** Pangenomics of the death cap mushroom *Amanita phalloides*, and of Agaricales, reveals dynamic evolution of toxin genes in an invasive range. *The ISME Journal* DOI: 10.1038/s41396-023-01432-x
- Drott MT, Rush TA, Satterlee TR, Giannone RJ, Abraham PE, Greco C, Venkatesh N, Skerker JM, Glass NL, Labbe JL, et al. 2021.** Microevolution in the pansecondary metabolome of *Aspergillus flavus* and its potential

- macroevolutionary implications for filamentous fungi. *Proceedings of the National Academy of Science USA* DOI: 10.1073/pnas.2021683118
- Dunk CW, Lebel T, Keane PJ. 2012.** Characterisation of ectomycorrhizal formation by the exotic fungus *Amanita muscaria* with *Nothofagus cunninghamii* in Victoria, Australia. *Mycorrhiza* DOI: 10.1007/s00572-011-0388-9
- Faulstich H, Cochet-Meilhac M. 1976.** Amatoxins in edible mushrooms. *FEBS Letters* DOI: 10.1016/0014-5793(76)80252-9
- Feng H, Meng P, Zhang S, Chen W, Wang H, Wang C. 2023.** Insights from comparative transcriptome analysis in the responses of Pb-tolerant fungi *Curvularia tsudae* to Pb stress. *Ecotoxicology and Environmental Safety* DOI: 10.1016/j.ecoenv.2022.114476
- Fuhrer BA. 1992.** *Rainforest fungi of Tasmania and south-east Australia*. East Melbourne, Vic. :: CSIRO Australia.
- Fujii I, Mori Y, Watanabe A, Kubo Y, Tsuji G, Ebizuka Y. 2000.** Enzymatic synthesis of 1,3,6,8-tetrahydroxynaphthalene solely from malonyl coenzyme A by a fungal iterative type I polyketide synthase PKS1. *Biochemistry* DOI: 10.1021/bi000644j
- Gao H, Zhou L, Zhang P, Wang Y, Qian X, Liu Y, Wu G. 2023.** Filamentous fungi-derived orsellinic acid-sesquiterpene meroterpenoids: Fungal sources, chemical structures, bioactivities, and biosynthesis. *Planta Medica* DOI: 10.1055/a-2099-4932
- Geml J, Laursen GA, O'Neill K, Nusbaum HC, Taylor DL. 2006.** Beringian origins and cryptic speciation events in the fly agaric (*Amanita muscaria*). *Molecular Ecology* DOI: 10.1111/j.1365-294X.2005.02799.x
- Geml J, Tulloss RE, Laursen GA, Sazanov NA, Taylor DL. 2008.** Evidence for strong inter- and intracontinental phylogeographic structure in *Amanita muscaria*, a wind-dispersed ectomycorrhizal basidiomycete. *Molecular Phylogenetics and Evolution* DOI: 10.1016/j.ympev.2008.04.029
- Gilchrist CLM, Booth TJ, van Wersch B, van Grieken L, Medema MH, Chooi Y-H. 2021.** cblaster: a remote search tool for rapid identification and visualization of homologous gene clusters. *Bioinformatics Advances* DOI: 10.1093/bioadv/vbab016
- Gillett J. 1962.** Pest pressure, an underestimated factor in evolution. *Systematics Association Publication* 4(37): 37-46.
- Global Biodiversity Information Facility (GBIF).** DOI: 10.15468/dl.4pqxfc
- Goldman GB, Gryzenhout M. 2019.** *Field Guide to Mushrooms and Other Fungi in South Africa*: Struik Nature.
- Gruber B, Unmack PJ, Berry OF, Georges A. 2018.** DARTR: an R package to facilitate analysis of SNP data generated from reduced representation genome sequencing. *Molecular Ecology Resources* DOI: 10.1111/1755-0998.12745
- Hayward J, Horton TR, Nuñez MA. 2015.** Ectomycorrhizal fungal communities coinvasive with Pinaceae host plants in Argentina: Gringos bajo el bosque. *New Phytologist* DOI: 10.1111/nph.13453
- He L, Diedrich J, Chu YY, Yates JR, 3rd. 2015.** Extracting accurate precursor information for Tandem Mass Spectra by RawConverter. *Analytical Chemistry* DOI: 10.1021/acs.analchem.5b02721

- Hoff KJ, Stanke M. 2013.** WebAUGUSTUS--a web service for training AUGUSTUS and predicting genes in eukaryotes. *Nucleic Acids Research* DOI: 10.1093/nar/gkt418
- Johnston GA. 2014.** Muscimol as an ionotropic GABA receptor agonist. *Neurochemical Research* DOI: 10.1007/s11064-014-1245-y
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.** ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* DOI: 10.1038/nmeth.4285
- Karst J, Marczak L, Jones MD, Turkington R. 2008.** The mutualism-parasitism continuum in ectomycorrhizas: a quantitative assessment using meta-analysis. *Ecology* DOI: 10.1890/07-0823.1
- Katoh K, Standley DM. 2013.** MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* DOI: 10.1093/molbev/mst010
- Keller NP. 2019.** Fungal secondary metabolism: regulation, function and drug discovery. *Nature Reviews Microbiology* DOI: 10.1038/s41579-018-0121-1
- Kohler A, Kuo A, Nagy LG, Morin E, Barry KW, Buscot F, Canback B, Choi C, Cichocki N, Clum A, et al. 2015.** Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in mycorrhizal mutualists. *Nature Genetics* DOI: 10.1038/ng.3223
- Korunes KL, Samuk K. 2021.** pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. *Molecular Ecology Resources* DOI: 10.1111/1755-0998.13326
- Lackner G, Bohnert M, Wick J, Hoffmeister D. 2013.** Assembly of melleolide antibiotics involves a polyketide synthase with cross-coupling activity. *Chemistry & Biology* DOI: 10.1016/j.chembiol.2013.07.009
- Lackner G, Misiek M, Braesel J, Hoffmeister D. 2012.** Genome mining reveals the evolutionary origin and biosynthetic potential of basidiomycete polyketide synthases. *Fungal Genetics and Biology* DOI: 10.1016/j.fgb.2012.09.009
- Le Maitre DC, van Wilgen BW, Gelderblom CM, Bailey C, Chapman RA, Nel JA. 2002.** Invasive alien trees and water resources in South Africa: case studies of the costs and benefits of management. *Forest Ecology and Management* DOI: 10.1016/s0378-1127(01)00474-1
- Lebel T, May TW, Cooper JA, Catcheside D, Catcheside P, Haska J. 2024.** Confirming the presence of five exotic species of *Amanita* in Australia and New Zealand. *Swainsona*. 1-44.
- Li H. 2013.** Seqtk: toolkit for Processing Sequences in FASTA/Q Formats. *GitHub*.
- Li H, Durbin R. 2009.** Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* DOI: 10.1093/bioinformatics/btp324
- Liu H, Stiling P. 2006.** Testing the enemy release hypothesis: a review and meta-analysis. *Biological Invasions* DOI: 10.1007/s10530-005-5845-y
- Lumpert M, Kreft S. 2016.** Catching flies with *Amanita muscaria*: traditional recipes from Slovenia and their efficacy in the extraction of ibotenic acid. *Journal of Ethnopharmacology* DOI: 10.1016/j.jep.2016.04.009

- Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. 2010.** Robust relationship inference in genome-wide association studies. *Bioinformatics* DOI: 10.1093/bioinformatics/btq559
- Manni M, Berkeley MR, Seppey M, Simao FA, Zdobnov EM. 2021.** BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution* DOI: 10.1093/molbev/msab199
- Márquez Parraguez CK. 2024.** *Ensayo de inducción de la simbiosis micorrícica entre el hongo exótico invasor Amanita muscaria y arboles nativos del género Nothofagus*. Undergraduate Thesis, Universidad de Concepción.
- McGinnis S, Madden TL. 2004.** BLAST: at the core of a powerful and diverse set of sequence analysis tools. *Nucleic Acids Res* 32(Web Server issue): W20-25.
- McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytzky A, Garimella K, Altshuler D, Gabriel S, Daly M, et al. 2010.** The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research* DOI: 10.1101/gr.107524.110
- Michelot D, Melendez-Howell LM. 2003.** *Amanita muscaria*: chemistry, biology, toxicology, and ethnomycology. *Mycological Research* DOI: 10.1017/s0953756203007305
- Mikola P 1970.** Mycorrhizal Inoculation in Afforestation. In: Romberger JA, Mikola P eds. *International Review of Forestry Research*: Elsevier, 123-196.
- Milani T, Hoeksema JD, Jobbágy EG, Rojas JA, Vilgalys R, Teste FP. 2022.** Co-invading ectomycorrhizal fungal succession in pine-invaded mountain grasslands. *Fungal Ecology* DOI: 10.1016/j.funeco.2022.101176
- Miller OK, Jenkins DT. 1978.** A taxonomic and nomenclatural study of the genus *Amanita* section *Amanita* for North America. *Mycologia* DOI: 10.2307/3759051
- Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020.** IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution* DOI: 10.1093/molbev/msaa015
- Mudbhari S, Lofgren L, Appidi MR, Vilgalys R, Hettich RL, Abraham PE. 2024.** Decoding the chemical language of *Suillus* fungi: genome mining and untargeted metabolomics uncover terpene chemical diversity. *mSystems* DOI: 10.1128/msystems.01225-23
- Navarro-Munoz JC, Selem-Mojica N, Mallowney MW, Kautsar SA, Tryon JH, Parkinson EI, De Los Santos ELC, Yeong M, Cruz-Morales P, Abubucker S, et al. 2020.** A computational framework to explore large-scale biosynthetic diversity. *Nature Chemical Biology* DOI: 10.1038/s41589-019-0400-9
- Neville P, Poumarat S. 2004.** *Amaniteae. Amanita, Limacella & Torrendia fungi Europaei* 9. Italy: Candusso Edizioni.
- Nickles GR, Oestereich B, Keller NP, Drott MT. 2023.** Mining for a new class of fungal natural products: the evolution, diversity, and distribution of isocyanide synthase biosynthetic gene clusters. *Nucleic Acids Research* DOI: 10.1093/nar/gkad573
- Núñez MA, Chiufo MC, Torres A, Paul T, Dimarco RD, Raal P, Policelli N, Moyano J, García RA, van Wilgen BW, et al. 2017.** Ecology and management of

- invasive Pinaceae around the world: progress and challenges. *Biological Invasions* DOI: 10.1007/s10530-017-1483-4
- Núñez MA, Horton TR, Simberloff D. 2009.** Lack of belowground mutualisms hinders Pinaceae invasions. *Ecology* DOI: 10.1890/08-2139.1
- Obermaier S, Müller M. 2020.** Ibotenic acid biosynthesis in the fly agaric is initiated by glutamate hydroxylation. *Angewandte Chemie International Edition* DOI: 10.1002/anie.202001870
- Oda T, Tanaka C, Tsuda M. 2004.** Molecular phylogeny and biogeography of the widely distributed *Amanita* species, *A. muscaria* and *A. pantherina*. *Mycological Research* DOI: 10.1017/s0953756204000620
- Peres MTLP, Mapeli A, Faccenda O, Gomes AT, Honda NK. 2009.** Allelopathic potential of orsellinic acid derivatives. *Brazilian Archives of Biology and Technology* DOI: Doi 10.1590/S1516-89132009000400027
- Pildain MB, Marchelli P, Azpilicueta MM, Starik C, Barroetavena C. 2021.** Understanding introduction history: genetic structure and diversity of the edible ectomycorrhizal fungus, *Suillus luteus*, in Patagonia (Argentina). *Mycologia* DOI: 10.1080/00275514.2021.1909449
- Policelli N, Bruns TD, Vilgalys R, Nunez MA. 2019.** Suilloid fungi as global drivers of pine invasions. *New Phytologist* DOI: 10.1111/nph.15660
- Policelli N, Hoeksema JD, Moyano J, Vilgalys R, Vivello S, Bhatnagar JM. 2023.** Global pine tree invasions are linked to invasive root symbionts. *New Phytologist* DOI: 10.1111/nph.18527
- Pringle A, Adams RI, Cross HB, Bruns TD. 2009.** The ectomycorrhizal fungus *Amanita phalloides* was introduced and is expanding its range on the west coast of North America. *Molecular Ecology* DOI: 10.1111/j.1365-294X.2008.04030.x
- Pringle A, Vellinga EC. 2006.** Last chance to know? Using literature to explore the biogeography and invasion biology of the death cap mushroom *Amanita phalloides* (Vaill. ex Fr. :Fr.) Link. *Biological Invasions* 8(5): 1131-1144.
- Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020.** Using SPAdes de novo assembler. *Current Protocols in Bioinformatics* DOI: 10.1002/cpbi.102
- Reid DA, Eicker A. 1991.** South African fungi: the genus *Amanita*. *Mycological Research* DOI: 10.1016/S0953-7562(09)81364-6
- Richardson DM, Higgins SI .1998.** Pines as invaders in the southern hemisphere. In: Richardson DM ed. *Ecology and biogeography of Pinus*. Cambridge: Cambridge University Press.
- Robey MT, Caesar LK, Drott MT, Keller NP, Kelleher NL. 2021.** An interpreted atlas of biosynthetic gene clusters from 1,000 fungal genomes. *Proceedings of the National Academy of Sciences* DOI: 10.1073/pnas.2020230118
- Rossman AY 2009.** The impact of invasive fungi on agricultural ecosystems in the United States. In: Langor DW, Sweeney J eds. *Ecological impacts of non-native invertebrates and fungi on terrestrial ecosystems*. Dordrecht: Springer Netherlands, 97-107.
- Rouget M, Richardson DM, Milton SJ, Polakow D. 2001.** Predicting invasion dynamics of four alien *Pinus* species in a highly fragmented semi-arid shrubland in South Africa. *Plant Ecology* DOI: 10.1023/A:1011412427075

- Sapsford SJ, Wakelin A, Peltzer DA, Dickie IA. 2022.** Pine invasion drives loss of soil fungal diversity. *Biological Invasions* DOI: 10.1007/s10530-021-02649-7
- Schaffer AA, Aravind L, Madden TL, Shavirin S, Spouge JL, Wolf YI, Koonin EV, Altschul SF. 2001.** Improving the accuracy of PSI-BLAST protein database searches with composition-based statistics and other refinements. *Nucleic Acids Research* DOI: 10.1093/nar/29.14.2994
- Schroeckh V, Scherlach K, Nuttmann HW, Shelest E, Schmidt-Heck W, Schuemann J, Martin K, Hertweck C, Brakhage AA. 2009.** Intimate bacterial-fungal interaction triggers biosynthesis of archetypal polyketides in *Aspergillus nidulans*. *Proceedings of the National Academy of Science USA* DOI: 10.1073/pnas.0901870106
- Schöffler A. 2018.** Secondary metabolites of basidiomycetes. In Anke T, Weber D. *Physiology and Genetics: Selected Basic and Applied Aspects*. New York: Springer. 231-275.
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. 2003.** Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Research* DOI: 10.1101/gr.1239303
- Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015.** BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* DOI: 10.1093/bioinformatics/btv351
- Simberloff D, Nuñez MA, Ledgard NJ, Pauchard A, Richardson DM, Sarasola M, Van Wilgen BW, Zalba SM, Zenni RD, Bustamante R, et al. 2010.** Spread and impact of introduced conifers in South America: Lessons from other southern hemisphere regions. *Austral Ecology* DOI: 10.1111/j.1442-9993.2009.02058.x
- Smith SE, Read DJ. 2008.** *Mycorrhizal Symbiosis*. London: Academic press.
- Staerck C, Vandeputte P, Gastebois A, Calenda A, Giraud S, Papon N, Bouchara JP, Fleury MJJ. 2018.** Enzymatic mechanisms involved in evasion of fungi to the oxidative stress: focus on *Scedosporium apiospermum*. *Mycopathologia* DOI: 10.1007/s11046-017-0160-6
- Stanke M, Diekhans M, Baertsch R, Haussler D. 2008.** Using native and syntenically mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* DOI: 10.1093/bioinformatics/btn013
- Størmer FC, Janak K, Koller GEB. 2004.** Ibotenic acid in *Amanita muscaria* spores and caps. *Mycologist* DOI: 10.1017/S0269915X04003039
- Su YT, Liu J, Yang DN, Cai Q, Yang ZL, Chen ZH. 2023.** Determination of ibotenic acid and muscimol in species of the genus *Amanita* section *Amanita* from China. *Toxicon* DOI: 10.1016/j.toxicon.2023.107257
- Tang MC, Fischer CR, Chari JV, Tan D, Suresh S, Chu A, Miranda M, Smith J, Zhang Z, Garg NK, et al. 2019.** Thioesterase-catalyzed aminoacylation and thiolation of polyketides in fungi. *Journal of the American Chemical Society* DOI: 10.1021/jacs.9b01083
- Tannous J, Cosetta CM, Drott MT, Rush TA, Abraham PE, Giannone RJ, Keller NP, Wolfe BE. 2022.** Fungal antibiotics control bacterial community diversity in the cheese rind microbiome. *bioRxiv* DOI: 10.1101/2022.11.26.518062

- Torchin ME, Lafferty KD, Dobson AP, McKenzie VJ, Kuris AM. 2003.** Introduced species and their missing parasites. *Nature* DOI: 10.1038/nature01346
- Treseder KK, Lennon JT. 2015.** Fungal traits that drive ecosystem dynamics on land. *Microbiology and Molecular Biology Reviews* DOI: 10.1128/MMBR.00001-15
- Tulloss R, Yang ZL. 2016.** Amanitaceae Studies.
- Tulloss RE, Rodríguez-Caycedo C, Hughes KW, Geml J, Kudzma, L.V. W, B.E., Arora D. 2015.** Nomenclatural changes in *Amanita* II. *Amanitaceae* 1(2): 1-6.
- Tuno N, Takahashi KH, Yamashita H, Osawa N, Tanaka C. 2007.** Tolerance of *Drosophila* flies to ibotenic acid poisons in mushrooms. *Journal of Chemical Ecology* DOI: 10.1007/s10886-006-9228-3
- van Wilgen BW, Richardson DM. 2012.** Three centuries of managing introduced conifers in South Africa: benefits, impacts, changing perceptions and conflict resolution. *Journal of Environmental Management* DOI: 10.1016/j.jenvman.2012.03.052
- Vargas N, Goncalves SC, Franco-Molano AE, Restrepo S, Pringle A. 2019.** In Colombia the Eurasian fungus *Amanita muscaria* is expanding its range into native, tropical *Quercus humboldtii* forests. *Mycologia* DOI: 10.1080/00275514.2019.1636608
- Vellinga EC, Wolfe BE, Pringle A. 2009.** Global patterns of ectomycorrhizal introductions. *New Phytologist* 181(4): 960-973.
- Walton J 2018.** The cyclic peptide toxins of *Amanita* and other poisonous mushrooms. Switzerland: Springer International Publishing AG.
- Wang YW, McKeon MC, Elmore H, Hess J, Golan J, Gage H, Mao W, Harrow L, Goncalves SC, Hull CM, et al. 2023.** Invasive Californian death caps develop mushrooms unisexually and bisexually. *Nature Communications* DOI: 10.1038/s41467-023-42317-z
- Wasson RG. 1969.** Soma divine mushroom of immortality. *Mycologia* DOI: 10.2307/3757481
- Wolfe BE, Tulloss RE, Pringle A. 2012.** The irreversible loss of a decomposition pathway marks the single origin of an ectomycorrhizal symbiosis. *PLoS One* DOI: 10.1371/journal.pone.0039597
- Won TH, Bok JW, Nadig N, Venkatesh N, Nickles G, Greco C, Lim FY, González JB, Turgeon BG, Keller NP, et al. 2022.** Copper starvation induces antimicrobial isocyanide integrated into two distinct biosynthetic pathways in fungi. *Nature Communications* DOI: 10.1038/s41467-022-32394-x
- Xu W, Chooi Y-H, Choi JW, Li S, Vederas JC, Da Silva NA, Tang Y. 2013.** LovG: the thioesterase required for dihydromonacolin L release and lovastatin nonaketide synthase turnover in lovastatin biosynthesis. *Angewandte Chemie International Edition* 52(25): 6472-6475.
- Yu GC, Smith DK, Zhu HC, Guan Y, Lam TTY. 2017.** GGTREE: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution* DOI: 10.1111/2041-210x.12628

1076 SUPPORTING INFORMATION

1077 Supporting Information may be found online in the Supporting Information section at the end of
1078 the article.

1079 **Table 1** Collection metadata for the specimens collected and analyzed.

1080 **Table 2** Summary of the bioactivity of *Amanita muscaria* extracts against Methicillin-resistant
1081 *Staphylococcus aureus* (MRSA), *Pseudomonas aeruginosa*, *Candida auris*, *C. elegans*, *Brugia*
1082 spp., *Musca domestica*, and *Aedes aegypti*.

1083 **Fig. 1** Records of *Amanita muscaria* in South Africa.

1084 **Fig. 2** A species reconstruction of 184 *Amanita muscaria* and 5 *Amanita pantherina* isolates.

1085 **Fig. 3** Specialized metabolism (SM) in *Amanita muscaria* genomes.

1086 **Fig. 4** Analysis of Type 1 polyketide synthase (PKS) gene duplication found in all *Amanita*
1087 *muscaria* samples.

1088 **Fig. 5** The metabolome of *Amanita muscaria* grouped by the geographic region of the sample.

1089 **Dataset S1:** Full metadata for *A. muscaria* data retrieved from prior publications.

1090 **Dataset S2:** Genome summary statistics

1091 **Dataset S3:** Kinship analyses of the sequenced genomes

1092 **Table S1** Summary of the isolates used in this study.

1093 **Table S2** Summary of heterozygosity in sequenced genomes.

1094 **Fig. S1:** Dried tissue of each sample prior to chemical analyses.

1095 **Fig. S2:** Determination of optimal cutoff to generate the PKS GCF predictions.

1096 **Fig. S3:** Determination of optimal cutoff to generate the Terpene GCF predictions.

1097 **Fig. S4:** Rooted phylogenies from the fully sequenced *Amanita muscaria* genomes.

1098 **Fig. S5:** Tree compatibility comparisons between different species reconstruction methods.

1099 **Fig. S6:** A cblaster analysis showing the presence and absence of genes in the ibotenic acid
1100 cluster in the *A. muscaria* and outgroup genomes.

1101 **Fig. S7:** Whole fungal kingdom phylogeny depicting the number of Reciprocal best-hit BLAST
1102 hits to genes in the ibotenic acid gene cluster.

1103 **Fig. S8:** Quantity of putative muscimol in every sample.

- 1104 **Fig. S9:** Comparison of polyketide synthases found in *Amanita* genomes.
- 1105 **Fig. S10:** Full GNPS output of every molecular family, including singletons
- 1106 **Fig. S11:** Bioassays of metabolite extracts against Methicillin-resistant *Staphylococcus aureus*L.
- 1107 **Fig. S12:** Bioassays of metabolite extracts against *Pseudomonas aeruginosa*L.
- 1108 **Fig. S13:** Bioassays of metabolite extracts against *Candida auris*.
- 1109 **Fig. S14:** Nematode bioassay results after treatment with several concentrations of extracts.
- 1110 **Fig. S15:** Viability assays in *Ae. aegypti* and *M. domestica*.
- 1111 **Fig. S16:** GO-Term map corresponding to genes that overlapped with 5 kb sliding windows
1112 containing at least 100 SNPs and where estimates of Fst corresponded to the right 5% tail of
1113 estimates.
- 1114 **Fig. S17:** Phylogenetic trees constructed using a codon-aware alignment of the *iboF* and *iboH*
1115 genes.
- 1116
- 1117
- 1118