### SCORE Methods Workshop

Causal Inference: Day 1

Benjamin Grant Purzycki<sup>a</sup> Joseph Bulbulia<sup>b</sup>

> Aarhus University<sup>a</sup> Victoria University<sup>b</sup>

> > 2025

# Introduction Outline

• Theory of causality and causal inference

- Theory of causality and causal inference
  - Counterfactuals and potential outcomes
  - EEESI workflow

- Theory of causality and causal inference
  - Counterfactuals and potential outcomes
  - EEESI workflow
- Causal directed acyclic graphs (DAGs)

- Theory of causality and causal inference
  - Counterfactuals and potential outcomes
  - EEESI workflow
- Causal directed acyclic graphs (DAGs)
  - Introduction to DAGs
  - DAGITTY workshop

- Theory of causality and causal inference
  - Counterfactuals and potential outcomes
  - EEESI workflow
- Causal directed acyclic graphs (DAGs)
  - Introduction to DAGs
  - DAGITTY workshop
- Simulating causal systems and estimating causal effects

- Theory of causality and causal inference
  - Counterfactuals and potential outcomes
  - EEESI workflow
- Causal directed acyclic graphs (DAGs)
  - Introduction to DAGs
  - DAGITTY workshop
- Simulating causal systems and estimating causal effects
  - Conditional vs. marginal effects
  - Cross-sectional and longitudinal
  - G-Methods
  - Sensitivity checks

Causation is inferred

• Intervention, counterfactuals, possible worlds, potential outcomes

control	treatment
eyes are fine	bleeding eyeballs

Fundamental problem of causal inference

"...one can never observe the potential outcome under the treatment state for those observed in the control state, and one can never observe the potential outcome under the control state for those observed in the treatment state. This impossibility implies that one can never calculate individual-level causal effects"

[Morgan and Winship, 2015]

Group	$Y^0$	$Y^1$
Control	observable	counterfactual
Treatment	counterfactual	observable

Average Treatment Effect

$$ATE = E[Y^{X=1}] - E[Y^{X=0}]$$

Rethinking how we use statistical tools

- give up reliance of point estimates
- effectively abandon significance tests (e.g., p-values)
- estimate uncertainty
- direct vs. total effects
- "control" for counfounding and colliding effects
- reduce near-endless model specifications
- think and work harder on the front end
- expose how under-specified theories are

# Theory of causality and causal inference Causal inference is EEESI [Bendixen and Purzycki, 2025]

• Estimand

# Theory of causality and causal inference Causal inference is EEESI [Bendixen and Purzycki, 2025]

- Estimand
  - What are you trying to estimate?

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?
- Estimator

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?
- Estimator
  - How will you estimate the Estimand?

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?
- Estimator
  - How will you estimate the Estimand?
  - statistical model


- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?
- Estimator
  - How will you estimate the Estimand?
  - statistical model
- Estimate

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?
- Estimator
  - How will you estimate the Estimand?
  - statistical model
- Estimate
  - Apply Estimator

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?
- Estimator
  - How will you estimate the Estimand?
  - statistical model
- Estimate
  - Apply Estimator
  - Interpret results in light of Estimand

- Estimand
  - What are you trying to estimate?
  - Intervention? Outcome?
  - What is the process that generates it?
- Estimator
  - How will you estimate the Estimand?
  - statistical model
- Estimate
  - Apply Estimator
  - Interpret results in light of Estimand
- SImulate

Variable types



Biases in estimates





Mediator




Collider




"collider"

A note on "controls" (prediction)



A note on "controls" (causal inference)




Controls: The Good, The Bad, and the Inconsequential [Cinelli et al., 2024]

```
n = 100
\beta = 0.5
X, Y, Z \sim \text{Normal}(0, 1)
error terms \sim \text{Normal}(0,1)
iter = 1,000
Z \leftarrow rnorm(n, 0, 1)
X <- beta * Z + err1
Y <- beta * X + beta * Z + err2
lm(Y \sim X)
lm(Y \sim X + Z)
```


n = 100

Controls: The Good, The Bad, and the Inconsequential [Cinelli et al., 2024]


```
\begin{split} \beta &= 0.5 \\ X,Y,Z &\sim \mathrm{Normal}(0,1) \\ \mathrm{error\ terms} &\sim \mathrm{Normal}(0,1) \\ \mathrm{iter} &= 1,000 \\ \\ Z &<- \ \mathrm{rnorm}(\mathrm{n,\ 0,\ 1}) \\ X &<- \ \mathrm{beta} \ * \ Z \ + \ \mathrm{err1} \\ Y &<- \ \mathrm{beta} \ * \ X \ + \ \mathrm{beta} \ * \ Z \ + \ \mathrm{err2} \\ \mathrm{lm}(Y \ \sim \ X) \\ \mathrm{lm}(Y \ \sim \ X \ + \ Z) \end{split}
```




Controls: The Good (Confounder)




Controls: The Conditionally Good (Instrumental)



Controls: The Conditionally Good (Mediator)




Note on Mediation




- Direct vs. total effects
  - Direct: M and  $U_2$
  - Total:  $U_1$  but not M


Controls: The Bad (Collider)



Controls: The Bad (Post-Treatment)



Controls: The Inconsequential



### Modeling Causal Systems

What to include in your DAG

• theoretically relevant variables

- theoretically relevant variables
- precedent relevant variables

- theoretically relevant variables
- precedent relevant variables
- meta-analytic variables


- theoretically relevant variables
- precedent relevant variables
- meta-analytic variables
- sources of bias (e.g., sampling, confounders, etc.)

- theoretically relevant variables
- precedent relevant variables
- meta-analytic variables
- sources of bias (e.g., sampling, confounders, etc.)
- common sense/gumption variables

- theoretically relevant variables
- precedent relevant variables
- meta-analytic variables
- sources of bias (e.g., sampling, confounders, etc.)
- common sense/gumption variables
- de-biasing or instrumental variables

- theoretically relevant variables
- precedent relevant variables
- meta-analytic variables
- sources of bias (e.g., sampling, confounders, etc.)
- common sense/gumption variables
- de-biasing or instrumental variables
- many are defined by the structure of your model!

# Simulating Causal Systems DAGITTY workshop




### Simulating Causal Systems

#### R workshop

```
fd <- function(n, beta) {
 e_rel <- rnorm(n, 0, 1) # noise
 e edu <- rnorm(n, 0, 1)
 e mat <- rnorm(n, 0, 1)
 e_qua <- rnorm(n, 0, 1)
 e sex <- rnorm(n, 0, 1)
 e age <- rnorm(n, 0, 1)
 e_kid <- rnorm(n, 0, 1)
 SEX <- rbinom(n, 1, .5) # sex
 XIA \leftarrow rbinom(n, 1, .5) # edu type
 PRI <- rbinom(n, 1, .5) # prime
 AGE <- rnorm(n, 0, 1)
 EDU <- beta * SEX + beta * AGE + e edu
 MAT <- beta * SEX * beta * EDU + beta * AGE + e mat
 KID <- beta * MAT + beta * AGE + e kid # children
 REL <- beta * EDU + beta * XIA + beta * MAT + beta * SEX + beta * AGE +
   beta * PRI + beta * KID + e rel # religiosity
 df <- data.frame(SEX, XIA, PRI, AGE, EDU, MAT, REL)</pre>
 open0 <- coef(lm(REL \sim EDU, dat = df))[2]
 open1 <- coef(lm(REL \sim EDU + SEX, dat = df))[2]
 open2 <- coef(lm(REL \sim EDU + SEX + MAT, dat = df))[2]
 open3 <- coef(lm(REL ~ EDU + SEX + MAT + AGE + KID, dat = df))[2]
 closed <- coef(lm(REL \sim EDU + SEX + MAT + AGE + KID + XIA, dat = df))[2]
 return(c(open0, open1, open2, open3, closed))
```


### Simulating Causal Systems

R workshop



### Directed Acyclic Graphs

Assumptions of experimental data



- Estimand obtained when:
  - $\bullet$  random assignment of X
  - no unmeasured confounders
  - no reverse causation
  - no measurement error
  - ATE =  $E[Y^{X=1}] E[Y^{X=0}]$

### Directed Acyclic Graphs


Assumptions of cross-sectional data



- Estimand obtained when:
  - $\bullet$  random assignment of X
  - no unmeasured confounders
  - no reverse causation
  - no measurement error
  - ATE =  $E[Y^{X=1}] E[Y^{X=0}]$
- Sensitivity analysis

### Directed Acyclic Graphs

Assumptions of longitudinal data



#### Estimand obtained when:

- $\bullet$  random assignment of X
- no unmeasured confounders
- no reverse causation
- no measurement error
- ATE =  $E[Y^{X=1}] E[Y^{X=0}]$

## Marginal vs. Conditional Effects

#### g-computation


- 1. Predict values where X=0
- 2. Predict values where X = 1
- 3. Calculate ATE
- 4. Bootstrap/Go Bayes


```
set.seed(777)
bZ <- 2
Z <- rnorm(n, 0, 0.5)
X \leftarrow rbinom(n, 1, plogis(0.5 + Z * bZ))
Y \leftarrow rnorm(n, 10 + X * bX + Z * bZ)
d <- data.frame(Y, X, Z)
model \leftarrow lm(Y \sim X + Z, data = d)
d$EX1 <- predict(model,
                  newdata = transform(d, X = 1)
d$EX0 <- predict(model.
                  newdata = transform(d, X = 0))
mean(d$EX1) - mean(d$EX0)
bootstrap.fun <- function(data, indices) {
  d <- data[indices, ] # resample
  model <- lm(Y ~ X + Z, data = d) # fit model
  d$EX1 <- predict(model, newdata = transform(d, X = 1))</pre>
  d$EX0 <- predict(model, newdata = transform(d, X = 0))</pre>
  ate <- with(d, mean(EX1) - mean(EX0)) # ate
bootstrap.res <- boot(data = d,
                            statistic = bootstrap.fun.
                            R = iter)
(bootstrap.sum <- boot.ci(bootstrap.res, type = c("norm")))
```

#### Instrumental variables revisited

#### Assumptions

- V is independent of U
- V is a cause of X
- V does not cause Y other than through X





$$\frac{\mathrm{E}[Y|V=1] - \mathrm{E}[Y|V=0]}{\mathrm{E}[X|V=1] - \mathrm{E}[Y|V=0]}$$

Inverse probability weighting (IPW)

- 1. Estimate probability of X = 1 as function of confound
- 2. Invert probs. and stabilize weights
- 3. Fit outcome model with weights
- 4. Bootstrap/Go Bayes

Treated:  $\frac{\Pr(X_i)}{\Pr(X_i|Z_i)}$ Untreated:  $\frac{1-\Pr(X_i)}{1-\Pr(X_i|Z_i)}$ 

Doubly robust estimation

-

### Acknowledgments

- Theiss Bendixen
- Joseph Bulbulia
- Daniel Major-Smith
- Richard McElreath
- Julia Rohrer
- Brooke, Don, and Mike

### SCORE Methods Workshop

Causal Inference: Day 2

Joseph Bulbulia<sup>a</sup> Benjamin Grant Purzycki<sup>b</sup>

> Victoria University<sup>a</sup> Aarhus University<sup>b</sup>

> > 2025

### Bibliography I

Bendixen, T. and Purzycki, B. G. (2025).

The Data Analyst's Guide to Cause and Effect: An Introduction to Applied Causal Inference.

Sage, London.

Cinelli, C., Forney, A., and Pearl, J. (2024). A crash course in good and bad controls. Sociological Methods & Research, 53(3):1071-1104.

Morgan, S. and Winship, C. (2015). Counterfactuals and causal inference. Cambridge University Press, Cambridge.