

Quadratic Programming Assignment

Gautam Singh

Abstract—This document contains the solution to a modification to Question 27 of Exercise 5 in Chapter 6 of the class 12 NCERT textbook.

1) Find the point on the curve

$$x^2 = 2y \quad (1)$$

which is nearest to the point $\mathbf{P} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$.

Solution: We need to find

$$\min_{\mathbf{x}} g(\mathbf{x}) = \|\mathbf{x} - \mathbf{P}\|^2 \quad (2)$$

$$\text{s.t. } h(\mathbf{x}) = \mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} = 0 \quad (3)$$

where

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 0 \\ -1 \end{pmatrix} \quad (4)$$

We find the required minima using constrained gradient descent in Fig. 1, plotted using the Python code `codes/grad_pits.py`. The gradient descent update equation is

$$\mathbf{x}_{n+1} = \mathbf{x}_n - \alpha \operatorname{sgn} \left((\nabla g(\mathbf{x}_n))^\top (\nabla h(\mathbf{x}_n)) \right) \nabla h(\mathbf{x}_n) \quad (5)$$

$$\mathbf{x}_{n+1} = \mathbf{x}_n - 2\alpha \operatorname{sgn} \left((\mathbf{x}_n - \mathbf{P})^\top (\mathbf{V} \mathbf{x}_n + \mathbf{u}) \right) (\mathbf{V} \mathbf{x}_n + \mathbf{u}) \quad (6)$$

where (6) follows from (3) and the fact that the signum function is used to choose the direction in which gradient descent is followed.




Fig. 1: Gradient descent for a convex optimization problem.