

Probability Assignment

Gautam Singh

Abstract—This document contains the solution to Question 17 of Exercise 1 in Chapter 13 of the class 12 NCERT textbook.

1) If A and B are events such that

$$\Pr(A|B) = \Pr(B|A) \quad (1)$$

then

- a) $A \subset B$ but $A \neq B$
- b) $A = B$
- c) $A \cap B = \phi$
- d) $\Pr(A) = \Pr(B)$

Solution: Using Bayes' Rule,

$$\Pr(A, B) = \Pr(A) \Pr(B|A) \quad (2)$$

$$= \Pr(B) \Pr(A|B) \quad (3)$$

Using (1) in (2) and (3), we get $\Pr(A) = \Pr(B)$.
We consider the options one by one.

- a) If $A \subset B$ and $A \neq B$, then clearly $\Pr(A|B) = 1$. Therefore, from (1),

$$\Pr(B|A) = \Pr(A|B) = 1 \quad (4)$$

This implies that $B \subset A$, which is a contradiction.

- b) We give a counterexample to show this is wrong. Consider A to denote the event that an even number shows on rolling a fair die and B denote the event that a prime number shows on rolling a fair die. Then, $\Pr(A|B) = \Pr(B|A) = \frac{1}{3}$ but $A \neq B$. See Table 1.
- c) The same example as before provides the required counterexample, as 2 is an even prime number.
- d) This is the correct answer, as discussed above.

	A	\bar{A}
B	$\frac{1}{6}$	$\frac{1}{3}$
\bar{B}	$\frac{1}{3}$	$\frac{1}{6}$

TABLE 1: Joint pmf for events A and B .