

Conic Assignment

Gautam Singh

Abstract—This document contains the solution to Question 14 of Exercise 4 in Chapter 11 of the class 11 NCERT textbook.

- 1) Find the equation of the hyperbola eccentricity is $e = \frac{4}{3}$ and whose vertices are

$$\mathbf{P}_1 = \begin{pmatrix} 7 \\ 0 \end{pmatrix}, \mathbf{P}_2 = \begin{pmatrix} -7 \\ 0 \end{pmatrix} \quad (1)$$

Solution: Let the equation of the conic with focus \mathbf{F} , directrix $\mathbf{n}^\top \mathbf{x} = c$ and eccentricity e be

$$\mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} + f = 0 \quad (2)$$

where

$$\mathbf{V} \triangleq \|\mathbf{n}\|^2 \mathbf{I} - e^2 \mathbf{n} \mathbf{n}^\top \quad (3)$$

$$\mathbf{u} \triangleq c e^2 \mathbf{n} - \|\mathbf{n}\|^2 \mathbf{F} \quad (4)$$

$$f \triangleq \|\mathbf{n}\|^2 \|\mathbf{F}\|^2 - c^2 e^2 \quad (5)$$

Since the conic is a hyperbola whose vertices are given by (1), the major axis is the x -axis and the directrix is parallel to the y -axis. Hence,

$$\mathbf{n} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (6)$$

Thus,

$$\mathbf{V} = \begin{pmatrix} 1 - e^2 & 0 \\ 0 & 1 \end{pmatrix} \quad (7)$$

$$\mathbf{u} = c e^2 \begin{pmatrix} 1 \\ 0 \end{pmatrix} - \mathbf{F} \quad (8)$$

$$f = \|\mathbf{F}\|^2 - c^2 e^2 \quad (9)$$

The centre of the conic is the midpoint of all chords that pass through the centre. We know that the centre lies on the x -axis. Hence, \mathbf{c} bisects $P_1 P_2$, and so

$$\mathbf{c} = \frac{\mathbf{P}_1 + \mathbf{P}_2}{2} = \mathbf{0} \quad (10)$$

The general expression of the centre of the

circle is given by

$$\mathbf{c} = -\mathbf{V}^{-1} \mathbf{u} \quad (11)$$

Since $\mathbf{c} = \mathbf{0}$ and $\mathbf{V}^{-1} \neq \mathbf{0}$, it follows from (11) that $\mathbf{u} = \mathbf{0}$. Thus, from (8),

$$\mathbf{F} = \begin{pmatrix} c e^2 \\ 0 \end{pmatrix} \quad (12)$$

and so,

$$f = c^2 e^2 (e^2 - 1) \quad (13)$$

Putting $\mathbf{x} = \mathbf{P}_1$ or $\mathbf{x} = \mathbf{P}_2$ in (2) and using (12) and (13),

$$(\pm 7 \ 0) \begin{pmatrix} 1 - e^2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ \pm 7 \end{pmatrix} + f = 0 \quad (14)$$

$$\implies 49e^2 - f = 49 \quad (15)$$

Since $e = \frac{4}{3}$, (15) implies

$$f = 49(e^2 - 1) = \frac{343}{9} \quad (16)$$

Therefore, the equation of the conic is

$$\mathbf{x}^\top \begin{pmatrix} -\frac{7}{9} & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} + \frac{343}{9} = 0 \quad (17)$$

The situation is illustrated in Fig. 1.




Fig. 1: Locus of the required hyperbola.