

Optimization Assignment

Gautam Singh

Abstract—This document contains the solution to Question 4 of Exercise 2 in Chapter 10 of the class 11 NCERT textbook.

- 1) Find the coordinates of the foot of perpendicular from the point

$$\mathbf{P} = \begin{pmatrix} -1 \\ 3 \end{pmatrix} \quad (1)$$

to the line

$$(3 \quad -4)\mathbf{x} = 16 \quad (2)$$

Solution: Any point on (2) is clearly of the form

$$\mathbf{Q} = \mathbf{A} + \lambda \mathbf{m} \quad (3)$$

where $\lambda \in \mathbb{R}$ and

$$\mathbf{A} = \begin{pmatrix} 0 \\ -4 \end{pmatrix}, \quad \mathbf{m} = \begin{pmatrix} 4 \\ 3 \end{pmatrix} \quad (4)$$

Thus,

$$f(\lambda) = \|\mathbf{Q} - \mathbf{P}\|^2 \quad (5)$$

$$= \|\mathbf{A} - \mathbf{P} + \lambda \mathbf{m}\|^2 \quad (6)$$

$$= \|\mathbf{m}\|^2 \lambda^2 + 2\mathbf{m}^\top (\mathbf{A} - \mathbf{P}) \lambda + \|\mathbf{A} - \mathbf{P}\|^2 \quad (7)$$

Since (7) is convex, we use the gradient descent function on λ to converge at the minimum of $f(\lambda)$.

$$\lambda_{n+1} = \lambda_n - \alpha f'(\lambda_n) \quad (8)$$

$$= (1 - 2\alpha \|\mathbf{m}\|^2) \lambda_n + 2\alpha \mathbf{m}^\top (\mathbf{A} - \mathbf{P}) \quad (9)$$

Taking the one-sided Z-transform on both sides of (9),

$$z\Lambda(z) = (1 - 2\alpha \|\mathbf{m}\|^2) \Lambda(z) - \frac{2\alpha \mathbf{m}^\top (\mathbf{A} - \mathbf{P})}{1 - z^{-1}} \quad (10)$$

Solving (10)

$$\Lambda(z) = -\frac{2\alpha \mathbf{m}^\top (\mathbf{A} - \mathbf{P}) z^{-1}}{(1 - z^{-1})(1 - (1 - 2\alpha \|\mathbf{m}\|^2) z^{-1})} \quad (11)$$

$$= -\frac{\mathbf{m}^\top (\mathbf{A} - \mathbf{P})}{\|\mathbf{m}\|^2} \left(\frac{1}{1 - z^{-1}} \right) \quad (12)$$

$$= -\frac{1}{1 - (1 - 2\alpha \|\mathbf{m}\|^2) z^{-1}} \quad (13)$$

$$= -\frac{\mathbf{m}^\top (\mathbf{A} - \mathbf{P})}{\|\mathbf{m}\|^2} \sum_{k=0}^{\infty} (1 - (1 - 2\alpha \|\mathbf{m}\|^2)^k) z^{-k} \quad (14)$$

From (11), the ROC is

$$|z| > \max\{1, 1 - 2\alpha \|\mathbf{m}\|^2\} \quad (15)$$

$$\implies 0 < 1 - 2\alpha \|\mathbf{m}\|^2 < 1 \quad (16)$$

$$\implies 0 < \alpha < \frac{1}{2 \|\mathbf{m}\|^2} \quad (17)$$

Thus, if α satisfies (17), then from (14), substituting from (4),

$$\lim_{n \rightarrow \infty} \lambda_n = -\frac{\mathbf{m}^\top (\mathbf{A} - \mathbf{P})}{\|\mathbf{m}\|^2} = \frac{17}{25} \quad (18)$$

We select the following parameters to arrive at the optimal λ , where N is the number of iterations and ϵ is the convergence limit. The gradient descent is demonstrated in Fig. 1, plotted by the Python code `codes/grad_desc.py`. The relevant parameters are shown in Table 1.

Parameter	Value
λ_0	0
α	0.1
N	1000000
ϵ	10^{-6}

TABLE 1: Parameters for Gradient Descent




Fig. 1: Gradient descent to get the optimal λ .