
CY104S1_Practical Exam-2022

Please read the questions carefully, there are 50 MCQ type questions, each carrying one mark and only one correct option. The duration of exam is 1:00 pm-2:00 pm (One hour), please submit the response before 2:00 pm.

1

Name of the student *

Garvit Shah

2

Admission number *

U21CS089

3

Division *

F

CY104S1_Practical Exam-2022

Please read the questions carefully, there are 50 MCQ type questions, each carrying one mark and only one correct option. The duration of exam is 1:00 pm-2:00 pm (One hour), please submit the response before 2:00 pm.

4

What is the role of adding conc. Sulphuric acid and conc. Nitric acid in the electrolysis beaker for the electrodeposition of Cu? *
(1 Point)

- ☐ To maintain the acidity of the solution.
- ☐ To oxidize the Cu atoms.
- ☐ To prevent the accumulation of gases near the electrodes.
- ☒ All the above.

5

Why a gauze type cathode is used in the electrodeposition of Cu? *
(1 Point)

- ☐ Because it is cheap and easily available.
- ☐ So that it can readily dissolve in the electrolytic solution.
- ☐ Because its weight will be minimal.
- ☒ To increase the surface area required for electrodeposition.

6

What kind of electrode is not advisable for cathode for studying electrodeposition of Cu? *
(1 Point)

- ☒ Copper
- ☐ Platinum

- ☐ Graphite
- ☐ None of the Above

7

Which of the following interferes in the electrodeposition of Cu? *
(1 Point)

- ☐ Nitrate
- ☐ Nitrite
- ☒ Silver
- ☐ All the above

8

What is the discharge potential of Cu? *
(1 Point)

- ☐ + 0.325 V
- ☐ -0.15 V
- ☐ -0.325 V
- ☒ + 0.35 V

9

Which of the following is not true about iodometric titration? *
(1 Point)

- ☐ It's not a direct iodine titration.
- ☐ It's a kind of redox titration.

- ☐ Sodium thiosulphate is generally used as titrant.
- ☒ None of the above.

10

In the Winkler method_____ *

(1 Point)

- ☐ Dissolved oxygen oxidizes iodine
- ☐ Dissolved oxygen oxidizes iodide
- ☐ Atmospheric oxygen oxidizes iodine
- ☒ Dissolved oxygen oxidizes manganese (+2) ion

11

During the titration in Winkler's method, what is the fate of thiosulphate ion? *

(1 Point)

- ☒ It forms tetrathionate ion.
- ☐ It forms colloidal solution of sulphur.
- ☐ It is converted to sulphur dioxide.
- ☐ It is converted to sulphate ion.

12

In Winkler method, what is the correct time to add starch? *

(1 Point)

- ☒ After titrating until pale yellow colour is obtained.
- ☐ just before starting the titration, when the colour is brownish yellow.

- ☐ before adding Manganese Sulphate.
- ☐ After titrating until colourless solution is obtained.

13

3.84 grams of dissolved oxygen will consume____ *

(1 Point)

- ☒ 119.12 grams of sodium thiosulphate pentahydrate.
- ☐ 75.89 grams of sodium thiosulphate pentahydrate.
- ☐ 15.36 grams of sodium thiosulphate pentahydrate.
- ☐ 29.78 grams of sodium thiosulphate pentahydrate.

14

The last line in the plot of Conductance v/s Volume of NaOH in all the titration is due to *

(1 Point)

- ☐ Decrease in the OH⁻ ions
- ☐ Increase in H⁺ ion
- ☒ Increase in the OH⁻ ions
- ☐ Decrease in H⁺ ion

15

The hydrogen ion has highest conductance due to____ *

(1 Point)

- ☐ High mobility
- ☐ Small size

☒ Both the above

☐ None of these

16

As temperature increases, electrolytic conduction____ *

(1 Point)

☐ Decreases

☒ Increases

☐ Remains unaffected

☐ None of the above

17

On dilution, the specific conductance____ *

(1 Point)

☐ Increase

☐ Remains same

☒ Decrease

☐ None of the mentioned

18

How will you prepare 0.1 N, 250 mL sodium thiosulphate solution *

(1 Point)

☐ By weighing 4.0 of sodium thiosulphate in 250 mL of water

☒ By weighing 6.2 of sodium thiosulphate in 250 mL of water

- ☐ By weighing 7.2 of sodium thiosulphate in 250 mL of water
- ☐ By weighing 10 g of sodium thiosulphate in 250 mL of water

19

The titrimetric determination of ascorbic acid is a type of ____ *

(1 Point)

- ☐ Iodometric titration
- ☒ Iodimetric titration
- ☐ Complexometric titration
- ☐ Acid-base titration

20

While the potassium iodide is a powerful reducing agent, *

(1 Point)

- ☐ Iodine is oxidised to iodide ion
- ☐ Iodine is reduced to iodide ion
- ☒ The iodide ion is oxidized to iodine
- ☐ The iodide ion is reduced to iodine

21

Starch-iodide paste is used as an external indicator in one of the following titrations. *

(1 Point)

- ☐ Iodometric titration of copper sulphate using sodium thiosulphate as titrant
- ☒ Iodimetric titration of ascorbic acid using iodine solution as titrant

- ☐ Potassium dichromate titration using sodium thiosulphate as titrant.
- ☐ Diazotisation titration of sulphadiazine using sodium nitrite as titrant

22

Ascorbic acid works as ____ *

(1 Point)

- ☒ Reducing agent
- ☐ Oxidizing agent
- ☐ Both the above
- ☐ None of the above

23

Which of the following cannot be used as secondary reference electrode? *

(1 Point)

- ☒ Glass electrode
- ☐ Mercury-mercury sulphate electrode
- ☐ Silver-silver chloride electrode
- ☐ Calomel electrode

24

Which of the following statement is not true? *

(1 Point)

- ☐ The saturated calomel electrode is a reference electrode
- ☐ The platinum electrode functions as a cathode

- ☒ Ceric ammonium sulfate act as a reducing agent in potentiometric titration
- ☐ The Indicator electrode is a platinum electrode

25

Calomel electrode can behave as _____ *

(1 Point)

- ☐ cathode only
- ☒ anode or cathode
- ☐ salt bridge
- ☐ anode only

26

In an electrochemical cell, reduction takes place at *

(1 Point)

- ☒ cathode
- ☐ anode
- ☐ salt bridge
- ☐ It depends on the potential applying

27

Which of the following is not the characteristic of a calomel electrode? *

(1 Point)

- ☒ The potential of the electrode is not temperature-dependent
- ☐ Value of potential or emf is higher for a lower concentration of KCl

- ☐ Preparation of electrode is easy
- ☐ Value of potential decreases with increasing concentration of KCl

28

Consider the titration of 30.0 mL of 0.20 M Acetic acid by adding 0.0500 M aqueous ammonia Hydroxide to it. The pH at the equivalence point is____ *

(1 Point)

- ☐ Is impossible to predict
- ☒ Less than 7
- ☐ Equal to 7
- ☐ Greater than 7

29

Consider a solution which is 0.10 M in CH_3COOH and 0.20 M in NaCH_3COO . Which of the following statements is true? *

(1 Point)

- ☐ If a small amount of NaOH is added, the pH decreases very slightly.
- ☐ If HCl is added, the H^+ ions react with CH_3COOH ions.
- ☐ If NaOH is added, the OH^- ions react with the CH_3COO^- ions.
- ☒ If a small amount of HCl is added, the pH decreases very slightly.

30

Which of the following statement is not true? *

(1 Point)

- ☒ The glass electrode is not sensitive to H^+ ion concentration

- ☐ The pH meter measures the total potential across the two electrodes
- ☐ The value of the Ionic product (K_w) is temperature-dependent
- ☐ Calomel electrode potential is constant

31

Why is the Standard hydrogen electrode called as the primary reference electrode? *

(1 Point)

- ☐ It has a known output potential
- ☐ Its output potential is independent of the composition of the solution
- ☐ It has a constant output potential
- ☒ Its output potential is zero volts

32

Choose the wrong statement for spectrometric analysis *

(1 Point)

- ☐ An absorbance of 0 at some wavelength means that no light of that particular wavelength has been absorbed
- ☐ On most of the diagrams you will come across, the absorbance ranges from 0 to 1, but it can go higher than that
- ☒ molar extinction coefficient is a measure of the probability of the electronic transition
- ☐ None of the above

33

Which of the following is not in the Beer Lambert's Law equation? *

(1 Point)

- ☐ concentration
- ☒ light wavelength
- ☐ cell path length
- ☐ molar absorptivity

34

Cell path length can be defined as____ *

(1 Point)

- ☐ distance between the light source and detector
- ☐ distance between light and object
- ☒ distance light travels through the substance
- ☐ the wavelength of the light

35

Unit of molar molar absorptivity is____ *

(1 Point)

- ☐ $L^{-1} mol\ cm^{-1}$
- ☒ $L\ mol^{-1}\ cm^{-2}$
- ☐ $L^{-1} mol^{-1}\ cm^{-1}$
- ☐ $L\ mol^{-1}\ cm^{-1}$

36

Beer Lambert's law gives the relation between which of the following? *

(1 Point)

- ☐ Reflected radiation and concentration
- ☐ Energy absorption and reflected radiation
- ☐ Scattered radiation and concentration
- ☒ Energy absorption and concentration

37

Lambert's law states that the intensity of light decreases with respect to *

(1 Point)

- ☐ Distance
- ☐ Volume
- ☒ Concentration
- ☐ Composition

38

_____ is the amount of oxygen required to oxidize only organic matter in sewage. *

(1 Point)

- ☐ DO
- ☐ Turbidity
- ☐ COD
- ☒ BOD

39

We can determine COD by_____ *

(1 Point)

- ☐ complexometric titration
- ☒ Back redox titration
- ☐ direct redox titration
- ☐ acid-base titration

40

Indicator that is used in determination of COD *

(1 Point)

- ☐ FAS
- ☒ Ferroin
- ☐ Phenolphthalein
- ☐ Starch

41

To test chemical oxygen demand (C.O.D.) of sewage, organic matter is oxidised by potassium dichromate in the presence of _____? *

(1 Point)

- ☐ Hydrochloric acid
- ☐ Citric acid
- ☒ Sulphuric acid
- ☐ Nitric acid

42

A 50 ml of a sewage water sample was refluxed with 10 ml of 0.25N $K_2Cr_2O_7$ solution of dil H_2SO_4 . The unreacted dichromate required 5 ml of 0.15 N ferrous ammonium sulphate. 10 ml of distilled water, under the same conditions as the sample, required 20 ml of 0.15 N ferrous ammonium sulphate. Calculate the COD of the sewage water sample. *

(1 Point)

- ☒ 360 ppm
- ☐ 370 ppm
- ☐ 300 ppm
- ☐ 350 ppm

43

The pH range is to be maintained in EDTA titration *

(1 Point)

- ☐ 3
- ☒ 10
- ☐ 7
- ☐ 8

44

Color change in the following reaction is-
Hard water + EBT
indicator complex

= Metal ion –

*

(1 Point)

- ☒ blue to red

- ☐ colorless to red
- ☐ Red to blue
- ☐ blue to colorless

45

In the determination of Cu in brass sample solution, the re-dissolution of precipitate is done by adding *

(1 Point)

- ☐ KI solution
- ☒ Acetic acid
- ☐ HNO₃
- ☐ Ammonium hydroxide

46

The salt _____ is a hardness producing substance *

(1 Point)

- ☐ Na_2SO_4
- ☒ $MgCl_2$
- ☐ $NaCl$
- ☐ Na_2CO_3

47

Naturally occurring ascorbic acid is found in _____ *

(1 Point)

- ☐ None of the above

- ☒ L form
- ☐ D and L both
- ☐ D form

48

Which one is used as primary Standard Solution in laboratory *
(1 Point)

- ☐ H_2SO_4
- ☐ $NaOH$
- ☐ HCl
- ☒ $K_2Cr_2O_7$

49

Which of the following acts as a self - indicator? *
(1 Point)

- ☐ $K_2Cr_2O_7$
- ☐ Oxalic acid
- ☐ Iodine
- ☒ $KMnO_4$

50

Ascorbic acid is a _____ *
(1 Point)

- ☐ cyclic ketone


- ☐ cyclic ether
- ☒ cyclic ester
- ☐ aldehyde

51

Erichrome black-T is a type of indicator *
(1 Point)


- ☒ Metal ion as well as acid-base indicator
- ☐ Metal ion indicator
- ☐ Acid-base indicator
- ☐ Redox indicator

52



The following pH-metric titration curve is the kind of curve expected for the titration of a ____ acid with a ____ base *
(1 Point)

- ☐ weak, weak
- ☐ weak, strong
- ☐ strong, weak

 strong, strong

53

Which indicator is used in potentiometric titration? *
(1 Point)

- ☐ Methyl orange
- ☐ Electrolyte solution
- ☐ EBT solution
- ☒ None of the above

This content is created by the owner of the form. The data you submit will be sent to the form owner. Microsoft is not responsible for the privacy or security practices of its customers, including those of this form owner. Never give out your password.

Powered by Microsoft Forms |

The owner of this form has not provided a privacy statement as to how they will use your response data. Do not provide personal or sensitive information.

| [Terms of use](#)