


Trig Cheat Sheet

Definition of the Trig Functions

Right triangle definition

For this definition we assume that

$$0 < \theta < \frac{\pi}{2} \text{ or } 0^\circ < \theta < 90^\circ.$$




$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}} \quad \csc \theta = \frac{\text{hypotenuse}}{\text{opposite}}$$

$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}} \quad \sec \theta = \frac{\text{hypotenuse}}{\text{adjacent}}$$

$$\tan \theta = \frac{\text{opposite}}{\text{adjacent}} \quad \cot \theta = \frac{\text{adjacent}}{\text{opposite}}$$

Unit circle definition

For this definition θ is any angle.



$$\sin \theta = \frac{y}{1} = y \quad \csc \theta = \frac{1}{y}$$

$$\cos \theta = \frac{x}{1} = x \quad \sec \theta = \frac{1}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

Facts and Properties

Domain

The domain is all the values of θ that can be plugged into the function.

$\sin \theta$, θ can be any angle

$\cos \theta$, θ can be any angle

$\tan \theta$, $\theta \neq \left(n + \frac{1}{2}\right)\pi$, $n = 0, \pm 1, \pm 2, \dots$

$\csc \theta$, $\theta \neq n\pi$, $n = 0, \pm 1, \pm 2, \dots$

$\sec \theta$, $\theta \neq \left(n + \frac{1}{2}\right)\pi$, $n = 0, \pm 1, \pm 2, \dots$

$\cot \theta$, $\theta \neq n\pi$, $n = 0, \pm 1, \pm 2, \dots$

Range

The range is all possible values to get out of the function.

$$-1 \leq \sin \theta \leq 1 \quad \csc \theta \geq 1 \text{ and } \csc \theta \leq -1$$

$$-1 \leq \cos \theta \leq 1 \quad \sec \theta \geq 1 \text{ and } \sec \theta \leq -1$$

$$-\infty < \tan \theta < \infty \quad -\infty < \cot \theta < \infty$$

Period

The period of a function is the number, T , such that $f(\theta + T) = f(\theta)$. So, if ω is a fixed number and θ is any angle we have the following periods.

$$\sin(\omega \theta) \rightarrow T = \frac{2\pi}{\omega}$$

$$\cos(\omega \theta) \rightarrow T = \frac{2\pi}{\omega}$$

$$\tan(\omega \theta) \rightarrow T = \frac{\pi}{\omega}$$

$$\csc(\omega \theta) \rightarrow T = \frac{2\pi}{\omega}$$

$$\sec(\omega \theta) \rightarrow T = \frac{2\pi}{\omega}$$

$$\cot(\omega \theta) \rightarrow T = \frac{\pi}{\omega}$$

Formulas and Identities

Tangent and Cotangent Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

Reciprocal Identities

$$\csc \theta = \frac{1}{\sin \theta} \quad \sin \theta = \frac{1}{\csc \theta}$$

$$\sec \theta = \frac{1}{\cos \theta} \quad \cos \theta = \frac{1}{\sec \theta}$$

$$\cot \theta = \frac{1}{\tan \theta} \quad \tan \theta = \frac{1}{\cot \theta}$$

Pythagorean Identities

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

Even/Odd Formulas

$$\sin(-\theta) = -\sin \theta \quad \csc(-\theta) = -\csc \theta$$

$$\cos(-\theta) = \cos \theta \quad \sec(-\theta) = \sec \theta$$

$$\tan(-\theta) = -\tan \theta \quad \cot(-\theta) = -\cot \theta$$

Periodic Formulas

If n is an integer.

$$\sin(\theta + 2\pi n) = \sin \theta \quad \csc(\theta + 2\pi n) = \csc \theta$$

$$\cos(\theta + 2\pi n) = \cos \theta \quad \sec(\theta + 2\pi n) = \sec \theta$$

$$\tan(\theta + \pi n) = \tan \theta \quad \cot(\theta + \pi n) = \cot \theta$$

Double Angle Formulas

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$$

$$= 2 \cos^2 \theta - 1$$

$$= 1 - 2 \sin^2 \theta$$

$$\tan(2\theta) = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

Degrees to Radians Formulas

If x is an angle in degrees and t is an angle in radians then

$$\frac{\pi}{180} = \frac{t}{x} \quad \Rightarrow \quad t = \frac{\pi x}{180} \quad \text{and} \quad x = \frac{180t}{\pi}$$

Half Angle Formulas (alternate form)

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}} \quad \sin^2 \theta = \frac{1}{2}(1 - \cos(2\theta))$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}} \quad \cos^2 \theta = \frac{1}{2}(1 + \cos(2\theta))$$

$$\tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}} \quad \tan^2 \theta = \frac{1 - \cos(2\theta)}{1 + \cos(2\theta)}$$

Sum and Difference Formulas

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$$

Product to Sum Formulas

$$\sin \alpha \sin \beta = \frac{1}{2}[\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

$$\cos \alpha \cos \beta = \frac{1}{2}[\cos(\alpha - \beta) + \cos(\alpha + \beta)]$$

$$\sin \alpha \cos \beta = \frac{1}{2}[\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$

$$\cos \alpha \sin \beta = \frac{1}{2}[\sin(\alpha + \beta) - \sin(\alpha - \beta)]$$

Sum to Product Formulas

$$\sin \alpha + \sin \beta = 2 \sin\left(\frac{\alpha + \beta}{2}\right) \cos\left(\frac{\alpha - \beta}{2}\right)$$

$$\sin \alpha - \sin \beta = 2 \cos\left(\frac{\alpha + \beta}{2}\right) \sin\left(\frac{\alpha - \beta}{2}\right)$$

$$\cos \alpha + \cos \beta = 2 \cos\left(\frac{\alpha + \beta}{2}\right) \cos\left(\frac{\alpha - \beta}{2}\right)$$

$$\cos \alpha - \cos \beta = -2 \sin\left(\frac{\alpha + \beta}{2}\right) \sin\left(\frac{\alpha - \beta}{2}\right)$$


Cofunction Formulas

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta \quad \cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$$

$$\csc\left(\frac{\pi}{2} - \theta\right) = \sec \theta \quad \sec\left(\frac{\pi}{2} - \theta\right) = \csc \theta$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta \quad \cot\left(\frac{\pi}{2} - \theta\right) = \tan \theta$$

Unit Circle



For any ordered pair on the unit circle (x, y) : $\cos \theta = x$ and $\sin \theta = y$

Example

$$\cos\left(\frac{5\pi}{3}\right) = \frac{1}{2} \quad \sin\left(\frac{5\pi}{3}\right) = -\frac{\sqrt{3}}{2}$$

Inverse Trig Functions

Definition

$y = \sin^{-1} x$ is equivalent to $x = \sin y$

$y = \cos^{-1} x$ is equivalent to $x = \cos y$

$y = \tan^{-1} x$ is equivalent to $x = \tan y$

Inverse Properties

$$\cos(\cos^{-1}(x)) = x \quad \cos^{-1}(\cos(\theta)) = \theta$$

$$\sin(\sin^{-1}(x)) = x \quad \sin^{-1}(\sin(\theta)) = \theta$$

$$\tan(\tan^{-1}(x)) = x \quad \tan^{-1}(\tan(\theta)) = \theta$$

Domain and Range

Function	Domain	Range
$y = \sin^{-1} x$	$-1 \leq x \leq 1$	$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$
$y = \cos^{-1} x$	$-1 \leq x \leq 1$	$0 \leq y \leq \pi$
$y = \tan^{-1} x$	$-\infty < x < \infty$	$-\frac{\pi}{2} < y < \frac{\pi}{2}$


Alternate Notation

$$\sin^{-1} x = \arcsin x$$

$$\cos^{-1} x = \arccos x$$

$$\tan^{-1} x = \arctan x$$

Law of Sines, Cosines and Tangents



Law of Sines

$$\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}$$

Law of Cosines

$$a^2 = b^2 + c^2 - 2bc \cos \alpha$$

$$b^2 = a^2 + c^2 - 2ac \cos \beta$$

$$c^2 = a^2 + b^2 - 2ab \cos \gamma$$

Law of Tangents

$$\frac{a-b}{a+b} = \frac{\tan \frac{1}{2}(\alpha - \beta)}{\tan \frac{1}{2}(\alpha + \beta)}$$

$$\frac{b-c}{b+c} = \frac{\tan \frac{1}{2}(\beta - \gamma)}{\tan \frac{1}{2}(\beta + \gamma)}$$

$$\frac{a-c}{a+c} = \frac{\tan \frac{1}{2}(\alpha - \gamma)}{\tan \frac{1}{2}(\alpha + \gamma)}$$

Mollweide's Formula

$$\frac{a+b}{c} = \frac{\cos \frac{1}{2}(\alpha - \beta)}{\sin \frac{1}{2}\gamma}$$

Trigonometric Identities & Formulas

Tutorial Services – Mission del Paso Campus

Reciprocal Identities

$$\sin x = \frac{1}{\csc x}$$

$$\csc x = \frac{1}{\sin x}$$

$$\cos x = \frac{1}{\sec x}$$

$$\sec x = \frac{1}{\cos x}$$

$$\tan x = \frac{1}{\cot x}$$

$$\cot x = \frac{1}{\tan x}$$

Ratio or Quotient Identities

$$\tan x = \frac{\sin x}{\cos x}$$

$$\cot x = \frac{\cos x}{\sin x}$$

$$\sin x = \cos x \tan x$$

$$\cos x = \sin x \cot x$$

Pythagorean Identities

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$1 + \cot^2 x = \csc^2 x$$

Note: there are only three, basic Pythagorean identities, the other forms are the same three identities, just arranged in a different order.

Pythagorean Identities in Radical Form

$$\sin x = \pm \sqrt{1 - \cos^2 x}$$

$$\tan x = \pm \sqrt{\sec^2 x - 1}$$

$$\cos x = \pm \sqrt{1 - \sin^2 x}$$

Confunction Identities

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x$$

$$\cos\left(\frac{\pi}{2} - x\right) = \sin x$$

$$\tan\left(\frac{\pi}{2} - x\right) = \cot x$$

$$\cot\left(\frac{\pi}{2} - x\right) = \tan x$$

$$\sec\left(\frac{\pi}{2} - x\right) = \csc x$$

$$\csc\left(\frac{\pi}{2} - x\right) = \sec x$$

Odd-Even Identities

Also called negative angle identities

$$\sin(-x) = -\sin x \quad \csc(-x) = -\csc x$$

$$\cos(-x) = \cos x \quad \sec(-x) = \sec x$$

$$\tan(-x) = -\tan x \quad \cot(-x) = -\cot x$$

$$\text{Phase Shift} = \frac{-c}{b}$$

$$\text{Period} = \frac{2\pi}{b}$$

Sum and Difference Formulas/Identities

$$\sin(u + v) = \sin u \cos v + \cos u \sin v$$

$$\sin(u - v) = \sin u \cos v - \cos u \sin v$$

$$\cos(u + v) = \cos u \cos v - \sin u \sin v$$

$$\cos(u - v) = \cos u \cos v + \sin u \sin v$$

$$\tan(u + v) = \frac{\tan u + \tan v}{1 - \tan u \tan v}$$

$$\tan(u - v) = \frac{\tan u - \tan v}{1 + \tan u \tan v}$$

How to Find Reference Angles

Step 1: Determine which quadrant the angle is in

Step 2: Use the appropriate formula

Quad I = is the angle itself

Quad II = $180 - \theta$ or $\pi - \theta$

Quad III = $\theta - 180$ or $\theta - \pi$

Quad IV = $360 - \theta$ or $2\pi - \theta$

Reciprocal Identities

$$\sin x = \frac{1}{\csc x} \quad \csc x = \frac{1}{\sin x}$$

$$\cos x = \frac{1}{\sec x} \quad \sec x = \frac{1}{\cos x}$$

$$\tan x = \frac{1}{\cot x} \quad \cot x = \frac{1}{\tan x}$$

Ratio or Quotient Identities

$$\tan x = \frac{\sin x}{\cos x} \quad \cot x = \frac{\cos x}{\sin x}$$

$$\sin x = \cos x \tan x \quad \cos x = \sin x \cot x$$

Pythagorean Identities

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$1 + \cot^2 x = \csc^2 x$$

Pythagorean Identities in Radical Form

$$\sin x = \pm \sqrt{1 - \cos^2 x}$$

$$\tan x = \pm \sqrt{\sec^2 x - 1}$$

Note: there are only three, basic Pythagorean identities, the other forms are the same three identities, just arranged in a different order.

Confunction Identities

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x \quad \cos\left(\frac{\pi}{2} - x\right) = \sin x$$

$$\tan\left(\frac{\pi}{2} - x\right) = \cot x \quad \cot\left(\frac{\pi}{2} - x\right) = \tan x$$

$$\sec\left(\frac{\pi}{2} - x\right) = \csc x \quad \csc\left(\frac{\pi}{2} - x\right) = \sec x$$

Odd-Even Identities

Also called negative angle identities

$$\sin(-x) = -\sin x \quad \csc(-x) = -\csc x$$

$$\cos(-x) = \cos x \quad \sec(-x) = \sec x$$

$$\tan(-x) = -\tan x \quad \cot(-x) = -\cot x$$

Sum and Difference Formulas - Identities

$$\sin(u + v) = \sin u \cos v + \cos u \sin v$$

$$\sin(u - v) = \sin u \cos v - \cos u \sin v$$

$$\cos(u + v) = \cos u \cos v - \sin u \sin v$$

$$\cos(u - v) = \cos u \cos v + \sin u \sin v$$

$$\tan(u + v) = \frac{\tan u + \tan v}{1 - \tan u \tan v}$$

$$\tan(u - v) = \frac{\tan u - \tan v}{1 + \tan u \tan v}$$


Right Triangle Definitions of Trigonometric Functions

Note: sin & cos are complementary angles, so are tan & cot and sec & csc, and the sum of complementary angles is 90 degrees.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{y}{r} \quad \csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{r}{y}$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{x}{r} \quad \sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{r}{x}$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{y}{x} \quad \cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{x}{y}$$



Adjacent = is the side adjacent to the angle in consideration. So if we are considering Angle A, then the adjacent side is CB

Trigonometric Values of Special Angles

Degrees	0°	30°	45°	60°	90°	180°	270°
Radians	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	π	$\frac{3\pi}{2}$
sinθ	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	0	-1
cosθ	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	-1	0
tanθ	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	undefined	0	undefined

To Convert Degrees to Radians, Multiply by $\frac{\pi \text{ rad}}{180 \text{ deg}}$

To Convert Radians to Degrees, Multiply by $\frac{180 \text{ deg}}{\pi \text{ rad}}$

Vocabulary

- Cotangent Angles - are two angles with the same terminal side
- Reference Angle - is an acute angle formed by terminal side of angle(α) with x-axis

$2ab$

Finding the Area of non-90degree Triangles

Area of an Oblique Triangle

$$area = \frac{1}{2}bc \sin A = \frac{1}{2}ab \sin C = \frac{1}{2}ac \sin B$$

Heron's Formula

Step 1: Find "s" $s = \frac{(a + b + c)}{2}$

Step 2: Use the formula $area = \sqrt{s(s-a)(s-b)(s-c)}$



Function Ranges:

$\sin(x)$	$-1 \leq y \leq 1$	$\arcsin(x)$	$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$
$\cos(x)$	$-1 \leq y \leq 1$	$\arccos(x)$	$0 \leq y \leq \pi$
$\tan(x)$	$-\infty < y < \infty$	$\arctan(x)$	$-\frac{\pi}{2} < y < \frac{\pi}{2}$
$\cot(x)$	$-\infty < y < \infty$	$\operatorname{arccot}(x)$	$0 < y < \pi$
$\csc(x)$	$-\infty < y \leq 1$ $\cup 1 \leq y < \infty$	$\operatorname{arccsc}(x)$	$0 \leq y < \frac{\pi}{2} \cup \pi \leq y < \frac{3\pi}{2}$
$\sec(x)$	$-\infty < y \leq 1 \cup$ $1 \leq y < \infty$	$\operatorname{arcsec}(x)$	$-\pi < y \leq -\frac{\pi}{2} \cup 0 < y < \frac{\pi}{2}$

Function Values:

	$\sin(x)$	$\cos(x)$	$\tan(x)$	$\cot(x)$
0	0	1	0	Undefined
$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	$\sqrt{3}$
$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	1
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{\sqrt{3}}{3}$
$\frac{\pi}{2}$	1	0	Undefined	0
$\frac{2\pi}{3}$	$\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$-\sqrt{3}$	$-\frac{\sqrt{3}}{3}$
$\frac{3\pi}{4}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{3}$	$-\sqrt{3}$
$\frac{5\pi}{6}$	$\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	-1	-1
π	0	-1	0	Undefined
$\frac{7\pi}{6}$	$-\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$		
$\frac{5\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$		
$\frac{4\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$		
$\frac{3\pi}{2}$	-1	0		
$\frac{5\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$\frac{1}{2}$		
$\frac{7\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$		
$\frac{11\pi}{6}$	$-\frac{1}{2}$	$\frac{\sqrt{3}}{2}$		

Triple Angle Identities

- $\sin(3x) = -\sin^3(x) + 3\cos^2(x)\sin(x)$
- $\sin(3x) = -4\sin^3(x) + 3\sin(x)$
- $\cos(3x) = \cos^3(x) - 3\sin^2(x)\cos(x)$
- $\cos(3x) = 4\cos^3(x) - 3\cos(x)$
- $\tan(3x) = \frac{3\tan(x) - \tan^3(x)}{1 - 3\tan^2(x)}$
- $\cot(3x) = \frac{3\cot(x) - \cot^3(x)}{1 - 3\cot^2(x)}$