Preprint typeset in JHEP style - HYPER VERSION

Lent Term, 2011 and 2012

Statistical Physics

University of Cambridge Part II Mathematical Tripos

Dr David Tong

Department of Applied Mathematics and Theoretical Physics,
Centre for Mathematical Sciences,
Wilberforce Road,
Cambridge, CB3 OBA, UK

http://www.damtp.cam.ac.uk/user/tong/statphys.html
d.tong@damtp.cam.ac.uk

Recommended Books and Resources

- Reif, *Fundamentals of Statistical and Thermal Physics*

A comprehensive and detailed account of the subject. It's solid. It's good. It isn't quirky.

- Kardar, *Statistical Physics of Particles*

A modern view on the subject which offers many insights. It's superbly written, if a little brief in places. A companion volume, "*The Statistical Physics of Fields*" covers aspects of critical phenomena. Both are available to download as lecture notes. Links are given on the course webpage

- Landau and Lifshitz, *Statistical Physics*

Russian style: terse, encyclopedic, magnificent. Much of this book comes across as remarkably modern given that it was first published in 1958.

- Mandl, *Statistical Physics*

This is an easy going book with very clear explanations but doesn't go into as much detail as we will need for this course. If you're struggling to understand the basics, this is an excellent place to look. If you're after a detailed account of more advanced aspects, you should probably turn to one of the books above.

- Pippard, *The Elements of Classical Thermodynamics*

This beautiful little book walks you through the rather subtle logic of classical thermodynamics. It's very well done. If Arnold Sommerfeld had read this book, he would have understood thermodynamics the first time round.

There are many other excellent books on this subject, often with different emphasis. I recommend "*States of Matter*" by David Goodstein which covers several topics beyond the scope of this course but offers many insights. For an entertaining yet technical account of thermodynamics that lies somewhere between a textbook and popular science, read "*The Four Laws*" by Peter Atkins.

A number of good lecture notes are available on the web. Links can be found on the course webpage: http://www.damtp.cam.ac.uk/user/tong/statphys.html

Contents

1. The Fundamentals of Statistical Mechanics **1**
- 1.1 Introduction 1
- 1.2 The Microcanonical Ensemble 2
  - 1.2.1 Entropy and the Second Law of Thermodynamics 5
  - 1.2.2 Temperature 8
  - 1.2.3 An Example: The Two State System 11
  - 1.2.4 Pressure, Volume and the First Law of Thermodynamics 14
  - 1.2.5 Ludwig Boltzmann (1844-1906) 16
- 1.3 The Canonical Ensemble 17
  - 1.3.1 The Partition Function 18
  - 1.3.2 Energy and Fluctuations 19
  - 1.3.3 Entropy 22
  - 1.3.4 Free Energy 25
- 1.4 The Chemical Potential 26
  - 1.4.1 Grand Canonical Ensemble 27
  - 1.4.2 Grand Canonical Potential 29
  - 1.4.3 Extensive and Intensive Quantities 29
  - 1.4.4 Josiah Willard Gibbs (1839-1903) 30

2. Classical Gases **32**
- 2.1 The Classical Partition Function 32
  - 2.1.1 From Quantum to Classical 33
- 2.2 Ideal Gas 34
  - 2.2.1 Equipartition of Energy 37
  - 2.2.2 The Sociological Meaning of Boltzmann's Constant 37
  - 2.2.3 Entropy and Gibbs's Paradox 39
  - 2.2.4 The Ideal Gas in the Grand Canonical Ensemble 40
- 2.3 Maxwell Distribution 42
  - 2.3.1 A History of Kinetic Theory 44
- 2.4 Diatomic Gas 45
- 2.5 Interacting Gas 48
  - 2.5.1 The Mayer f Function and the Second Virial Coefficient 50
  - 2.5.2 van der Waals Equation of State 53
  - 2.5.3 The Cluster Expansion 55

2.6 Screening and the Debye-Hückel Model of a Plasma 60

3. Quantum Gases 62
3.1 Density of States 62
3.1.1 Relativistic Systems 63
3.2 Photons: Blackbody Radiation 64
3.2.1 Planck Distribution 66
3.2.2 The Cosmic Microwave Background Radiation 68
3.2.3 The Birth of Quantum Mechanics 69
3.2.4 Max Planck (1858-1947) 70
3.3 Phonons 70
3.3.1 The Debye Model 70
3.4 The Diatomic Gas Revisited 75
3.5 Bosons 77
3.5.1 Bose-Einstein Distribution 78
3.5.2 A High Temperature Quantum Gas is (Almost) Classical 81
3.5.3 Bose-Einstein Condensation 82
3.5.4 Heat Capacity: Our First Look at a Phase Transition 86
3.6 Fermions 90
3.6.1 Ideal Fermi Gas 91
3.6.2 Degenerate Fermi Gas and the Fermi Surface 92
3.6.3 The Fermi Gas at Low Temperature 93
3.6.4 A More Rigorous Approach: The Sommerfeld Expansion 97
3.6.5 White Dwarfs and the Chandrasekhar limit 100
3.6.6 Pauli Paramagnetism 102
3.6.7 Landau Diamagnetism 104

4. Classical Thermodynamics 108
4.1 Temperature and the Zeroth Law 109
4.2 The First Law 111
4.3 The Second Law 113
4.3.1 The Carnot Cycle 115
4.3.2 Thermodynamic Temperature Scale and the Ideal Gas 117
4.3.3 Entropy 120
4.3.4 Adiabatic Surfaces 123
4.3.5 A History of Thermodynamics 126
4.4 Thermodynamic Potentials: Free Energies and Enthalpy 128
4.4.1 Enthalpy 131

4.4.2 Maxwell's Relations 131
4.5 The Third Law 133

5. Phase Transitions 135
5.1 Liquid-Gas Transition 135
5.1.1 Phase Equilibrium 137
5.1.2 The Clausius-Clapeyron Equation 140
5.1.3 The Critical Point 142
5.2 The Ising Model 147
5.2.1 Mean Field Theory 149
5.2.2 Critical Exponents 152
5.2.3 Validity of Mean Field Theory 154
5.3 Some Exact Results for the Ising Model 155
5.3.1 The Ising Model in $d = 1$ Dimensions 156
5.3.2 2d Ising Model: Low Temperatures and Peierls Droplets 157
5.3.3 2d Ising Model: High Temperatures 162
5.3.4 Kramers-Wannier Duality 165
5.4 Landau Theory 170
5.4.1 Second Order Phase Transitions 172
5.4.2 First Order Phase Transitions 175
5.4.3 Lee-Yang Zeros 176
5.5 Landau-Ginzburg Theory 180
5.5.1 Correlations 182
5.5.2 Fluctuations 183

Acknowledgements

These lecture notes are far from original. They borrow heavily both from the books described above and the online resources listed on the course webpage. I benefited a lot from the lectures by Mehran Kardar and by Chetan Nayak. This course is built on the foundation of previous courses given in Cambridge by Ron Horgan and Matt Wingate. I am also grateful to Ray Goldstein for help in developing the present syllabus. I am supported by the Royal Society and Alex Considine.

1. The Fundamentals of Statistical Mechanics

> "Ludwig Boltzmann, who spent much of his life studying statistical mechanics, died in 1906 by his own hand. Paul Ehrenfest, carrying on the work, died similarly in 1933. Now it is our turn to study statistical mechanics."
>
> *David Goodstein*

1.1 Introduction

Statistical mechanics is the art of turning the microscopic laws of physics into a description of Nature on a macroscopic scale.

Suppose you've got theoretical physics cracked. Suppose you know all the fundamental laws of Nature, the properties of the elementary particles and the forces at play between them. How can you turn this knowledge into an understanding of the world around us? More concretely, if I give you a box containing 10^{23} particles and tell you their mass, their charge, their interactions, and so on, what can you tell me about the stuff in the box?

There's one strategy that definitely won't work: writing down the Schrödinger equation for 10^{23} particles and solving it. That's typically not possible for 23 particles, let alone 10^{23}. What's more, even if you could find the wavefunction of the system, what would you do with it? The positions of individual particles are of little interest to anyone. We want answers to much more basic, almost childish, questions about the contents of the box. Is it wet? Is it hot? What colour is it? Is the box in danger of exploding? What happens if we squeeze it, pull it, heat it up? How can we begin to answer these kind of questions starting from the fundamental laws of physics?

The purpose of this course is to introduce the dictionary that allows you translate from the microscopic world where the laws of Nature are written to the everyday macroscopic world that we're familiar with. This will allow us to begin to address very basic questions about how matter behaves.

We'll see many examples. For centuries — from the 1600s to the 1900s — scientists were discovering "laws of physics" that govern different substances. There are many hundreds of these laws, mostly named after their discovers. Boyle's law and Charles's law relate pressure, volume and temperature of gases (they are usually combined into the ideal gas law); the Stefan-Boltzmann law tells you how much energy a hot object emits; Wien's displacement law tells you the colour of that hot object; the Dulong-Petit

law tells you how much energy it takes to heat up a lump of stuff; Curie's law tells you how a magnet loses its magic if you put it over a flame; and so on and so on. Yet we now know that these laws aren't fundamental. In some cases they follow simply from Newtonian mechanics and a dose of statistical thinking. In other cases, we need to throw quantum mechanics into the mix as well. But in all cases, we're going to see how derive them from first principles.

A large part of this course will be devoted to figuring out the interesting things that happen when you throw 10^{23} particles together. One of the recurring themes will be that $10^{23} \neq 1$. More is different: there are key concepts that are not visible in the underlying laws of physics but emerge only when we consider a large collection of particles. One very simple example is temperature. This is not a fundamental concept: it doesn't make sense to talk about the temperature of a single electron. But it would be impossible to talk about physics of the everyday world around us without mention of temperature. This illustrates the fact that the language needed to describe physics on one scale is very different from that needed on other scales. We'll see several similar emergent quantities in this course, including the phenomenon of phase transitions where the smooth continuous laws of physics conspire to give abrupt, discontinuous changes in the structure of matter.

Historically, the techniques of statistical mechanics proved to be a crucial tool for understanding the deeper laws of physics. Not only is the development of the subject intimately tied with the first evidence for the existence of atoms, but quantum mechanics itself was discovered by applying statistical methods to decipher the spectrum of light emitted from hot objects. (We will study this derivation in Section 3). However, physics is not a finished subject. There are many important systems in Nature – from high temperature superconductors to black holes – which are not yet understood at a fundamental level. The information that we have about these systems concerns their macroscopic properties and our goal is to use these scant clues to deconstruct the underlying mechanisms at work. The tools that we will develop in this course will be crucial in this task.

1.2 The Microcanonical Ensemble

> "Anyone who wants to analyze the properties of matter in a real problem might want to start by writing down the fundamental equations and then try to solve them mathematically. Although there are people who try to use such an approach, these people are the failures in this field..."
>
> *Richard Feynman, sugar coating it.*

We'll start by considering an isolated system with fixed energy, E. For the purposes of the discussion we will describe our system using the language of quantum mechanics, although we should keep in mind that nearly everything applies equally well to classical systems.

In your first two courses on quantum mechanics you looked only at systems with a few degrees of freedom. These are defined by a Hamiltonian, $\hat{H}$, and the goal is usually to solve the time independent Schrödinger equation

$$\hat{H}|\psi\rangle = E|\psi\rangle$$

In this course, we will still look at systems that are defined by a Hamiltonian, but now with a very large number of degrees of freedom, say $N \sim 10^{23}$. The energy eigenstates $|\psi\rangle$ are very complicated objects since they contain information about what each of these particles is doing. They are called *microstates*.

In practice, it is often extremely difficult to write down the microstate describing all these particles. But, more importantly, it is usually totally uninteresting. The wavefunction for a macroscopic system very rarely captures the relevant physics because real macroscopic systems are not described by a single pure quantum state. They are in contact with an environment, constantly buffeted and jostled by outside influences. Each time the system is jogged slightly, it undergoes a small perturbation and there will be a probability that it transitions to another state. If the perturbation is very small, then the transitions will only happen to states of equal (or very nearly equal) energy. But with 10^{23} particles, there can be many many microstates all with the same energy E. To understand the physics of these systems, we don't need to know the intimate details of any one state. We need to know the crude details of all the states.

It would be impossibly tedious to keep track of the dynamics which leads to transitions between the different states. Instead we will resort to statistical methods. We will describe the system in terms of a probability distribution over the quantum states. In other words, the system is in a *mixed state* rather than a pure state. Since we have fixed the energy, there will only be a non-zero probability for states which have the specified energy E. We will denote a basis of these states as $|n\rangle$ and the probability that the systems sits in a given state as $p(n)$. Within this probability distribution, the expectation value of any operator $\hat{\mathcal{O}}$ is

$$\langle\hat{\mathcal{O}}\rangle = \sum_n p(n)\langle n|\hat{\mathcal{O}}|n\rangle$$

Our immediate goal is to understand what probability distribution $p(n)$ is appropriate for large systems.

Firstly, we will greatly restrict the kind of situations that we can talk about. We will only discuss systems that have been left alone for some time. This ensures that the energy and momentum in the system has been redistributed among the many particles and any memory of whatever special initial conditions the system started in has long been lost. Operationally, this means that the probability distribution is independent of time which ensures that the expectation values of the macroscopic observables are also time independent. In this case, we say that the system is in *equilibrium*. Note that just because the system is in equilibrium does not mean that all the components of the system have stopped moving; a glass of water left alone will soon reach equilibrium but the atoms inside are still flying around.

We are now in a position to state the fundamental assumption of statistical mechanics. It is the idea that we should take the most simple minded approach possible and treat all states the same. Or, more precisely:

For an isolated system in equilibrium, all accessible microstates are equally likely.

Since we know nothing else about the system, such a democratic approach seems eminently reasonable. Notice that we've left ourselves a little flexibility with the inclusion of the word "accessible". This refers to any state that can be reached due to the small perturbations felt by the system. For the moment, we will take it mean all states that have the same energy E. Later, we shall see contexts where we add further restrictions on what it means to be an accessible state.

Let us introduce some notation. We define

$$\Omega(E) = \text{Number of states with energy } E$$

The probability that the system with fixed energy E is in a given state $|n\rangle$ is simply

$$p(n) = \frac{1}{\Omega(E)} \tag{1.1}$$

The probability that the system is in a state with some different energy $E' \neq E$ is zero. This probability distribution, relevant for systems with fixed energy, is known as the *microcanonical ensemble*. Some comments:

- $\Omega(E)$ is a usually ridiculously large number. For example, suppose that we have $N \sim 10^{23}$ particles, each of which can only be in one of two quantum states – say "spin up" and "spin down". Then the total number of microstates of the system is $2^{10^{23}}$. This is a silly number. In some sense, numbers this large can never have

any physical meaning! They only appear in combinatoric problems, counting possible eventualities. They are never answers to problems which require you to count actual existing physical objects. One, slightly facetious, way of saying this is that numbers this large can't have physical meaning because they are the same no matter what units they have. (If you don't believe me, think of $2^{10^{23}}$ as a distance scale: it is effectively the same distance regardless of whether it is measured in microns or lightyears. Try it!).

- In quantum systems, the energy levels will be discrete. However, with many particles the energy levels will be finely spaced and can be effectively treated as a continuum. When we say that $\Omega(E)$ counts the number of states with energy E we implicitly mean that it counts the number of states with energy between E and $E + \delta E$ where δE is small compared to the accuracy of our measuring apparatus but large compared to the spacing of the levels.

- We phrased our discussion in terms of quantum systems but everything described above readily carries over the classical case. In particular, the probabilities $p(n)$ have nothing to do with quantum indeterminacy. They are due entirely to our ignorance.

1.2.1 Entropy and the Second Law of Thermodynamics

We define the *entropy* of the system to be

$$S(E) = k_B \log \Omega(E) \tag{1.2}$$

Here k_B is a fundamental constant, known as *Boltzmann's constant* . It has units of Joules per Kelvin.

$$k_B \approx 1.381 \times 10^{-23} JK^{-1} \tag{1.3}$$

The log in (1.2) is the natural logarithm (base e, not base 10). Why do we take the log in the definition? One reason is that it makes the numbers less silly. While the number of states is of order $\Omega \sim e^N$, the entropy is merely proportional to the number of particles in the system, $S \sim N$. This also has the happy consequence that the entropy is an additive quantity. To see this, consider two non-interacting systems with energies E_1 and E_2 respectively. Then the total number of states of both systems is

$$\Omega(E_1, E_2) = \Omega_1(E_1)\Omega(E_2)$$

while the entropy for both systems is

$$S(E_1, E_2) = S_1(E_1) + S_2(E_2)$$

The Second Law

Suppose we take the two, non-interacting, systems mentioned above and we bring them together. We'll assume that they can exchange energy, but that the energy levels of each individual system remain unchanged. (These are actually contradictory assumptions! If the systems can exchange energy then there must be an interaction term in their Hamiltonian. But such a term would shift the energy levels of each system. So what we really mean is that these shifts are negligibly small and the only relevant effect of the interaction is to allow the energy to move between systems).

The energy of the combined system is still $E_{\text{total}} = E_1 + E_2$. But the first system can have any energy $E \leq E_{\text{total}}$ while the second system must have the remainder $E_{\text{total}} - E$. In fact, there is a slight caveat to this statement: in a quantum system we can't have any energy at all: only those discrete energies E_i that are eigenvalues of the Hamiltonian. So the number of available states of the combined system is

$$\begin{aligned}\Omega(E_{\text{total}}) &= \sum_{\{E_i\}} \Omega_1(E_i)\Omega_2(E_{\text{total}} - E_i) \\ &= \sum_{\{E_i\}} \exp\left(\frac{S_1(E_i)}{k_B} + \frac{S_2(E_{\text{total}} - E_i)}{k_B}\right) \end{aligned} \tag{1.4}$$

There is a slight subtlety in the above equation. Both system 1 and system 2 have discrete energy levels. How do we know that if E_i is an energy of system 1 then $E_{\text{total}} - E_i$ is an energy of system 2. In part this goes back to the comment made above about the need for an interaction Hamiltonian that shifts the energy levels. In practice, we will just ignore this subtlety. In fact, for most of the systems that we will discuss in this course, the discreteness of energy levels will barely be important since they are so finely spaced that we can treat the energy E of the first system as a continuous variable and replace the sum by an integral. We will see many explicit examples of this in the following sections.

At this point, we turn again to our fundamental assumption — all states are equally likely — but now applied to the combined system. This has fixed energy E_{total} so can be thought of as sitting in the microcanonical ensemble with the distribution (1.1) which means that the system has probability $p = 1/\Omega(E_{\text{total}})$ to be in each state. Clearly, the entropy of the combined system is greater or equal to that of the original system,

$$S(E_{\text{total}}) \equiv k_B \log \Omega(E_{\text{total}}) \geq S_1(E_1) + S_2(E_2) \tag{1.5}$$

which is true simply because the states of the two original systems are a subset of the total number of possible states.

While (1.5) is true for any two systems, there is a useful approximation we can make to determine $S(E_{\rm total})$ which holds when the number of particles, N, in the game is very large. We have already seen that the entropy scales as $S \sim N$. This means that the expression (1.4) is a sum of exponentials of N, which is itself an exponentially large number. Such sums are totally dominated by their maximum value. For example, suppose that for some energy, $E_\star$, the exponent has a value that's twice as large as any other E. Then this term in the sum is larger than all the others by a factor of e^N. And that's a very large number. All terms but the maximum are completely negligible. (The equivalent statement for integrals is that they can be evaluated using the saddle point method). In our case, the maximum value, $E = E_\star$, occurs when

$$\frac{\partial S_1(E_\star)}{\partial E} - \frac{\partial S_2(E_{\rm total} - E_\star)}{\partial E} = 0 \tag{1.6}$$

where this slightly cumbersome notation means (for the first term) $\partial S_1/\partial E$ evaluated at $E = E_\star$. The total entropy of the combined system can then be very well approximated by

$$S(E_{\rm total}) \approx S_1(E_\star) + S_2(E_{\rm total} - E_\star) \geq S_1(E_1) + S_2(E_2)$$

It's worth stressing that there is no a priori reason why the first system should have a fixed energy once it is in contact with the second system. But the large number of particles involved means that it is overwhelmingly likely to be found with energy $E_\star$ which maximises the number of states of the combined system. Conversely, once in this bigger set of states, it is highly unlikely that the system will ever be found back in a state with energy E_1 or, indeed, any other energy different from $E_\star$.

Figure 1: Arthur Eddington

It is this simple statement that is responsible for all the irreversibility that we see in the world around us. This is the *second law of thermodynamics*. As a slogan, "entropy increases". When two systems are brought together — or, equivalently, when constraints on a system are removed — the total number of available states available is vastly enlarged.

It is sometimes stated that second law is the most sacred in all of physics. Arthur Eddington's rant, depicted in the cartoon, is one of the more famous acclamations of the law.

And yet, as we have seen above, the second law hinges on probabilistic arguments. We said, for example, that it is "highly unlikely" that the system will return to its initial configuration. One might think that this may allow us a little leeway. Perhaps, if probabilities are underlying the second law, we can sometimes get lucky and find counterexamples. While, it is most likely to find system 1 to have energy $E_\star$, surely occasionally one sees it in a state with a different energy? In fact, this never happens. The phrase "highly unlikely" is used only because the English language does not contain enough superlatives to stress how ridiculously improbable a violation of the second law would be. The silly number of possible states in a macroscopic systems means that violations happen only on silly time scales: exponentials of exponentials. This is a good operational definition of the word "never".

1.2.2 Temperature

We next turn to a very familiar quantity, albeit viewed in an unfamiliar way. The temperature, T, of a system is defined as

$$\frac{1}{T} = \frac{\partial S}{\partial E} \tag{1.7}$$

This is an extraordinary equation. We have introduced it as the definition of temperature. But why is this a good definition? Why does it agree with the idea of temperature that your mum has? Why is this the same T that makes mercury rise (the element, not the planet...that's a different course). Why is it the same T that makes us yell when we place our hand on a hot stove?

First, note that T has the right units, courtesy of Boltzmann's constant (1.3). But that was our merely a choice of convention: it doesn't explain why T has the properties that we expect of temperature. To make progress, we need to think more carefully about the kind of properties that we do expect. We will describe this in some detail in Section 4. For now it will suffice to describe the key property of temperature, which is the following: suppose we take two systems, each in equilibrium and each at the same temperature T, and place them in contact so that they can exchange energy. Then...nothing happens.

It is simple to see that this follows from our definition (1.7). We have already done the hard work above where we saw that two systems, brought into contact in this way,

will maximize their entropy. This is achieved when the first system has energy $E_\star$ and the second energy $E_{\text{total}} - E_\star$, with $E_\star$ determined by equation (1.6). If we want nothing noticeable to happen when the systems are brought together, then it must have been the case that the energy of the first system was already at $E_1 = E_\star$. Or, in other words, that equation (1.6) was obeyed before the systems were brought together,

$$\frac{\partial S_1(E_1)}{\partial E} = \frac{\partial S_2(E_2)}{\partial E} \tag{1.8}$$

From our definition (1.7), this is the same as requiring that the initial temperatures of the two systems are equal: $T_1 = T_2$.

Suppose now that we bring together two systems at slightly different temperatures. They will exchange energy, but conservation ensures that what the first system gives up, the second system receives and vice versa. So $\delta E_1 = -\delta E_2$. If the change of entropy is small, it is well approximated by

$$\begin{aligned}
\delta S &= \frac{\partial S_1(E_1)}{\partial E}\,\delta E_1 + \frac{\partial S_2(E_2)}{\partial E}\,\delta E_2 \\
&= \left(\frac{\partial S_1(E_1)}{\partial E} - \frac{\partial S_2(E_2)}{\partial E}\right)\delta E_1 \\
&= \left(\frac{1}{T_1} - \frac{1}{T_2}\right)\delta E_1
\end{aligned}$$

The second law tells us that entropy must increase: $\delta S > 0$. This means that if $T_1 > T_2$, we must have $\delta E_1 < 0$. In other words, the energy flows in the way we would expect: from the hotter system to colder.

To summarise: the equilibrium argument tell us that $\partial S/\partial E$ should have the interpretation as *some* function of temperature; the heat flowing argument tell us that it should be a monotonically decreasing function. But why $1/T$ and not, say, $1/T^2$? To see this, we really need to compute T for a system that we're all familiar with and see that it gives the right answer. Once we've got the right answer for one system, the equilibrium argument will ensure that it is right for all systems. Our first business in Section 2 will be to compute the temperature T for an ideal gas and confirm that (1.7) is indeed the correct definition.

Heat Capacity

The heat capacity, C, is defined by

$$C = \frac{\partial E}{\partial T} \tag{1.9}$$

We will later introduce more refined versions of the heat capacity (in which various, yet-to-be-specified, external parameters are held constant or allowed to vary and we are more careful about the mode of energy transfer into the system). The importance of the heat capacity is that it is defined in terms of things that we can actually measure! Although the key theoretical concept is entropy, if you're handed an experimental system involving 10^{23} particles, you can't measure the entropy directly by counting the number of accessible microstates. You'd be there all day. But you can measure the heat capacity: you add a known quantity of energy to the system and measure the rise in temperature. The result is C^{-1}.

There is another expression for the heat capacity that is useful. The entropy is a function of energy, $S = S(E)$. But we could invert the formula (1.7) to think of energy as a function of temperature, $E = E(T)$. We then have the expression

$$\frac{\partial S}{\partial T} = \frac{\partial S}{\partial E} \cdot \frac{\partial E}{\partial T} = \frac{C}{T}$$

This is a handy formula. If we can measure the heat capactiy of the system for various temperatures, we can get a handle on the function $C(T)$. From this we can then determine the entropy of the system. Or, more precisely, the entropy difference

$$\Delta S = \int_{T_1}^{T_2} \frac{C(T)}{T}\, dT \tag{1.10}$$

Thus the heat capacity is our closest link between experiment and theory.

The heat capacity is always proportional to N, the number of particles in the system. It is common to define the *specific heat capacity*, which is simply the heat capacity divided by the mass of the system and is independent of N.

There is one last point to make about heat capacity. Differentiating (1.7) once more, we have

$$\frac{\partial^2 S}{\partial E^2} = -\frac{1}{T^2 C} \tag{1.11}$$

Nearly all systems you will meet have $C > 0$. (There is one important exception: a black hole has negative heat capacity!). Whenever $C > 0$, the system is said to be *thermodynamically stable*. The reason for this language goes back to the previous discussion concerning two systems which can exchange energy. There we wanted to maximize the entropy and checked that we had a stationary point (1.6), but we forgot to check whether this was a maximum or minimum. It is guaranteed to be a maximum if the heat capacity of both systems is positive so that $\partial^2 S/\partial E^2 < 0$.

1.2.3 An Example: The Two State System

Consider a system of N non-interacting particles. Each particle is fixed in position and can sit in one of two possible states which, for convenience, we will call "spin up" $|\uparrow\rangle$ and "spin down" $|\downarrow\rangle$. We take the energy of these states to be,

$$E_\downarrow = 0 \quad , \quad E_\uparrow = \epsilon$$

which means that the spins want to be down; you pay an energy cost of ϵ for each spin which points up. If the system has $N_\uparrow$ particles with spin up and $N_\downarrow = N - N_\uparrow$ particles with spin down, the energy of the system is

$$E = N_\uparrow \epsilon$$

We can now easily count the number of states $\Omega(E)$ of the total system which have energy E. It is simply the number of ways to pick $N_\uparrow$ particles from a total of N,

$$\Omega(E) = \frac{N!}{N_\uparrow!(N - N_\uparrow)!}$$

and the entropy is given by

$$S(E) = k_B \log\left(\frac{N!}{N_\uparrow!(N - N_\uparrow)!}\right)$$

An Aside: Stirling's Formula

For large N, there is a remarkably accurate approximation to the factorials that appear in the expression for the entropy. It is known as Stirling's formula,

$$\log N! = N \log N - N + \tfrac{1}{2}\log 2\pi N + \mathcal{O}(1/N)$$

You will prove this on the first problem sheet. However, for our purposes we will only need the first two terms in this expansion and these can be very quickly derived by looking at the expression

$$\log N! = \sum_{p=1}^{N} \log p \approx \int_1^N dp \;\; \log p = N \log N - N + 1$$

where we have approximated the sum by the integral as shown in the figure. You can also see from the figure that integral gives a lower bound on the sum which is confirmed by checking the next terms in Stirling's formula.

Figure 2:

Back to the Physics

Using Stirling's approximation, we can write the entropy as

$$\begin{aligned} S(E) &= k_B \left[N \log N - N - N_\uparrow \log N_\uparrow + N_\uparrow - (N - N_\uparrow) \log(N - N_\uparrow) + (N - N_\uparrow) \right] \\ &= -k_B \left[(N - N_\uparrow) \log \left(\frac{N - N_\uparrow}{N} \right) + N_\uparrow \log \left(\frac{N_\uparrow}{N} \right) \right] \\ &= -k_B N \left[\left(1 - \frac{E}{N\epsilon} \right) \log \left(1 - \frac{E}{N\epsilon} \right) + \frac{E}{N\epsilon} \log \left(\frac{E}{N\epsilon} \right) \right] \end{aligned} \tag{1.12}$$

A sketch of $S(E)$ plotted against E is shown in Figure 3. The entropy vanishes when $E = 0$ (all spins down) and $E = N\epsilon$ (all spins up) because there is only one possible state with each of these energies. The entropy is maximal when $E = N\epsilon/2$ where we have $S = Nk_B \log 2$.

If the system has energy E, its temperature is

$$\frac{1}{T} = \frac{\partial S}{\partial E} = \frac{k_B}{\epsilon} \log \left(\frac{N\epsilon}{E} - 1 \right)$$

We can also invert this expression. If the system has temperature T, the fraction of particles with spin up is given by

$$\frac{N_\uparrow}{N} = \frac{E}{N\epsilon} = \frac{1}{e^{\epsilon/k_B T} + 1} \tag{1.13}$$

Note that as $T \to \infty$, the fraction of spins $N_\uparrow/N \to 1/2$. In the limit of infinite temperature, the system sits at the peak of the curve in Figure 3.

Figure 3: Entropy of the two-state system

What happens for energies $E > N\epsilon/2$, where $N_\uparrow/N > 1/2$? From the definition of temperature as $1/T = \partial S/\partial E$, it is clear that we have entered the realm of *negative temperatures*. This should be thought of as hotter than infinity! (This is simple to see in the variables $1/T$ which tends towards zero and then just keeps going to negative values). Systems with negative temperatures have the property that the number of microstates decreases as we add energy. They can be realised in laboratories, at least temporarily, by instantaneously flipping all the spins in a system.

Heat Capacity and the Schottky Anomaly

Finally, we can compute the heat capacity, which we choose to express as a function of temperature (rather than energy) since this is more natural when comparing to experiment. We then have

$$C = \frac{dE}{dT} = \frac{N\epsilon^2}{k_B T^2} \frac{e^{\epsilon/k_B T}}{(e^{\epsilon/k_B T} + 1)^2} \tag{1.14}$$

Note that C is of order N, the number of particles in the system. This property extends to all other examples that we will study. A sketch of C vs T is shown in Figure 4. It starts at zero, rises to a maximum, then drops off again. We'll see a lot of graphs in this course that look more or less like this. Let's look at some of the key features in this case. Firstly, the maximum is around $T \sim \epsilon/k_B$. In other words, the maximum point sits at the characteristic energy scale in the system.

Figure 4: Heat Capacity of the two state system

As $T \to 0$, the specific heat drops to zero exponentially quickly. (Recall that $e^{-1/x}$ is a function which tends towards zero faster than any power x^n). The reason for this fast fall-off can be traced back to the existence of an *energy gap*, meaning that the first excited state is a finite energy above the ground state. The heat capacity also drops off as $T \to \infty$, but now at a much slower power-law pace. This fall-off is due to the fact that all the states are now occupied.

The contribution to the heat capacity from spins is not the dominant contribution in most materials. It is usually dwarfed by the contribution from phonons and, in metals, from conduction electrons, both of which we will calculate later in the course. Nonetheless, in certain classes of material — for example, paramagnetic salts — a spin contribution of the form (1.14) can be seen at low temperatures where it appears as a small bump in the graph and is referred to as the *Schottky anomaly*. (It is "anomalous" because most materials have a heat capacity which decreases monotonically as temperature is reduced). In Figure 5, the Schottky contribution has been isolated from the phonon

Figure 5:

contribution[1]. The open circles and dots are both data (interpreted in mildly different ways); the solid line is theoretical prediction, albeit complicated slightly by the presence of a number of spin states. The deviations are most likely due to interactions between the spins.

The two state system can also be used as a model for defects in a lattice. In this case, the "spin down" state corresponds to an atom sitting in the lattice where it should be with no energy cost. The "spin up" state now corresponds to a missing atom, ejected from its position at energy cost ϵ.

1.2.4 Pressure, Volume and the First Law of Thermodynamics

We will now start to consider other external parameters which can affect different systems. We will see a few of these as we move through the course, but the most important one is perhaps the most obvious – the volume V of a system. This didn't play a role in the two-state example because the particles were fixed. But as soon as objects are free to move about, it becomes crucial to understand how far they can roam.

We'll still use the same notation for the number of states and entropy of the system, but now these quantities will be functions of both the energy and the volume,

$$S(E,V) = k_B \log \Omega(E,V)$$

The temperature is again given by $1/T = \partial S/\partial E$, where the partial derivative implicitly means that we keep V fixed when we differentiate. But now there is a new, natural quantity that we can consider — the differentiation with respect to V. This also gives a quantity that you're all familiar with — *pressure*, p. Well, almost. The definition is

$$p = T\frac{\partial S}{\partial V} \tag{1.15}$$

To see that this is a sensible definition, we can replay the arguments of Section 1.2.2. Imagine two systems in contact through a moveable partition as shown in the figure above, so that the total volume remains fixed, but system 1 can expand at the expense of system 2 shrinking. The same equilibrium arguments that previously lead to (1.8) now tell us that the volumes of the systems don't change as long as $\partial S/\partial V$ is the same for both systems. Or, in other words, as long as the pressures are equal.

Despite appearances, the definition of pressure actually has little to do with entropy. Roughly speaking, the S in the derivative cancels the factor of S sitting in T. To make

[1]The data is taken from Chirico and Westrum Jr., J. Chem. Thermodynamics 12 (1980), 311, and shows the spin contribution to the heat capacity of $Tb(OH)_3$

this mathematically precise, consider a system with entropy $S(E, V)$ that undergoes a small change in energy and volume. The change in entropy is

$$dS = \frac{\partial S}{\partial E} dE + \frac{\partial S}{\partial V} dV$$

Rearranging, and using our definitions (1.7) and (1.15), we can write

$$dE = TdS - pdV \tag{1.16}$$

The left-hand side is the change in energy of the system. It is easy to interpret the second term on the right-hand side: it is the work done on the system. To see this, consider the diagram on the right. Recall that pressure is force per area. The change of volume in the set-up depicted is $dV = \text{Area} \times dx$. So the work done on the system is Force $\times\, dx = (pA)dx = pdV$. To make sure that we've got the minus signs right, remember that if $dV < 0$, we're exerting a force to squeeze the system, increasing its energy. In contrast, if $dV > 0$, the system itself is doing the work and hence losing energy.

Figure 7: Work Done

Alternatively, you may prefer to run this argument in reverse: if you're happy to equate squeezing the system by dV with doing work, then the discussion above is sufficient to tell you that pressure as defined in (1.15) has the interpretation of force per area.

What is the interpretation of the first term on the right-hand side of (1.16)? It must be some form of energy transferred to the system. It turns out that the correct interpretation of TdS is the amount of *heat* the system absorbs from the surroundings. Much of Section 4 will be concerned with understanding why this is right way to think about TdS and we postpone a full discussion until then.

Equation (1.16) expresses the conservation of energy for a system at finite temperature. It is known as the *First Law of Thermodynamics*. (You may have noticed that we're not doing these laws in order! This too will be rectified in Section 4).

As a final comment, we can now give a slightly more refined definition of the heat capacity (1.9). In fact, there are several different heat capacities which depend on which other variables are kept fixed. Throughout most of these lectures, we will be interested in the heat capacity at fixed volume, denoted C_V,

$$C_V = \left.\frac{\partial E}{\partial T}\right|_V \tag{1.17}$$

Using the first law of thermodynamics (1.16), we see that something special happens when we keep volume constant: the work done term drops out and we have

$$C_V = T \left.\frac{\partial S}{\partial T}\right|_V \tag{1.18}$$

This form emphasises that, as its name suggests, the heat capacity measures the ability of the system to absorb *heat* TdS as opposed to any other form of energy. (Although, admittedly, we still haven't really defined what heat is. As mentioned above, this will have to wait until Section 4).

The equivalence of (1.17) and (1.18) only followed because we kept volume fixed. What is the heat capacity if we keep some other quantity, say pressure, fixed? In this case, the correct definition of heat capacity is the expression analogous to (1.18). So, for example, the heat capacity at constant pressure C_p is defined by

$$C_p = T \left.\frac{\partial S}{\partial T}\right|_p$$

For the next few Sections, we'll only deal with C_V. But we'll return briefly to the relationship between C_V and C_p in Section 4.4.

1.2.5 Ludwig Boltzmann (1844-1906)

> "My memory for figures, otherwise tolerably accurate, always lets me down when I am counting beer glasses"
>
> *Boltzmann Counting*

Ludwig Boltzmann was born into a world that doubted the existence of atoms[2]. The cumulative effect of his lifetime's work was to change this. No one in the 1800s ever thought we could see atoms directly and Boltzmann's strategy was to find indirect, yet overwhelming, evidence for their existence. He developed much of the statistical machinery that we have described above and, building on the work of Maxwell, showed that many of the seemingly fundamental laws of Nature — those involving heat and gases in particular — were simply consequences of Newton's laws of motion when applied to very large systems.

[2]If you want to learn more about his life, I recommend the very enjoyable biography, *Boltzmann's Atom* by David Lindley. The quote above is taken from a travel essay that Boltzmann wrote recounting a visit to California. The essay is reprinted in a drier, more technical, biography by Carlo Cercignani.

It is often said that Boltzmann's great insight was the equation which is now engraved on his tombstone, $S = k_B \log \Omega$, which lead to the understanding of the second law of thermodynamics in terms of microscopic disorder. Yet these days it is difficult to appreciate the boldness of this proposal simply because we rarely think of any other definition of entropy. We will, in fact, meet the older thermodynamic notion of entropy and the second law in Section 4 of this course. In the meantime, perhaps Boltzmann's genius is better reflected in the surprising equation for temperature: $1/T = \partial S/\partial E$.

Boltzmann gained prominence during his lifetime, holding professorships at Graz, Vienna, Munich and Leipzig (not to mention a position in Berlin that he somehow failed to turn up for). Nonetheless, his work faced much criticism from those who would deny the existence of atoms, most notably Mach. It is not known whether these battles contributed to the depression Boltzmann suffered later in life, but the true significance of his work was only appreciated after his body was found hanging from the rafters of a guest house near Trieste.

1.3 The Canonical Ensemble

The microcanonical ensemble describes systems that have a fixed energy E. From this, we deduce the equilibrium temperature T. However, very often this is not the best way to think about a system. For example, a glass of water sitting on a table has a well defined average energy. But the energy is constantly fluctuating as it interacts with the environment. For such systems, it is often more appropriate to think of them as sitting at fixed temperature T, from which we then deduce the average energy.

To model this, we will consider a system — let's call it S — in contact with a second system which is a large *heat reservoir* – let's call it R. This reservoir is taken to be at some equilibrium temperature T. The term "reservoir" means that the energy of S is negligible compared with that of R. In particular, S can happily absorb or donate energy from or to the reservoir without changing the ambient temperature T.

How are the energy levels of S populated in such a situation? We label the states of S as $|n\rangle$, each of which has energy E_n. The number of microstates of the combined systems S and R is given by the sum over all states of S,

$$\Omega(E_{\text{total}}) = \sum_n \Omega_R(E_{\text{total}} - E_n) \equiv \sum_n \exp\left(\frac{S_R(E_{\text{total}} - E_n)}{k_B}\right)$$

I stress again that the sum above is over all the states of S, rather than over the energy levels of S. (If we'd written the latter, we would have to include a factor of $\Omega_S(E_n)$ in the sum to take into account the degeneracy of states with energy E_n). The fact that

R is a reservoir means that $E_n \ll E_{\text{total}}$. This allows us to Taylor expand the entropy, keeping just the first two terms,

$$\Omega(E_{\text{total}}) \approx \sum_n \exp\left(\frac{S_R(E_{\text{total}})}{k_B} - \frac{\partial S_R}{\partial E_{\text{total}}}\frac{E_n}{k_B}\right)$$

But we know that $\partial S_R/\partial E_{\text{total}} = 1/T$, so we have

$$\Omega(E_{\text{total}}) = e^{S_R(E_{\text{total}})/k_B} \sum_n e^{-E_n/k_B T}$$

We now apply the fundamental assumption of statistical mechanics — that all accessible energy states are equally likely — to the combined system + reservoir. This means that each of the $\Omega(E_{\text{total}})$ states above is equally likely. The number of these states for which the system sits in $|n\rangle$ is $e^{S_R/k_B}e^{-E_n/k_B T}$. So the probabilty that the system sits in state $|n\rangle$ is just the ratio of this number of states to the total number of states,

$$p(n) = \frac{e^{-E_n/k_B T}}{\sum_m e^{-E_m/k_B T}} \tag{1.19}$$

This is the *Boltzmann distribution*, also known as the *canonical ensemble*. Notice that the details of the reservoir have dropped out. We don't need to know $S_R(E)$ for the reservoir; all that remains of its influence is the temperature T.

The exponential suppression in the Boltzmann distribution means that it is very unlikely that any of the states with $E_n \gg k_B T$ are populated. However, all states with energy $E_n \leq k_B T$ have a decent chance of being occupied. Note that as $T \to 0$, the Boltzmann distribution forces the system into its ground state (i.e. the state with lowest energy); all higher energy states have vanishing probability at zero temperature.

1.3.1 The Partition Function

Since we will be using various quantities a lot, it is standard practice to introduce new notation. Firstly, the inverse factor of the temperature is universally denoted,

$$\beta \equiv \frac{1}{k_B T} \tag{1.20}$$

And the normalization factor that sits in the denominator of the probability is written,

$$Z = \sum_n e^{-\beta E_n} \tag{1.21}$$

In this notation, the probability for the system to be found in state $|n\rangle$ is

$$p(n) = \frac{e^{-\beta E_n}}{Z} \tag{1.22}$$

Rather remarkably, it turns out that the most important quantity in statistical mechanics is Z. Although this was introduced as a fairly innocuous normalization factor, it actually contains all the information we need about the system. We should think of Z, as defined in (1.21), as a function of the (inverse) temperature β. When viewed in this way, Z is called the *partition function*.

We'll see lots of properties of Z soon. But we'll start with a fairly basic, yet important, point: for independent systems, Z's multiply. This is easy to prove. Suppose that we have two systems which don't interact with each other. The energy of the combined system is then just the sum of the individual energies. The partition function for the combined system is (in, hopefully, obvious notation)

$$\begin{aligned} Z &= \sum_{n,m} e^{-\beta(E_n^{(1)}+E_m^{(2)})} \\ &= \sum_{n,m} e^{-\beta E_n^{(1)}} e^{-\beta E_m^{(2)}} \\ &= \sum_n e^{-\beta E_n^{(1)}} \sum_m e^{-\beta E_m^{(2)}} = Z_1 Z_2 \end{aligned} \tag{1.23}$$

A Density Matrix for the Canonical Ensemble

In statistical mechanics, the inherent probabilities of the quantum world are joined with probabilities that arise from our ignorance of the underlying state. The correct way to describe this is in term of a density matrix, $\hat{\rho}$. The canonical ensemble is really a choice of density matrix,

$$\hat{\rho} = \frac{e^{-\beta \hat{H}}}{Z} \tag{1.24}$$

If we make a measurement described by an operator $\hat{\mathcal{O}}$, then the probability that we find ourselves in the eigenstate $|\phi\rangle$ is given by

$$p(\phi) = \langle \phi | \hat{\rho} | \phi \rangle$$

For energy eigenstates, this coincides with our earlier result (1.22). We won't use the language of density matrices in this course, but it is an elegant and conceptually clear framework to describe more formal results.

1.3.2 Energy and Fluctuations

Let's see what information is contained in the partition function. We'll start by thinking about the energy. In the microcanonical ensemble, the energy was fixed. In the

canonical ensemble, that is no longer true. However, we can happily compute the average energy,

$$\langle E \rangle = \sum_n p(n) E_n = \sum_n \frac{E_n e^{-\beta E_n}}{Z}$$

But this can be very nicely expressed in terms of the partition function by

$$\langle E \rangle = -\frac{\partial}{\partial \beta} \log Z \tag{1.25}$$

We can also look at the spread of energies about the mean — in other words, about fluctuations in the probability distribution. As usual, this spread is captured by the variance,

$$\Delta E^2 = \langle (E - \langle E \rangle)^2 \rangle = \langle E^2 \rangle - \langle E \rangle^2$$

This too can be written neatly in terms of the partition function,

$$\Delta E^2 = \frac{\partial^2}{\partial \beta^2} \log Z = -\frac{\partial \langle E \rangle}{\partial \beta} \tag{1.26}$$

There is another expression for the fluctuations that provides some insight. Recall our definition of the heat capacity (1.9) in the microcanonical ensemble. In the canonical ensemble, where the energy is not fixed, the corresponding definition is

$$C_V = \left. \frac{\partial \langle E \rangle}{\partial T} \right|_V$$

Then, since $\beta = 1/k_B T$, the spread of energies in (1.26) can be expressed in terms of the heat capacity as

$$\Delta E^2 = k_B T^2 C_V \tag{1.27}$$

There are two important points hiding inside this small equation. The first is that the equation relates two rather different quantities. On the left-hand side, ΔE describes the probabilistic fluctuations in the energy of the system. On the right-hand side, the heat capacity C_V describes the ability of the system to absorb energy. If C_V is large, the system can take in a lot of energy without raising its temperature too much. The equation (1.27) tells us that the fluctuations of the systems are related to the ability of the system to dissipate, or absorb, energy. This is the first example of a more general result known as the *fluctuation-dissipation theorem*.

The other point to take away from (1.27) is the size of the fluctuations as the number of particles N in the system increases. Typically $E \sim N$ and $C_V \sim N$. Which means that the relative size of the fluctuations scales as

$$\frac{\Delta E}{E} \sim \frac{1}{\sqrt{N}} \tag{1.28}$$

The limit $N \to \infty$ in known as the *thermodynamic limit*. The energy becomes peaked closer and closer to the mean value $\langle E \rangle$ and can be treated as essentially fixed. But this was our starting point for the microcanonical ensemble. In the thermodynamic limit, the microcanonical and canonical ensembles coincide.

All the examples that we will discuss in the course will have a *very* large number of particles, N, and we can consider ourselves safely in the thermodynamic limit. For that reason, even in the canonical ensemble, we will often write E for the average energy rather than $\langle E \rangle$.

An Example: The Two State System Revisited

We can rederive our previous results for the two state system using the canonical ensemble. It is marginally simpler. For a single particle with two energy levels, 0 and ϵ, the partition function is given by

$$Z_1 = \sum_n e^{-\beta E_n} = 1 + e^{-\beta\epsilon} = 2e^{-\beta\epsilon/2}\cosh(\beta\epsilon/2)$$

We want the partition function for N such particles. But we saw in (1.23) that if we have independent systems, then we simply need to multiply their partition functions together. We then have

$$Z = 2^N e^{-N\beta\epsilon/2}\cosh^N(\beta\epsilon/2)$$

from which we can easily compute the average energy

$$\langle E \rangle = -\frac{\partial}{\partial\beta}\log Z = \frac{N\epsilon}{2}\left(1 - \tanh(\beta\epsilon/2)\right)$$

A bit of algebra will reveal that this is the same expression that we derived in the microcanonical ensemble (1.13). We could now go on to compute the heat capacity and reproduce the result (1.14).

Notice that, unlike in the microcanonical ensemble, we didn't have to solve any combinatoric problem to count states. The partition function has done all that work for us. Of course, for this simple two state system, the counting of states wasn't difficult but in later examples, where the counting gets somewhat tricker, the partition function will be an invaluable tool to save us the work.

1.3.3 Entropy

Recall that in the microcanonical ensemble, the entropy counts the (log of the) number of states with fixed energy. We would like to define an analogous quantity in the canonical ensemble where we have a probability distribution over states with different energies. How to proceed? Our strategy will be to again return to the microcanonical ensemble, now applied to the combined system + reservoir.

In fact, we're going to use a little trick. Suppose that we don't have just one copy of our system S, but instead a large number, W, of identical copies. Each system lives in a particular state $|n\rangle$. If W is large enough, the number of systems that sit in state $|n\rangle$ must be simply $p(n)W$. We see that the trick of taking W copies has translated the probabilities into eventualities. To determine the entropy we can treat the whole collection of W systems as sitting in the microcanonical ensemble to which we can apply the familiar Boltzmann definition of entropy (1.2). We must only figure out how many ways there are of putting $p(n)W$ systems into state $|n\rangle$ for each $|n\rangle$. That's a simple combinatoric problem: the answer is

$$\Omega = \frac{W!}{\prod_n (p(n)W)!}$$

And the entropy is therefore

$$S = k_B \log \Omega = -k_B W \sum_n p(n) \log p(n) \tag{1.29}$$

where we have used Stirling's formula to simplify the logarithms of factorials. This is the entropy for all W copies of the system. But we also know that entropy is additive. So the entropy for a single copy of the system, with probability distribution $p(n)$ over the states is

$$S = -k_B \sum_n p(n) \log p(n) \tag{1.30}$$

This beautiful formula is due to Gibbs. It was rediscovered some decades later in the context of information theory where it goes by the name of Shannon entropy for classical systems or von Neumann entropy for quantum systems. In the quantum context, it is sometimes written in terms of the density matrix (1.24) as

$$S = -k_B \mathrm{Tr}\, \hat{\rho} \log \hat{\rho}$$

When we first introduced entropy in the microcanonical ensemble, we viewed it as a function of the energy E. But (1.30) gives a very different viewpoint on the entropy:

it says that we should view S as a function of a probability distribution. There is no contradiction with the microcanonical ensemble because in that simple case, the probability distribution is itself determined by the choice of energy E. Indeed, it is simple to check (and you should!) that the Gibbs entropy (1.30) reverts to the Boltzmann entropy in the special case of $p(n) = 1/\Omega(E)$ for all states $|n\rangle$ of energy E.

Meanwhile, back in the canonical ensemble, the probability distribution is entirely determined by the choice of temperature T. This means that the entropy is naturally a function of T. Indeed, substituting the Boltzmann distribution $p(n) = e^{-\beta E_n}/Z$ into the expression (1.30), we find that the entropy in the canonical ensemble is given by

$$\begin{aligned} S &= -\frac{k_B}{Z}\sum_n e^{-\beta E_n}\log\left(\frac{e^{-\beta E_n}}{Z}\right) \\ &= \frac{k_B\beta}{Z}\sum_n E_n e^{-\beta E_n} + k_B\log Z \end{aligned}$$

As with all other important quantities, this can be elegantly expressed in terms of the partition function by

$$S = k_B\frac{\partial}{\partial T}(T\log Z) \tag{1.31}$$

A Comment on the Microcanonical vs Canonical Ensembles

The microcanonical and canonical ensembles are different probability distributions. This means, using the definition (1.30), that they generally have different entropies. Nonetheless, in the limit of a large number of particles, $N \to \infty$, all physical observables — including entropy — coincide in these two distributions. We've already seen this when we computed the variance of energy (1.28) in the canonical ensemble. Let's take a closer look at how this works.

The partition function in (1.21) is a sum over all states. We can rewrite it as a sum over energy levels by including a degeneracy factor

$$Z = \sum_{\{E_i\}}\Omega(E_i)e^{-\beta E_i}$$

The degeneracy factor $\Omega(E)$ factor is typically a rapidly rising function of E, while the Boltzmann suppression $e^{-\beta E}$ is rapidly falling. But, for both the exponent is proportional to N which is itself exponentially large. This ensures that the sum over energy levels is entirely dominated by the maximum value, $E_\star$, defined by the requirement

$$\frac{\partial}{\partial E}\Big(\Omega(E)e^{-\beta E}\Big)_{E=E_\star} = 0$$

and the partition function can be well approximated by

$$Z \approx \Omega(E_\star)e^{-\beta E_\star}$$

(This is the same kind of argument we used in (1.2.1) in our discussion of the Second Law). With this approximation, we can use (1.25) to show that the most likely energy $E_\star$ and the average energy $\langle E \rangle$ coincide:

$$\langle E \rangle = E_\star$$

(We need to use the result (1.7) in the form $\partial \log \Omega(E_\star)/\partial E_\star = \beta$ to derive this). Similarly, using (1.31), we can show that the entropy in the canonical ensemble is given by

$$S = k_B \log \Omega(E_\star)$$

Maximizing Entropy

There is actually a unified way to think about the microcanonical and canonical ensembles in terms of a variational principle: the different ensembles have the property that they maximise the entropy subject to various constraints. The only difference between them is the constraints that are imposed.

Let's start with the microcanonical ensemble, in which we fix the energy of the system so that we only allow non-zero probabilities for those states which have energy E. We could then compute the entropy using the Gibbs formula (1.30) for any probability distribution, including systems away from equilibrium. We need only insist that all the probabilities add up to one: $\sum_n p(n) = 1$. We can maximise S subject to this condition by introducing a Lagrange multiplier α and maximising $S + \alpha k_B(\sum_n p(n) - 1)$,

$$\frac{\partial}{\partial p(n)}\left(-\sum_n p(n)\log p(n) + \alpha \sum_n p(n) - \alpha\right) = 0 \qquad \Rightarrow \qquad p(n) = e^{\alpha - 1}$$

We learn that all states with energy E are equally likely. This is the microcanonical ensemble.

In the examples sheet, you will be asked to show that the canonical ensemble can be viewed in the same way: it is the probability distribution that maximises the entropy subject to the constraint that the *average* energy is fixed.

1.3.4 Free Energy

We've left the most important quantity in the canonical ensemble to last. It is called the *free energy*,

$$F = \langle E \rangle - TS \tag{1.32}$$

There are actually a number of quantities all vying for the name "free energy", but the quantity F is the one that physicists usually work with. When necessary to clarify, it is sometimes referred to as the Helmholtz free energy. The word "free" here doesn't mean "without cost". Energy is never free in that sense. Rather, it should be interpreted as the "available" energy.

Heuristically, the free energy captures the competition between energy and entropy that occurs in a system at constant temperature. Immersed in a heat bath, energy is not necessarily at a premium. Indeed, we saw in the two-state example that the ground state plays little role in the physics at non-zero temperature. Instead, the role of entropy becomes more important: the existence of many high energy states can beat a few low-energy ones.

The fact that the free energy is the appropriate quantity to look at for systems at fixed temperature is also captured by its mathematical properties. Recall, that we started in the microcanonical ensemble by defining entropy $S = S(E, V)$. If we invert this expression, then we can equally well think of energy as a function of entropy and volume: $E = E(S, V)$. This is reflected in the first law of thermodynamics (1.16) which reads $dE = TdS - pdV$. However, if we look at small variations in F, we get

$$dF = d\langle E \rangle - d(TS) = -SdT - pdV \tag{1.33}$$

This form of the variation is telling us that we should think of the free energy as a function of temperature and volume: $F = F(T, V)$. Mathematically, F is a Legendre transform of E.

Given the free energy, the variation (1.33) tells us how to get back the entropy,

$$S = -\left.\frac{\partial F}{\partial T}\right|_V \tag{1.34}$$

Similarly, the pressure is given by

$$p = -\left.\frac{\partial F}{\partial V}\right|_T \tag{1.35}$$

The free energy is the most important quantity at fixed temperature. It is also the quantity that is most directly related to the partition function Z:

$$F = -k_B T \log Z \tag{1.36}$$

This relationship follows from (1.25) and (1.31). Using the identity $\partial/\partial\beta = -k_B T^2 \partial/\partial T$, these expressions allow us to write the free energy as

$$\begin{aligned} F = E - TS &= k_B T^2 \frac{\partial}{\partial T} \log Z - k_B T \frac{\partial}{\partial T}(T \log Z) \\ &= -k_B T \log Z \end{aligned}$$

as promised.

1.4 The Chemical Potential

Before we move onto applications, there is one last bit of formalism that we will need to introduce. This arises in situations where there is some other conserved quantity which restricts the states that are accessible to the system. The most common example is simply the number of particles N in the system. Another example is the electric charge Q. For the sake of definiteness, we will talk about particle number below but all the comments apply to any conserved quantity.

In both the microcanonical and canonical ensembles, we should only consider states that have a fixed value of N. We already did this when we discussed the two state system — for example, the expression for entropy (1.12) depends explicitly on the number of particles N. We will now make this dependence explicit and write

$$S(E, V, N) = k_B \log \Omega(E, V, N)$$

The entropy leads us to the temperature as $1/T = \partial S/\partial E$ and the pressure as $p = T\partial S/\partial V$. But now we have another option: we can differentiate with respect to particle number N. The resulting quantity is called the *chemical potential*,

$$\mu = -T \frac{\partial S}{\partial N} \tag{1.37}$$

Using this definition, we can re-run the arguments given in Section 1.2.2 for systems which are allowed to exchange particles. Such systems are in equilibrium only if they have equal chemical potential μ. This condition is usually referred to as chemical equilibrium.

To get a feel for the meaning of the chemical potential, we can look again at the first law of thermodynamics (1.16), now allowing for a change in particle number as well. Writing $dS = \ldots$ and rearranging, we have,

$$dE = TdS - pdV + \mu dN \tag{1.38}$$

This tells us the meaning of the chemical potential: it is the energy cost to add one more particle to the system while keeping both S and V fixed. (Strictly speaking, an infinitesimal amount of particle, but if we're adding one more to 10^{23} that effectively counts as infinitesimal). If we're interested in electric charge Q, rather than particle number, the chemical potential is the same thing as the familiar electrostatic potential of the system that you met in your first course in Electromagnetism.

There's actually a subtle point in the above derivation that is worth making explicit. It's the kind of thing that will crop up a lot in thermodynamics where you typically have many variables and need to be careful about which ones are kept fixed. We defined the chemical potential as $\mu = -T\, \partial S/\partial N|_{E,V}$. But the first law is telling us that we can also think of the chemical potential as $\mu = \partial E/\partial N|_{S,V}$. Why is this the same thing? This follows from a general formula for partial derivatives. If you have three variables, x, y and z, with a single constraint between them, then

$$\left.\frac{\partial x}{\partial y}\right|_z \left.\frac{\partial y}{\partial z}\right|_x \left.\frac{\partial z}{\partial x}\right|_y = -1$$

Applying this general formula to E, S and N gives us the required result

$$\left.\frac{\partial E}{\partial N}\right|_{S,V} = -\left.\frac{\partial S}{\partial N}\right|_{E,v} \left.\frac{\partial E}{\partial S}\right|_{N,V} = \mu$$

If we work at constant temperature rather than constant energy, the relevant function is the free energy $F(T, V, N) = E - TS$. Small changes are given by

$$dF = -SdT - pdV + \mu dN$$

from which we see that the chemical potential can also be defined as

$$\mu = \left.\frac{\partial F}{\partial N}\right|_{T,V}$$

1.4.1 Grand Canonical Ensemble

When we made the transition from microcanonical to canonical ensemble, we were no longer so rigid in our insistence that the system has a fixed energy. Rather it could freely

exchange energy with the surrounding reservoir, which was kept at a fixed temperature. We could now imagine the same scenario with any other conserved quantity. For example, if particles are free to move between the system and the reservoir, then N is no longer fixed. In such a situation, we will require that the reservoir sits at fixed chemical potential μ as well as fixed temperature T.

The probability distribution that we need to use in this case is called the *grand canonical ensemble*. The probability of finding the system in a state $|n\rangle$ depends on both the energy E_n and the particle number N_n. (Notice that because N is conserved, the quantum mechanical operator necessarily commutes with the Hamiltonian so there is no difficulty in assigning both energy and particle number to each state). We introduce the *grand canonical partition function*

$$\mathcal{Z}(T,\mu,V) = \sum_n e^{-\beta(E_n-\mu N_n)} \tag{1.39}$$

Re-running the argument that we used for the canonical ensemble, we find the probability that the system is in state $|n\rangle$ to be

$$p(n) = \frac{e^{-\beta(E_n-\mu N_n)}}{\mathcal{Z}}$$

In the canonical ensemble, all the information that we need is contained within the partition function Z. In the grand canonical ensemble it is contained within $\mathcal{Z}$. The entropy (1.30) is once again given by

$$S = k_B \frac{\partial}{\partial T}(T \log \mathcal{Z}) \tag{1.40}$$

while differentiating with respect to β gives us

$$\langle E\rangle - \mu\langle N\rangle = -\frac{\partial}{\partial\beta}\log\mathcal{Z} \tag{1.41}$$

The average particle number $\langle N\rangle$ in the system can then be separately extracted by

$$\langle N\rangle = \frac{1}{\beta}\frac{\partial}{\partial\mu}\log\mathcal{Z} \tag{1.42}$$

and its fluctuations,

$$\Delta N^2 = \frac{1}{\beta^2}\frac{\partial^2}{\partial\mu^2}\log\mathcal{Z} = \frac{1}{\beta}\frac{\partial\langle N\rangle}{\partial\mu} \tag{1.43}$$

Just as the average energy is determined by the temperature in the canonical ensemble, here the average particle number is determined by the chemical potential. The grand canonical ensemble will simplify several calculations later, especially when we come to discuss Bose and Fermi gases in Section 3.

The relative size of these fluctuations scales in the same way as the energy fluctuations, $\Delta N/\langle N\rangle \sim 1/\sqrt{\langle N\rangle}$, and in the thermodynamic limit $N \to \infty$ results from all three ensembles coincide. For this reason, we will drop the averaging brackets $\langle\cdot\rangle$ from our notation and simply refer to the average particle number as N.

1.4.2 Grand Canonical Potential

The grand canonical potential Φ is defined by

$$\Phi = F - \mu N$$

Φ is a Legendre transform of F, from variable N to μ. This is underlined if we look at small variations,

$$d\Phi = -SdT - pdV - Nd\mu \tag{1.44}$$

which tells us that Φ should be thought of as a function of temperature, volume and chemical potential, $\Phi = \Phi(T, V, \mu)$.

We can perform the same algebraic manipulations that gave us F in terms of the canonical partition function Z, this time using the definitions (1.40) and (1.41)) to write Φ as

$$\Phi = -k_B T \log \mathcal{Z} \tag{1.45}$$

1.4.3 Extensive and Intensive Quantities

There is one property of Φ that is rather special and, at first glance, somewhat surprising. This property actually follows from very simple considerations of how different variables change as we look at bigger and bigger systems.

Suppose we have a system and we double it. That means that we double the volume V, double the number of particles N and double the energy E. What happens to all our other variables? We have already seen back in Section 1.2.1 that entropy is additive, so S also doubles. More generally, if we scale V, N and E by some amount λ, the entropy must scale as

$$S(\lambda E, \lambda V, \lambda N) = \lambda S(E, V, N)$$

Quantities such as E, V, N and S which scale in this manner are called *extensive*. In contrast, the variables which arise from differentiating the entropy, such as temperature $1/T = \partial S/\partial E$ and pressure $p = T\partial S/\partial V$ and chemical potential $\mu = T\partial S/\partial N$ involve the ratio of two extensive quantities and so do not change as we scale the system: they are called *intensive* quantities.

What now happens as we make successive Legendre transforms? The free energy $F = E - TS$ is also extensive (since E and S are extensive while T is intensive). So it must scale as

$$F(T, \lambda V, \lambda N) = \lambda F(T, V, N) \tag{1.46}$$

Similarly, the grand potential $\Phi = F - \mu N$ is extensive and scales as

$$\Phi(T, \lambda V, \mu) = \lambda \Phi(T, V, \mu) \tag{1.47}$$

But there's something special about this last equation, because Φ only depends on a single extensive variable, namely V. While there are many ways to construct a free energy F which obeys (1.46) (for example, any function of the form $F \sim V^{n+1}/N^n$ will do the job), there is only one way to satisfy (1.47): Φ must be proportional to V. But we've already got a name for this proportionality constant: it is pressure. (Actually, it is minus the pressure as you can see from (1.44)). So we have the equation

$$\Phi(T, V, \mu) = -p(T, \mu)V \tag{1.48}$$

It looks as if we got something for free! If F is a complicated function of V, where do these complications go after the Legendre transform to Φ? The answer is that the complications go into the pressure $p(T, \mu)$ when expressed as a function of T and μ. Nonetheless, equation (1.48) will prove to be an extremely economic way to calculate the pressure of various systems.

1.4.4 Josiah Willard Gibbs (1839-1903)

> "Usually, Gibbs' prose style conveys his meaning in a sufficiently clear way, using no more than twice as many words as Poincaré or Einstein would have used to say the same thing."
>
> *E.T.Jaynes on the difficulty of reading Gibbs*

Gibbs was perhaps the first great American theoretical physicist. Many of the developments that we met in this chapter are due to him, including the free energy, the chemical potential and, most importantly, the idea of ensembles. Even the name "statistical mechanics" was invented by Gibbs.

Gibbs provided the first modern rendering of the subject in a treatise published shortly before his death. Very few understood it. Lord Rayleigh wrote to Gibbs suggesting that the book was "too condensed and too difficult for most, I might say all, readers". Gibbs disagreed. He wrote back saying the book was only "too long".

There do not seem to be many exciting stories about Gibbs. He was an undergraduate at Yale. He did a PhD at Yale. He became a professor at Yale. Apparently he rarely left New Haven. Strangely, he did not receive a salary for the first ten years of his professorship. Only when he received an offer from John Hopkins of $3000 dollars a year did Yale think to pay America's greatest physicist. They made a counter-offer of $2000 dollars and Gibbs stayed.

2. Classical Gases

Our goal in this section is to use the techniques of statistical mechanics to describe the dynamics of the simplest system: a gas. This means a bunch of particles, flying around in a box. Although much of the last section was formulated in the language of quantum mechanics, here we will revert back to classical mechanics. Nonetheless, a recurrent theme will be that the quantum world is never far behind: we'll see several puzzles, both theoretical and experimental, which can only truly be resolved by turning on $\hbar$.

2.1 The Classical Partition Function

For most of this section we will work in the canonical ensemble. We start by reformulating the idea of a partition function in classical mechanics. We'll consider a simple system – a single particle of mass m moving in three dimensions in a potential $V(\vec{q})$. The classical Hamiltonian of the system[3] is the sum of kinetic and potential energy,

$$H = \frac{\vec{p}^2}{2m} + V(\vec{q})$$

We earlier defined the partition function (1.21) to be the sum over all quantum states of the system. Here we want to do something similar. In classical mechanics, the state of a system is determined by a point in *phase space*. We must specify both the position and momentum of each of the particles — only then do we have enough information to figure out what the system will do for all times in the future. This motivates the definition of the partition function for a single classical particle as the integration over phase space,

$$Z_1 = \frac{1}{h^3} \int d^3q d^3p \; e^{-\beta H(p,q)} \tag{2.1}$$

The only slightly odd thing is the factor of $1/h^3$ that sits out front. It is a quantity that needs to be there simply on dimensional grounds: Z should be dimensionless so h must have dimension (length $\times$ momentum) or, equivalently, Joules-seconds (Js). The actual value of h won't matter for any physical observable, like heat capacity, because we always take $\log Z$ and then differentiate. Despite this, there is actually a correct value for h: it is Planck's constant, $h = 2\pi\hbar \approx 6.6 \times 10^{-34}\ Js$.

It is very strange to see Planck's constant in a formula that is supposed to be classical. What's it doing there? In fact, it is a vestigial object, like the male nipple. It is redundant, serving only as a reminder of where we came from. And the classical world came from the quantum.

[3] If you haven't taken the Classical Dynamics course, you should think of the Hamiltonian as the energy of the system expressed in terms of the position and momentum of the particle.

2.1.1 From Quantum to Classical

It is possible to derive the classical partition function (2.1) directly from the quantum partition function (1.21) without resorting to hand-waving. It will also show us why the factor of $1/h$ sits outside the partition function. The derivation is a little tedious, but worth seeing. (Similar techniques are useful in later courses when you first meet the path integral). To make life easier, let's consider a single particle moving in one spatial dimension. It has position operator $\hat{q}$, momentum operator $\hat{p}$ and Hamiltonian,

$$\hat{H} = \frac{\hat{p}^2}{2m} + V(\hat{q})$$

If $|n\rangle$ is the energy eigenstate with energy E_n, the quantum partition function is

$$Z_1 = \sum_n e^{-\beta E_n} = \sum_n \langle n|e^{-\beta \hat{H}}|n\rangle \tag{2.2}$$

In what follows, we'll make liberal use of the fact that we can insert the identity operator anywhere in this expression. Identity operators can be constructed by summing over any complete basis of states. We'll need two such constructions, using the position eigenvectors $|q\rangle$ and the momentum eigenvectors $|p\rangle$,

$$\mathbf{1} = \int dq\, |q\rangle\langle q| \quad , \quad \mathbf{1} = \int dp\, |p\rangle\langle p|$$

We start by inserting two copies of the identity built from position eigenstates,

$$\begin{aligned} Z_1 &= \sum_n \langle n| \int dq\, |q\rangle\langle q|e^{-\beta\hat{H}} \int dq'|q'\rangle\langle q'|n\rangle \\ &= \int dq dq'\, \langle q|e^{-\beta\hat{H}}|q'\rangle \sum_n \langle q'|n\rangle\langle n|q\rangle \end{aligned}$$

But now we can replace $\sum_n |n\rangle\langle n|$ with the identity matrix and use the fact that $\langle q'|q\rangle = \delta(q'-q)$, to get

$$Z_1 = \int dq\, \langle q|e^{-\beta\hat{H}}|q\rangle \tag{2.3}$$

We see that the result is to replace the sum over energy eigenstates in (2.2) with a sum (or integral) over position eigenstates in (2.3). If you wanted, you could play the same game and get the sum over any complete basis of eigenstates of your choosing. As an aside, this means that we can write the partition function in a basis independent fashion as

$$Z_1 = \mathrm{Tr}\, \mathrm{e}^{-\beta\hat{\mathrm{H}}}$$

So far, our manipulations could have been done for any quantum system. Now we want to use the fact that we are taking the classical limit. This comes about when we try to factorize $e^{-\beta\hat{H}}$ into a momentum term and a position term. The trouble is that this isn't always possible when there are matrices (or operators) in the exponent. Recall that,

$$e^{\hat{A}}e^{\hat{B}} = e^{\hat{A}+\hat{B}+\frac{1}{2}[\hat{A},\hat{B}]+\ldots}$$

For us $[\hat{q},\hat{p}] = i\hbar$. This means that if we're willing to neglect terms of order $\hbar$ — which is the meaning of taking the classical limit — then we can write

$$e^{-\beta\hat{H}} = e^{-\beta\hat{p}^2/2m}\, e^{-\beta V(\hat{q})} + \mathcal{O}(\hbar)$$

We can now start to replace some of the operators in the exponent, like $V(\hat{q})$, with functions $V(q)$. (The notational difference is subtle, but important, in the expressions below!),

$$\begin{aligned} Z_1 &= \int dq\, \langle q|e^{-\beta\hat{p}^2/2m}e^{-\beta V(\hat{q})}|q\rangle \\ &= \int dq\, e^{-\beta V(q)}\langle q|e^{-\beta\hat{p}^2/2m}|q\rangle \\ &= \int dqdpdp' e^{-\beta V(q)}\langle q|p\rangle\langle p|e^{-\beta\hat{p}^2/2m}|p'\rangle\langle p'|q\rangle \\ &= \frac{1}{2\pi\hbar}\int dqdp\, e^{-\beta H(p,q)} \end{aligned}$$

where, in the final line, we've used the identity

$$\langle q|p\rangle = \frac{1}{\sqrt{2\pi\hbar}}e^{ipq/\hbar}$$

This completes the derivation.

2.2 Ideal Gas

The first classical gas that we'll consider consists of N particles trapped inside a box of volume V. The gas is "ideal". This simply means that the particles do not interact with each other. For now, we'll also assume that the particles have no internal structure, so no rotational or vibrational degrees of freedom. This situation is usually referred to as the monatomic ideal gas. The Hamiltonian for each particle is simply the kinetic energy,

$$H = \frac{\vec{p}^2}{2m}$$

And the partition function for a single particle is

$$Z_1(V,T) = \frac{1}{(2\pi\hbar)^3}\int d^3q d^3p\, e^{-\beta\vec{p}^2/2m} \tag{2.4}$$

The integral over position is now trivial and gives $\int d^3q = V$, the volume of the box. The integral over momentum is also straightforward since it factorizes into separate integrals over p_x, p_y and p_z, each of which is a Gaussian of the form,

$$\int dx\, e^{-ax^2} = \sqrt{\frac{\pi}{a}}$$

So we have

$$Z_1 = V\left(\frac{mk_BT}{2\pi\hbar^2}\right)^{3/2}$$

We'll meet the combination of factors in the brackets a lot in what follows, so it is useful to give it a name. We'll write

$$Z_1 = \frac{V}{\lambda^3} \tag{2.5}$$

The quantity λ goes by the name of the *thermal de Broglie wavelength*,

$$\lambda = \sqrt{\frac{2\pi\hbar^2}{mk_BT}} \tag{2.6}$$

λ has the dimensions of length. We will see later that you can think of λ as something like the average de Broglie wavelength of a particle at temperature T. Notice that it is a quantum object – it has an $\hbar$ sitting in it – so we expect that it will drop out of any genuinely classical quantity that we compute. The partition function itself (2.5) is counting the number of these thermal wavelengths that we can fit into volume V.

Z_1 is the partition function for a single particle. We have N, non-interacting, particles in the box so the partition function of the whole system is

$$Z(N,V,T) = Z_1^N = \frac{V^N}{\lambda^{3N}} \tag{2.7}$$

(Full disclosure: there's a slightly subtle point that we're brushing under the carpet here and this equation isn't quite right. This won't affect our immediate discussion and we'll explain the issue in more detail in Section 2.2.3.)

Figure 8: Deviations from ideal gas law at sensible densities

Figure 9: Deviations from ideal gas law at extreme densities

Armed with the partition function Z, we can happily calculate anything that we like. Let's start with the pressure, which can be extracted from the partition function by first computing the free energy (1.36) and then using (1.35). We have

$$\begin{aligned} p &= -\frac{\partial F}{\partial V} \\ &= \frac{\partial}{\partial V}(k_B T \log Z) \\ &= \frac{N k_B T}{V} \end{aligned} \tag{2.8}$$

This equation is an old friend – it is the ideal gas law, $pV = Nk_BT$, that we all met in kindergarten. Notice that the thermal wavelength λ has indeed disappeared from the discussion as expected. Equations of this form, which link pressure, volume and temperature, are called *equations of state*. We will meet many throughout this course.

As the plots above show[4], the ideal gas law is an extremely good description of gases at low densities. Gases deviate from this ideal behaviour as the densities increase and the interactions between atoms becomes important. We will see how this comes about from the viewpoint of microscopic forces in Section 2.5.

It is worth pointing out that this derivation should calm any lingering fears that you had about the definition of temperature given in (1.7). The object that we call T really does coincide with the familiar notion of temperature applied to gases. But the key property of the temperature is that if two systems are in equilibrium then they have the same T. That's enough to ensure that equation (1.7) is the right definition of temperature for *all* systems because we can always put any system in equilibrium with an ideal gas.

[4]Both figures are taken from the web textbook "*General Chemistry*" and credited to John Hutchinson.

2.2.1 Equipartition of Energy

The partition function (2.7) has more in store for us. We can compute the average energy of the ideal gas,

$$E = -\frac{\partial}{\partial \beta} \log Z = \frac{3}{2} N k_B T \tag{2.9}$$

There's an important, general lesson lurking in this formula. To highlight this, it is worth repeating our analysis for an ideal gas in arbitrary number of spatial dimensions, D. A simple generalization of the calculations above shows that

$$Z = \frac{V^N}{\lambda^{DN}} \qquad \Rightarrow \qquad E = \frac{D}{2} N k_B T$$

Each particle has D degrees of freedom (because it can move in one of D spatial directions). And each particle contributes $\frac{1}{2} D k_B T$ towards the average energy. This is a general rule of thumb, which holds for all classical systems: the average energy of each free degree of freedom in a system at temperature T is $\frac{1}{2} k_B T$. This is called the *equipartition of energy*. As stated, it holds only for degrees of freedom in the absence of a potential. (There is a modified version if you include a potential). Moreover, it holds only for classical systems or quantum systems at suitably high temperatures.

We can use the result above to see why the thermal de Broglie wavelength (2.6) can be thought of as roughly equal to the average de Broglie wavelength of a particle. Equating the average energy (2.9) to the kinetic energy $E = p^2/2m$ tells us that the average (root mean square) momentum carried by each particle is $p \sim \sqrt{m k_B T}$. In quantum mechanics, the de Broglie wavelength of a particle is $\lambda_{dB} = h/p$, which (up to numerical factors of 2 and π) agrees with our formula (2.6).

Finally, returning to the reality of $d = 3$ dimensions, we can compute the heat capacity for a monatomic ideal gas. It is

$$C_V = \left. \frac{\partial E}{\partial T} \right|_V = \frac{3}{2} N k_B \tag{2.10}$$

2.2.2 The Sociological Meaning of Boltzmann's Constant

We introduced Boltzmann's constant k_B in our original the definition of entropy (1.2). It has the value,

$$k_B = 1.381 \times 10^{-23} \ JK^{-1}$$

In some sense, there is no deep physical meaning to Boltzmann's constant. It is merely a conversion factor that allows us to go between temperature and energy, as reflected

in (1.7). It is necessary to include it in the equations only for historical reasons: our ancestors didn't realise that temperature and energy were closely related and measured them in different units.

Nonetheless, we could ask why does k_B have the value above? It doesn't seem a particularly natural number. The reason is that both the units of temperature (Kelvin) and energy (Joule) are picked to reflect the conditions of human life. In the everyday world around us, measurements of temperature and energy involve fairly ordinary numbers: room temperature is roughly $300\,K$; the energy required to lift an apple back up to the top of the tree is a few Joules. Similarly, in an everyday setting, all the measurable quantities — p, V and T — in the ideal gas equation are fairly normal numbers when measured in SI units. The only way this can be true is if the combination Nk_B is a fairly ordinary number, of order one. In other words the number of atoms must be huge,

$$N \sim 10^{23} \tag{2.11}$$

This then is the real meaning of the value of Boltzmann's constant: *atoms are small.*

It's worth stressing this point. Atoms aren't just small: they're really really small. 10^{23} is an astonishingly large number. The number of grains of sand in all the beaches in the world is around 10^{18}. The number of stars in our galaxy is about 10^{11}. The number of stars in the entire visible Universe is probably around 10^{22}. And yet the number of water molecules in a cup of tea is more than 10^{23}.

Chemist Notation

While we're talking about the size of atoms, it is probably worth reminding you of the notation used by chemists. They too want to work with numbers of order one. For this reason, they define a *mole* to be the number of atoms in one gram of Hydrogen. (Actually, it is the number of atoms in 12 grams of Carbon-12, but this is roughly the same thing). The mass of Hydrogen is $1.6 \times 10^{-27}\ Kg$, so the number of atoms in a mole is Avogadro's number,

$$N_A \approx 6 \times 10^{23}$$

The number of moles in our gas is then $n = N/N_A$ and the ideal gas law can be written as

$$pV = nRT$$

where $R = N_A k_B$ is the called the Universal gas constant. Its value is a nice sensible number with no silly power in the exponent: $R \approx 8\ JK^{-1}\text{mol}^{-1}$.

2.2.3 Entropy and Gibbs's Paradox

> "It has always been believed that Gibbs's paradox embodied profound thought. That it was intimately linked up with something so important and entirely new could hardly have been foreseen."
>
> *Erwin Schrödinger*

We said earlier that the formula for the partition function (2.7) isn't quite right. What did we miss? We actually missed a subtle point from quantum mechanics: quantum particles are indistinguishable. If we take two identical atoms and swap their positions, this doesn't give us a new state of the system – it is the same state that we had before. (Up to a sign that depends on whether the atoms are bosons or fermions – we'll discuss this aspect in more detail in Sections 3.5 and 3.6). However, we haven't taken this into account – we wrote the expression $Z = Z_1^N$ which would be true if all the N particles in the were distinguishable — for example, if each of the particles were of a different type. But this naive partition function overcounts the number of states in the system when we're dealing with indistinguishable particles.

It is a simple matter to write down the partition function for N indistinguishable particles. We simply need to divide by the number of ways to permute the particles. In other words, for the ideal gas the partition function is

$$Z_{\text{ideal}}(N, V, T) = \frac{1}{N!} Z_1^N = \frac{V^N}{N! \lambda^{3N}} \tag{2.12}$$

The extra factor of $N!$ doesn't change the calculations of pressure or energy since, for each, we had to differentiate $\log Z$ and any overall factor drops out. However, it does change the entropy since this is given by,

$$S = \frac{\partial}{\partial T}(k_B T \log Z_{\text{ideal}})$$

which includes a factor of $\log Z$ without any derivative. Of course, since the entropy is counting the number of underlying microstates, we would expect it to know about whether particles are distinguishable or indistinguishable. Using the correct partition function (2.12) and Stirling's formula, the entropy of an ideal gas is given by,

$$S = N k_B \left[\log \left(\frac{V}{N \lambda^3} \right) + \frac{5}{2} \right] \tag{2.13}$$

This result is known as the *Sackur-Tetrode* equation. Notice that not only is the entropy sensitive to the indistinguishability of the particles, but it also depends on λ. However, the entropy is not directly measurable classically. We can only measure entropy differences by the integrating the heat capacity as in (1.10).

The benefit of adding an extra factor of $N!$ was noticed before the advent of quantum mechanics by Gibbs. He was motivated by the change in entropy of mixing between two gases. Suppose that we have two different gases, say red and blue. Each has the same number of particles N and sits in a volume V, separated by a partition. When the partition is removed the gases mix and we expect the entropy to increase. But if the gases are of the same type, removing the partition shouldn't change the macroscopic state of the gas. So why should the entropy increase? This is referred to as the *Gibb's paradox*. Including the factor of $N!$ in the partition function ensures that the entropy does not increase when identical atoms are mixed[5]

2.2.4 The Ideal Gas in the Grand Canonical Ensemble

It is worth briefly looking at the ideal gas in the grand canonical ensemble. Recall that in such an ensemble, the gas is free to exchange both energy and particles with the outside reservoir. You could think of the system as some fixed subvolume inside a much larger gas. If there are no walls to define this subvolume then particles, and hence energy, can happily move in and out. We can ask how many particles will, on average, be inside this volume and what fluctuations in particle number will occur. More importantly, we can also start to gain some intuition for this strange quantity called the chemical potential, μ.

The grand partition function (1.39) for the ideal gas is

$$\mathcal{Z}_{\text{ideal}}(\mu, V, T) = \sum_{N=0}^{\infty} e^{\beta\mu N} Z_{\text{ideal}}(N, V, T) = \exp\left(\frac{e^{\beta\mu}V}{\lambda^3}\right)$$

From this we can determine the average particle number,

$$N = \frac{1}{\beta}\frac{\partial}{\partial\mu}\log\mathcal{Z} = \frac{e^{\beta\mu}V}{\lambda^3}$$

Which, rearranging, gives

$$\mu = k_B T \log\left(\frac{\lambda^3 N}{V}\right) \tag{2.14}$$

If $\lambda^3 < V/N$ then the chemical potential is negative. Recall that λ is roughly the average de Broglie wavelength of each particle, while V/N is the average volume taken

[5]Be warned however: a closer look shows that the Gibbs paradox is rather toothless and, in the classical world, there is no real necessity to add the $N!$. A clear discussion of these issues can be found in E.T. Jaynes' article "*The Gibbs Paradox*" which you can download from the course website.

up by each particle. But whenever the de Broglie wavelength of particles becomes comparable to the inter-particle separation, then quantum effects become important. In other words, to trust our classical calculation of the ideal gas, we must have $\lambda^3 \ll V/N$ and, correspondingly, $\mu < 0$.

At first sight, it is slightly strange that μ is negative. When we introduced μ in Section 1.4.1, we said that it should be thought of as the energy cost of adding an extra particle to the system. Surely that energy should be positive! To see why this isn't the case, we should look more closely at the definition. From the energy variation (1.38), we have

$$\mu = \left.\frac{\partial E}{\partial N}\right|_{S,V}$$

So the chemical potential should be thought of as the energy cost of adding an extra particle *at fixed entropy and volume*. But adding a particle will give more ways to share the energy around and so increase the entropy. If we insist on keeping the entropy fixed, then we will need to reduce the energy when we add an extra particle. This is why we have $\mu < 0$ for the classical ideal gas.

There are situations where $\mu > 0$. This can occur if we have a suitably strong repulsive interaction between particles so that there's a large energy cost associated to throwing in one extra. We also have $\mu > 0$ for fermion systems at low temperatures as we will see in Section 3.6.

We can also compute the fluctuation in the particle number,

$$\Delta N^2 = \frac{1}{\beta^2}\frac{\partial^2}{\partial \mu^2}\log \mathcal{Z}_{\text{ideal}} = N$$

As promised in Section 1.4.1, the relative fluctuations $\Delta N/\langle N\rangle = 1/\sqrt{N}$ are vanishingly small in the thermodynamic $N \to \infty$ limit.

Finally, it is very easy to compute the equation of state in the grand canonical ensemble because (1.45) and (1.48) tell us that

$$pV = k_B T \log \mathcal{Z} = k_B T \frac{e^{\beta\mu} V}{\lambda^3} = k_B T N \tag{2.15}$$

which gives us back the ideal gas law.

Figure 10: Maxwell distribution for Noble gases: He, Ne, Ar and Xe.

2.3 Maxwell Distribution

Our discussion above focusses on understanding macroscopic properties of the gas such as pressure or heat capacity. But we can also use the methods of statistical mechanics to get a better handle on the microscopic properties of the gas. Like everything else, the information is hidden in the partition function. Let's return to the form of the single particle partition function (2.4) before we do the integrals. We'll still do the trivial spatial integral $\int d^3q = V$, but we'll hold off on the momentum integral and instead change variables from momentum to velocity, $\vec{p} = m\vec{v}$. Then the single particle partition function is

$$Z_1 = \frac{m^3 V}{(2\pi\hbar)^3} \int d^3v\, e^{-\beta m\vec{v}^2/2} = \frac{4\pi m^3 V}{(2\pi\hbar)^3} \int dv\, v^2 e^{-\beta m v^2/2}$$

We can compare this to the original definition of the partition function: the sum over states of the probability of that state. But here too, the partition function is written as a sum, now over speeds. The integrand must therefore have the interpretation as the probability distribution over speeds. The probability that the atom has speed between v and $v + dv$ is

$$f(v)dv = \mathcal{N} v^2 e^{-mv^2/2k_BT}\, dv \tag{2.16}$$

where the normalization factor $\mathcal{N}$ can be determined by insisting that probabilities sum to one, $\int_0^\infty f(v)\, dv = 1$, which gives

$$\mathcal{N} = 4\pi \left(\frac{m}{2\pi k_B T}\right)^{3/2}$$

This is the *Maxwell distribution*. It is sometimes called the Maxwell-Boltzmann distribution. Figure 10 shows this distribution for a variety of gases with different masses at the same temperature, from the slow heavy Xenon (purple) to light, fast Helium (blue). We can use it to determine various average properties of the speeds of atoms in a gas. For example, the mean square speed is

$$\langle v^2 \rangle = \int_0^\infty dv\ v^2 f(v) = \frac{3k_B T}{m}$$

This is in agreement with the equipartition of energy: the average kinetic energy of the gas is $E = \frac{1}{2}m\langle v^2 \rangle = \frac{3}{2}k_B T$.

Maxwell's Argument

The above derivation tells us the distribution of velocities in a non-interacting gas of particles. Remarkably, the Maxwell distribution also holds in the presence of any interactions. In fact, Maxwell's original derivation of the distribution makes no reference to any properties of the gas. It is very slick!

Let's first think about the distribution of velocities in the x direction; we'll call this distribution $\phi(v_x)$. Rotational symmetry means that we must have the same distribution of velocities in both the y and z directions. However, rotational invariance also requires that the full distribution can't depend on the direction of the velocity; it can only depend on the speed $v = \sqrt{v_x^2 + v_y^2 + v_z^2}$. This means that we need to find functions $F(v)$ and $\phi(v_x)$ such that

$$F(v)\, dv_x dv_y dv_z = \phi(v_x)\phi(v_y)\phi(v_z)\, dv_x dv_y dv_z$$

It doesn't look as if we possibly have enough information to solve this equation for both F and ϕ. But, remarkably, there is only one solution. The only function which satisfies this equation is

$$\phi(v_x) = Ae^{-Bv_x^2}$$

for some constants A and B. Thus the distribution over speeds must be

$$F(v)\, dv_x dv_y dv_z = 4\pi v^2 F(v)\, dv = 4\pi A^3 v^2 e^{-Bv^2} dv$$

We see that the functional form of the distribution arises from rotational invariance alone. To determine the coefficient $B = m/2k_B T$ we need the more elaborate techniques of statistical mechanics that we saw above. (In fact, one can derive it just from equipartition of energy).

2.3.1 A History of Kinetic Theory

The name *kinetic theory* refers to the understanding the properties of gases through their underlying atomic constituents. The discussion given above barely scratches the surface of this important subject.

Kinetic theory traces its origin to the work of Daniel Bernoulli in 1738. He was the first to argue that the phenomenon that we call pressure is due to the constant bombardment of tiny atoms. His calculation is straightforward. Consider a cubic box with sides of length L. Suppose that an atom travelling with momentum v_x in the x direction bounces elastically off a wall so that it returns with velocity $-v_x$. The particle experiences a change in momentum is $\Delta p_x = 2mv_x$. Since the particle is trapped in a box, it will next hit the wall at a time $\Delta t = 2L/v_x$ later. This means that the force on the wall due to this atom is

$$F = \frac{\Delta p_x}{\Delta t} = \frac{mv_x^2}{L}$$

Summing over all the atoms which hit the wall, the force is

$$F = \frac{Nm\langle v_x^2\rangle}{L}$$

where $\langle v_x^2\rangle$ is the average velocity in the x-direction. Using the same argument as we gave in Maxwell's derivation above, we must have $\langle v_x^2\rangle = \langle v^2\rangle/3$. Thus $F = Nm\langle v\rangle^2/3L$ and the pressure, which is force per area, is given be

$$p = \frac{Nm\langle v^2\rangle}{3L^3} = \frac{Nm\langle v^2\rangle}{3V}$$

If this equation is compared to the ideal gas law (which, at the time, had only experimental basis) one concludes that the phenomenon of temperature must arise from the kinetic energy of the gas. Or, more precisely, one finds the equipartition result that we derived previously: $\frac{1}{2}m\langle v^2\rangle = \frac{3}{2}k_BT$.

After Bernoulli's pioneering work, kinetic theory languished. No one really knew what to do with his observation nor how to test the underlying atomic hypothesis. Over the next century, Bernouilli's result was independently rediscovered by a number of people, all of whom were ignored by the scientific community. One of the more interesting attempts was by John Waterson, a Scottish engineer and naval instructor working for the East India Company in Bombay. Waterson was considered a crackpot. His 1843 paper was rejected by the Royal Society as "nothing but nonsense" and he wrote up his results in a self-published book with the wonderfully crackpot title "Thoughts on Mental Functions".

The results of Bernouilli and Waterson finally became accepted only after they were re-rediscovered by more established scientists, most notably Rudolph Clausius who, in 1857, extended these ideas to rotating and vibrating molecules. Soon afterwards, in 1859, Maxwell gave the derivation of the distribution of velocities that we saw above. This is often cited as the first statistical law of physics. But Maxwell was able to take things further. He used kinetic theory to derive the first genuinely new prediction of the atomic hypothesis: that the viscosity of a gas is independent of its density. Maxwell himself wrote,

> *"Such a consequence of the mathematical theory is very startling and the only experiment I have met with on the subject does not seem to confirm it."*

Maxwell decided to rectify the situation. With help from his wife, he spent several years constructing an experimental apparatus in his attic which was capable of providing the first accurate measurements of viscosity of gases[6]. His surprising theoretical prediction was confirmed by his own experiment.

There are many further developments in kinetic theory which we will not cover in this course. Perhaps the most important is the Boltzmann equation. This describes the evolution of a particle's probability distribution in position and momentum space as it collides with other particles. Stationary, unchanging, solutions bring you back to the Maxwell-Boltzmann distribution, but the equation also provides a framework to go beyond the equilibrium description of a gas. You can read about this in the lecture notes on Kinetic Theory.

2.4 Diatomic Gas

> "I must now say something about these internal motions, because the greatest difficulty which the kinetic theory of gases has yet encountered belongs to this part of the subject".
>
> *James Clerk Maxwell*, 1875

Consider a molecule that consists of two atoms in a bound state. We'll construct a very simple physicist's model of this molecule: two masses attached to a spring. As well as the translational degrees of freedom, there are two further ways in which the molecule can move

[6]You can see the original apparatus down the road in the corridor of the Cavendish lab. Or, if you don't fancy the walk, you can simply click here:
http://www-outreach.phy.cam.ac.uk/camphy/museum/area1/exhibit1.htm

- Rotation: the molecule can rotate rigidly about the two axes perpendicular to the axis of symmetry, with moment of inertia I. (For now, we will neglect the rotation about the axis of symmetry. It has very low moment of inertia which will ultimately mean that it is unimportant).

- Vibration: the molecule can oscillate along the axis of symmetry

We'll work under the assumption that the rotation and vibration modes are independent. In this case, the partition function for a single molecule factorises into the product of the translation partition function $Z_{\rm trans}$ that we have already calculated (2.5) and the rotational and vibrational contributions,

$$Z_1 = Z_{\rm trans} Z_{\rm rot} Z_{\rm vib}$$

We will now deal with $Z_{\rm rot}$ and $Z_{\rm vib}$ in turn.

Rotation

The Lagrangian for the rotational degrees of freedom is[7]

$$L_{\rm rot} = \frac{1}{2} I(\dot\theta^2 + \sin^2\theta \dot\phi^2) \tag{2.17}$$

The conjugate momenta are therefore

$$p_\theta = \frac{\partial L_{\rm rot}}{\partial \dot\theta} = I\dot\theta \quad , \quad p_\phi = \frac{\partial L_{\rm rot}}{\partial \dot\phi} = I\sin^2\theta\,\dot\phi$$

from which we get the Hamiltonian for the rotating diatomic molecule,

$$H_{\rm rot} = \dot\theta p_\theta + \dot\phi p_\phi - L = \frac{p_\theta^2}{2I} + \frac{p_\phi^2}{2I\sin^2\theta} \tag{2.18}$$

The rotational contribution to the partition function is then

$$\begin{aligned} Z_{\rm rot} &= \frac{1}{(2\pi\hbar)^2}\int d\theta d\phi dp_\theta dp_\phi\, e^{-\beta H_{\rm rot}} \\ &= \frac{1}{(2\pi\hbar)^2}\sqrt{\frac{2\pi I}{\beta}}\int_0^\pi d\theta \sqrt{\frac{2\pi I\sin^2\theta}{\beta}}\int_0^{2\pi} d\phi \\ &= \frac{2Ik_BT}{\hbar^2} \end{aligned} \tag{2.19}$$

[7]See, for example, Section 3.6 of the lecture notes on Classical Dynamics

From this we can compute the average rotational energy of each molecule,

$$E_{\rm rot} = k_B T$$

If we now include the translational contribution (2.5), the partition function for a diatomic molecule that can spin and move, but can't vibrate, is given by $Z_1 = Z_{\rm trans} Z_{\rm rot} \sim (k_B T)^{5/2}$, and the partition function for a gas of these object $Z = Z_1^N/N!$, from which we compute the energy $E = \frac{5}{2} N k_B T$ and the heat capacity,

$$C_V = \frac{5}{2} k_B N$$

In fact we can derive this result simply from equipartition of energy: there are 3 translational modes and 2 rotational modes, giving a contribution of $5N \times \frac{1}{2} k_B T$ to the energy.

Vibrations

The Hamiltonian for the vibrating mode is simply a harmonic oscillator. We'll denote the displacement away from the equilibrium position by ζ. The molecule vibrates with some frequency ω which is determined by the strength of the atomic bond. The Hamiltonian is then

$$H_{\rm vib} = \frac{p_\zeta^2}{2m} + \frac{1}{2} m \omega^2 \zeta^2$$

from which we can compute the partition function

$$Z_{\rm vib} = \frac{1}{2\pi\hbar} \int d\zeta dp_\zeta e^{-\beta H_{\rm vib}} = \frac{k_B T}{\hbar\omega} \tag{2.20}$$

The average vibrational energy of each molecule is now

$$E_{\rm vib} = k_B T$$

(You may have anticipated $\frac{1}{2} k_B T$ since the harmonic oscillator has just a single degree of freedom, but equipartition works slightly differently when there is a potential energy. You will see another example on the problem sheet from which it is simple to deduce the general form).

Putting together all the ingredients, the contributions from translational motion, rotation and vibration give the heat capacity

$$C_V = \frac{7}{2} N k_B$$

This result depends on neither the moment of inertia, I, nor the stiffness of the molecular bond, ω. A molecule with large I will simply spin more slowly so that the average rotational kinetic energy is k_BT; a molecule attached by a stiff spring with high ω will vibrate with smaller amplitude so that the average vibrational energy is k_BT. This ensures that the heat capacity is constant.

Figure 11: The heat capacity of Hydrogen gas H_2. The graph was created by P. Eyland.

Great! So the heat capacity of a diatomic gas is $\frac{7}{2}Nk_BT$. Except it's not! An idealised graph of the heat capacity for H_2, the simplest diatomic gas, is shown in Figure 11. At suitably high temperatures, around $5000K$, we do see the full heat capacity that we expect. But at low temperatures, the heat capacity is that of monatomic gas. And, in the middle, it seems to rotate, but not vibrate. What's going on? Towards the end of the nineteenth century, scientists were increasingly bewildered about this behaviour.

What's missing in the discussion above is something *very* important: $\hbar$. The successive freezing out of vibrational and rotational modes as the temperature is lowered is a quantum effect. In fact, this behaviour of the heat capacities of gases was the first time that quantum mechanics revealed itself in experiment. We're used to thinking of quantum mechanics as being relevant on small scales, yet here we see that affects the physics of gases at temperatures of $2000\,K$. But then, that is the theme of this course: how the microscopic determines the macroscopic. We will return to the diatomic gas in Section 3.4 and understand its heat capacity including the relevant quantum effects.

2.5 Interacting Gas

Until now, we've only discussed free systems; particles moving around unaware of each other. Now we're going to turn on interactions. Here things get much more interesting.

And much more difficult. Many of the most important unsolved problems in physics are to do with the interactions between large number of particles. Here we'll be gentle. We'll describe a simple approximation scheme that will allow us to begin to understand the effects of interactions between particles.

We'll focus once more on the monatomic gas. The ideal gas law is exact in the limit of no interactions between atoms. This is a good approximation when the density of atoms N/V is small. Corrections to the ideal gas law are often expressed in terms of a density expansion, known as the *virial expansion*. The most general equation of state is,

$$\frac{p}{k_B T} = \frac{N}{V} + B_2(T)\frac{N^2}{V^2} + B_3(T)\frac{N^3}{V^3} + \ldots \tag{2.21}$$

where the functions $B_j(T)$ are known as virial coefficients.

Our goal is to compute the virial coefficients from first principles, starting from a knowledge of the underlying potential energy $U(r)$ between two neutral atoms separated by a distance r. This potential has two important features:

- An attractive $1/r^6$ force. This arises from fluctuating dipoles of the neutral atoms. Recall that two permanent dipole moments, p_1 and p_2, have a potential energy which scales as $p_1 p_2/r^3$. Neutral atoms don't have permanent dipoles, but they can acquire a temporary dipole due to quantum fluctuations. Suppose that the first atom has an instantaneous dipole p_1. This will induce an electric field which is proportional to $E \sim p_1/r^3$ which, in turn, will induce a dipole of the second atom $p_2 \sim E \sim p_1/r^3$. The resulting potential energy between the atoms scales as $p_1 p_2/r^3 \sim 1/r^6$. This is sometimes called the *van der Waals* interaction.

- A rapidly rising repulsive interaction at short distances, arising from the Pauli exclusion principle that prevents two atoms from occupying the same space. For our purposes, the exact form of this repulsion is not so relevant: just as long as it's big. (The Pauli exclusion principle is a quantum effect. If the exact form of the potential is important then we really need to be dealing with quantum mechanics all along. We will do this in the next section).

One very common potential that is often used to model the force between atoms is the *Lennard-Jones potential*,

$$U(r) \sim \left(\frac{r_0}{r}\right)^{12} - \left(\frac{r_0}{r}\right)^{6} \tag{2.22}$$

The exponent 12 is chosen only for convenience: it simplifies certain calculations because $12 = 2 \times 6$.

An even simpler form of the potential incorporates a *hard core repulsion*, in which the particles are simply forbidden from closer than a fixed distance by imposing an infinite potential,

$$U(r) = \begin{cases} \infty & r < r_0 \\ -U_0 \left(\frac{r_0}{r}\right)^6 & r \geq r_0 \end{cases} \tag{2.23}$$

Figure 12:

The hard-core potential with van der Waals attraction is sketched to the right. We will see shortly that the virial coefficients are determined by increasingly difficult integrals involving the potential $U(r)$. For this reason, it's best to work with a potential that's as simple as possible. When we come to do some actual calculations we will use the form (2.23).

2.5.1 The Mayer f Function and the Second Virial Coefficient

We're going to change notation and call the positions of the particles $\vec{r}$ instead of $\vec{q}$. (The latter notation was useful to stress the connection to quantum mechanics at the beginning of this Section, but we've now left that behind!). The Hamiltonian of the gas is

$$H = \sum_{i=1}^{N} \frac{p_i^2}{2m} + \sum_{i>j} U(r_{ij})$$

where $r_{ij} = |\vec{r}_i - \vec{r}_j|$ is the separation between particles. The restriction $i > j$ on the final sum ensures that we sum over each pair of particles exactly once. The partition function is then

$$\begin{aligned} Z(N,V,T) &= \frac{1}{N!}\frac{1}{(2\pi\hbar)^{3N}} \int \prod_{i=1}^{N} d^3p_i d^3r_i \; e^{-\beta H} \\ &= \frac{1}{N!}\frac{1}{(2\pi\hbar)^{3N}} \left[\int \prod_i d^3p_i \; e^{-\beta \sum_j p_j^2/2m}\right] \times \left[\int \prod_i d^3r_i \; e^{-\beta \sum_{j<k} U(r_{jk})}\right] \\ &= \frac{1}{N!\lambda^{3N}} \int \prod_i d^3r_i \; e^{-\beta \sum_{j<k} U(r_{jk})} \end{aligned}$$

where λ is the thermal wavelength that we met in (2.6). We still need to do the integral over positions. And that looks hard! The interactions mean that the integrals don't

factor in any obvious way. What to do? One obvious way thing to try is to Taylor expand (which is closely related to the so-called *cumulant expansion* in this context)

$$e^{-\beta\sum_{j<k}U(r_{jk})} = 1 - \beta\sum_{j<k}U(r_{jk}) + \frac{\beta^2}{2}\sum_{j<k,l<m}U(r_{jk})U(r_{lm}) + \ldots$$

Unfortunately, this isn't so useful. We want each term to be smaller than the preceding one. But as $r_{ij} \to 0$, the potential $U(r_{ij}) \to \infty$, which doesn't look promising for an expansion parameter.

Instead of proceeding with the naive Taylor expansion, we will instead choose to work with the following quantity, usually called the *Mayer f function*,

$$f(r) = e^{-\beta U(r)} - 1 \tag{2.24}$$

This is a nicer expansion parameter. When the particles are far separated at $r \to \infty$, $f(r) \to 0$. However, as the particles come close and $r \to 0$, the Mayer function approaches $f(r) \to -1$. We'll proceed by trying to construct a suitable expansion in terms of f. We define

$$f_{ij} = f(r_{ij})$$

Then we can write the partition function as

$$\begin{aligned} Z(N,V,T) &= \frac{1}{N!\lambda^{3N}}\int\prod_i d^3r_i\prod_{j>k}(1+f_{jk}) \\ &= \frac{1}{N!\lambda^{3N}}\int\prod_i d^3r_i\left(1+\sum_{j>k}f_{jk}+\sum_{j>k,l>m}f_{jk}f_{lm}+\ldots\right) \end{aligned} \tag{2.25}$$

The first term simply gives a factor of the volume V for each integral, so we get V^N. The second term has a sum, each element of which is the same. They all look like

$$\int\prod_{i=1}^N d^3r_i\ f_{12} = V^{N-2}\int d^3r_1 d^3r_2\ f(r_{12}) = V^{N-1}\int d^3r\ f(r)$$

where, in the last equality, we've simply changed integration variables from $\vec{r}_1$ and $\vec{r}_2$ to the centre of mass $\vec{R} = \frac{1}{2}(\vec{r}_1 + \vec{r}_2)$ and the separation $\vec{r} = \vec{r}_1 - \vec{r}_2$. (You might worry that the limits of integration change in the integral over $\vec{r}$, but the integral over $f(r)$ only picks up contributions from atomic size distances and this is only actually a problem close to the boundaries of the system where it is negligible). There is a term

like this for each pair of particles – that is $\frac{1}{2}N(N-1)$ such terms. For $N \sim 10^{23}$, we can just call this a round $\frac{1}{2}N^2$. Then, ignoring terms quadratic in f and higher, the partition function is approximately

$$\begin{aligned} Z(N,V,T) &= \frac{V^N}{N!\lambda^{3N}}\left(1+\frac{N^2}{2V}\int d^3r\ f(r)+\ldots\right) \\ &= Z_{\text{ideal}}\left(1+\frac{N}{2V}\int d^3r\ f(r)+\ldots\right)^N \end{aligned}$$

where we've used our previous result that $Z_{\text{ideal}} = V^N/N!\lambda^{3N}$. We've also engaged in something of a sleight of hand in this last line, promoting one power of N from in front of the integral to an overall exponent. Massaging the expression in this way ensures that the free energy is proportional to the number of particles as one would expect:

$$F = -k_BT\log Z = F_{\text{ideal}} - Nk_BT\log\left(1+\frac{N}{2V}\int d^3r\ f(r)\right) \tag{2.26}$$

However, if you're uncomfortable with this little trick, it's not hard to convince yourself that the result (2.27) below for the equation of state doesn't depend on it. We will also look at the expansion more closely in the following section and see how all the higher order terms work out.

From the expression (2.26) for the free energy, it is clear that we are indeed performing an expansion in density of the gas since the correction term is proportional to N/V. This form of the free energy will give us the second virial coefficient $B_2(T)$.

We can be somewhat more precise about what it means to be at low density. The exact form of the integral $\int d^3rf(r)$ depends on the potential, but for both the Lennard-Jones potential (2.22) and the hard-core repulsion (2.23), the integral is approximately $\int d^3rf(r) \sim r_0^3$, where r_0 is roughly the minimum of the potential. (We'll compute the integral exactly below for the hard-core potential). For the expansion to be valid, we want each term with an extra power of f to be smaller than the preceding one. (This statement is actually only approximately true. We'll be more precise below when we develop the cluster expansion). That means that the second term in the argument of the log should be smaller than 1. In other words,

$$\frac{N}{V} \ll \frac{1}{r_0^3}$$

The left-hand side is the density of the gas. The right-hand side is atomic density. Or, equivalently, the density of a substance in which the atoms are packed closely together. But we have a name for such substances – we call them liquids! Our expansion is valid for densities of the gas that are much lower than that of the liquid state.

2.5.2 van der Waals Equation of State

We can use the free energy (2.26) to compute the pressure of the gas. Expanding the logarithm as $\log(1+x) \approx x$ we get

$$p = -\frac{\partial F}{\partial V} = \frac{Nk_BT}{V}\left(1 - \frac{N}{2V}\int d^3r f(r) + \ldots\right)$$

As expected, the pressure deviates from that of an ideal gas. We can characterize this by writing

$$\frac{pV}{Nk_BT} = 1 - \frac{N}{2V}\int d^3r\ f(r) \tag{2.27}$$

To understand what this is telling us, we need to compute $\int d^3r f(r)$. Firstly let's look at two trivial examples:

Repulsion: Suppose that $U(r) > 0$ for all separations r with $U(r=\infty) = 0$. Then $f = e^{-\beta U} - 1 < 0$ and the pressure increases, as we'd expect for a repulsive interaction.

Attraction: If $U(r) < 0$, we have $f > 0$ and the pressure decreases, as we'd expect for an attractive interaction.

What about a more realistic interaction that is attractive at long distances and repulsive at short? We will compute the equation of state of a gas using the hard-core potential with van der Waals attraction (2.23). The integral of the Mayer f function is

$$\int d^3r\ f(r) = \int_0^{r_0} d^3r(-1) + \int_{r_0}^{\infty} d^3r\ (e^{+\beta U_0(r_0/r)^6} - 1) \tag{2.28}$$

We'll approximate the second integral in the high temperature limit, $\beta U_0 \ll 1$, where $e^{+\beta U_0(r_0/r)^6} \approx 1 + \beta U_0(r_0/r)^6$. Then

$$\begin{aligned}\int d^3r\ f(r) &= -4\pi\int_0^{r_0} dr\, r^2 + \frac{4\pi U_0}{k_BT}\int_{r_0}^{\infty} dr\ \frac{r_0^6}{r^4} \\ &= \frac{4\pi r_0^3}{3}\left(\frac{U_0}{k_BT} - 1\right)\end{aligned} \tag{2.29}$$

Inserting this into (2.27) gives us an expression for the equation of state,

$$\frac{pV}{Nk_BT} = 1 - \frac{N}{V}\left(\frac{a}{k_BT} - b\right)$$

We recognise this expansion as capturing the second virial coefficient in (2.21) as promised. The constants a and b are defined by

$$a = \frac{2\pi r_0^3 U_0}{3} \quad , \quad b = \frac{2\pi r_0^3}{3}$$

It is actually slightly more useful to write this in the form $k_B T = \ldots$. We can multiply through by $k_B T$ then, rearranging we have

$$k_B T = \frac{V}{N}\left(p + \frac{N^2}{V^2}a\right)\left(1 + \frac{N}{V}b\right)^{-1}$$

Since we're working in an expansion in density, N/V, we're at liberty to Taylor expand the last bracket, keeping only the first two terms. We get

$$k_B T = \left(p + \frac{N^2}{V^2}a\right)\left(\frac{V}{N} - b\right) \tag{2.30}$$

This is the famous *van der Waals equation of state* for a gas. We stress again the limitations of our analysis: it is valid only at low densities and (because of our approximation when performing the integral (2.28)) at high temperatures.

We will return to the van der Waals equation in Section 5 where we'll explore many of its interesting features. For now, we can get a feeling for the physics behind this equation of state by rewriting it in yet another way,

$$p = \frac{N k_B T}{V - bN} - a\frac{N^2}{V^2} \tag{2.31}$$

The constant a contains a factor of U_0 and so capures the effect of the attractive interaction at large distances. We see that its role is to reduce the pressure of the gas. The reduction in pressure is proportional to the density squared because this is, in turn, proportional to the number of *pairs* of particles which feel the attractive force. In contrast, b only contains r_0 and arises due to the hard-core repulsion in the potential. Its effect is the reduce the effective volume of the gas because of the space taken up by the particles.

Figure 13:

It is worth pointing out where some quizzical factors of two come from in $b = 2\pi r_0^3/3$. Recall that r_0 is the minimum distance that two atoms can approach. If we think of the each atom as a hard sphere, then they have radius $r_0/2$ and volume $4\pi(r_0/2)^3/3$. Which isn't equal to b. However, as illustrated in the figure, the excluded volume around each

atom is actually $\Omega = 4\pi r_0^3/3 = 2b$. So why don't we have Ω sitting in the denominator of the van der Waals equation rather than $b = \Omega/2$? Think about adding the atoms one at a time. The first guy can move in volume V; the second in volume $V - \Omega$; the third in volume $V - 2\Omega$ and so on. For $\Omega \ll V$, the total configuration space available to the atoms is

$$\frac{1}{N!}\prod_{m=1}^{N}(V - m\Omega) \approx \frac{V^N}{N!}\left(1 - \frac{N^2}{2}\frac{\Omega}{V} + \ldots\right) \approx \frac{1}{N!}\left(V - \frac{N\Omega}{2}\right)^N$$

And there's that tricky factor of 1/2.

Above we computed the equation of state for the dipole van der Waals interaction with hard core potential. But our expression (2.27) can seemingly be used to compute the equation of state for any potential between atoms. However, there are limitations. Looking back to the integral (2.29), we see that a long-range force of the form $1/r^n$ will only give rise to a convergent integral for $n \geq 4$. This means that the techniques described above do not work for *long-range* potentials with fall-off $1/r^3$ or slower. This includes the important case of $1/r$ Coulomb interactions.

2.5.3 The Cluster Expansion

Above we computed the leading order correction to the ideal gas law. In terms of the virial expansion (2.21) this corresponds to the second virial coefficient B_2. We will now develop the full expansion and explain how to compute the higher virial coefficients.

Let's go back to equation (2.25) where we first expressed the partition function in terms of f,

$$\begin{aligned} Z(N,V,T) &= \frac{1}{N!\lambda^{3N}}\int\prod_i d^3r_i \prod_{j>k}(1+f_{jk}) \\ &= \frac{1}{N!\lambda^{3N}}\int\prod_i d^3r_i\left(1+\sum_{j>k}f_{jk}+\sum_{j>k,l>m}f_{jk}f_{lm}+\ldots\right) \end{aligned} \tag{2.32}$$

Above we effectively related the second virial coefficient to the term linear in f: this is the essence of the equation of state (2.27). One might think that terms quadratic in f give rise to the third virial coefficient and so on. But, as we'll now see, the expansion is somewhat more subtle than that.

The expansion in (2.32) includes terms of the form $f_{ij}f_{kl}f_{mn}\ldots$ where the indices denote pairs of atoms, (i,j) and (k,l) and so on. These pairs may have atoms in common or they may all be different. However, the same pair never appears twice in a given term as you may check by going back to the first line in (2.32). We'll introduce a diagrammatic method to keep track of all the terms in the sum. To each term of the form $f_{ij}f_{kl}f_{mn}\ldots$ we associate a picture using the following rules

- Draw N atoms. (This gets tedious for $N \sim 10^{23}$ but, as we'll soon see, we will actually only need pictures with small subset of atoms).

- Draw a line between each pair of atoms that appear as indices. So for $f_{ij}f_{kl}f_{mn}\ldots$, we draw a line between atom i and atom j; a line between atom k and atom l; and so on.

For example, if we have just $N = 4$, we have the following pictures for different terms in the expansion,

$f_{12} =$

$f_{12}f_{34} =$

$f_{12}f_{23} =$

$f_{21}f_{23}f_{31} =$

We call these diagrams *graphs*. Each possible graph appears exactly once in the partition function (2.32). In other words, the partition function is a sum over all graphs. We still have to do the integrals over all positions $\vec{r}_i$. We will denote the integral over graph G to be $W[G]$. Then the partition function is

$$Z(N,V,T) = \frac{1}{N!\lambda^{3N}} \sum_G W[G]$$

Nearly all the graphs that we can draw will have disconnected components. For example, those graphs that correspond to just a single f_{ij} will have two atoms connected and the remaining $N-2$ sitting alone. Those graphs that correspond to $f_{ij}f_{kl}$ fall into two categories: either they consist of two pairs of atoms (like the second example above) or, if (i,j) shares an atom with (k,l), there are three linked atoms (like the third example above). Importantly, the integral over positions $\vec{r}_i$ then factorises into a product of integrals over the positions of atoms in disconnected components. This is illustrated by an example with $N = 5$ atoms,

$W\Bigg[$

$\Bigg] = \left(\int d^3r_1 d^3r_2 d^3r_3 f_{12}f_{23}f_{31}\right)\left(\int d^3r_4 d^3r_5 f_{45}\right)$

We call the disconnected components of the graph *clusters*. If a cluster has l atoms, we will call it an l-cluster. The $N = 5$ example above has a single 3-cluster and a single 2-cluster. In general, a graph G will split into m_l l-clusters. Clearly, we must have

$$\sum_{l=1}^{N} m_l l = N \tag{2.33}$$

Of course, for a graph with only a few lines and lots of atoms, nearly all the atoms will be in lonely 1-clusters.

We can now make good on the promise above that we won't have to draw all $N \sim 10^{23}$ atoms. The key idea is that we can focus on clusters of l-atoms. We will organise the expansion in such a way that the $(l+1)$-clusters are less important than the l-clusters. To see how this works, let's focus on 3-clusters for now. There are four different ways that we can have a 3-cluster,

Each of these 3-clusters will appear in a graph with any other combination of clusters among the remaining $N-3$ atoms. But since clusters factorise in the partition function, we know that Z must include a factor

$$U_3 \equiv \int d^3r_1 d^3r_2 d^3r_3 \Bigg(\quad + \quad + \quad + \quad \Bigg)$$

U_3 contains terms of order f^2 and f^3. It turns out that this is the correct way to arrange the expansion: not in terms of the number of lines in the diagram, which is equal to the power of f, but instead in terms of the number of atoms that they connect. The partition function will similarly contain factors associated to all other l-clusters. We define the corresponding integrals as

$$U_l \equiv \int \prod_{i=1}^{l} d^3r_i \sum_{G \in \{l\text{-cluster}\}} G \tag{2.34}$$

Notice that U_1 is simply the integral over space, namely $U_1 = V$. The full partition function must be a product of U_l's. The tricky part is to get all the combinatoric factors right to make sure that you count each graph exactly once. The sum over graphs G that appears in the partition function turns out to be

$$\sum_G W[G] = N! \sum_{\{m_l\}} \prod_l \frac{U_l^{m_l}}{(l!)^{m_l} m_l!} \tag{2.35}$$

The product $N!/\prod_l m_l!(l!)^{m_l}$ counts the number of ways to split the particles into m_l l-clusters, while ignoring the different ways to internally connect each cluster. This is the right thing to do since the different internal connections are taken into account in the integral U_l.

Combinatoric arguments are not always transparent. Let's do a couple of checks to make sure that this is indeed the right answer. Firstly, consider $N = 4$ atoms split into two 2-clusters (i.e $m_2 = 2$). There are three such diagrams, $f_{12}f_{34} =$, $f_{13}f_{24} =$, and $f_{14}f_{23} =$. Each of these gives the same answer when integrated, namely U_2^2 so the final result should be $3U_2^2$. We can check this against the relevant terms in (2.35) which are $4!U_2^2/2!^2 2! = 3U_2^2$ as expected.

Another check: $N = 5$ atoms with $m_2 = m_3 = 1$. All diagrams come in the combinations

$$U_3U_2 = \int \prod_{i=1}^{5} d^3r_i \Bigg($$ + + + $\Bigg)$

together with graphs that are related by permutations. The permutations are fully determined by the choice of the two atoms that sit in the pair: there are 10 such choices. The answer should therefore be $10U_3U_2$. Comparing to (2.35), we have $5!U_3U_2/3!2! = 10U_3U_2$ as required.

Hopefully you are now convinced that (2.35) counts the graphs correctly. The end result for the partition function is therefore

$$Z(N,V,T) = \frac{1}{\lambda^{3N}} \sum_{\{m_l\}} \prod_l \frac{U_l^{m_l}}{(l!)^{m_l} m_l!}$$

The problem with computing this sum is that we still have to work out the different ways that we can split N atoms into different clusters. In other words, we still have to obey the constraint (2.33). Life would be very much easier if we didn't have to worry about this. Then we could just sum over any m_l, regardless. Thankfully, this is exactly what we can do if we work in the grand canonical ensemble where N is not fixed! The grand canonical ensemble is

$$\mathcal{Z}(\mu,V,T) = \sum_N e^{\beta\mu N} Z(N,V,T)$$

We define the *fugacity* as $z = e^{\beta\mu}$. Then we can write

$$\mathcal{Z}(\mu,V,T) = \sum_N z^n Z(N,V,T) = \sum_{m_l=0}^{\infty} \prod_{l=1}^{\infty} \left(\frac{z}{\lambda^3}\right)^{m_l l} \frac{1}{m_l!} \left(\frac{U_l}{l!}\right)^{m_l} = \prod_{l=1}^{\infty} \exp\left(\frac{U_l z^l}{\lambda^{3l} l!}\right)$$

One usually defines

$$b_l = \frac{\lambda^3}{V}\frac{U_l}{l!\lambda^{3l}} \tag{2.36}$$

Notice in particular that $U_1 = V$ so this definition gives $b_1 = 1$. Then we can write the grand partition function as

$$\mathcal{Z}(\mu, V, T) = \prod_{l=1}^{\infty} \exp\left(\frac{V}{\lambda^3} b_l z^l\right) = \exp\left(\frac{V}{\lambda^3}\sum_{l=1}^{\infty} b_l z^l\right) \tag{2.37}$$

Something rather cute happened here. The sum over all diagrams got rewritten as the exponential over the sum of all *connected* diagrams, meaning all clusters. This is a general lesson which also carries over to quantum field theory where the diagrams in question are Feynman diagrams.

Back to the main plot of our story, we can now compute the pressure

$$\frac{pV}{k_B T} = \log \mathcal{Z} = \frac{V}{\lambda^3}\sum_{l=1}^{\infty} b_l z^l$$

and the number of particles

$$\frac{N}{V} = \frac{z}{V}\frac{\partial}{\partial z}\log \mathcal{Z} = \frac{1}{\lambda^3}\sum_{l=1}^{\infty} l b_l z^l \tag{2.38}$$

Dividing the two gives us the equation of state,

$$\frac{pV}{N k_B T} = \frac{\sum_l b_l z^l}{\sum_l l b_l z^l} \tag{2.39}$$

The only downside is that the equation of state is expressed in terms of z. To massage it into the form of the virial expansion (2.21), we need to invert (2.38) to get z in terms of the particle density N/V. Equating (2.39) with (2.21) (and defining $B_1 = 1$), we have

$$\begin{aligned}
\sum_{l=1}^{\infty} b_l z^l &= \sum_{l=1}^{\infty} B_l \left(\frac{N}{V}\right)^{l-1} \sum_{m=1}^{\infty} m b_m z^m \\
&= \sum_{l=1}^{\infty} \frac{B_l}{\lambda^{3(l-1)}} \left(\sum_{n=1}^{\infty} n b_n z^n\right)^{l-1} \sum_{m=1}^{\infty} m b_m z^m \\
&= \left[1 + \frac{B_2}{\lambda^3}(z + 2b_2 z^2 + 3b_3 z^3 + \ldots) + \frac{B_3}{\lambda^6}(z + 2b_2 z^2 + 3b_3 z^3 + \ldots)^2 + \ldots\right] \\
&\qquad \times \left[z + 2b_2 z^2 + 3b_3 z^3 + \ldots\right]
\end{aligned}$$

where we've used both $B_1 = 1$ and $b_1 = 1$. Expanding out the left- and right-hand sides to order z^3 gives

$$z + b_2 z^2 + b_3 z^3 + \ldots = z + \left(\frac{B_2}{\lambda^3} + 2b_2\right) z^2 + \left(3b_3 + \frac{4b_2 B_2}{\lambda^3} + \frac{B_3}{\lambda^3}\right) z^3 + \ldots$$

Comparing terms, and recollecting the definitions of b_l (2.36) in terms of U_l (2.34) in terms of graphs, we find the second virial coefficient is given by

$$B_2 = -\lambda^3 b_2 = -\frac{U_2}{2V} = -\frac{1}{2V}\int d^3r_1 d^3r_2 f(\vec{r}_1 - \vec{r}_2) = -\frac{1}{2}\int d^3r f(r)$$

which reproduces the result (2.27) that we found earlier using slightly simpler methods. We now also have an expression for the third coefficient,

$$B_3 = \lambda^6(4b_2^2 - 2b_3)$$

although admittedly we still have a nasty integral to do before we have a concrete result. More importantly, the cluster expansion gives us the technology to perform a systematic perturbation expansion to any order we wish.

2.6 Screening and the Debye-Hückel Model of a Plasma

There are many other applications of the classical statistical methods that we saw in this chapter. Here we use them to derive the important phenomenon of *screening*. The problem we will consider, which sometimes goes by the name of a "one-component plasma", is the following: a gas of electrons, each with charge $-q$, moves in a fixed background of uniform positive charge density $+q\rho$. The charge density is such that the overall system is neutral which means that ρ is also the average charge density of the electrons. This is the Debye-Hückel model.

In the absence of the background charge density, the interaction between electons is given by the Coulomb potential

$$U(r) = \frac{q^2}{r}$$

where we're using units in which $4\pi\epsilon_0 = 1$. How does the fixed background charge affect the potential between electrons? The clever trick of the Debye-Hückel model is to use statistical methods to figure out the answer to this question. Consider placing one electron at the origin. Let's try to work out the electrostatic potential $\phi(\vec{r})$ due to this electron. It is not obvious how to do this because ϕ will also depend on the positions of all the other electrons. In general we can write,

$$\nabla^2\phi(\vec{r}) = -4\pi\left(-q\delta(\vec{r}) + q\rho - q\rho g(\vec{r})\right) \tag{2.40}$$

where the first term on the right-hand side is due to the electron at the origin; the second term is due to the background positive charge density; and the third term is due to the other electrons whose average charge density close to the first electron is $\rho g(\vec{r})$. The trouble is that we don't know the function g. If we were sitting at zero temperature, the electrons would try to move apart as much as possible. But at non-zero temperatures, their thermal energy will allow them to approach each other. This is the clue that we need. The energy cost for an electron to approach the origin is, of course, $E(\vec{r}) = -q\phi(\vec{r})$. We will therefore assume that the charge density near the origin is given by the Boltzmann factor,

$$g(\vec{r}) \approx e^{\beta q \phi(\vec{r})}$$

For high temperatures, $\beta q \phi \ll 1$, we can write $e^{\beta q \phi} \approx 1 + \beta q \phi$ and the Poisson equation (2.40) becomes

$$\left(\nabla^2 + \frac{1}{\lambda_D^2}\right) \phi(\vec{r}) = 4\pi q \delta(\vec{r})$$

where $\lambda_D^2 = 1/4\pi\beta\rho q^2$. This equation has the solution,

$$\phi(\vec{r}) = -\frac{q e^{-r/\lambda_D}}{r} \tag{2.41}$$

which immediately translates into an effective potential energy between electrons,

$$U_{\text{eff}}(r) = \frac{q^2 e^{-r/\lambda_D}}{r}$$

We now see that the effect of the plasma is to introduce the exponential factor in the numerator, causing the potential to decay very quickly at distances $r > \lambda_D$. This effect is called *screening* and λ_D is known as the *Debye screening length*. The derivation of (2.41) is self-consistent if we have a large number of electrons within a distance λ_D of the origin so that we can happily talk about average charge density. This means that we need $\rho \lambda_D^3 \gg 1$.

3. Quantum Gases

In this section we will discuss situations where quantum effects are important. We'll still restrict attention to gases — meaning a bunch of particles moving around and barely interacting — but one of the first things we'll see is how versatile the idea of a gas can be in the quantum world. We'll use it to understand not just the traditional gases that we met in the previous section but also light and, ironically, certain properties of solids. In the latter part of this section, we will look at what happens to gases at low temperatures where their behaviour is dominated by quantum statistics.

3.1 Density of States

We start by introducing the important concept of the density of states. To illustrate this, we'll return once again to the ideal gas trapped in a box with sides of length L and volume $V = L^3$. Viewed quantum mechanically, each particle is described by a wavefunction. We'll impose periodic boundary conditions on this wavefunction (although none of the physics that we'll discuss in this course will be sensitive to the choice of boundary condition). If there are no interactions between particles, the energy eigenstates are simply plane waves,

$$\psi = \frac{1}{\sqrt{V}}\, e^{i\vec{k}\cdot\vec{x}}$$

Boundary conditions require that the wavevector $\vec{k} = (k_1, k_2, k_3)$ is quantized as

$$k_i = \frac{2\pi n_i}{L} \quad \text{with} \quad n_i \in \mathbf{Z}$$

and the energy of the particle is

$$E_{\vec{n}} = \frac{\hbar^2 k^2}{2m} = \frac{4\pi^2\hbar^2}{2mL^2}\,(n_1^2 + n_2^2 + n_3^2)$$

with $k = |\vec{k}|$. The quantum mechanical single particle partition function (1.21) is given by the sum over all energy eigenstates,

$$Z_1 = \sum_{\vec{n}} e^{-\beta E_{\vec{n}}}$$

The question is: how do we do the sum? The simplest way is to approximate it by an integral. Recall from the previous section that the thermal wavelength of the particle is defined to be

$$\lambda = \sqrt{\frac{2\pi\hbar^2}{mk_BT}}$$

The exponents that appear in the sum are all of the form $\sim \lambda^2 n^2/L^2$, up to some constant factors. For any macroscopic size box, $\lambda \ll L$ (a serious understatement! Actually $\lambda \lll L$) which ensures that there are many states with $E_{\vec{n}} \leq k_B T$ all of which contribute to the sum. (There will be an exception to this at very low temperatures which will be the focus of Section 3.5.3). We therefore lose very little by approximating the sum by an integral. We can write the measure of this integral as

$$\sum_{\vec{n}} \approx \int d^3n \;=\; \frac{V}{(2\pi)^3}\int d^3k = \frac{4\pi V}{(2\pi)^3}\int_0^\infty dk\; k^2$$

where, in the last equality, we have integrated over the angular directions to get 4π, the area of the 2-sphere, leaving an integration over the magnitude $k = |\vec{k}|$ and the Jacobian factor k^2. For future applications, it will prove more useful to change integration variables at this stage. We work instead with the energy,

$$E = \frac{\hbar^2 k^2}{2m} \quad \Rightarrow \quad dE = \frac{\hbar^2 k}{m}\, dk$$

We can now write out integral as

$$\frac{4\pi V}{(2\pi)^3}\int dk\; k^2 \;=\; \frac{V}{2\pi^2}\int dE\; \sqrt{\frac{2mE}{\hbar^2}}\frac{m}{\hbar^2} \;\equiv\; \int dE\; g(E) \tag{3.1}$$

where

$$g(E) = \frac{V}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2} E^{1/2} \tag{3.2}$$

is the *density of states*: $g(E)dE$ counts the number of states with energy between E and $E + dE$. Notice that we haven't actually done the integral over E in (3.1); instead this is to be viewed as a measure which we can integrate over any function $f(E)$ of our choosing.

There is nothing particularly quantum mechanical about the density of states. Indeed, in the derivation above we have replaced the quantum sum with an integral over momenta which actually looks rather classical. Nonetheless, as we encounter more and more different types of gases, we'll see that the density of states appears in all the calculations and it is a useful quantity to have at our disposal.

3.1.1 Relativistic Systems

Relativistic particles moving in $d = 3 + 1$ spacetime dimensions have kinetic energy

$$E = \sqrt{\hbar^2 k^2 c^2 + m^2 c^4} \tag{3.3}$$

Repeating the steps above, we find the density of states is given by

$$g(E) = \frac{VE}{2\pi^2\hbar^3c^3}\sqrt{E^2 - m^2c^4} \tag{3.4}$$

In particular, for massless particles, the density of states is

$$g(E) = \frac{VE^2}{2\pi^2\hbar^3c^3} \tag{3.5}$$

3.2 Photons: Blackbody Radiation

> "It was an act of desperation. For six years I had struggled with the blackbody theory. I knew the problem was fundamental and I knew the answer. I had to find a theoretical explanation at any cost, except for the inviolability of the two laws of thermodynamics"
>
> *Max Planck*

We now turn to our first truly quantum gas: light. We will consider a gas of photons — the quanta of the electromagnetic field — and determine a number of its properties, including the distribution of wavelengths. Or, in other words, its colour.

Below we will describe the colour of light at a fixed temperature. But this also applies (with a caveat) to the colour of any object at the same temperature. The argument for this is as follows: consider bathing the object inside the gas of photons. In equilibrium, the object sits at the same temperature as the photons, emitting as many photons as it absorbs. The colour of the object will therefore mimic that of the surrounding light.

For a topic that's all about colour, a gas of photons is usually given a rather bland name — *blackbody radiation*. The reason for this is that any real object will exhibit absorption and emission lines due to its particular atomic make-up (this is the caveat mentioned above). We're not interested in these details; we only wish to compute the spectrum of photons that a body emits because it's hot. For this reason, one sometimes talks about an idealised body that absorbs photons of any wavelength and reflects none. At zero temperature, such an object would appear black: this is the blackbody of the title. We would like to understand its colour as we turn up the heat.

To begin, we need some facts about photons. The energy of a photon is determined by its wavelength λ or, equivalently, by its frequency $\omega = 2\pi c/\lambda$ to be

$$E = \hbar\omega$$

This is a special case of the relativistic energy formula (3.3) for massless particles, $m = 0$. The frequency is related to the (magnitude of the) wavevector by $\omega = kc$.

Photons have two polarization states (one for each dimension transverse to the direction of propagation). To account for this, the density of states (3.5) should be multiplied by a factor of two. The number of states available to a single photon with energy between E and $E + dE$ is therefore

$$g(E)dE = \frac{VE^2}{\pi^2\hbar^3c^3}dE$$

Equivalently, the number of states available to a single photon with frequency between ω and $\omega + d\omega$ is

$$g(E)dE = g(\omega)d\omega = \frac{V\omega^2}{\pi^2c^3}d\omega \tag{3.6}$$

where we've indulged in a slight abuse of notation since $g(\omega)$ is not the same function as $g(E)$ but is instead defined by the equation above. It is also worth pointing out an easy mistake to make when performing these kinds of manipulations with densities of states: you need to remember to rescale the interval dE to $d\omega$. This is most simply achieved by writing $g(E)dE = g(\omega)d\omega$ as we have above. If you miss this then you'll get $g(\omega)$ wrong by a factor of $\hbar$.

The final fact that we need is important: photons are not conserved. If you put six atoms in a box then they will still be there when you come back a month later. This isn't true for photons. There's no reason that the walls of the box can't absorb one photon and then emit two. The number of photons in the world is not fixed. To demonstrate this, you simply need to turn off the light.

Because photon number is not conserved, we're unable to define a chemical potential for photons. Indeed, even in the canonical ensemble we must already sum over states with different numbers of photons because these are all "accessible states". (It is sometimes stated that we should work in the grand canonical ensemble at $\mu = 0$ which is basically the same thing). This means that we should consider states with any number N of photons.

We'll start by looking at photons with a definite frequency ω. A state with N such photons has energy $E = N\hbar\omega$. Summing over all N gives us the partition function for photons at fixed frequency,

$$Z_\omega = 1 + e^{-\beta\hbar\omega} + e^{-2\beta\hbar\omega} + \ldots = \frac{1}{1 - e^{-\beta\hbar\omega}} \tag{3.7}$$

We now need to sum over all possible frequencies. As we've seen a number of times, independent partition functions multiply, which means that the logs add. We only need

Figure 14: The Planck Distribution function (Source: E. Schubert, *Light Emitting Diodes*).

to know how many photon states there are with some frequency ω. But this is what the density of states (3.6) tells us. We have

$$\log Z = \int_0^\infty d\omega \; g(w) \; \log Z_\omega = -\frac{V}{\pi^2 c^3} \int_0^\infty d\omega \; \omega^2 \log\left(1 - e^{-\beta\hbar\omega}\right) \tag{3.8}$$

3.2.1 Planck Distribution

From the partition function (3.8) we can calculate all interesting quantities for a gas of light. For example, the energy density stored in the photon gas is

$$E = -\frac{\partial}{\partial\beta} \log Z = \frac{V\hbar}{\pi^2 c^3} \int_0^\infty d\omega \; \frac{\omega^3}{e^{\beta\hbar\omega} - 1} \tag{3.9}$$

However, before we do the integral over frequency, there's some important information contained in the integrand itself: it tells us the amount of energy carried by photons with frequency between ω and $\omega + d\omega$

$$E(\omega)d\omega = \frac{V\hbar}{\pi^2 c^3} \frac{\omega^3}{e^{\beta\hbar\omega} - 1} d\omega \tag{3.10}$$

This is the *Planck distribution*. It is plotted above for various temperatures. As you can see from the graph, for hot gases the maximum in the distribution occurs at a lower wavelength or, equivalently, at a higher frequency. We can easily determine where this maximum occurs by finding the solution to $dE(\omega)/d\omega = 0$. It is

$$\omega_{\max} = \zeta \frac{k_B T}{\hbar}$$

where $\zeta \approx 2.822$ solves $3 - \zeta = 3e^{-\zeta}$. The equation above is often called *Wien's displacement law*. Roughly speaking, it tells you the colour of a hot object.

To compute the total energy in the gas of photons, we need to do the integration in (3.9). To highlight how the total energy depends on temperature, it is useful to perform the rescaling $x = \beta\hbar\omega$, to get

$$E = \frac{V}{\pi^2 c^3}\frac{(k_B T)^4}{\hbar^3}\int_0^\infty \frac{x^3 dx}{e^x - 1}$$

The integral $I = \int dx\, x^3/(e^x - 1)$ is tricky but doable. It turns out to be $I = \pi^4/15$. (We will effectively prove this fact later in the course when we consider a more general class of integrals (3.27) which can be manipulated into the sum (3.28). The net result of this is to express the integral I above in terms of the Gamma function and the Riemann zeta function: $I = \Gamma(4)\zeta(4) = \pi^4/15$). We learn that the energy density $\mathcal{E} = E/V$ in a gas of photons scales is proportional to T^4,

$$\mathcal{E} = \frac{\pi^2 k_B^4}{15\hbar^3 c^3}\, T^4$$

Stefan-Boltzmann Law

The expression for the energy density above is closely related to the *Stefan-Boltzmann law* which describes the energy emitted by an object at temperature T. That energy flux is defined as the rate of transfer of energy from the surface per unit area. It is given by

$$\text{Energy Flux} = \frac{\mathcal{E}c}{4} \equiv \sigma T^4 \tag{3.11}$$

where

$$\sigma = \frac{\pi^2 k_B^4}{60\hbar^3 c^2} = 5.67 \times 10^{-8}\ Js^{-1}m^{-2}K^{-4}$$

is the Stefan constant.

The factor of the speed of light in the middle equation of (3.11) appears because the flux is the *rate* of transfer of energy. The factor of 1/4 comes because we're not considering the flux emitted by a point source, but rather by an actual object whose size is bigger than the wavelength of individual photons. This means that the photon are only emitted in one direction: away from the object, not into it. Moreover, we only care about the velocity perpendicular to the object, which is $(c\cos\theta)$ where θ is the angle the photon makes with the normal. This means that rather than filling out a sphere of area 4π surrounding the object, the actual flux of photons from any point on the object's surface is given by

$$\frac{1}{4\pi}\int_0^{2\pi} d\phi \int_0^{\pi/2} d\theta\, \sin\theta\, (c\cos\theta) = \frac{c}{4}$$

Radiation Pressure and Other Stuff

All other quantities of interest can be computed from the free energy,

$$\begin{aligned} F &= -k_B T \log Z \\ &= \frac{V k_B T}{\pi^2 c^3} \int_o^\infty d\omega \; \omega^2 \log\left(1 - e^{-\beta\hbar\omega}\right) \end{aligned}$$

We can remove the logarithm through an integration by parts to get,

$$\begin{aligned} F &= -\frac{V\hbar}{3\pi^2 c^3} \int_0^\infty d\omega \; \frac{\omega^3 e^{-\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} \\ &= -\frac{V\hbar}{3\pi^2 c^3} \frac{1}{\beta^4 \hbar^4} \int_0^\infty dx \; \frac{x^3}{e^x - 1} \\ &= -\frac{V\pi^2}{45\hbar^3 c^3} \left(k_B T\right)^4 \end{aligned}$$

From this we can compute the pressure due to electromagnetic radiation,

$$p = -\left.\frac{\partial F}{\partial V}\right|_T = \frac{E}{3V} = \frac{4\sigma}{3c} T^4$$

This is the equation of state for a gas of photons. The middle equation tells us that the pressure of photons is one third of the energy density — a fact which will be important in the *Cosmology* course.

We can also calculate the entropy S and heat capacity C_V. They are both most conveniently expressed in terms of the Stefan constant which hides most of the annoying factors,

$$S = -\left.\frac{\partial F}{\partial T}\right|_V = \frac{16V\sigma}{3c} T^3 \quad , \quad C_V = \left.\frac{\partial E}{\partial T}\right|_V = \frac{16V\sigma}{c} T^3$$

3.2.2 The Cosmic Microwave Background Radiation

The cosmic microwave background, or CMB, is the afterglow of the big bang, a uniform light that fills the Universe. The intensity of this light was measured accurately by the FIRAS (far infrared absolute spectrophotometer) instrument on the COBE satellite in the early 1990s. The result is shown on the right, together with the theoretical curve for a blackbody spectrum at $T = 2.725\ K$. It may look as if the error bars are large, but this is only because they have been multiplied by a factor of 400. If the error bars were drawn at the correct size, you wouldn't be able to to see them.

This result is totally astonishing. The light has been traveling for 13.7 billion years, almost since the beginning of time itself. And yet we can understand it with ridiculous accuracy using such a simple calculation. If you're not awed by this graph then you have no soul.

3.2.3 The Birth of Quantum Mechanics

The key conceptual input in the derivation of Planck's formula (3.10) is that light of frequency ω comes in packets of energy $E = \hbar\omega$. Historically, this was the first time that the idea of quanta arose in theoretical physics.

Let's see what would happen in a classical world where light of frequency ω can have arbitrarily low intensity and, correspondingly, arbitrarily low energy. This is effectively what happens in the regime $h\omega \ll k_B T$ of the Planck distribution where the minimum energy $\hbar\omega$ is completely swamped by the temperature. There we can approximate

$$\frac{1}{e^{\beta\hbar\omega} - 1} \approx \frac{1}{\beta\hbar\omega}$$

and Planck's distribution formula (3.10) reduces to

$$E(\omega) = \frac{V}{2\pi^2 c^3}\,\omega^2 k_B T$$

Notice that all hints of the quantum $\hbar$ have vanished. This is the *Rayleigh-Jeans law* for the distribution of classical radiation. It has a serious problem if we try to extrapolate it to high frequencies since the total energy, $E = \int_0^\infty E(\omega)d\omega$, diverges. This was referred to as the ultra-violet catastrophe. In contrast, in Planck's formula (3.10) there is an exponential suppression at high frequencies. This arises because when $\hbar\omega \gg k_B T$, the

temperature is not high enough to create even a single photon. By imposing a minimum energy on these high frequency modes, quantum mechanics ensures that they become frozen out.

3.2.4 Max Planck (1858-1947)

> "A new scientific truth does not triumph by convincing its opponents and making them see the light, but rather because its opponents eventually die"
>
> *Max Planck*

Planck was educated in Munich and, after a brief spell in Kiel, moved to take a professorship in Berlin. (The same position that Boltzmann had failed to turn up for).

For much of his early career, Planck was adamantly against the idea of atoms. In his view the laws of thermodynamics were inviolable. He could not accept that they sat on a foundation of probability. In 1882, he wrote "atomic theory, despite its great successes, will ultimately have to be abandoned".

Twenty years later, Planck had changed his mind. In 1900, he applied Boltzmann's statistical methods to photons to arrive at the result we derived above, providing the first hints of quantum mechanics. However, the key element of the derivation — that light comes in quanta — was not emphasised by Planck. Later, when quantum theory was developed, Planck refused once more to accept the idea of probability underlying physics. This time he did not change his mind.

3.3 Phonons

It is hard to imagine substances much more different than a gas and a solid. It is therefore quite surprising that we can employ our study of gases to accurately understand certain properties of solids. Consider a crystal of atoms of the type shown in the figure. The individual atoms are stuck fast in position: they certainly don't act like a gas. But the vibrations of the atoms — in other words, sound waves — can be treated using the same formalism that we introduced for photons.

Figure 15:

3.3.1 The Debye Model

Quantum mechanics turns electromagnetic waves into discrete packets of energy called photons. In exactly the same way, sounds waves in solids also come in discrete packets. They are called *phonons*. We'll take the energy of a phonon to again be of the form

$$E = \hbar\omega = \hbar k c_s \tag{3.12}$$

where c_s is now the speed of sound rather than the speed of light.

The density of states for phonons is the same as that of photons (3.6) with two exceptions: we must replace the speed of light c with the speed of sound c_s; and phonons have three polarization states rather than two. There are two transverse polarizations (like the photon) but also a longitudinal mode. The density of states is therefore

$$g(\omega)d\omega = \frac{3V}{2\pi^2 c_s^2}\omega^2 d\omega$$

There is one further important difference between phonons and photons. While light waves can have arbitrarily high frequency, sound waves cannot. This is because high frequency waves have small wavelengths, $\lambda = 2\pi c_s/\omega$. But there is a minimum wavelength in the problem set by the spacing between atoms. it is not possible for sound waves to propagate through a solid with wavelength smaller than the atomic spacing because there's nothing in the middle there to shake.

We will denote the maximum allowed phonon frequency as ω_D. The minimum wavelength, λ_D, should be somewhere around the lattice spacing between atoms, which is $(V/N)^{1/3}$, so we expect that $\omega_D \sim (N/V)^{1/3}c_s$. But how can we work out the coefficient? There is a rather clever argument to determine ω_D due to Debye. (So clever that he gets his initial on the frequency and his name on the section heading). We start by counting the number of single phonon states,

$$\int_0^{\omega_D} d\omega\ g(w) = \frac{V\omega_D^3}{2\pi^2 c_s^3}$$

The clever bit of the argument is to identify this with the number of degrees of freedom in the solid. This isn't immediately obvious. The number of degrees of freedom in a lattice of N atoms is $3N$ since each atom can move in three directions. But the number of single phonon states is counting collective vibrations of all the atoms together. Why should they be equal?

To really see what's going on, one should compute the correct energy eigenstates of the lattice and just count the number of single phonon modes with wavevectors inside the first Brillouin zone. (You will learn about Brillouin zones in the *Applications of Quantum Mechanics* course). But to get the correct intuition, we can think in the following way: in general the solid will have many phonons inside it and each of these phonons could sit in any one of the single-phonon states that we counted above. Suppose that there are three phonons sitting in the same state. We say that this state is occupied three times. In this language, each of the states above can be occupied

an arbitrary number of times, restricted only by the energy available. If you want to describe the total state of the system, you need to say how many phonons are in the first state, and how many are in the second state and so on. The number of one-phonon states is then playing the role of the number of degrees of freedom: it is the number of things you can excite.

The net result of this argument is to equate

$$3N = \frac{V\omega_D^3}{2\pi^2 c_s^3} \quad \Rightarrow \quad \omega_D = \left(\frac{6\pi^2 N}{V}\right)^{1/3} c_s$$

We see that ω_D is related to the atomic spacing $(V/N)^{1/3}$ as we anticipated above, but now we have the coefficient too. However, in some sense, the argument of Debye is "answer analysis". It was constructed to ensure that we have the right high-temperature behaviour that we will see below.

From the maximum frequency ω_D we can construct an associated energy scale, $\hbar\omega_D$, and temperature scale,

$$T_D = \frac{\hbar\omega_D}{k_B}$$

This is known as the Debye temperature. It provides a way of characterising all solids: it is the temperature at which the highest frequency phonon starts to become excited. T_D ranges from around $100K$ for very soft materials such as lead through to $2000K$ for hard materials such as diamond. Most materials have Debye temperatures around room temperature ($\pm 100K$ or so).

Heat Capacity of Solids

All that remains is to put the pieces together. Like photons, the number of phonons is not conserved. The partition function for phonons of a fixed frequency, ω, is the same as for photons (3.7),

$$Z_\omega = 1 + e^{-\beta\hbar\omega} + e^{-2\beta\hbar\omega} + \ldots = \frac{1}{1 - e^{-\beta\hbar\omega}}$$

Summing over all frequencies, the partition function is then

$$\log Z_{\text{phonon}} = \int_0^{\omega_D} d\omega \; g(\omega) \log Z_\omega$$

where the partition function Z_ω for a single phonon of frequency ω is the same as that of a photon (3.7). The total energy in sound waves is therefore

$$E = \int_0^{\omega_D} d\omega \; \frac{\hbar\omega g(\omega)}{e^{\beta\hbar\omega} - 1} = \frac{3V\hbar}{2\pi^2 c_s^3} \int_0^{\omega_D} d\omega \frac{\omega^3}{e^{\beta\hbar\omega} - 1}$$

Figure 16: Experimental data for heat capacities. The solid line is the Debye prediction. (Source: D. Schroeder *An Introduction to Thermal Physics*)

We again rescale the integration variable to $x = \beta\hbar\omega$ so the upper limit of the integral becomes $x_D = T_D/T$. Then we have

$$E = \frac{3V}{2\pi^2(\hbar c_s)^3}(k_B T)^4 \int_0^{T_D/T} dx \; \frac{x^3}{e^x - 1}$$

The integral is a function of T_D/T. It has no analytic expression. However, we can look in the two extremes. Firstly, for $T \ll T_D$ we can replace the upper limit of the integral by infinity. We're then left with the same definite integral that we appeared for photons, $I = \int^\infty dx \; x^3/(e^x - 1) = \pi^4/15$. In this low-temperature regime, the heat capacity is proportional to T^3,

$$C_V = \frac{\partial E}{\partial T} = \frac{2\pi^2 V k_B^4}{5\hbar^3 c_s^3} T^3 \qquad (T \ll T_D) \tag{3.13}$$

It is often expressed in terms of the Debye temperature T_D, so it reads

$$C_V = N k_B \frac{12\pi^4}{5} \left(\frac{T}{T_D}\right)^3 \tag{3.14}$$

In contrast, at temperatures $T \gg T_D$ we only integrate over small values of x, allowing us to Taylor expand the integrand,

$$\int_0^{T_D/T} dx \; \frac{x^3}{e^x - 1} = \int_0^{T_D/T} dx \; \left(x^2 + \ldots\right) = \frac{1}{3}\left(\frac{T_D}{T}\right)^3 + \ldots$$

This ensures that the energy grows linearly with T and the heat capacity tends towards a constant value

$$C_V = \frac{V k_B^4 T_D^3}{2\pi^2 \hbar^3 c_s^3} = 3Nk_B \qquad (T \gg T_D) \tag{3.15}$$

This high-temperature behaviour has been known experimentally since the early 1800's. It is called the *Dulong-Petit law*. Debye's argument for the value of ω_D was basically constructed to reproduce the coefficient $3N$ in the formula above. This was known experimentally, but also from an earlier model of vibrations in a solid due to Einstein. (You met the Einstein model in the first problem sheet). Historically, the real success of the Debye model was the correct prediction of the T^3 behaviour of C_V at low temperatures.

In most materials the heat capacity is dominated by the phonon contribution. (In metals there is an additional contribution from conduction electrons that we will calculate in Section 3.6). The heat capacity of three materials is shown in Figure 16, together with the predictions from the Debye model. As you can see, it works very well! The deviation from theory at high temperatures is due to differences between C_V and C_p, the heat capacity at constant pressure.

What's Wrong with the Debye Model?

As we've seen, the Debye model is remarkably accurate in capturing the heat capacities of solids. Nonetheless, it is a caricature of the physics. The most glaring problem is our starting point (3.12). The relationship $E = \hbar\omega$ between energy and frequency is fine; the mistake is the relationship between frequency ω and wavevector (or momentum) k, namely $\omega = kc_s$. Equations of this type, relating energy and momentum, are called *dispersion relations*. It turns out that that the dispersion relation for phonons is a little more complicated.

It is not hard to compute the dispersion relation for phonons. (You will, in fact, do this calculation in the Applications of Quantum Mechanics course). For simplicity, we'll work with a one dimensional periodic lattice of N atoms as shown in the figure. The equilibrium position of each atom is $x_l = la$ and we impose periodic boundary conditions by insisting that $x_{N+1} \equiv x_1$. Let u_l be the deviation from equilibrium, $u_l = x_l - la$. If we approximate the bonds joining the atoms as har-

Figure 17:

monic oscillators, the Hamiltonian governing the vibrations is

$$H = \frac{1}{2m}\sum_i \dot{u}_i^2 + \frac{\alpha}{2}\sum_i (u_i - u_{i+1})^2$$

where α is a parameter governing the strength of the bonds between atoms. The equation of motion is

$$\ddot{u}_i = \frac{\alpha}{m}(2u_i - u_{i+1} - u_{i-1})$$

This is easily solved by the discrete Fourier transform. We make the ansatz

$$u_l = \frac{1}{\sqrt{N}}\sum_k \tilde{u}_k e^{i(kla - \omega_k t)}$$

Plugging this into the equation of motion gives the dispersion relation

$$\omega_k = 2\sqrt{\frac{\alpha}{m}}\left|\sin\left(\frac{ka}{2}\right)\right|$$

To compute the partition function correctly in this model, we would have to revisit the density of states using the new dispersion relation $E(k) = \hbar\omega_k$. The resulting integrals are messy. However, at low temperatures only the smallest frequency modes are excited and, for small ka, the sin function is approximately linear. This means that we get back to the dispersion relation that we used in the Debye model, $\omega = kc_s$, with the speed of sound given by $c_s = a\sqrt{\alpha/m}$. Moreover, at very high temperatures it is simple to check that this model gives the Dulong-Petit law as expected. It deviates from the Debye model only at intermediate temperatures and, even here, this deviation is mostly negligible.

3.4 The Diatomic Gas Revisited

With a bit of quantum experience under our belt, we can look again at the diatomic gas that we discussed in Section 2.4. Recall that the classical prediction for the heat capacity — $C_V = \frac{7}{2}Nk_B$ — only agrees with experiment at very high temperatures. Instead, the data suggests that as the temperature is lowered, the vibrational modes and the rotational modes become frozen out. But this is exactly the kind of behaviour that we expect for a quantum system where there is a minimum energy necessary to excite each degree of freedom. Indeed, this "freezing out" of modes saved us from ultra-violet catastrophe in the case of blackbody radiation and gave rise to a reduced heat capacity at low temperatures for phonons.

Let's start with the rotational modes, described by the Hamiltonian (2.18). Treating this as a quantum Hamiltonian, it has energy levels

$$E = \frac{\hbar^2}{2I} j(j+1) \qquad j = 0, 1, 2, \ldots$$

The degeneracy of each energy level is $2j+1$. Thus the rotational partition function for a single molecule is

$$Z_{\rm rot} = \sum_{j=0}^{\infty} (2j+1) e^{-\beta\hbar^2 j(j+1)/2I}$$

When $T \gg \hbar^2/2Ik_B$, we can approximate the sum by the integral to get

$$Z_{\rm rot} \approx \int_0^\infty dx\ (2x+1) e^{-\beta\hbar^2 x(x+1)/2I} = \frac{2I}{\beta\hbar^2}$$

which agrees with our result for the classical partition function (2.19).

In contrast, for $T \ll \hbar^2/2Ik_B$ all states apart from $j=0$ effectively decouple and we have simply $Z_{\rm rot} \approx 1$. At these temperatures, the rotational modes are frozen at temperatures accessible in experiment so only the translational modes contribute to the heat capacity.

This analysis also explains why there is no rotational contribution to the heat capacity of a monatomic gas. One could try to argue this away by saying that atoms are point particles and so can't rotate. But this simply isn't true. The correct argument is that the moment of inertia I of an atom is very small and the rotational modes are frozen. Similar remarks apply to rotation about the symmetry axis of a diatomic molecule.

The vibrational modes are described by the harmonic oscillator. You already computed the partition function for this on the first examples sheet (and, in fact, implicitly in the photon and phonon calculations above). The energies are

$$E = \hbar\omega(n + \frac{1}{2})$$

and the partition function is

$$Z_{\rm vib} = \sum_n e^{-\beta\hbar\omega(n+\frac{1}{2})} = e^{-\beta\hbar\omega/2} \sum_n e^{-\beta\hbar\omega n} = \frac{e^{-\beta\hbar\omega/2}}{1 - e^{-\beta\hbar\omega}} = \frac{1}{2\sinh(\beta\hbar\omega/2)}$$

At high temperatures $\beta\hbar\omega \ll 1$, we can approximate the partition function as $Z_{\rm vib} \approx 1/\beta\hbar\omega$ which again agrees with the classical result (2.20). At low temperatures $\beta\hbar\omega \gg$

1, the partition function becomes $Z_{\rm vib} \approx e^{-\beta\hbar\omega/2}$. This is a contribution from the zero-point energy of the harmonic oscillator. It merely gives the expected additive constant to the energy per particle,

$$E_{\rm vib} = -\frac{\partial}{\partial\beta}\log Z_{\rm vib} \approx \frac{\hbar\omega}{2}$$

and doesn't contribute the heat capacity. Once again, we see how quantum effects explain the observed behaviour of the heat capacity of the diatomic gas. The end result is a graph that looks like that shown in Figure 11.

3.5 Bosons

For the final two topics of this section, we will return again to the simple monatomic ideal gas. The classical treatment that we described in Section 2.2 has limitations. As the temperature decreases, the thermal de Broglie wavelength,

$$\lambda = \sqrt{\frac{2\pi\hbar^2}{mk_BT}}$$

gets larger. Eventually it becomes comparable to the inter-particle separation, $(V/N)^{1/3}$. At this point, quantum effects become important. If the particles are non-interacting, there is really only one important effect that we need to consider: quantum statistics.

Recall that in quantum mechanics, particles come in two classes: bosons and fermions. Which class a given particle falls into is determined by its spin, courtesy of the spin-statistics theorem. Integer spin particles are bosons. This means that any wavefunction must be symmetric under the exchange of two particles,

$$\psi(\vec{r}_1,\vec{r}_2) = \psi(\vec{r}_2,\vec{r}_1)$$

Particles with $\frac{1}{2}$-integer spin are fermions. They have an anti-symmetrized wavefunction,

$$\psi(\vec{r}_1,\vec{r}_2) = -\psi(\vec{r}_2,\vec{r}_1)$$

At low temperatures, the behaviour of bosons and fermions is very different. All familiar fundamental particles such as the electron, proton and neutron are fermions. But an atom that contains an even number of fermions acts as a boson as long as we do not reach energies large enough to dislodge the constituent particles from their bound state. Similarly, an atom consisting of an odd number of electrons, protons and neutrons will be a fermion. (In fact, the proton and neutron themselves are not fundamental: they

are fermions because they contain three constituent quarks, each of which is a fermion. If the laws of physics were different so that four quarks formed a bound state rather than three, then both the proton and neutron would be bosons and, as we will see in the next two sections, nuclei would not exist!).

We will begin by describing the properties of bosons and then turn to a discussion of fermions to Section 3.6.

3.5.1 Bose-Einstein Distribution

We'll change notation slightly from earlier sections and label the single particle quantum states of the system by $|r\rangle$. (We used $|n\rangle$ previously, but n will be otherwise occupied for most of this section). The single particle energies are then E_r and we'll assume that our particles are non-interacting. In that case, you might think that to specify the state of the whole system, you would need to say which state particle 1 is in, and which state particle 2 is in, and so on. But this is actually too much information because particle 1 and particle 2 are indistinguishable. To specify the state of the whole system, we don't need to attach labels to each particle. Instead, it will suffice to say how many particles are in state 1 and how many particles are in state 2 and so on.

We'll denote the number of particles in state $|r\rangle$ as n_r. If we choose to work in the canonical ensemble, we must compute the partition function,

$$Z = \sum_{\{n_r\}} e^{-\beta n_r E_r}$$

where the sum is over all possible ways of partitioning N particles into sets $\{n_r\}$ subject to the constraint that $\sum_r n_r = N$. Unfortunately, the need to impose this constraint makes the sums tricky. This means that the canonical ensemble is rather awkward when discussing indistinguishable particles. It turns out to be *much* easier to work in the grand canonical ensemble where we introduce a chemical potential μ and allow the total number of particles N to fluctuate.

Life is simplest it we think of each state $|r\rangle$ in turn. In the grand canonical ensemble, a given state can be populated by an arbitrary number of particles. The grand partition function for this state is

$$\mathcal{Z}_r = \sum_{n_r} e^{-\beta n_r (E_r - \mu)} = \frac{1}{1 - e^{-\beta(E_r - \mu)}}$$

Notice that we've implicitly assumed that the sum above converges, which is true only if $(E_r - \mu) > 0$. But this should be true for all states E_r. We will set the ground state

to have energy $E_0 = 0$, so the grand partition function for a Bose gas only makes sense if

$$\mu < 0 \tag{3.16}$$

Now we use the fact that all the occupation of one state is independent of any other. The full grand partition function is then

$$\mathcal{Z} = \prod_r \frac{1}{1 - e^{-\beta(E_r - \mu)}}$$

From this we can compute the average number of particles,

$$N = \frac{1}{\beta}\frac{\partial}{\partial \mu} \log \mathcal{Z} = \sum_r \frac{1}{e^{\beta(E_r - \mu)} - 1} \equiv \sum_r \langle n_r \rangle$$

Here $\langle n_r \rangle$ denotes the average number of particles in the state $|r\rangle$,

$$\langle n_r \rangle = \frac{1}{e^{\beta(E_r - \mu)} - 1} \tag{3.17}$$

This is the *Bose-Einstein distribution*. In what follows we will always be interested in the thermodynamic limit where fluctuations around the average are negligible. For this reason, we will allow ourselves to be a little sloppy in notation and we write the average number of particles in $|r\rangle$ as n_r instead of $\langle n_r \rangle$.

Notice that we've seen expressions of the form (3.17) already in the calculations for photons and phonons — see, for example, equation (3.7). This isn't coincidence: both photons and phonons are bosons and the origin of this term in both calculations is the same: it arises because we sum over the number of particles in a given state rather than summing over the states for a single particle. As we mentioned in the Section on blackbody radiation, it is not really correct to think of photons or phonons in the grand canonical ensemble because their particle number is not conserved. Nonetheless, the equations are formally equivalent if one just sets $\mu = 0$.

In what follows, it will save ink if we introduce the *fugacity*

$$z = e^{\beta\mu} \tag{3.18}$$

Since $\mu < 0$, we have $0 < z < 1$.

Ideal Bose Gas

Let's look again at a gas of non-relativistic particles, now through the eyes of quantum mechanics. The energy is

$$E = \frac{\hbar^2 k^2}{2m}$$

As explained in Section 3.1, we can replace the sum over discrete momenta with an integral over energies as long as we correctly account for the density of states. This was computed in (3.2) and we reproduce the result below:

$$g(E) = \frac{V}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2} E^{1/2} \tag{3.19}$$

From this, together with the Bose-Einstein distribution, we can easily compute the total number of particles in the gas,

$$N = \int dE\ \frac{g(E)}{z^{-1}e^{\beta E} - 1}\ , \tag{3.20}$$

There is an obvious, but important, point to be made about this equation. If we can do the integral (and we will shortly) we will have an expression for the number of particles in terms of the chemical potential and temperature: $N = N(\mu, T)$. That means that if we keep the chemical potential fixed and vary the temperature, then N will change. But in most experimental situations, N is fixed and we're working in the grand canonical ensemble only because it is mathematically simpler. But nothing is for free. The price we pay is that we will have to invert equation (3.20) to get it in the form $\mu = \mu(N, T)$. Then when we change T, keeping N fixed, μ will change too. We have already seen an example of this in the ideal gas where the chemical potential is given by (2.14).

The average energy of the Bose gas is,

$$E = \int dE\ \frac{Eg(E)}{z^{-1}e^{\beta E} - 1}\ , \tag{3.21}$$

And, finally, we can compute the pressure. In the grand canonical ensemble this is,

$$pV = \frac{1}{\beta}\log \mathcal{Z} = -\frac{1}{\beta}\int dE\, g(E) \log\left(1 - ze^{-\beta E}\right)$$

We can manipulate this last expression using an integration by parts. Because $g(E) \sim E^{1/2}$, this becomes

$$pV = \frac{2}{3}\int dE\ \frac{Eg(E)}{z^{-1}e^{\beta E} - 1} = \frac{2}{3}E \tag{3.22}$$

This is implicitly the equation of state. But we still have a bit of work to do. Equation (3.21) gives us the energy as a function of μ and T. And, by the time we have inverted (3.20), it gives us μ as a function of N and T. Substituting both of these into (3.22) will give the equation of state. We just need to do the integrals...

3.5.2 A High Temperature Quantum Gas is (Almost) Classical

Unfortunately, the integrals (3.20) and (3.21) look pretty fierce. Shortly, we will start to understand some of their properties, but first we look at a particularly simple limit. We will expand the integrals (3.20), (3.21) and (3.22) in the limit $z = e^{\beta\mu} \ll 1$. We'll figure out the meaning of this expansion shortly (although there's a clue in the title of this section if you're impatient). Let's look at the particle density (3.20),

$$\begin{aligned}
\frac{N}{V} &= \frac{1}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\int_0^\infty dE\ \frac{E^{1/2}}{z^{-1}e^{\beta E}-1} \\
&= \frac{1}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\int_0^\infty dE\ \frac{ze^{-\beta E}E^{1/2}}{1-ze^{-\beta E}} \\
&= \frac{1}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\frac{z}{\beta^{3/2}}\int_0^\infty dx\ \sqrt{x}e^{-x}(1+ze^{-x}+\ldots)
\end{aligned}$$

where we made the simple substitution $x = \beta E$. The integrals are all of the Gaussian type and can be easily evaluated by making one further substitution $x = u^2$. The final answer can be conveniently expressed in terms of the thermal wavelength λ,

$$\frac{N}{V} = \frac{z}{\lambda^3}\left(1+\frac{z}{2\sqrt{2}}+\ldots\right) \tag{3.23}$$

Now we've got the answer, we can ask what we've done! What kind of expansion is $z \ll 1$? From the above expression, we see that the expansion is consistent only if $\lambda^3 N/V \ll 1$, which means that the thermal wavelength is much less than the interparticle spacing. But this is true at high temperatures: the expansion $z \ll 1$ is a high temperature expansion.

At first glance, it is surprising that $z = e^{\beta\mu} \ll 1$ corresponds to high temperatures. When $T \to \infty$, $\beta \to 0$ so naively it looks as if $z \to 1$. But this is too naive. If we keep the particle number N fixed in (3.20) then μ must vary as we change the temperature, and it turns out that $\mu \to -\infty$ faster than $\beta \to 0$. To see this, notice that to leading order we need z/λ^3 to be constant, so $z \sim T^{-3/2}$.

High Temperature Equation of State of a Bose Gas

We now wish to compute the equation of state. We know from (3.22) that $pV = \frac{2}{3}E$. So we need to compute E using the same $z \ll 1$ expansion as above. From (3.21), the energy density is

$$\begin{aligned}\frac{E}{V} &= \frac{1}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\int_0^\infty dE\ \frac{E^{3/2}}{z^{-1}e^{\beta E}-1} \\ &= \frac{1}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\frac{z}{\beta^{5/2}}\int_0^\infty dx\ x^{3/2}e^{-x}(1+ze^{-x}+\ldots) \\ &= \frac{3z}{2\lambda^3\beta}\left(1+\frac{z}{4\sqrt{2}}+\ldots\right) \qquad (3.24)\end{aligned}$$

The next part's a little fiddly. We want to eliminate z from the expression above in favour of N/V. To do this we invert (3.23), remembering that we're working in the limit $z \ll 1$ and $\lambda^3 N/V \ll 1$. This gives

$$z = \frac{\lambda^3 N}{V}\left(1-\frac{1}{2\sqrt{2}}\frac{\lambda^3 N}{V}+\ldots\right)$$

which we then substitute into (3.24) to get

$$E = \frac{3}{2}\frac{N}{\beta}\left(1-\frac{1}{2\sqrt{2}}\frac{\lambda^3 N}{V}+\ldots\right)\left(1+\frac{1}{4\sqrt{2}}\frac{\lambda^3 N}{V}+\ldots\right)$$

and finally we substitute this into $pV = \frac{2}{3}E$ to get the equation of state of an ideal Bose gas at high temperatures,

$$pV = Nk_BT\left(1-\frac{\lambda^3 N}{4\sqrt{2}V}+\ldots\right) \qquad (3.25)$$

which reproduces the classical ideal gas, together with a term that we can identify as the second virial coefficient in the expansion (2.21). However, this correction to the pressure hasn't arisen from any interactions among the atoms: it is solely due to quantum statistics. We see that the effect of bosonic statistics on the high temperature gas is to reduce the pressure.

3.5.3 Bose-Einstein Condensation

We now turn to the more interesting question: what happens to a Bose gas at low temperatures? Recall that for convergence of the grand partition function we require $\mu < 0$ so that $z = e^{\beta\mu} \in (0,1)$. Since the high temperature limit is $z \to 0$, we'll anticipate that quantum effects and low temperatures come about as $z \to 1$.

Recall that the number density is given by (3.20) which we write as

$$\frac{N}{V} = \frac{1}{4\pi^2}\left(\frac{2mk_BT}{\hbar^2}\right)^{3/2}\int_0^\infty dx\ \frac{x^{1/2}}{z^{-1}e^x - 1} \equiv \frac{1}{\lambda^3}\, g_{3/2}(z) \tag{3.26}$$

where the function $g_{3/2}(z)$ is one of a class of functions which appear in the story of Bose gases,

$$g_n(z) = \frac{1}{\Gamma(n)}\int_0^\infty dx\ \frac{x^{n-1}}{z^{-1}e^x - 1} \tag{3.27}$$

These functions are also known as polylogarithms and sometimes denoted as $Li_n(z) = g_n(z)$. The function $g_{3/2}$ is relevant for the particle number and the function $g_{5/2}$ appears in the calculation of the energy in (3.21). For reference, the gamma function has value $\Gamma(3/2) = \sqrt{\pi}/2$. (Do not confuse these functions $g_n(z)$ with the density of states $g(E)$. They are not related; they just share a similar name).

In Section 3.5.2 we looked at (3.26) in the $T \to \infty$ limit. There we saw that $\lambda \to 0$ but the function $g_{3/2}(z) \to 0$ in just the right way to compensate and keep N/V fixed. What now happens in the other limit, $T \to 0$ and $\lambda \to \infty$. One might harbour the hope that $g_{3/2}(z) \to \infty$ and again everything balances nicely with N/V remaining constant. We will now show that this can't happen.

A couple of manipulations reveals that the integral (3.27) can be expressed in terms of a sum,

$$\begin{aligned}
g_n(z) &= \frac{1}{\Gamma(n)}\int dx\ \frac{zx^{n-1}e^{-x}}{1 - ze^{-x}} \\
&= \frac{1}{\Gamma(n)} z \int dx\ x^{n-1}e^{-x}\sum_{m=0}^{\infty} z^m e^{-mx} \\
&= \frac{1}{\Gamma(n)}\sum_{m=1}^{\infty} z^m \int dx\ x^{n-1}e^{-mx} \\
&= \frac{1}{\Gamma(n)}\sum_{m=1}^{\infty}\frac{z^m}{m^n}\int du\ u^{n-1}e^{-u}
\end{aligned}$$

But the integral that appears in the last line above is nothing but the definition of the gamma function $\Gamma(n)$. This means that we can write

$$g_n(z) = \sum_{m=1}^{\infty}\frac{z^m}{m^n} \tag{3.28}$$

We see that $g_n(z)$ is a monotonically increasing function of z. Moreover, at $z = 1$, it is equal to the Riemann zeta function

$$g_n(1) = \zeta(n)$$

For our particular problem it will be useful to know that $\zeta(3/2) \approx 2.612$.

Let's now return to our story. As we decrease T in (3.26), keeping N/V fixed, z and hence $g_{3/2}(z)$ must both increase. But z can't take values greater than 1. When we reach $z = 1$, let's denote the temperature as $T = T_c$. We can determine T_c by setting $z = 1$ in equation (3.26),

$$T_c = \left(\frac{2\pi\hbar^2}{k_B m}\right)\left(\frac{1}{\zeta(3/2)}\frac{N}{V}\right)^{2/3} \tag{3.29}$$

What happens if we try to decrease the temperature below T_c? Taken at face value, equation (3.26) tells us that the number of particles should decrease. But that's a stupid conclusion! It says that we can make particles disappear by making them cold. Where did they go? We may be working in the grand canonical ensemble, but in the thermodynamic limit we expect $\Delta N \sim 1/\sqrt{N}$ which is too small to explain our missing particles. Where did we misplace them? It looks as if we made a mistake in the calculation.

In fact, we did make a mistake in the calculation. It happened right at the beginning. Recall that back in Section 3.1 we replaced the sum over states with an integral over energies,

$$\sum_{\vec{k}} \approx \frac{V(2m)^{3/2}}{4\pi^2\hbar^3}\int dE\ E^{1/2}$$

Because of the weight $\sqrt{E}$ in the integrand, the ground state with $E = 0$ doesn't contribute to the integral. We can use the Bose-Einstein distribution (3.17) to compute the number of states that we expect to be missing because they sit in this $E = 0$ ground state,

$$n_0 = \frac{1}{z^{-1} - 1} \tag{3.30}$$

For most values of $z \in (0, 1)$, there are just a handful of particles sitting in this lowest state and it doesn't matter if we miss them. But as z gets *very very* close to 1 (meaning $z \approx 1 - 1/N$) then we get a macroscopic number of particles occupying the ground state.

It is a simple matter to redo our calculation taking into account the particles in the ground state. Equation (3.26) is replaced by

$$N = \frac{V}{\lambda^3}\, g_{3/2}(z) + \frac{z}{1-z}$$

Now there's no problem keeping N fixed as we take z close to 1 because the additional term diverges. This means that if we have finite N, then as we decrease T we can never get to $z = 1$. Instead, z must level out around $z \approx 1 - 1/N$ as $T \to 0$.

For $T < T_c$, the number of particles sitting in the ground state is $\mathcal{O}(\mathcal{N})$. Some simple algebra allows us to determine that fraction of particles in the ground state is

$$\frac{n_0}{N} = 1 - \frac{V}{N\lambda^3}\,\zeta(3/2) = 1 - \left(\frac{T}{T_c}\right)^{3/2} \tag{3.31}$$

At temperatures $T < T_c$, a macroscopic number of atoms discard their individual identities and merge into a communist collective, a single quantum state so large that it can be seen with the naked eye. This is known as the *Bose-Einstein condensate*. It provides an exquisitely precise playground in which many quantum phenomena can be tested.

Bose-Einstein Condensates in the Lab

Figure 18: UFO=BEC

Bose-Einstein condensates (often shortened to BECs) of weakly interacting atoms were finally created in 1995, some 70 years after they were first predicted. These first BECs were formed of Rubidium, Sodium or Lithium and contained between $N \sim 10^4 \to 10^7$ atoms. The transition temperatures needed to create these condensates are extraordinarily small, around $T_c \sim 10^{-7}K$. Figure 19 shows the iconic colour enhanced plots that reveal the existence of the condensate. To create these plots, the atoms are stored in a magnetic trap which is then turned off. A picture is taken of the atom cloud a short time t later when the atoms have travelled a distance $\hbar kt/m$. The grey UFO-like smudges above are the original pictures. From the spread of atoms, the momentum distribution of the cloud is inferred and this is what is shown in Figure 19. The peak that appears in the last two plots reveals that a substantial number of atoms were indeed sitting in the momentum ground state. (This is not a $k = 0$ state because of the finite trap and the Heisenberg uncertainty relation). The initial discoverers of BECs, Eric Cornell and Carl Wieman from Boulder and Wolfgang Ketterle from MIT, were awarded the 2001 Nobel prize in physics.

Figure 19: The velocity distribution of Rubidium atoms, taken from the Ketterle lab at MIT. The left-hand picture shows $T > T_c$, just before the condensate forms. The middle and right-hand pictures both reveal the existence of the condensate.

Low Temperature Equation of State of a Bose Gas

The pressure of the ideal Bose gas was computed in (3.22). We can express this in terms of our new favourite functions (3.27) as

$$p = \frac{2}{3}\frac{E}{V} = \frac{k_B T}{\lambda^3}\, g_{5/2}(z) \tag{3.32}$$

Formally there is also a contribution from the ground state, but it is $\log(1-z)/V$ which is a factor of N smaller than the term above and can be safely ignored. At low temperatures, $T < T_c$, we have $z \approx 1$ and

$$p = \frac{k_B T}{\lambda^3}\zeta(5/2)$$

So at low temperatures, the equation of state of the ideal Bose gas is very different from the classical, high temperature, behaviour. The pressure scales as $p \sim T^{5/2}$ (recall that there is a factor of $T^{3/2}$ lurking in the λ). More surprisingly, the pressure is independent of the density of particles N/V.

3.5.4 Heat Capacity: Our First Look at a Phase Transition

Let's try to understand in more detail what happens as we pass through the critical temperature T_c. We will focus on how the heat capacity behaves on either side of the critical temperature.

We've already seen in (3.32) that we can express the energy in terms of the function $g_{5/2}(z)$,

$$\frac{E}{V} = \frac{3}{2}\frac{k_B T}{\lambda^3}\, g_{5/2}(z)$$

so the heat capacity becomes

$$\frac{C_V}{V} = \frac{1}{V}\frac{dE}{dT} = \frac{15k_B}{4\lambda^3} g_{5/2}(z) + \frac{3}{2}\frac{k_B T}{\lambda^3}\frac{dg_{5/2}}{dz}\frac{dz}{dT} \tag{3.33}$$

The first term gives a contribution both for $T < T_c$ and for $T > T_c$. However, the second term includes a factor of dz/dT and z is a very peculiar function of temperature: for $T > T_c$, it is fairly smooth, dropping off at $T = T_c$. However, as $T \to T_c$, the fugacity rapidly levels off to the value $z \approx 1 - 1/N$. For $T < T_c$, z doesn't change very much at all. The net result of this is that the second term only contributes when $T > T_c$. Our goal here is to understand how this contribution behaves as we approach the critical temperature.

Let's begin with the easy bit. Below the critical temperature, $T < T_c$, only the first term in (3.33) contributes and we may happily set $z = 1$. This gives the heat capacity,

$$C_V = \frac{15V}{4\lambda^3}\zeta(5/2) \sim T^{3/2} \tag{3.34}$$

Now we turn to $T > T_c$. Here we have $z < 1$, so $g_{5/2}(z) < g_{5/2}(1)$. We also have $dz/dT < 0$. This means that the heat capacity decreases for $T > T_c$. But we know that for $T < T_c$, $C_V \sim T^{3/2}$ so the heat capacity must have a maximum at $T = T_c$. Our goal in this section is to understand a little better what the function C_V looks like in this region.

To compute the second term in (3.33) we need to understand both $g'_{5/2}$ and how z changes with T as we approach T_c from above. The first calculation is easy is we use our expression (3.28),

$$g_n(z) = \sum_{m=1}^{\infty} \frac{z^m}{m^n} \quad \Rightarrow \quad \frac{d}{dz} g_n(z) = \frac{1}{z} g_{n-1}(z) \tag{3.35}$$

As $T \to T_c$ from above, $dg_{5/2}/dT \to \zeta(3/2)$, a constant. All the subtleties lie in the remaining term, dz/dT. After all, this is the quantity which is effectively vanishing for $T < T_c$. What's it doing at $T > T_c$? To figure this out is a little more involved. We start with our expression (3.26),

$$g_{3/2}(z) = \frac{N\lambda^3}{V} \qquad T > T_c \tag{3.36}$$

and we'll ask what happens to the function $g_{3/2}(z)$ as $z \to 1$, keeping N fixed. We know that exactly at $z = 1$, $g_{3/2}(1) = \zeta(3/2)$. But how does it approach this value? To answer this, it is actually simplest to look at the derivative $dg_{3/2}/dz = g_{1/2}/z$, where

$$g_{1/2}(z) = \frac{1}{\Gamma(1/2)} \int_0^\infty dx \; \frac{x^{-1/2}}{z^{-1}e^x - 1}$$

The reason for doing this is that $g_{1/2}$ diverges as $z \to 1$ and it is generally easier to isolate divergent parts of a function than some finite piece. Indeed, we can do this straightforwardly for $g_{1/2}$ by looking at the integral very close to $x = 0$, we can write

$$\begin{aligned} g_{1/2}(z) &= \frac{1}{\Gamma(1/2)} \int_0^\epsilon dx \; \frac{x^{-1/2}}{z^{-1}(1+x) - 1} + \text{finite} \\ &= \frac{z}{\Gamma(1/2)} \int_0^\epsilon dx \; \frac{x^{-1/2}}{(1-z) + x} + \ldots \\ &= \frac{2z}{\sqrt{1-z}} \frac{1}{\Gamma(1/2)} \int_0^\epsilon du \; \frac{1}{1+u^2} + \ldots \end{aligned}$$

where, in the last line, we made the substution $u = \sqrt{x/(1-z)}$. So we learn that as $z \to 1$, $g_{1/2}(z) \to z(1-z)^{-1/2}$. But this is enough information to tell us how $g_{3/2}$ approaches its value at $z = 1$: it must be

$$g_{3/2}(z) \approx \zeta(3/2) + A(1-z)^{1/2} + \ldots$$

for some constant A. Inserting this into our equation (3.36) and rearranging, we find that as $T \to T_c$ from above,

$$\begin{aligned} z &\approx 1 - \frac{1}{A^2} \left(\zeta(3/2) - \frac{N\lambda^3}{V} \right)^2 \\ &= 1 - \frac{\zeta(3/2)^2}{A^2} \left(\left(\frac{T}{T_c} \right)^{3/2} - 1 \right)^2 \\ &\approx 1 - B \left(\frac{T - T_c}{T_c} \right)^2 \end{aligned}$$

where, in the second line, we used the expression of the critical temperature (3.29). B is some constant that we could figure out with a little more effort, but it won't be important for our story. From the expression above, we can now determine dz/dT as $T \to T_c$. We see that it vanishes linearly at $T = T_c$.

Putting all this together, we can determine the expression for the heat capacity (3.33) when $T > T_c$. We're not interested in the coefficients, so we'll package a bunch of numbers of order 1 into a constant b and the end result is

$$C_V = \frac{15Vk_B}{4\lambda^3} g_{5/2}(z) - b\left(\frac{T - T_c}{T_c}\right)$$

The first term above goes smoothly over to the expression (3.34) for C_V when $T < T_c$. But the second term is only present for $T > T_c$. Notice that it goes to zero as $T \to T_c$, which ensures that the heat capacity is continuous at this point. But the derivative is *not* continuous. A sketch of the heat capacity is shown in the figure.

Figure 20: Heat Capacity for a BEC

Functions in physics are usually nice and smooth. How did we end up with a discontinuity in the derivative? In fact, if we work at finite N, strictly speaking everything is nice and smooth. There is a similar contribution to dz/dT even at $T < T_c$. We can see that by looking again at the expressions (3.30) and (3.31), which tell us

$$z = \left(1 + \frac{1}{n_0}\right)^{-1} = \left(1 + \frac{1}{N}\frac{1}{1 - (T/T_c)^{3/2}}\right)^{-1} \qquad (T < T_c)$$

The difference is that while dz/dT is of order one above T_c, it is of order $1/N$ below T_c. In the thermodynamic limit, $N \to \infty$, this results in the discontinuity that we saw above. This is a general lesson: phase transitions with their associated discontinuities can only arise in strictly infinite systems. There are no phase transitions in finite systems.

Superfluid Helium-4

A similar, but much more pronounced, discontinuity is seen in Helium-4 as it becomes a superfluid, a transition which occurs at 2.17 K. The atom contains two protons, two neutrons and two electrons and is therefore a boson. (In contrast, Helium-3 contains just a single neutron and is a fermion). The experimental data for the heat capacity of Helium-4 is shown on the right. The successive graphs are zooming in on the phase transistion: the scales are (from left to right) Kelvin, milliKelvin and microKelvin. The discontinuity is often called the *lambda* transition on account of the shape of this graph.

There is a close connection between Bose-Einstein condensation described above and superfluids: strictly speaking a non-interacting Bose-Einstein condensate is not a superfluid but superfluidity is a consequence of arbitrarily weak repulsive interactions between the atoms. However, in He-4, the interactions between atoms are strong and the system cannot be described using the simple techniques developed above.

Figure 21: ^4He.

Something very similar to Bose condensation also occurs in superconductivity and superfluidity of Helium-3. Now the primary characters are fermions rather than bosons (electrons in the case of superconductivity). As we will see in the next section, fermions cannot condense. But they may form bound states due to interactions and these effective bosons can then undergo condensation.

3.6 Fermions

For our final topic, we will discuss the fermion gases. Our analysis will focus solely on non-interacting fermions. Yet this simple model provides a (surprisingly) good first approximation to a wide range of systems, including electrons in metals at low temperatures, liquid Helium-3 and white dwarfs and neutron stars.

Fermions are particles with $\frac{1}{2}$-integer spin. By the spin-statistics theorem, the wavefunction of the system is required to pick up a minus sign under exchange of any particle,

$$\psi(\vec{r}_1, \vec{r}_2) = -\psi(\vec{r}_2, \vec{r}_1)$$

As a corollory, the wavefunction vanishes if you attempt to put two identical fermions in the same place. This is a reflection of the *Pauli exclusion principle* which states that fermions cannot sit in the same state. We will see that the low-energy physics of a gas of fermions is entirely dominated by the exclusion principle.

We work again in the grand canonical ensemble. The grand partition function for a single state $|r\rangle$ is very easy: the state is either occupied or it is not. There is no other option.

$$\mathcal{Z}_r = \sum_{n=0,1} e^{-\beta n(E_r - \mu)} = 1 + e^{-\beta(E_r - \mu)}$$

So, the grand partition function for all states is $\mathcal{Z} = \prod_r \mathcal{Z}_r$, from which we can compute the average number of particles in the system

$$N = \sum_r \frac{1}{e^{\beta(E_r-\mu)}+1} \equiv \sum_r n_r$$

where the average number of particles in the state $|r\rangle$ is

$$n_r = \frac{1}{e^{\beta(E_r-\mu)}+1} \tag{3.37}$$

This is the *Fermi-Dirac distribution*. It differs from the Bose-Einstein distribution only by the sign in the denominator. Note however that we had no convergence issues in defining the partition function. Correspondingly, the chemical potential μ can be either positive or negative for fermions.

3.6.1 Ideal Fermi Gas

We'll look again at non-interacting, non-relativistic particles with $E = \hbar^2 k^2/2m$. Since fermions necessarily have $\frac{1}{2}$-integer spin, s, there is always a degeneracy factor when counting the number of states given by

$$g_s = 2s + 1$$

For example, electrons have spin $\frac{1}{2}$ and, correspondingly have a degeneracy of $g_s = 2$ which just accounts for "spin up" and "spin down" states. We saw similar degeneracy factors when computing the density of states for photons (which have two polarizations) and phonons (which had three). For non-relativistic fermions, the density of states is

$$g(E) = \frac{g_s V}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2} E^{1/2}$$

We'll again use the notation of fugacity, $z = e^{\beta\mu}$. The particle number is

$$N = \int dE\ \frac{g(E)}{z^{-1}e^{\beta E}+1} \tag{3.38}$$

The average energy is

$$E = \int dE\ \frac{Eg(E)}{z^{-1}e^{\beta E}+1}$$

And the pressure is

$$pV = \frac{1}{\beta}\int dE\ g(E)\log\left(1+ze^{-\beta E}\right) = \frac{2}{3}E \tag{3.39}$$

At high temperatures, it is simple to repeat the steps of Section 3.5.2. (This is one of the questions on the problem sheet). Ony a few minus signs differ along the way and one again finds that for $z \ll 1$, the equation of state reduces to that of a classical gas,

$$pV = Nk_BT\left(1 + \frac{\lambda^3 N}{4\sqrt{2}g_s V} + \ldots\right) \tag{3.40}$$

Notice that the minus signs filter down to the final answer: the first quantum correction to a Fermi gas increases the pressure.

3.6.2 Degenerate Fermi Gas and the Fermi Surface

In the extreme limit $T \to 0$, the Fermi-Dirac distribution becomes very simple: a state is either filled or empty.

$$\frac{1}{e^{\beta(E-\mu)}+1} \longrightarrow \begin{cases} 1 & \text{for } E < \mu \\ 0 & \text{for } E > \mu \end{cases}$$

It's simple to see what's going on here. Each fermion that we throw into the system settles into the lowest available energy state. These are successively filled until we run out of particles. The energy of the last filled state is called the *Fermi energy* and is denoted as E_F. Mathematically, it is the value of the chemical potential at $T = 0$,

$$\mu(T = 0) = E_F \tag{3.41}$$

Filling up energy states with fermions is just like throwing balls into a box. With one exception: the energy states of free particles are not localised in position space; they are localised in momentum space. This means that successive fermions sit in states with ever-increasing momentum. In this way, the fermions fill out a ball in momentum space. The momentum of the final fermion is called the *Fermi momentum* and is related to the Fermi energy in the usual way: $\hbar k_F = (2mE_F)^{1/2}$. All states with wavevector $|\vec{k}| \leq k_F$ are filled and are said to form the *Fermi sea* or *Fermi sphere*. Those states with $|\vec{k}| = k_F$ lie on the edge of the Fermi sea. They are said to form the *Fermi surface*. The concept of a Fermi surface is extremely important in later applications to condensed matter physics.

We can derive an expression for the Fermi energy in terms of the number of particles N in the system. To do this, we should appreciate that we've actually indulged in a slight abuse of notation when writing (3.41). In the grand canonical ensemble, T and μ are independent variables: they're not functions of each other! What this equation really means is that if we want to keep the average particle number N in the system

fixed (which we do) then as we vary T we will have to vary μ to compensate. So a slightly clearer way of defining the Fermi energy is to write it directly in terms of the particle number

$$N = \int_0^{E_F} dE\, g(E) = \frac{g_s V}{6\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2} E_F^{3/2} \tag{3.42}$$

Or, inverting,

$$E_F = \frac{\hbar^2}{2m}\left(\frac{6\pi^2}{g_s}\frac{N}{V}\right)^{2/3} \tag{3.43}$$

The Fermi energy sets the energy scale for the system. There is an equivalent temperature scale, $T_F = E_F/k_B$. The high temperature expansion that resulted in the equation of state (3.40) is valid at temperatures $T > T_F$. In contrast, temperatures $T < T_F$ are considered "low" temperatures for systems of fermions. Typically, these low temperatures do not have to be too low: for electrons in a metal, $T_F \sim 10^4 K$; for electrons in a white dwarf, $T_F > 10^7 K$.

While E_F is the energy of the last occupied state, the average energy of the system can be easily calculated. It is

$$E = \int_0^{E_F} dE\; Eg(E) = \frac{3}{5}NE_F \tag{3.44}$$

Similarly, the pressure of the degenerate Fermi gas can be computed using (3.39),

$$pV = \frac{2}{5}NE_F \tag{3.45}$$

Even at zero temperature, the gas has non-zero pressure, known as *degeneracy pressure*. It is a consequence of the Pauli exclusion principle and is important in the astrophysics of white dwarf stars and neutron stars. (We will describe this application in Section 3.6.5). The existence of this residual pressure at $T = 0$ is in stark contrast to both the classical ideal gas (which, admittedly, isn't valid at zero temperature) and the bosonic quantum gas.

3.6.3 The Fermi Gas at Low Temperature

We now turn to the low-temperature behaviour of the Fermi gas. As mentioned above, "low" means $T \ll T_F$, which needn't be particularly low in everyday terms. The number of particles N and the average energy E are given by,

$$N = \int_0^\infty dE\; \frac{g(E)}{z^{-1}e^{\beta E} + 1} \tag{3.46}$$

Figure 22: The Fermi-Dirac Distribution function at $T = 0$ and very small T. The distribution differs from the $T = 0$ ground state only for a range of energies $k_B T$ around E_F.

and

$$E = \int_0^\infty dE \; \frac{Eg(E)}{z^{-1}e^{\beta E} + 1} \tag{3.47}$$

where, for non-relativistic fermions, the density of states is

$$g(E) = \frac{g_s V}{4\pi^2} \left(\frac{2m}{\hbar^2}\right)^{3/2} E^{1/2}$$

Our goal is to firstly understand how the chemical potential μ, or equivalently the fugacity $z = e^{\beta\mu}$, changes with temperature when N is held fixed. From this we can determine how E changes with temperature when N is held fixed.

There are two ways to proceed. The first is a direct approach on the problem by Taylor expanding (3.46) and (3.47) for small T. But it turns out that this is a little delicate because the starting point at $T = 0$ involves the singular distribution shown on the left-hand side of Figure 22 and it's easy for the physics to get lost in a morass of integrals. For this reason, we start by developing a heuristic understanding of how the integrals (3.46) and (3.47) behave at low temperatures which will be enough to derive the required results. We will then give the more rigorous expansion – sometimes called the *Sommerfeld expansion* – and confirm that our simpler derivation is indeed correct.

The Fermi-Dirac distribution (3.37) at small temperatures is sketched on the right-hand side of Figure 22. The key point is that only states with energy within $k_B T$ of the Fermi surface are affected by the temperature. We want to insist that as we vary the temperature, the number of particles stays fixed which means that $dN/dT = 0$. I claim that this holds if, to leading order, the chemical potential is independent of temperature, so

$$\left.\frac{d\mu}{dT}\right|_{T=0} = 0 \tag{3.48}$$

Let's see why this is the case. The change in particle number can be written as

$$\begin{aligned}\frac{dN}{dT} &= \frac{d}{dT}\int_0^\infty dE\ \frac{g(E)}{e^{\beta(E-\mu)}+1} \\ &= \int_0^\infty dE\ g(E)\frac{d}{dT}\left(\frac{1}{e^{\beta(E-\mu)}+1}\right) \\ &\approx\ g(E_F)\int_0^\infty dE\ \frac{\partial}{\partial T}\left(\frac{1}{e^{\beta(E-E_F)}+1}\right)\end{aligned}$$

There are two things going on in the step from the second line to the third. Firstly, we are making use of the fact that, for $k_BT \ll E_F$, the Fermi-Dirac distribution only changes significantly in the vicinity of E_F as shown in the right-hand side of Figure 22. This means that the integral in the middle equation above is only receiving contributions in the vicinity of E_F and we have used this fact to approximate the density of states $g(E)$ with its value at $g(E_F)$. Secondly, we have used our claimed result (3.48) to replace the total derivative d/dT (which acts on the chemical potential) with the partial derivative $\partial/\partial T$ (which doesn't) and μ is replaced with its zero temperature value E_F.

Explicitly differentiating the Fermi-Dirac distribution in the final line, we have

$$\frac{dN}{dT} \approx g(E_F)\int_0^\infty dE\ \left(\frac{E-E_F}{k_BT^2}\right)\frac{1}{4\cosh^2(\beta(E-E_F)/2)} \approx 0$$

This integral vanishes because $(E-E_F)$ is odd around E_F while the cosh function is even. (And, as already mentioned, the integral only receives contributions in the vicinity of E_F).

Let's now move on to compute the change in energy with temperature. This, of course, is the heat capacity. Employing the same approximations that we made above, we can write

$$\begin{aligned}C_V &= \left.\frac{\partial E}{\partial T}\right|_{N,V} = \int_0^\infty dE\ Eg(E)\,\frac{\partial}{\partial T}\left(\frac{1}{e^{\beta(E-\mu)}+1}\right) \\ &\approx \int_0^\infty dE\ \left[E_Fg(E_F)+\frac{3}{2}g(E_F)(E-E_f)\right]\frac{\partial}{\partial T}\left(\frac{1}{e^{\beta(E-E_F)}+1}\right)\end{aligned}$$

However, this time we have not just replaced $Eg(E)$ by $E_Fg(E_F)$, but instead Taylor expanded to include a term linear in $(E-E_F)$. (The factor of 3/2 comes about because $Eg(E) \sim E^{3/2}$. The first $E_Fg(E_F)$ term in the square brackets vanishes by the same even/odd argument that we made before. But the $(E-E_F)$ term survives.

Writing $x = \beta(E - E_F)$, the integral becomes

$$C_V \approx \frac{3}{2} g(E_F) T \int_{-\infty}^{\infty} dx \; \frac{x^2}{4\cosh^2(x/2)}$$

where we've extended the range of the integral from $-\infty$ to $+\infty$, safe in the knowledge that only the region near E_F (or $x = 0$) contributes anyway. More importantly, however, this integral only gives an overall coefficient which we won't keep track of. The final result for the heat capacity is

$$C_V \sim T\, g(E_F)$$

There is a simple way to intuitively understand this linear behaviour. At low temperatures, only fermions within $k_B T$ of E_F are participating in the physics. The number of these particles is $\sim g(E_F) k_B T$. If each picks up energy $\sim k_B T$ then the total energy of the system scales as $E \sim g(E_F)(k_B T)^2$, resulting in the linear heat capacity found above.

Finally, the heat capacity is often re-expressed in a slightly different form. Using (3.42) we learn that $N \sim E_F^{3/2}$ which allows us to write,

$$C_V \sim N k_B \left(\frac{T}{T_F}\right) \tag{3.49}$$

Heat Capacity of Metals

One very important place that we can apply the theory above it to metals. We can try to view the conduction electrons — those which are free to move through the lattice — as an ideal gas. At first glance, this seems very unlikely to work. This is because we are neglecting the Coulomb interaction between electrons. Nonetheless, the ideal gas approximation for electrons turns out to work remarkably well.

From what we've learned in this section, we would expect two contributions to the heat capacity of a metal. The vibrations of the lattice gives a phonon contribution which goes as T^3 (see equation (3.13)). If the conduction electrons act as an ideal gas, they should give a linear term. The low-temperature heat capacity can therefore be written as

$$C_V = \gamma T + \alpha T^3$$

Experimental data is usually plotted as C_V/T vs T^2 since this gives a straight line which looks nice. The intercept with the C_V axis tells us the electron contribution. The heat capacity for copper is shown in the figure.

We can get a rough idea of when the phonon and electron contributions to heat capacity are comparable. Equating (3.14) with (3.56), and writing $24\pi^2/5 \approx 50$, we have the the two contributions are equal when $T^2 \sim T_D^3/50T_F$. Ballpark figures are $T_D \sim 10^2 K$ and $T_F \sim 10^4 K$ which tells us that we have to get down to temperatures of around $1K$ or so to see the electron contribution.

Figure 23: The heat capacity of copper (taken from A. Tari "*The Specific Heat of Matter at Low Temperatures*")

For many metals, the coefficient γ of the linear heat capacity is fairly close to the ideal gas value (within, say, 20% or so). Yet the electrons in a metal are far from free: the Coulomb energy from their interations is at least as important as the kinetic energy. So why does the ideal gas approximation work so well? This was explained in the 1950s by Landau and the resulting theory — usually called Landau's Fermi liquid theory — is the basis for our current understanding of electron systems.

3.6.4 A More Rigorous Approach: The Sommerfeld Expansion

The discussion in Section 3.6.3 uses some approximations but highlights the physics of the low-temperature fermi gas. For completeness, here we present a more rigorous derivation of the low-temperature expansion.

To start, we introduce some new notation for the particle number and energy integrals, (3.46) and (3.47). We write,

$$\begin{aligned} \frac{N}{V} &= \frac{g_s}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\int_0^\infty dE\ \frac{E^{1/2}}{z^{-1}e^{\beta E}+1} \\ &= \frac{g_s}{\lambda^3}\, f_{3/2}(z) \end{aligned} \tag{3.50}$$

and

$$\begin{aligned} \frac{E}{V} &= \frac{g_s}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\int_0^\infty dE\ \frac{E^{3/2}}{z^{-1}e^{\beta E}+1} \\ &= \frac{3}{2}\frac{g_s}{\lambda^3}k_BT\, f_{5/2}(z) \end{aligned} \tag{3.51}$$

where $\lambda = \sqrt{2\pi\hbar^2/mk_BT}$ is the familiar thermal wavelength and the functions $f_n(z)$ are the fermionic analogs of g_n defined in (3.27),

$$f_n(z) = \frac{1}{\Gamma(n)}\int_0^\infty dx\ \frac{x^{n-1}}{z^{-1}e^x+1} \tag{3.52}$$

where it is useful to recall $\Gamma(3/2) = \sqrt{\pi}/2$ and $\Gamma(5/2) = 3\sqrt{\pi}/4$ (which follows, of course, from the property $\Gamma(n+1) = n\Gamma(n)$). This function is an example of a polylogarithm and is sometimes written as $Li_n(-z) = -f_n(z)$.

Expanding $f_n(z)$

We will now derive the large z expansion of $f_n(z)$, sometimes called the *Sommerfeld expansion*. The derivation is a little long. We begin by splitting the dx integral into two parts,

$$\begin{aligned}
\Gamma(n) f_n(z) &= \int_0^{\beta\mu} dx\ \frac{x^{n-1}}{z^{-1}e^x + 1} + \int_{\beta\mu}^{\infty} dx\ \frac{x^{n-1}}{z^{-1}e^x + 1} \\
&= \int_0^{\beta\mu} dx\ x^{n-1}\left(1 - \frac{1}{1 + ze^{-x}}\right) + \int_{\beta\mu}^{\infty} dx\ \frac{x^{n-1}}{z^{-1}e^x + 1} \\
&= \frac{(\log z)^n}{n} - \int_0^{\beta\mu} dx\ \frac{x^{n-1}}{1 + ze^{-x}} + \int_{\beta\mu}^{\infty} dx\ \frac{x^{n-1}}{z^{-1}e^x + 1}
\end{aligned}$$

We now make a simple change of variable, $\eta_1 = \beta\mu - x$ for the first integral and $\eta_2 = x - \beta\mu$ for the second,

$$\Gamma(n) f_n(z) = \frac{(\log z)^n}{n} - \int_0^{\beta\mu} d\eta_1\ \frac{(\beta\mu - \eta_1)^{n-1}}{1 + e^{\eta_1}} + \int_0^{\infty} d\eta_2\ \frac{(\beta\mu + \eta_2)^{n-1}}{1 + e^{\eta_2}}$$

So far we have only massaged the integral into a useful form. We now make use of the approximation $\beta\mu \gg 1$. Firstly, we note that the first integral has a e^{η_1} suppression in the denominator. If we replace the upper limit of the integral by ∞, the expression changes by a term of order $e^{-\beta\mu} = z^{-1}$ which is a term we're willing to neglect. With this approximation, we can combine the two integrals into one,

$$\Gamma(n) f_n(z) = \frac{(\log z)^n}{n} + \int_0^{\infty} d\eta\ \frac{(\beta\mu + \eta)^{n-1} - (\beta\mu - \eta)^{n-1}}{1 + e^{\eta}}$$

We now Taylor expand the numerator,

$$\begin{aligned}
(\beta\mu + \eta)^{n-1} - (\beta\mu - \eta)^{n-1} &= (\beta\mu)^{n-1}\left[(1 + \eta/\beta\mu)^{n-1} - (1 - \eta/\beta\mu)^{n-1}\right] \\
&= (\beta\mu)^{n-1}\left[1 + (n-1)\frac{\eta}{\beta\mu} + \ldots - 1 + (n-1)\frac{\eta}{\beta\mu} + \ldots\right] \\
&= 2(n-1)(\beta\mu)^{n-2}\eta + \ldots
\end{aligned}$$

From which we get

$$\Gamma(n) f_n(z) = \frac{(\log z)^n}{n} + 2(n-1)(\log z)^{n-2} \int_0^{\infty} d\eta\ \frac{\eta}{e^{\eta} + 1}$$

We're left with a fairly simple integral.

$$\begin{aligned}\int_0^\infty d\eta\ \frac{\eta}{e^\eta+1} &= \int_0^\infty d\eta\ \frac{\eta e^{-\eta}}{1+e^{-\eta}} \\ &= \int_0^\infty d\eta\ \eta \sum_{m=1}^\infty e^{-m\eta}(-1)^{m+1} \\ &= \sum_{m=1}^\infty \frac{(-1)^{m+1}}{m^2} \int_0^\infty du\ ue^{-u}\end{aligned}$$

The integral in the last line is very simple: $\int_0^\infty du\ ue^{-u} = 1$. We have only to do the sum. There are a number of ways to manipulate this. We chose to write

$$\begin{aligned}1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} \ldots &= \left(1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \ldots\right) - 2\left(\frac{1}{2^2} + \frac{1}{4^2} + \frac{1}{6^2} + \ldots\right) \\ &= \left(1 - \frac{2}{2^2}\right) \cdot \left(1 + \frac{1}{2^2} + \frac{1}{3^2} + \ldots\right) \qquad (3.53)\end{aligned}$$

This final sum is well known. It is $\sum 1/m^2 = \zeta(2) = \pi^2/6$. (The original proof of this, by Euler, proceeds by Taylor expanding $\sin x/x$, writing resulting sum as a product of roots, and then equating the x^2 term).

After all this work, we finally have the result we want. The low temperature expansion of $f_n(z)$ is an expansion in $1/\log z = 1/\beta\mu$,

$$f_n(z) = \frac{(\log z)^n}{\Gamma(n+1)}\left(1 + \frac{\pi^2}{6}\frac{n(n-1)}{(\log z)^2} + \ldots\right) \qquad (3.54)$$

Back to Physics: Heat Capacity of a Fermi Gas Again

The expansion (3.54) is all we need to determine the leading order effects of temperature on the Fermi gas. The number of particles (3.50) can be expressed in terms of the chemical potential as

$$\frac{N}{V} = \frac{g_s}{6\pi^2\hbar^3}(2m\mu)^{3/2}\left(1 + \frac{\pi^2}{8}\left(\frac{k_BT}{\mu}\right)^2 + \ldots\right) \qquad (3.55)$$

But the number of particles determines the Fermi energy through (3.43). For fixed particle number N, we can therefore express the chemical potential at finite temperature in terms of the Fermi energy. After a small amount of algebra, we find

$$\mu = E_F\left(1 - \frac{\pi^2}{12}\left(\frac{k_BT}{E_F}\right)^2 + \ldots\right)$$

We see that the chemical potential is a maximum at $T = 0$ and decreases at higher T. This is to be expected: recall that by the time we get to the classical gas, the chemical potential must be negative. Moreover, note that the leading order correction is quadratic in temperature rather than linear, in agreement with our earlier claim (3.48).

The energy density can be computed from (3.51) and (3.54). It is

$$\frac{E}{V} = \frac{g_s}{10\pi^2\hbar^3}(2m)^{3/2}\mu^{5/2}\left(1 + \frac{5\pi^2}{8}\left(\frac{k_B T}{\mu}\right)^2 + \dots\right)$$

Our real interest is in the heat capacity. However, with fixed particle number N, the chemical potential also varies with temperature at quadratic order. We can solve this problem by dividing by (3.55) to get

$$\frac{E}{N} = \frac{3E_F}{5}\left(1 + \frac{5\pi^2}{12}\left(\frac{k_B T}{E_F}\right)^2 + \dots\right)$$

Now the only thing that depends on temperature on the right-hand side is T itself. From this we learn that the heat capacity of a low temperature Fermi gas is linear in T,

$$C_V = \left.\frac{\partial E}{\partial T}\right|_{N,V} = N k_B \frac{\pi^2}{2}\frac{T}{T_F} \tag{3.56}$$

But we now have the correct coefficient that completes our earlier result (3.49).

3.6.5 White Dwarfs and the Chandrasekhar limit

When stars exhaust their fuel, the temperature $T \to 0$ and they have to rely on the Pauli exclusion principle to support themselves through the degeneracy pressure (3.45). Such stars supported by electron degeneracy pressure are called white dwarfs. In addition to the kinetic energy E_{kinetic} of fermions given in (3.43) the system has gravitational energy. If we make the further assumption that the density in a star of radius R is uniform, then

$$E_{\text{grav}} = -\frac{3}{5}\frac{G_N M^2}{R}$$

where G_N is Newton's constant. In the problem set, you will minimise $E_{\text{grav}} + E_{\text{kinetic}}$ to find the relationship between the radius and mass of the star,

$$R \sim M^{-1/3}$$

This is unusual. Normally if you throw some stuff on a pile, the pile gets bigger. Not so for stars supported by degeneracy pressure. The increased gravitational attraction wins and causes them to shrink.

As the star shrinks, the Fermi energy (3.43) grows. Eventually it becomes comparable to the electron mass m_e and our non-relativistic approximation breaks down. We can redo our calculations using the relativistic formula for the density of states (3.4) (which needs to be multiplied by a factor of 2 to account for the spin degeneracy). In the opposite regime of ultra-relativistic electrons, where $E \gg m_e$, then we can expand the density of states as

$$g(E) = \frac{V}{\pi^2\hbar^3c^3}\left(E^2 - \frac{m^2c^4}{2} + \dots\right)$$

from which we can determine the kinetic energy due to fermions, replacing the non-relativistic result (3.44) by

$$E_{\text{kinetic}} = \frac{V}{\pi^2\hbar^3c^3}\left(\frac{1}{4}E_F^4 - \frac{m^2c^4}{4}E_F^2 + \dots\right)$$

The Fermi energy can be expressed in terms of the particle number by

$$N = \frac{V}{\pi^2\hbar^3c^3}\left(\frac{1}{3}E_F^3 - \frac{m^2c^4}{2}E_F + \dots\right)$$

The total energy of the system is then given by,

$$E_{\text{grav}} + E_F = \left[\frac{3\hbar c}{4}\left(\frac{9\pi M^4}{4m_p^4}\right)^{1/3} - \frac{3}{5}G_N M^2\right]\frac{1}{R} + \mathcal{O}(R)$$

where m_p is the proton mass (and we're taking $M = Nm_p$ as a good approximation to the full mass of the star). If the term above is positive then the star once again settles into a minimum energy state, where the $1/R$ term balances the term that grows linearly in R. But if the $1/R$ term is negative, we have a problem: the star is unstable to gravitational collapse and will shrink to smaller and smaller values of R. This occurs when the mass exceeds the *Chandrasekhar limit*, $M > M_C$. Neglecting factors of 2 and π, this limit is

$$M_C \sim \left(\frac{\hbar c}{G_N}\right)^{3/2}\frac{1}{m_p^2} \tag{3.57}$$

This is roughly 1.5 times the mass of the Sun. Stars that exceed this bound will not end their life as white dwarfs. Their collapse may ultimately be stopped by degeneracy pressure of neutrons so that they form a neutron star. Or it may not be stopped at all in which case they form a black hole.

The factor in brackets in (3.57) is an interesting mix of fundamental constants associated to quantum theory, relativity and gravity. In this combination, they define a mass scale called the Planck mass,

$$M_{pl}^2 = \frac{\hbar c}{G_N}$$

When energy densities reach the Planck scale, quantum gravity effects are important meaning that we have to think of quantising spacetime itself. We haven't quite reached this limit in white dwarf stars. (We're shy by about 25 orders of magnitude or so!). Nonetheless the presence of the Planck scale in the Chandra limit is telling us that both quantum effects and gravitational effects are needed to understand white dwarf stars: the matter is quantum; the gravity is classical.

3.6.6 Pauli Paramagnetism

> "With heavy heart, I have decided that Fermi-Dirac, not Einstein, is the correct statistics, and I have decided to write a short note on paramagnetism"
>
> *Wolfgang Pauli, in a letter to Schrödinger, reluctantly admitting that electrons are fermions*

We now describe a gas of electrons in a background magnetic field $\vec{B}$. There are two effects that are important: the Lorentz force $\vec{v} \times B$ on the motion of electrons and the coupling of the electron spin to the magnetic field. We first discuss the spin coupling and postpone the Lorentz force for the next section.

An electron can have two spin states, "up" and "down". We'll denote the spin state by a discrete label: $s = 1$ for spin up; $s = -1$ for spin down. In a background magnetic field $\vec{B}$, the kinetic energy picks up an extra term

$$E_{\text{spin}} = \mu_B B s \tag{3.58}$$

where $\mu_B = |e|\hbar/2m$ is the Bohr magneton. (It has nothing to do with the chemical potential. Do not confuse them!)

Since spin up and spin down electrons have different energies, their occupation numbers differ. Equation (3.50) is now replaced by

$$\frac{N_\uparrow}{V} = \frac{1}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\int_0^\infty dE\ \frac{E^{1/2}}{e^{\beta(E+\mu_B B-\mu)}+1} = \frac{1}{\lambda^3} f_{3/2}(ze^{-\beta\mu_B B})$$
$$\frac{N_\downarrow}{V} = \frac{1}{4\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\int_0^\infty dE\ \frac{E^{1/2}}{e^{\beta(E-\mu_B B-\mu)}+1} = \frac{1}{\lambda^3} f_{3/2}(ze^{+\beta\mu_B B})$$

Our interest is in computing the *magnetization*, which is a measure of how the energy of the system responds to a magnetic field,

$$M = -\frac{\partial E}{\partial B} \tag{3.59}$$

Since each spin has energy (3.58), the magnetization is simply the difference in the in the number of up spins and down spins,

$$M = -\mu_B(N_\uparrow - N_\downarrow) = -\frac{\mu_B V}{\lambda^3}\left[f_{3/2}(ze^{-\beta\mu_B B}) - f_{3/2}(ze^{\beta\mu_B B})\right]$$

At high temperatures, and suitably low magnetic fields, we can approximate $f_{3/2}(z) \approx z$ as $z \to 0$, so that

$$M \approx \frac{2\mu_B V z}{\lambda^3}\sinh(\beta\mu_B B)$$

We can eliminate the factor of z in favour of the particle number,

$$N = N_\uparrow + N_\downarrow \approx \frac{2Vz}{\lambda^3}\cosh(\beta\mu_B B)$$

From which we can determine the high temperature magnetization,

$$M \approx \mu_B N \tanh(\beta\mu_B B)$$

This is the same result that you computed in the first problem sheet using the simple two-state model. We see that, once again, at high temperatures the quantum statistics are irrelevant. The magnetic susceptibility is a measure of how easy it is to magnetise a substance. It is defined as

$$\chi \equiv \frac{\partial M}{\partial B} \tag{3.60}$$

Evaluating χ in the limit of vanishing magnetic field, we find that the magnetization is inversely proportional to the temperature,

$$\chi(B=0) = \frac{N\mu_B^2}{k_B T}$$

This $1/T$ behaviour is known as *Curie's law*. (Pierre, not Marie).

The above results hold in the high temperature limit. In contrast, at low temperatures the effects of the Fermi surface become important. We need only keep the leading order term in (3.54) to find

$$M \approx \frac{\mu_B V}{6\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2}\left[(E_F+\mu_B B)^{3/2}-(E_F-\mu_B B)^{3/2}\right] \approx \frac{\mu_B^2 V}{2\pi^2}\left(\frac{2m}{\hbar^2}\right)^{3/2} E_F^{1/2} B$$

where we're written the final result in the limit $\mu_B B \ll E_F$ and expanded to linear order in B. We could again replace some of the factors for the electron number N. However, we can emphasise the relevant physics slightly better if we write the result above in terms of the density of states (3.19). Including the factor of 2 for spin degeneracy, we have

$$M \approx \mu_B^2 g(E_F) B \tag{3.61}$$

We see that at low temperatures, the magnetic susceptibility no longer obeys Curie's law, but saturates to a constant

$$\chi \equiv \frac{\partial M}{\partial B} = \mu_B^2 g(E_F)$$

What's happening here is that most electrons are embedded deep within the Fermi surface and the Pauli exclusion principle forbids them from flipping their spins in response to a magnetic field. Only those electrons that sit on the Fermi surface itself — all $g(E_F)$ of them — have the luxury to save energy by aligning their spins with the magnetic field.

Notice that $\chi > 0$. Such materials are called paramagnetic: in the presence of a magnetic field, the magnetism increases. Materials that are paramagnetic are attracted to applied magnetic fields. The effect described above is known as *Pauli paramagnetism*.

3.6.7 Landau Diamagnetism

For charged fermions, such as electrons, there is a second consequence of placing them in a magnetic field: they move in circles. The Hamiltonian describing a particle of charge $-e$ in a magnetic field $\vec{B} = \nabla \times \vec{A}$ is,

$$H = \frac{1}{2m}\left(\vec{p} + e\vec{A}(\vec{r})\right)^2$$

This has an immediate consequence: a magnetic field cannot affect the statistical physics of a classical gas! This follows simply from substituting the above Hamiltonian into the classical partition function (2.1). A simple shift of variables, $\vec{p}' = \vec{p} + e\vec{A}$ shows that the classical partition function does not depend on $\vec{B}$. This result (which was also mentioned in a slightly different way in the Classical Dynamics course) is known as the Bohr-van Leeuwen theorem: it states that there can be no classical magnetism.

Let's see how quantum effects change this conclusion. Our first goal is to understand the one-particle states for an electron in a constant background magnetic field. This is a problem that you will also solve in the Applications of Quantum Mechanics course. It is the problem of Landau levels.

Landau Levels

We will consider an electron moving in a constant magnetic field pointing in the $\hat{z}$ direction: $\vec{B} = (0, 0, B)$. There are many different ways to pick $\vec{A}$ such that $\vec{B} = \nabla \times \vec{A}$ gives the right magnetic field. These are known as *gauge choices*. Each choice will give rise to the same physics but, usually, very different equations in the intervening steps. Here we make the following gauge choice,

$$\vec{A} = (-By, 0, 0) \tag{3.62}$$

Let's place our particle in a cubic box with sides of length L. Solutions to the Schrödinger equation $\hat{H}\psi = E\psi$ are simply plane waves in the x and z directions,

$$\psi(\vec{r}) = e^{i(k_x x + k_z z)} f(y)$$

But the wavefunction $f(y)$ must satisfy,

$$\left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial y^2} + \frac{1}{2m}\left(\hbar k_x - eBy\right)^2 \right] f(y) = E' f(y) \tag{3.63}$$

But this is a familiar equation! If we define the *cyclotron frequency*,

$$\omega_c = \frac{eB}{m}$$

Then we can rewrite (3.63) as

$$\left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial y^2} + \frac{1}{2}m\omega_c^2\left(y - y_0\right)^2 \right] f(y) = E' f(y)$$

where $y_0 = \hbar k_x / eB$. This is simply the Schrödinger equation for a harmonic oscillator, jiggling with frequency ω_c and situated at position y_0.

One might wonder why the harmonic oscillator appeared. The cyclotron frequency ω_c is the frequency at which electrons undergo classical Larmor orbits, so that kind of makes sense. But why have these become harmonic oscillators sitting at positions y_0 which depend on k_x? The answer is that there is no deep meaning to this at all! it is merely an artefact of our gauge choice (3.62). Had we picked a different $\vec{A}$ giving rise to

the same $\vec{B}$, we would find ourselves trying to answer a different question! The lesson here is that when dealing with gauge potentials, the intermediate steps of a calculation do not always have physical meaning. Only the end result is important. Let's see what this is. The energy of the particle is

$$E = E' + \frac{\hbar^2 k_z^2}{2m}$$

where E' is the energy of the harmonic oscillator,

$$E' = \left(n + \frac{1}{2}\right)\hbar\omega_c \qquad n \in \mathbf{Z}$$

These discrete energy levels are known as *Landau levels*. They are highly degenerate. To see this, note that k_x is quantized in unit of $\Delta k_x = 2\pi/L$. This means that we can have a harmonic oscillator located every $\Delta y_0 = 2\pi\hbar/eBL$. The number of such oscillators that we can fit into the box of size L is $L/\Delta y_0$. This is the degeneracy of each level,

$$\frac{eBL^2}{2\pi\hbar} \equiv \frac{\Phi}{\Phi_0}$$

where $\Phi = L^2 B$ is the total flux through the system and $\Phi_0 = 2\pi\hbar/e$ is known as the *flux quantum*. (Note that the result above does not include a factor of 2 for the spin degeneracy of the electron).

Back to the Diamagnetic Story

We can now compute the grand partition function for non-interacting electrons in a magnetic field. Including the factor of 2 from electron spin, we have

$$\log \mathcal{Z} = \frac{L}{2\pi}\int dk_z \sum_{n=0}^{\infty} \frac{2L^2 B}{\Phi_0} \log\left[1 + z\exp\left(-\frac{\beta\hbar^2 k_z^2}{2m} - \beta\hbar\omega_c(n+1/2)\right)\right]$$

To perform the sum, we use the Euler summation formula which states that for any function $h(x)$,

$$\sum_{n=0}^{\infty} h(n+1/2) = \int_0^{\infty} h(x)dx + \frac{1}{24}h'(0) + \ldots$$

We'll apply the Euler summation formula to the function,

$$h(x) = \int dk_z \log\left[1 + \exp\left(-\frac{\beta\hbar^2 k_z^2}{2m} + \beta x\right)\right]$$

So the grand partition function becomes

$$\begin{aligned}
\log \mathcal{Z} &= \frac{VB}{\pi\Phi_0}\sum_{n=0}^{\infty} h(\mu - \hbar\omega_c(n+1/2)) \\
&= \frac{VB}{\pi\Phi_0}\int_0^{\infty} h(\mu - \hbar\omega_c x)\,dx - \frac{VB}{\pi\Phi_0}\frac{\hbar\omega_c}{24}\frac{dh(\mu)}{d\mu} + \ldots \\
&= \frac{Vm}{2\pi^2\hbar^2}\left[\int_{-\infty}^{\mu} h(y)\,dy - \frac{(\hbar\omega_c)^2}{24}\int_{-\infty}^{+\infty} dk\ \frac{\beta}{e^{\beta(\hbar^2k^2/2m-\mu)}+1}\right] + \ldots \qquad (3.64)
\end{aligned}$$

The first term above does not depend on B. In fact it is simply a rather perverse way of writing the partition function of a Fermi gas when $B = 0$. However, our interest here is in the magnetization which is again defined as (3.59). In the grand canonical ensemble, this is simply

$$M = \frac{1}{\beta}\frac{\partial \log \mathcal{Z}}{\partial B}$$

The second term in (3.64) is proportional to B^2 (which is hiding in the ω_c^2 term). Higher order terms in the Euler summation formula will carry higher powers of B and, for small B, the expression above will suffice.

At $T = 0$, the integrand is simply 1 for $|k| < k_F$ and zero otherwise. To compare to Pauli paramagnetism, we express the final result in terms of the Bohr magneton $\mu_B = |e|\hbar/2mc$. We find

$$M = -\frac{\mu_B^2}{3}g(E_F)B$$

This is comparable to the Pauli contribution (3.61). But it has opposite sign. Substances whose magnetization is opposite to the magnetic field are called *diamagnetic*: they are repelled by applied magnetic fields. The effect that we have derived above is known as *Landau diamagnetism*.

4. Classical Thermodynamics

> "Thermodynamics is a funny subject. The first time you go through it, you don't understand it at all. The second time you go through it, you think you understand it, except for one or two small points. The third time you go through it, you know you don't understand it, but by that time you are used to it, so it doesn't bother you any more."
>
> *Arnold Sommerfeld, making excuses*

So far we've focussed on a statistical mechanics, studying systems in terms of their microscopic constituents. In this section, we're going to take a step back and look at classical thermodynamics. This is a theory that cares nothing for atoms and microscopics. Instead it describes relationships between the observable macroscopic phenomena that we see directly.

In some sense, returning to thermodynamics is a retrograde step. It is certainly not as fundamental as the statistical description. Indeed, the "laws" of thermodynamics that we describe below can all be derived from statistical physics. Nonetheless, there are a number of reasons for developing classical thermodynamics further.

First, pursuing classical thermodynamics will give us a much deeper understanding of some of the ideas that briefly arose in Section 1. In particular, we will focus on how energy flows due to differences in temperature. Energy transferred in this way is called *heat*. Through a remarkable series of arguments involving heat, one can deduce the existence of a quantity called entropy and its implications for irreversibility in the Universe. This definition of entropy is entirely equivalent to Boltzmann's later definition $S = k_B \log \Omega$ but makes no reference to the underlying states.

Secondly, the weakness of thermodynamics is also its strength. Because the theory is ignorant of the underlying nature of matter, it is limited in what it can tell us. But this means that the results we deduce from thermodynamics are not restricted to any specific system. They will apply equally well in any circumstance, from biological systems to quantum gravity. And you can't say that about a lot of theories!

In Section 1, we briefly described the first and second laws of thermodynamics as consequences of the underlying principles of statistical physics. Here we instead place ourselves in the shoes of Victorian scientists with big beards, silly hats and total ignorance of atoms. We will present the four laws of thermodynamics as axioms on which the theory rests.

4.1 Temperature and the Zeroth Law

We need to start with a handful of definitions:

- A system that is completely isolated from all outside influences is said to be contained in *adiabatic walls*. We will also refer to such systems as *insulated*.

- Walls that are not adiabatic are said to be *diathermal* and two systems separated by a diathermal wall are said to be in *thermal contact*. A diathermal wall is still a wall which means that it neither moves, nor allows particles to transfer from one system to the other. However, it is not in any other way special and it will allow heat (to be defined shortly) to be transmitted between systems. If in doubt, think of a thin sheet of metal.

- An isolated system, when left alone for a suitably long period of time, will relax to a state where no further change is noticeable. This state is called *equilibrium*

Since we care nothing for atoms and microstates, we must use macroscopic variables to describe any system. For a gas, the only two variables that we need to specify are pressure p and volume V: if you know the pressure and volume, then all other quantities — colour, smell, viscosity, thermal conductivity — are fixed. For other systems, further (or different) variables may be needed to describe their macrostate. Common examples are surface tension and area for a film; magnetic field and magnetization for a magnet; electric field and polarization for a dielectric. In what follows we'll assume that we're dealing with a gas and use p and V to specify the state. Everything that we say can be readily extended to more general settings.

So far, we don't have a definition of temperature. This is provided by the zeroth law of thermodynamics which states that equilibrium is a transitive property,

Zeroth Law: If two systems, A and B, are each in equilibrium with a third body C, then they are also in equilibrium with each other

Let's see why this allows us to define the concept of temperature. Suppose that system A is in state (p_1, V_1) and C is in state (p_3, V_3). To test if the two systems are in equilibrium, we need only place them in thermal contact and see if their states change. For generic values of pressure and volume, we will find that the systems are not in equilibrium. Equilibrium requires some relationship between the(p_1, V_1) and (p_3, V_3). For example, suppose that we choose p_1, V_1 and p_3, then there will be a special value of V_3 for which nothing happens when the two systems are brought together.

We'll write the constraint that determines when A and C are in equilibrium as

$$F_{AC}(p_1, V_1; p_3, V_3) = 0$$

which can be solved to give

$$V_3 = f_{AC}(p_1, V_1; p_3)$$

Since systems B and C are also in equilibrium, we also have a constraint,

$$F_{BC}(p_2, V_2; p_3, V_3) = 0 \quad \Rightarrow \quad V_3 = f_{BC}(p_2, V_2; p_3)$$

These two equilibrium conditions give us two different expressions for the volume V_3,

$$f_{AC}(p_1, V_1; p_3) = f_{BC}(p_2, V_2; p_3) \tag{4.1}$$

At this stage we invoke the zeroth law, which tells us that systems A and B must also be in equilibrium, meaning that (4.1) must be equivalent to a constraint

$$F_{AB}(p_1, V_1; p_2, V_2) = 0 \tag{4.2}$$

Equation (4.1) implies (4.2), but the latter does not depend on p_3. That means that p_3 must appear in (4.1) in such a way that it can just be cancelled out on both sides. When this cancellation is performed, (4.1) tells us that there is a relationship between the states of system A and system B.

$$\theta_A(p_1, V_1) = \theta_B(p_2, V_2)$$

The value of $\theta(p, V)$ is called the *temperature* of the system. The function $T = \theta(p, V)$ is called the *equation of state*.

The above argument really only tells us that there exists a property called temperature. There's nothing yet to tell us why we should pick $\theta(p, V)$ as temperature rather than, say $\sqrt{\theta(p, V)}$. We will shortly see that there is, in fact, a canonical choice of temperature that is defined through the second law of thermodynamics and a construct called the Carnot cycle. However, in the meantime it will suffice to simply pick a reference system to define temperature. The standard choice is the ideal gas equation of state (which, as we have seen, is a good approximation to real gases at low densities),

$$T = \frac{pV}{Nk_B}$$

4.2 The First Law

The first law is simply the statement of the conservation of energy, together with the tacit acknowledgement that there's more than one way to change the energy of the system. It is usually expressed as something along the lines of

First Law: The amount of work required to change an *isolated* system from state 1 to state 2 is independent of how the work is performed.

This rather cumbersome sentence is simply telling us that there is another function of state of the system, $E(P, V)$. This is the energy. We could do an amount of work W on an isolated system in any imaginative way we choose: squeeze it, stir it, place a wire and resistor inside with a current passing through it. The method that we choose does not matter: in all cases, the change of the energy is $\Delta E = W$.

However, for systems that are not isolated, the change of energy is *not* equal to the amount of work done. For example, we could take two systems at different temperatures and place them in thermal contact. We needn't do any work, but the energy of each system will change. We're forced to accept that there are ways to change the energy of the system other than by doing work. We write

$$\Delta E = Q + W \tag{4.3}$$

where Q is the amount of energy that was transferred to the system that can't be accounted for by the work done. This transfer of energy arises due to temperature differences. It is called *heat*.

Heat is *not* a type of energy. It is a process — a mode of transfer of energy. There is no sense in which we can divide up the energy $E(p, V)$ of the system into heat and work. We can't write "$E = Q + W$" because neither Q nor W are functions of state.

Quasi-Static Processes

In the discussion above, the transfer of energy can be as violent as you like. There is no need for the system to be in equilibrium while the energy is being added: the first law as expressed in (4.3) refers only to energy at the beginning and end.

From now on, we will be more gentle. We will add or subtract energy to the system very slowly, so that at every stage of the process the system is effectively in equilibrium and can be described by the thermodynamic variables p and V. Such a process is called *quasi-static*.

For quasi-static processes, it is useful to write (4.3) in infinitesimal form. Unfortunately, this leads to a minor notational headache. The problem is that we want to retain the distinction between $E(p, V)$, which is a function of state, and Q and W, which are not. This means that an infinitesimal change in the energy is a total derivative,

$$dE = \frac{\partial E}{\partial p}\, dp + \frac{\partial E}{\partial V}\, dV$$

while an infinitesimal amount of work or heat has no such interpretation: it is merely something small. To emphasise this, it is common to make up some new notation[8]. A small amount of heat is written $đQ$ and a small amount of work is written $đW$. The first law of thermodynamics in infinitesimal form is then

$$dE = đQ + đW \tag{4.4}$$

Although we introduced the first law as applying to all types of work, from now on the discussion is simplest if we just restrict to a single method to applying work to a system: squeezing. We already saw in Section 1 that the infinitesimal work done on a system is

$$đW = -pdV$$

which is the same thing as "force $\times$ distance". Notice the sign convention. When $đW > 0$, we are doing work on the system by squeezing it so that $dV < 0$. However, when the system expands, $dV > 0$ so $đW < 0$ and the system is performing work.

Expressing the work as $đW = -pdV$ also allows us to underline the meaning of the new symbol $đ$. There is no function $W(p, V)$ which has "$dW = -pdV$". (For example, you could try $W = -pV$ but that gives $dW = -pdV - Vdp$ which isn't what we want). The notation $đW$ is there to remind us that work is not an exact differential.

Figure 24:

Suppose now that we vary the state of a system through two different quasi-static paths as shown in the figure. The change in energy is independent of the path taken: it is $\int dE = E(p_2, V_2) - E(p_1, V_1)$. In contrast, the work done $\int đW = -\int pdV$ depends on the path taken. This simple observation will prove important for our next discussion.

[8]In a more sophisticated language, dE, $đW$ and $đQ$ are all one-forms on the state space of the system. dE is exact; $đW$ and $đQ$ are not.

4.3 The Second Law

> "Once or twice I have been provoked and have asked company how many of them could describe the Second Law of Thermodynamics, the law of entropy. The response was cold: it was also negative. Yet I was asking something which is about the scientific equivalent of: 'Have you read a work of Shakespeare?' "
>
> *C.P.Snow (1959)*

C.P. Snow no doubt had in mind the statement that entropy increases. Yet this is a consequence of the second law; it is not the axiom itself. Indeed, we don't yet even have a thermodynamic definition of entropy.

The essence of the second law is that there is a preferred direction of time. There are many macroscopic processes in Nature which cannot be reversed. Things fall apart. The lines on your face only get deeper. Words cannot be unsaid. The second law summarises all such observations in a single statements about the motion of heat.

Reversible Processes

Before we state the second law, it will be useful to first focus on processes which can happily work in both directions of time. These are a special class of quasi-static processes that can be run backwards. They are called *reversible*

A reversible process must lie in equilibrium at each point along the path. This is the quasi-static condition. But now there is the further requirement that there is no friction involved.

For reversible processes, we can take a round trip as shown to the right. Start in state (p_1, V_1), take the lower path to (p_2, V_2) and then the upper path back to (p_1, V_1). The energy is unchanged because $\oint dE = 0$. But the total work done is non-zero: $\oint pdV \neq 0$. By the first law of thermodynamics (4.4), the work performed by the system during the cycle must be equal to the heat absorbed by the system, $\oint \text{đ}Q = \oint pdV$. If we go one way around the cycle, the system does work and absorbs heat from the surroundings; the other way round work in done on the system which then emits energy as heat.

Figure 25:

Processes which move in a cycle like this, returning to their original starting point, are interesting. Run the right way, they convert heat into work. But that's very very useful. The work can be thought of as a piston which can be used to power a steam train. Or a playstation. Or an LHC.

The Statement of the Second Law

The second law is usually expressed in one of two forms. The first tells us when energy can be fruitfully put to use. The second emphasises the observation that there is an arrow of time in the macroscopic world: heat flows from hot to cold. They are usually stated as

Second Law à la Kelvin: No process is possible whose sole effect is to extract heat from a hot reservoir and convert this entirely into work

Second Law à la Clausius: No process is possible whose sole effect is the transfer of heat from a colder to hotter body

It's worth elaborating on the meaning of these statements. Firstly, we all have objects in our kitchens which transfer heat from a cold environment to a hot environment: this is the purpose of a fridge. Heat is extracted from inside the fridge where it's cold and deposited outside where it's warm. Why doesn't this violate Clausius' statement? The reason lies in the words "sole effect". The fridge has an extra effect which is to make your electricity meter run up. In thermodynamic language, the fridge operates because we're supplying it with "work". To get the meaning of Clausius' statement, think instead of a hot object placed in contact with a cold object. Energy always flows from hot to cold; never the other way round.

The statements by Kelvin and Clausius are equivalent. Suppose, for example, that we build a machine that violates Kelvin's statement by extracting heat from a hot reservoir and converting it entirely into work. We can then use this work to power a fridge, extracting heat from a cold source and depositing it back into a hot source. The combination of the two machines then violates Clausius's statement. It is not difficult to construct a similar argument to show that "not Clausius" $\Rightarrow$ "not Kelvin".

Figure 26:

Our goal in this Section is to show how these statements of the second law allow us to define a quantity called "entropy".

Figure 27: The Carnot cycle in cartoon.

4.3.1 The Carnot Cycle

Kelvin's statement of the second law is that we can't extract heat from a hot reservoir and turn it entirely into work. Yet, at first glance, this appears to be in contrast with what we know about reversible cycles. We have just seen that these necessarily have $\oint \text{đ}Q = \oint \text{đ}W$ and so convert heat to work. Why is this not in contradiction with Kelvin's statement?

The key to understanding this is to appreciate that a reversible cycle does more than just extract heat from a hot reservoir. It also, by necessity, deposits some heat elsewhere. The energy available for work is the difference between the heat extracted and the heat lost. To illustrate this, it's very useful to consider a particular kind of reversible cycle called a *Carnot engine*. This is series of reversible processes, running in a cycle, operating between two temperatures T_H and T_C. It takes place in four stages shown in cartoon in Figures 27 and 28.

- Isothermal expansion AB at a constant hot temperature T_H. The gas pushes against the side of its container and is allowed to slowly expand. In doing so, it can be used to power your favourite electrical appliance. To keep the temperature constant, the system will need to absorb an amount of heat Q_H from the surroundings

- Adiabatic expansion BC. The system is now isolated, so no heat is absorbed. But the gas is allowed to continue to expand. As it does so, both the pressure and temperature will decrease.

- Isothermal contraction CD at constant temperature T_C. We now start to restore the system to its original state. We do work on the system by compressing the

Figure 28: The Carnot cycle, shown in the $p - V$ plane and the $T - S$ plane.

gas. If we were to squeeze an isolated system, the temperature would rise. But we keep the system at fixed temperature, so it dumps heat Q_C into its surroundings.

- Adiabatic contraction DA. We isolate the gas from its surroundings and continue to squeeze. Now the pressure and temperature both increase. We finally reach our starting point when the gas is again at temperature T_H.

At the end of the four steps, the system has returned to its original state. The net heat absorbed is $Q_H - Q_C$ which must be equal to the work performed by the system W. We define the *efficiency* η of an engine to be the ratio of the work done to the heat absorbed from the hot reservoir,

$$\eta = \frac{W}{Q_H} = \frac{Q_H - Q_C}{Q_H} = 1 - \frac{Q_C}{Q_H}$$

Ideally, we would like to take all the heat Q_H and convert it to work. Such an engine would have efficiency $\eta = 1$ but would be in violation of Kelvin's statement of the second law. We can see the problem in the Carnot cycle: we have to deposit some amount of heat Q_C back to the cold reservoir as we return to the original state. And the following result says that the Carnot cycle is the best we can do:

Carnot's Theorem: Carnot is the best. Or, more precisely: Of all engines operating between two heat reservoirs, a reversible engine is the most efficient. As a simple corollary, all reversible engines have the same efficiency which depends only on the temperatures of the reservoirs $\eta(T_H, T_C)$.

Proof: Let's consider a second engine — call it Ivor — operating between the same two temperatures T_H and T_C. Ivor also performs work W but, in contrast to Carnot, is not reversible. Suppose that Ivor absorbs Q'_H from the hot reservoir and deposits Q'_C into the cold. Then we can couple Ivor to our original Carnot engine set to reverse.

Figure 29:

The work W performed by Ivor now goes into driving Carnot. The net effect of the two engines is to extract $Q'_H - Q_H$ from the hot reservoir and, by conservation of energy, to deposit the same amount $Q'_C - Q_C = Q'_H - Q_H$ into the cold. But Clausius's statement tells us that we must have $Q'_H \geq Q_H$; if this were not true, energy would be moved from the colder to hotter body. Performing a little bit of algebra then gives

$$Q'_C - Q'_H = Q_C - Q_H \quad \Rightarrow \quad \eta_{\text{Ivor}} = 1 - \frac{Q'_C}{Q'_H} = \frac{Q_H - Q_C}{Q'_H} \leq \frac{Q_H - Q_C}{Q_H} = \eta_{\text{Carnot}}$$

The upshot of this argument is the result that we wanted, namely

$$\eta_{\text{Carnot}} \geq \eta_{\text{Ivor}}$$

The corollary is now simple to prove. Suppose that Ivor was reversible after all. Then we could use the same argument above to prove that $\eta_{\text{Ivor}} \geq \eta_{\text{Carnot}}$, so it must be true that $\eta_{\text{Ivor}} = \eta_{\text{Carnot}}$ if Ivor is reversible. This means that for all reversible engines operating between T_H and T_C have the same efficiency. Or, said another way, the ratio Q_H/Q_C is the same for all reversible engines. Moreover, this efficiency must be a function only of the temperatures, $\eta_{\text{Carnot}} = \eta(T_H, T_C)$, simply because they are the only variables in the game. $\square$.

4.3.2 Thermodynamic Temperature Scale and the Ideal Gas

Recall that the zeroth law of thermodynamics showed that there was a function of state, which we call temperature, defined so that it takes the same value for any two systems in equilibrium. But at the time there was no canonical way to decide between different definitions of temperature: $\theta(p, V)$ or $\sqrt{\theta(p, V)}$ or any other function are all equally good choices. In the end we were forced to pick a reference system — the ideal gas — as a benchmark to define temperature. This was a fairly arbitrary choice. We can now do better.

Since the efficiency of a Carnot engine depends only on the temperatures T_H and T_C, we can use this to define a temperature scale that is independent of any specific material. (Although, as we shall see, the resulting temperature scale turns out to be equivalent to the ideal gas temperature). Let's now briefly explain how we can define a temperature through the Carnot cycle.

The key idea is to consider two Carnot engines. The first operates between two temperature reservoirs $T_1 > T_2$; the second engine operates between two reservoirs $T_2 > T_3$. If the first engine extracts heat Q_1 then it must dump heat Q_2 given by

$$Q_2 = Q_1 \left(1 - \eta(T_1, T_2)\right)$$

where the arguments above tell us that $\eta = \eta_{\text{Carnot}}$ is a function only of T_1 and T_2. If the second engine now takes this same heat Q_2, it must dump heat Q_3 into the reservoir at temperature T_3, given by

$$Q_3 = Q_2 \left(1 - \eta(T_2, T_3)\right) = Q_1 \left(1 - \eta(T_1, T_2)\right) \left(1 - \eta(T_2, T_3)\right)$$

But we can also consider both engines working together as a Carnot engine in its own right, operating between reservoirs T_1 and T_3. Such an engine extracts heat Q_1, dumps heat Q_3 and has efficiency $\eta(T_1, T_3)$, so that

$$Q_3 = Q_1 \left(1 - \eta(T_1, T_3)\right)$$

Combining these two results tells us that the efficiency must be a function which obeys the equation

$$1 - \eta(T_1, T_3) = \left(1 - \eta(T_1, T_2)\right) \left(1 - \eta(T_2, T_3)\right)$$

The fact that T_2 has to cancel out on the right-hand side is enough to tell us that

$$1 - \eta(T_1, T_2) = \frac{f(T_2)}{f(T_1)}$$

for some function $f(T)$. At this point, we can use the ambiguity in the definition of temperature to simply pick a nice function, namely $f(T) = T$. Hence, we *define* the thermodynamic temperature to be such that the efficiency is given by

$$\eta = 1 - \frac{T_2}{T_1} \tag{4.5}$$

The Carnot Cycle for an Ideal Gas

We now have two ways to specify temperature. The first arises from looking at the equation of state of a specific, albeit simple, system: the ideal gas. Here temperature is defined to be $T = pV/Nk_B$. The second definition of temperature uses the concept of Carnot cycles. We will now show that, happily, these two definitions are equivalent by explicitly computing the efficiency of a Carnot engine for the ideal gas.

We deal first with the isothermal changes of the ideal gas. We know that the energy in the gas depends only on the temperature[9],

$$E = \frac{3}{2}Nk_BT \tag{4.6}$$

So $dT = 0$ means that $dE = 0$. The first law then tells us that $đQ = -đW$. For the motion along the line AB in the Carnot cycle, we have

$$Q_H = \int_A^B đQ = -\int_A^B đW = \int_A^B pdV = \int_A^B \frac{Nk_BT_H}{V}dV = Nk_BT_H \log\left(\frac{V_B}{V_A}\right) \tag{4.7}$$

Similarly, the heat given up along the line CD in the Carnot cycle is

$$Q_C = -Nk_BT_C \log\left(\frac{V_D}{V_C}\right) \tag{4.8}$$

Next we turn to the adiabatic change in the Carnot cycle. Since the system is isolated, $đQ = 0$ and all work goes into the energy, $dE = -pdV$. Meanwhile, from (4.6), we can write the change of energy as $dE = C_VdT$ where $C_V = \frac{3}{2}Nk_B$, so

$$C_VdT = -\frac{Nk_BT}{V}dV \quad \Rightarrow \quad \frac{dT}{T} = -\left(\frac{Nk_B}{C_V}\right)\frac{dV}{V}$$

After integrating, we have

$$TV^{2/3} = \text{constant}$$

[9] A confession: strictly speaking, I'm using some illegal information in the above argument. The result $E = \frac{3}{2}Nk_BT$ came from statistical mechanics and if we're really pretending to be Victorian scientists we should discuss the efficiency of the Carnot cycle without this knowledge. Of course, we could just measure the heat capacity $C_V = \partial E/\partial T|_V$ to determine $E(T)$ experimentally and proceed. Alternatively, and more mathematically, we could note that it's not necessary to use this exact form of the energy to carry through the argument: we need only use the fact that the energy is a function of temperature only: $E = E(T)$. The isothermal parts of the Carnot cycle are trivially the same and we reproduce (4.7) and (4.8). The adiabatic parts cannot be solved exactly without knowledge of $E(T)$ but you can still show that $V_A/V_B = V_D/V_C$ which is all we need to derive the efficiency (4.9).

Applied to the line BC and DA in the Carnot cycle, this gives

$$T_H V_B^{2/3} = T_C V_C^{2/3} \quad , \quad T_C V_D^{2/3} = T_H V_A^{2/3}$$

which tells us that $V_A/V_B = V_D/V_C$. But this means that the factors of $\log(V/V)$ cancel when we take the ratio of heats. The efficiency of a Carnot engine for an ideal gas — and hence for any system — is given by

$$\eta_{\text{carnot}} = 1 - \frac{Q_C}{Q_H} = 1 - \frac{T_C}{T_H} \tag{4.9}$$

We see that the efficiency using the ideal gas temperature coincides with our thermodynamic temperature (4.5) as advertised.

4.3.3 Entropy

The discussion above was restricted to Carnot cycles: reversible cycles operating between two temperatures. The second law tells us that we can't turn all the extracted heat into work. We have to give some back. To generalize, let's change notation slightly so that Q always denotes the energy absorbed by the system. If the system releases heat, then Q is negative. In terms of our previous notation, $Q_1 = Q_H$ and $Q_2 = -Q_C$. Similarly, $T_1 = T_H$ and $T_2 = T_C$. Then, for all Carnot cycles

Figure 30:

$$\sum_{i=1}^{2} \frac{Q_i}{T_i} = 0$$

Now consider the reversible cycle shown in the figure in which we cut the corner of the original cycle. From the original Carnot cycle $ABCD$, we know that

$$\frac{Q_{AB}}{T_H} + \frac{Q_{CD}}{T_C} = 0$$

Meanwhile, we can view the square $EBGF$ as a mini-Carnot cycle so we also have

$$\frac{Q_{GF}}{T_{FG}} + \frac{Q_{EB}}{T_H} = 0$$

What if we now run along the cycle $AEFGCD$? Clearly $Q_{AB} = Q_{AE} + Q_{EB}$. But we also know that the heat absorbed along the segment FG is equal to that dumped along

the segment GF when we ran the mini-Carnot cycle. This follows simply because we're taking the same path but in the opposite direction and tells us that $Q_{FG} = -Q_{GF}$. Combining these results with the two equations above gives us

$$\frac{Q_{AE}}{T_H} + \frac{Q_{FG}}{T_{FG}} + \frac{Q_{CD}}{T_C} = 0$$

By cutting more and more corners, we can consider any reversible cycle as constructed of (infinitesimally) small isothermal and adiabatic segments. Summing up all contributions Q/T along the path, we learn that the total heat absorbed in any reversible cycle must obey

$$\oint \frac{đQ}{T} = 0$$

But this is a very powerful statement. It means that if we reversibly change our system from state A to state B, then the quantity $\int_A^B đQ/T$ is independent of the path taken. Either of the two paths shown in the figure will give the same result:

$$\int_{\text{Path I}} \frac{đQ}{T} = \int_{\text{Path II}} \frac{đQ}{T}$$

Figure 31:

Given some reference state O, this allows us to define a new function of state. It is called *entropy*, S

$$S(A) = \int_0^A \frac{đQ}{T} \tag{4.10}$$

Entropy depends only on the state of the system: $S = S(p, V)$. It does not depend on the path we took to get to the state. We don't even have to take a reversible path: as long as the system is in equilibrium, it has a well defined entropy (at least relative to some reference state).

We have made no mention of microstates in defining the entropy. Yet it is clearly the same quantity that we met in Section 1. From (4.10), we can write $dS = đQ/T$, so that the first law of thermodynamics (4.4) is written in the form of (1.16)

$$dE = TdS - pdV \tag{4.11}$$

Irreversibility

What can we say about paths that are not reversible? By Carnot's theorem, we know that an irreversible engine that operates between two temperatures T_H and T_C is less efficient than the Carnot cycle. We use the same notation as in the proof of Carnot's theorem; the Carnot engine extracts heat Q_H and dumps heat Q_C; the irreversible engine extracts heat Q'_H and dumps Q'_C. Both do the same amount of work $W = Q_H - Q_C = Q'_H - Q'_C$. We can then write

$$\begin{aligned}\frac{Q'_H}{T_H} - \frac{Q'_C}{T_C} &= \frac{Q_H}{T_H} - \frac{Q_C}{T_C} + (Q'_H - Q_H)\left(\frac{1}{T_H} - \frac{1}{T_C}\right) \\ &= (Q'_H - Q_H)\left(\frac{1}{T_H} - \frac{1}{T_C}\right) \leq 0\end{aligned}$$

In the second line, we used $Q_H/T_H = Q_C/T_C$ for a Carnot cycle, and to derive the inequality we used the result of Carnot's theorem, namely $Q'_H \geq Q_H$ (together with the fact that $T_H > T_C$).

The above result holds for any engine operating between two temperatures. But by the same method of cutting corners off a Carnot cycle that we used above, we can easily generalise the statement to any path, reversible or irreversible. Putting the minus signs back in so that heat dumped has the opposite sign to heat absorbed, we arrive at a result is known as the *Clausius inequality*,

$$\oint \frac{đQ}{T} \leq 0$$

We can express this in slightly more familiar form. Suppose that we have two possible paths between states A and B as shown in the figure. Path I is irreversible while path II is reversible. Then Clausius's inequality tells us that

Figure 32:

$$\oint \frac{đQ}{T} = \int_I \frac{đQ}{T} - \int_{II} \frac{đQ}{T} \leq 0$$

$$\Rightarrow \quad \int_I \frac{đQ}{T} \leq S(B) - S(A) \tag{4.12}$$

Suppose further that path I is adiabatic, meaning that it is isolated from the environment. Then $đQ = 0$ and we learn that the entropy of any isolated system never decreases,

$$S(B) \geq S(A) \tag{4.13}$$

Moreover, if an adiabatic process is reversible, then the resulting two states have equal entropy.

The second law, as expressed in (4.13), is responsible for the observed arrow of time in the macroscopic world. Isolated systems can only evolve to states of equal or higher entropy. This coincides with the statement of the second law that we saw back in Section 1.2.1 using Boltzmann's definition of the entropy.

4.3.4 Adiabatic Surfaces

The primary consequence of the second law is that there exists a new function of state, entropy. Surfaces of constant entropy are called *adiabatic surfaces*. The states that sit on a given adiabatic surface can all be reached by performing work on the system while forbidding any heat to enter or leave. In other words, they are the states that can be reached by adiabatic processes with $đQ = 0$ which is equivalent to $dS = 0$.

In fact, for the simplest systems such as the ideal gas which require only two variables p and V to specify the state, we do not need the second law to infer to the existence of an adiabatic surface. In that case, the adiabatic surface is really an adiabatic line in the two-dimensional space of states. The existence of this line follows immediately from the first law. To see this, we write the change of energy for an adiabatic process using (4.4) with $đQ = 0$,

$$dE + pdV = 0 \tag{4.14}$$

Let's view the energy itself as a function of p and V so that we can write

$$dE = \frac{\partial E}{\partial p} dP + \frac{\partial E}{\partial V} dV$$

Then the condition for an adiabatic process (4.14) becomes

$$\frac{\partial E}{\partial p} dp + \left(\frac{\partial E}{\partial V} + p \right) dV = 0$$

Which tells us the slope of the adiabatic line is given by

$$\frac{dp}{dV} = - \left(\frac{\partial E}{\partial V} + p \right) \left(\frac{\partial E}{\partial p} \right)^{-1} \tag{4.15}$$

The upshot of this calculation is that if we sit in a state specified by (p, V) and transfer work but no heat to the system then we necessarily move along a line in the space of states determined by (4.15). If we want to move off this line, then we have to add heat to the system.

However, the analysis above does not carry over to more complicated systems where more than two variables are needed to specify the state. Suppose that the state of the system is specified by three variables. The first law of thermodynamics is now gains an extra term, reflecting the fact that there are more ways to add energy to the system,

$$dE = \text{đ}Q - pdV - ydX$$

We've already seen examples of this with $y = -\mu$, the chemical potential, and $X = N$, the particle number. Another very common example is $y = -M$, the magnetization, and $X = H$, the applied magnetic field. For our purposes, it won't matter what variables y and X are: just that they exist. We need to choose three variables to specify the state. Any three will do, but we will choose p, V and X and view the energy as a function of these: $E = E(p, V, X)$. An adiabatic process now requires

$$dE + pdV + ydX = 0 \quad \Rightarrow \quad \frac{\partial E}{\partial p}dp + \left(\frac{\partial E}{\partial V} + p\right)dV + \left(\frac{\partial E}{\partial X} + y\right)dX = 0 \tag{4.16}$$

But this equation does not necessarily specify a surface in $\mathbf{R}^3$. To see that this is not sufficient, we can look at some simple examples. Consider $\mathbf{R}^3$, parameterised by z_1, z_2 and z_3. If we move in an infinitesimal direction satisfying

$$z_1 dz_1 + z_2 dz_2 + z_3 dz_3 = 0$$

then it is simple to see that we can integrate this equation to learn that we are moving on the surface of a sphere,

$$z_1^2 + z_2^2 + z_3^2 = \text{constant}$$

In contrast, if we move in an infinitesimal direction satisfying the condition

$$z_2 dz_1 + dz_2 + dz_3 = 0 \tag{4.17}$$

Then there is no associated surface on which we're moving. Indeed, you can convince yourself that if you move in a such a way as to always obey (4.17) then you can reach any point in $\mathbf{R}^3$ from any other point.

In general, an infinitesimal motion in the direction

$$Z_1 dz_1 + Z_2 dz_2 + Z_3 dz_3 = 0$$

has the interpretation of motion on a surface only if the functions Z_i obey the condition

$$Z_1\left(\frac{\partial Z_2}{\partial z_3} - \frac{\partial Z_3}{\partial z_2}\right) + Z_2\left(\frac{\partial Z_3}{\partial z_1} - \frac{\partial Z_1}{\partial z_3}\right) + Z_3\left(\frac{\partial Z_1}{\partial z_2} - \frac{\partial Z_2}{\partial z_1}\right) = 0 \tag{4.18}$$

So for systems with three or more variables, the existence of an adiabatic surface is not guaranteed by the first law alone. We need the second law. This ensures the existence of a function of state S such that adiabatic processes move along surfaces of constant S. In other words, the second law tells us that (4.16) satisfies (4.18).

In fact, there is a more direct way to infer the existence of adiabatic surfaces which uses the second law but doesn't need the whole rigmarole of Carnot cycles. We will again work with a system that is specified by three variables, although the argument will hold for any number. But we choose our three variables to be V, X and the internal energy E. We start in state A shown in the figure. We will show that Kelvin's statement of the second law implies that it is not possible to reach both states B and C through reversible adiabatic processes. The key feature of these states is that they have the same values of V and X and differ only in their energy E.

Figure 33:

To prove this statement, suppose the converse: i.e. we can indeed reach both B and C from A through means of reversible adiabatic processes. Then we can start at A and move to B. Since the energy is lowered, the system performs work along this path but, because the path is adiabatic, no heat is exchanged. Now let's move from B to C. Because $dV = dX = 0$ on this trajectory, we do no work but the internal energy E changes so the system must absorb heat Q from the surroundings. Now finally we do work on the system to move from C back to A. However, unlike in the Carnot cycle, we don't return any heat to the environment on this return journey because, by assumption, this second path is also adiabatic. The net result is that we have extracted heat Q and employed this to undertake work $W = Q$. This is in contradiction with Kelvin's statement of the second law.

The upshot of this argument is that the space of states can be foliated by adiabatic surfaces such that each vertical line at constant V and X intersects the surface only once. We can describe these surfaces by some function $S(E, V, X) =$ constant. This function is the entropy.

The above argument shows that Kelvin's statement of the second law implies the existence of adiabatic surfaces. One may wonder if we can run the argument the other way around and use the existence of adiabatic surfaces as the basis of the second law, dispensing with the Kelvin and Clausius postulates all together. In fact, we can almost do this. From the discussion above it should already be clear that the existence of

adiabatic surfaces implies that the addition of heat is proportional to the change in entropy $đQ \sim dS$. However, it remains to show that the integrating factor, relating the two, is temperature so $đQ = TdS$. This can be done by returning to the zeroth law of thermodynamics. A fairly simple description of the argument can be found at the end of Chapter 4 of Pippard's book. This motivates a mathematically concise statement of the second law due to Carathéodory.

Second Law à la Carathéodory: Adiabatic surfaces exist. Or, more poetically: if you want to be able to return, there are places you cannot go through work alone. Sometimes you need a little heat.

What this statement is lacking is perhaps the most important aspect of the second law: an arrow of time. But this is easily remedied by providing one additional piece of information telling us which side of a surface can be reached by irreversible processes. To one side of the surface lies the future, to the other the past.

4.3.5 A History of Thermodynamics

The history of heat and thermodynamics is long and complicated, involving wrong turns, insights from disparate areas of study such as engineering and medicine, and many interesting characters, more than one of which find reason to change their name at some point in the story[10].

Although ideas of "heat" date back to pre-history, a good modern starting point is the 1787 *caloric* theory of Lavoisier. This postulates that heat is a conserved fluid which has a tendency to repel itself, thereby flowing from hot bodies to cold bodies. It was, for its time, an excellent theory, explaining many of the observed features of heat. Of course, it was also wrong.

Lavoisier's theory was still prominent 30 years later when the French engineer Sadi Carnot undertook the analysis of steam engines that we saw above. Carnot understood all of his processes in terms of caloric. He was inspired by mechanics of waterwheels and saw the flow of caloric from hot to cold bodies as analogous to the fall of water from high to low. This work was subsequently extended and formalised in a more mathematical framework by another French physicist, Émile Clapeyron. By the 1840s, the properties of heat were viewed by nearly everyone through the eyes of Carnot-Clapeyron caloric theory.

[10] A longer description of the history of heat can be found in Michael Fowler's lectures from the University of Virginia: http://galileo.phys.virginia.edu/classes/152.mf1i.spring02/HeatIndex.htm

Yet the first cracks in caloric theory has already appeared before the turn of 19th century due to the work of Benjamin Thompson. Born in the English colony of Massachusetts, Thompson's CV includes turns as mercenary, scientist and humanitarian. He is the inventor of thermal underwear and pioneer of the soup kitchen for the poor. By the late 1700s, Thompson was living in Munich under the glorious name "Count Rumford of the Holy Roman Empire" where he was charged with overseeing artillery for the Prussian Army. But his mind was on loftier matters. When boring cannons, Rumford was taken aback by the amount of heat produced by friction. According to Lavoisier's theory, this heat should be thought of as caloric fluid squeezed from the brass cannon. Yet is seemed inexhaustible: when a cannon was bored a second time, there was no loss in its ability to produce heat. Thompson/Rumford suggested that the cause of heat could not be a conserved caloric. Instead he attributed heat correctly, but rather cryptically, to "motion".

Having put a big dent in Lavoisier's theory, Rumford rubbed salt in the wound by marrying his widow. Although, in fairness, Lavoisier was beyond caring by this point. Rumford was later knighted by Britain, reverting to Sir Benjamin Thompson, where he founded the Royal Institution.

The journey from Thompson's observation to an understanding of the first law of thermodynamics was a long one. Two people in particular take the credit.

In Manchester, England, James Joule undertook a series of extraordinarily precise experiments. He showed how different kinds of work — whether mechanical or electrical – could be used to heat water. Importantly, the amount by which the temperature was raised depended only on the amount of work, not the manner in which it was applied. His 1843 paper "The Mechanical Equivalent of Heat" provided compelling quantitative evidence that work could be readily converted into heat.

But Joule was apparently not the first. A year earlier, in 1842, the German physician Julius von Mayer came to the same conclusion through a very different avenue of investigation: blood letting. Working on a ship in the Dutch East Indies, Mayer noticed that the blood in sailors veins was redder in Germany. He surmised that this was because the body needed to burn less fuel to keep warm. Not only did he essentially figure out how the process of oxidation is responsible for supplying the body's energy but, remarkably, he was able to push this to an understanding of how work and heat are related. Despite limited physics training, he used his intuition, together with known experimental values of the heat capacities C_p and C_V of gases, to determine essentially the same result as Joule had found through more direct means.

The results of Thompson, Mayer and Joule were synthesised in an 1847 paper by Hermann von Helmholtz, who is generally credited as the first to give a precise formulation of the first law of thermodynamics. (Although a guy from Swansea called William Grove has a fairly good, albeit somewhat muddled, claim from a few years earlier). It's worth stressing the historical importance of the first law: this was the first time that the conservation of energy was elevated to a key idea in physics. Although it had been known for centuries that quantities such as "$\frac{1}{2}mv^2 + V$" were conserved in certain mechanical problems, this was often viewed as a mathematical curiosity rather than a deep principle of Nature. The reason, of course, is that friction is important in most processes and energy does not appear to be conserved. The realisation that there is a close connection between energy, work and heat changed this. However, it would still take more than half a century before Emmy Noether explained the true reason behind the conservation of tenergy.

With Helmholtz, the first law was essentially nailed. The second remained. This took another two decades, with the pieces put together by a number of people, notably William Thomson and Rudolph Clausius.

William Thomson was born in Belfast but moved to Glasgow at the age of 10. He came to Cambridge to study, but soon returned to Glasgow and stayed there for the rest of his life. After his work as a scientist, he gained fame as an engineer, heavily involved in laying the first trans-atlantic cables. For this he was made Lord Kelvin, the name chosen for the River Kelvin which flows nearby Glasgow University. He was the first to understand the importance of absolute zero and to define the thermodynamic temperature scale which now bears his favourite river's name. We presented Kelvin's statement of the second law of thermodynamics earlier in this Section.

In Germany, Rudolph Clausius was developing the same ideas as Kelvin. But he managed to go further and, in 1865, presented the subtle thermodynamic argument for the existence of entropy that we saw in Section 4.3.3. Modestly, Clausius introduced the unit "Clausius" (symbol **Cl**) for entropy. It didn't catch on.

4.4 Thermodynamic Potentials: Free Energies and Enthalpy

We now have quite a collection of thermodynamic variables. The state of the system is dictated by pressure p and volume V. From these, we can define temperature T, energy E and entropy S. We can also mix and match. The state of the system can just as well be labelled by T and V; or E and V; or T and p; or V and S ...

While we're at liberty to pick any variables we like, certain quantities are more naturally expressed in terms of some variables instead of others. We've already seen examples both in Section 1 and in this section. If we're talking about the energy E, it is best to label the state in terms of S and V, so $E = E(S, V)$. In these variables the first law has the nice form (4.11).

Equivalently, inverting this statement, the entropy should be thought of as a function of E and V, so $S = S(E, V)$. It is not just mathematical niceties underlying this: it has physical meaning too for, as we've seen above, at fixed energy the second law tells us that entropy can never decrease.

What is the natural object to consider at constant temperature T, rather than constant energy? In fact we already answered this way back in Section 1.3 where we argued that one should minimise the Helmholtz free energy,

$$F = E - TS$$

The arguments that we made back in Section 1.3 were based on a microscopic viewpoint of entropy. But, with our thermodynamic understanding of the second law, we can easily now repeat the argument without mention of probability distributions. We consider our system in contact with a heat reservoir such that the total energy, E_{total} of the combined system and reservoir is fixed. The combined entropy is then,

$$\begin{aligned} S_{\text{total}}(E_{\text{total}}) &= S_R(E_{\text{total}} - E) + S(E) \\ &\approx S_R(E_{\text{total}}) - \frac{\partial S_R}{\partial E_{\text{total}}} E + S(E) \\ &= S_R(E_{\text{total}}) - \frac{F}{T} \end{aligned}$$

The total entropy can never decrease; the free energy of the system can never increase.

One interesting situation that we will meet in the next section is a system which, at fixed temperature and volume, has two different equilibrium states. Which does it choose? The answer is the one that has lower free energy, for random thermal fluctuations will tend to take us to this state, but very rarely bring us back.

We already mentioned in Section 1.3 that the free energy is a Legendre transformation of the energy; it is most naturally thought of as a function of T and V, which is reflected in the infinitesimal variation,

$$dF = -SdT - pdV \quad \Rightarrow \quad \left.\frac{\partial F}{\partial T}\right|_V = -S \quad , \quad \left.\frac{\partial F}{\partial V}\right|_T = -p \tag{4.19}$$

We can now also explain what's free about this energy. Consider taking a system along a reversible isotherm, from state A to state B. Because the temperature is constant, the change in free energy is $dF = -pdV$, so

$$F(B) - F(A) = -\int_A^B pdV = -W$$

where W is the work done by the system. The free energy is a measure of the amount of energy free to do work at finite temperature.

Gibbs Free Energy

We can also consider systems that don't live at fixed volume, but instead at fixed pressure. To do this, we will once again imagine the system in contact with a reservoir at temperature T. The volume of each can fluctuate, but the total volume V_{total} of the combined system and reservoir is fixed. The total entropy is

$$\begin{aligned} S_{\text{total}}(E_{\text{total}}, V_{\text{total}}) &= S_R(E_{\text{total}} - E, V_{\text{total}} - V) + S(E, V) \\ &\approx S_R(E_{\text{total}}, V_{\text{total}}) - \frac{\partial S_R}{\partial E_{\text{total}}} E - \frac{\partial S_R}{\partial V_{\text{total}}} V + S(E, V) \\ &= S_R(V_{\text{total}}) - \frac{E + pV - TS}{T} \end{aligned}$$

At fixed temperature and pressure we should minimise the *Gibbs Free Energy*,

$$G = F + pV = E + pV - TS \tag{4.20}$$

This is a Legendre transform of F, this time swapping volume for pressure: $G = G(T, p)$. The infinitesimal variation is

$$dG = -SdT + Vdp$$

In our discussion we have ignored the particle number N. Yet both F and G implicitly depend on N (as you may check by re-examining the many examples of F that we computed earlier in the course). If we also consider changes dN then each variations gets the additional term μdN, so

$$dF = -SdT - pdV + \mu dN \quad \text{and} \quad dG = -SdT + Vdp + \mu dN \tag{4.21}$$

While F can have an arbitrarily complicated dependence on N, the Gibbs free energy G has a very simple dependence. To see this, we simply need to look at the extensive properties of the different variables and make the same kind of argument that we've already seen in Section 1.4.1. From its definition (4.20), we see that the Gibbs free

energy G is extensive. It a function of p, T and N, of which only N is extensive. Therefore,

$$G(p,T,N) = \mu(p,T)N \tag{4.22}$$

where the fact that the proportionality coefficient is μ follows from variation (4.21) which tells us that $\partial G/\partial N = \mu$.

The Gibbs free energy is frequently used by chemists, for whom reactions usually take place at constant pressure rather than constant volume. (When a chemist talks about "free energy", they usually mean G. For a physicist, "free energy" usually means F). We'll make use of the result (4.22) in the next section when we discuss first order phase transitions.

4.4.1 Enthalpy

There is one final combination that we can consider: systems at fixed energy and pressure. Such systems are governed by the *enthalpy*,

$$H = E + pV \quad \Rightarrow \quad dH = TdS + Vdp$$

The four objects E, F, G and H are sometimes referred to as *thermodynamic potentials*.

4.4.2 Maxwell's Relations

Each of the thermodynamic potentials has an interesting present for us. Let's start by considering the energy. Like any function of state, it can be viewed as a function of any of the other two variables which specify the system. However, the first law of thermodynamics (4.11) suggests that it is most natural to view energy as a function of entropy and volume: $E = E(S,V)$. This has the advantage that the partial derivatives are familiar quantities,

$$\left.\frac{\partial E}{\partial S}\right|_V = T \quad , \quad \left.\frac{\partial E}{\partial V}\right|_S = -p$$

We saw both of these results in Section 1. It is also interesting to look at the double mixed partial derivative, $\partial^2 E/\partial S\partial V = \partial^2 E/\partial V\partial S$. This gives the relation

$$\left.\frac{\partial T}{\partial V}\right|_S = -\left.\frac{\partial p}{\partial S}\right|_V \tag{4.23}$$

This result is mathematically trivial. Yet physically it is far from obvious. It is the first of four such identities, known as the *Maxwell Relations*.

The other Maxwell relations are derived by playing the same game with F, G and H. From the properties (4.19), we see that taking mixed partial derivatives of the free energy gives us,

$$\left.\frac{\partial S}{\partial V}\right|_T = \left.\frac{\partial p}{\partial T}\right|_V \tag{4.24}$$

The Gibbs free energy gives us

$$\left.\frac{\partial S}{\partial p}\right|_T = -\left.\frac{\partial V}{\partial T}\right|_p \tag{4.25}$$

While the enthalpy gives

$$\left.\frac{\partial T}{\partial p}\right|_S = \left.\frac{\partial V}{\partial S}\right|_p \tag{4.26}$$

The four Maxwell relations (4.23), (4.24), (4.25) and (4.26) are remarkable in that they are mathematical identities that hold for any system. They are particularly useful because they relate quantities which are directly measurable with those which are less easy to determine experimentally, such as entropy.

It is not too difficult to remember the Maxwell relations. Cross-multiplication always yields terms in pairs: TS and pV, which follows essentially on dimensional grounds. The four relations are simply the four ways to construct such equations. The only tricky part is to figure out the minus signs.

Heat Capacities Revisted

By taking further derivatives of the Maxwell relations, we can derive yet more equations which involve more immediate quantities. You will be asked to prove a number of these on the examples sheet, including results for the heat capacity at constant volume, $C_V = T\,\partial S/\partial T|_V$, as well as the heat capacity at capacity at constant pressure $C_p = T\,\partial S/\partial T|_p$. Useful results include,

$$\left.\frac{\partial C_V}{\partial V}\right|_T = T\left.\frac{\partial^2 p}{\partial T^2}\right|_V \quad , \quad \left.\frac{\partial C_p}{\partial p}\right|_T = -T\left.\frac{\partial^2 V}{\partial T^2}\right|_p$$

You will also prove a relationship between these two heat capacities,

$$C_p - C_V = T\left.\frac{\partial V}{\partial T}\right|_p \left.\frac{\partial p}{\partial T}\right|_V$$

This last expression has a simple consequence. Consider, for example, an ideal gas obeying $pV = Nk_BT$. Evaluating the right-hand side gives us

$$C_p - C_v = Nk_B$$

There is an intuitive reason why C_p is greater than C_V. At constant volume, if you dump heat into a system then it all goes into increasing the temperature. However, at constant pressure some of this energy will cause the system to expand, thereby doing work. This leaves less energy to raise the temperature, ensuring that $C_p > C_V$.

4.5 The Third Law

The second law only talks about entropy differences. We can see this in (4.10) where the entropy is defined with respect to some reference state. The third law, sometimes called *Nernst's postulate*, provides an absolute scale for the entropy. It is usually taken to be

$$\lim_{T\to 0} S(T) = 0$$

In fact we can relax this slightly to allow a finite entropy, but vanishing entropy density S/N. We know from the Boltzmann definition that, at $T = 0$, the entropy is simply the logarithm of the degeneracy of the ground state of the system. The third law really requires $S/N \to 0$ as $T \to 0$ and $N \to \infty$. This then says that the ground state entropy shouldn't grow extensively with N.

The third law doesn't quite have the same teeth as its predecessors. Each of the first three laws provided us with a new function of state of the system: the zeroth law gave us temperature; the first law energy; and the second law entropy. There is no such reward from the third law.

One immediate consequence of the third law is that heat capacities must also tend to zero as $T \to 0$. This follows from the equation (1.10)

$$S(B) - S(A) = \int_A^B dT\, \frac{C_V}{T}$$

If the entropy at zero temperature is finite then the integral must converge which tells us that $C_V \to T^n$ for some $n \geq 1$ or faster. Looking back at the various examples of heat capacities, we can check that this is always true. (The case of a degenerate Fermi gas is right on the borderline with $n = 1$). However, in each case the fact that the heat capacity vanishes is due to quantum effects freezing out degrees of freedom.

In contrast, when we restrict to the classical physics, we typically find constant heat capacities such as the classical ideal gas (2.10) or the Dulong-Petit law (3.15). These would both violate the third law. In that sense, the third law is an admission that the low temperature world is not classical. It is quantum.

Thinking about things quantum mechanically, it is very easy to see why the third law holds. A system that violates the third law would have a very large – indeed, an extensive – number of ground states. But large degeneracies do not occur naturally in quantum mechanics. Moreover, even if you tune the parameters of a Hamiltonian so that there are a large number of ground states, then any small perturbation will lift the degeneracy, introducing an energy splitting between the states. From this perspective, the third law is a simple a consequence of the properties of the eigenvalue problem for large matrices.

5. Phase Transitions

A *phase transition* is an abrupt, discontinuous change in the properties of a system. We've already seen one example of a phase transition in our discussion of Bose-Einstein condensation. In that case, we had to look fairly closely to see the discontinuity: it was lurking in the derivative of the heat capacity. In other phase transitions — many of them already familiar — the discontinuity is more manifest. Examples include steam condensing to water and water freezing to ice.

In this section we'll explore a couple of phase transitions in some detail and extract some lessons that are common to all transitions.

5.1 Liquid-Gas Transition

Recall that we derived the van der Waals equation of state for a gas (2.31) in Section 2.5. We can write the van der Waals equation as

$$p = \frac{k_B T}{v - b} - \frac{a}{v^2} \tag{5.1}$$

where $v = V/N$ is the volume per particle. In the literature, you will also see this equation written in terms of the particle density $\rho = 1/v$.

On the right we fix T at different values and sketch the graph of p vs. V determined by the van der Waals equation. These curves are *isotherms* — line of constant temperature. As we can see from the diagram, the isotherms take three different shapes depending on the value of T. The top curve shows the isotherm for large values of T. Here we can effectively ignore the $-a/v^2$ term. (Recall that v cannot take values smaller than b, reflecting the fact that atoms cannot approach to arbitrarily closely). The result is a monotonically decreasing function, essentially the same as we would get for an ideal gas. In contrast, when T is low enough, the second term in (5.1) can compete with the first term. Roughly speaking, this happens when $k_B T \sim a/v$ is in the allowed region $v > b$. For these low value of the temperature, the isotherm has a wiggle.

Figure 34:

At some intermediate temperature, the wiggle must flatten out so that the bottom curve looks like the top one. This happens when the maximum and minimum meet to form an inflection point. Mathematically, we are looking for a solution to $dp/dv = d^2p/dv^2 = 0$. it is simple to check that these two equations only have a solution at the *critical temperature* $T = T_c$ given by

$$k_B T_c = \frac{8a}{27b} \tag{5.2}$$

Let's look in more detail at the $T < T_c$ curve. For a range of pressures, the system can have three different choices of volume. A typical, albeit somewhat exagerated, example of this curve is shown in the figure below. What's going on? How should we interpret the fact that the system can seemingly live at three different densities $\rho = 1/v$?

Figure 35:

First look at the middle solution. This has some fairly weird properties. We can see from the graph that the gradient is positive: $dp/dv|_T > 0$. This means that if we apply a force to the container to squeeze the gas, the pressure decreases. The gas doesn't push back; it just relents. But if we expand the gas, the pressure increases and the gas pushes harder. Both of these properties are telling us that the gas in that state is unstable. If we were able to create such a state, it wouldn't hand around for long because any tiny perturbation would lead to a rapid, explosive change in its density. If we want to find states which we are likely to observe in Nature then we should look at the other two solutions.

The solution to the left on the graph has v slightly bigger than b. But, recall from our discussion of Section 2.5 that b is the closest that the atoms can get. If we have $v \sim b$, then the atoms are very densely packed. Moreover, we can also see from the graph that $|dp/dv|$ is very large for this solution which means that the state is very difficult to compress: we need to add a great deal of pressure to change the volume only slightly. We have a name for this state: it is a *liquid*.

You may recall that our original derivation of the van der Waals equation was valid only for densities much lower than the liquid state. This means that we don't really trust (5.1) on this solution. Nonetheless, it is interesting that the equation predicts the existence of liquids and our plan is to gratefully accept this gift and push ahead to explore what the van der Waals tells us about the liquid-gas transition. We will see that it captures many of the qualitative features of the phase transition.

The last of the three solutions is the one on the right in the figure. This solution has $v \gg b$ and small $|dp/dv|$. It is the *gas state*. Our goal is to understand what happens in between the liquid and gas state. We know that the naive, middle, solution given to us by the van der Waals equation is unstable. What replaces it?

5.1.1 Phase Equilibrium

Throughout our derivation of the van der Waals equation in Section 2.5, we assumed that the system was at a fixed density. But the presence of two solutions — the liquid and gas state — allows us to consider more general configurations: part of the system could be a liquid and part could be a gas.

How do we figure out if this indeed happens? Just because both liquid and gas states can exist, doesn't mean that they can cohabit. It might be that one is preferred over the other. We already saw some conditions that must be satisfied in order for two systems to sit in equilibrium back in Section 1. Mechanical and thermal equilibrium are guaranteed if two systems have the same pressure and temperature respectively. But both of these are already guaranteed by construction for our two liquid and gas solutions: the two solutions sit on the same isotherm and at the same value of p. We're left with only one further requirement that we must satisfy which arises because the two systems can exchange particles. This is the requirement of chemical equilibrium,

$$\mu_{\text{liquid}} = \mu_{\text{gas}} \tag{5.3}$$

Because of the relationship (4.22) between the chemical potential and the Gibbs free energy, this is often expressed as

$$g_{\text{liquid}} = g_{\text{gas}} \tag{5.4}$$

where $g = G/N$ is the Gibbs free energy per particle.

Notice that all the equilibrium conditions involve only intensive quantities: p, T and μ. This means that if we have a situation where liquid and gas are in equilibrium, then we can have any number N_{liquid} of atoms in the liquid state and any number N_{gas} in the gas state. But how can we make sure that chemical equilibrium (5.3) is satisfied?

Maxwell Construction

We want to solve $\mu_{\text{liquid}} = \mu_{\text{gas}}$. We will think of the chemical potential as a function of p and T: $\mu = \mu(p, T)$. Importantly, we won't assume that $\mu(p, T)$ is single valued since that would be assuming the result we're trying to prove! Instead we will show that if we fix T, the condition (5.3) can only be solved for a very particular value of pressure

p. To see this, start in the liquid state at some fixed value of p and T and travel along the isotherm. The infinitesimal change in the chemical potential is

$$d\mu = \left.\frac{\partial \mu}{\partial p}\right|_T dp$$

However, we can get an expression for $\partial\mu/\partial p$ by recalling that arguments involving extensive and intensive variables tell us that the chemical potential is proportional to the Gibbs free energy: $G(p,T,N) = \mu(p,T)N$ (4.22). Looking back at the variation of the Gibbs free energy (4.21) then tells us that

$$\left.\frac{\partial G}{\partial p}\right|_{N,T} = \left.\frac{\partial \mu}{\partial p}\right|_T N = V \tag{5.5}$$

Integrating along the isotherm then tells us the chemical potential of any point on the curve,

$$\mu(p,T) = \mu_{\text{liquid}} + \int_{p_{\text{liquid}}}^{p} dp' \, \frac{V(p',T)}{N}$$

When we get to gas state at the same pressure $p = p_{\text{liquid}}$ that we started from, the condition for equilibrium is $\mu = \mu_{\text{liquid}}$. Which means that the integral has to vanish. Graphically this is very simple to describe: the two shaded areas in the graph must have equal area. This condition, known as the *Maxwell construction*, tells us the pressure at which gas and liquid can co-exist.

Figure 36:

I should confess that there's something slightly dodgy about the Maxwell construction. We already argued that the part of the isotherm with $dp/dv > 0$ suffers an instability and is unphysical. But we needed to trek along that part of the curve to derive our result. There are more rigorous arguments that give the same answer.

For each isotherm, we can determine the pressure at which the liquid and gas states are in equilibrium. The gives us the *co-existence curve*, shown by the dotted line in Figure 37. Inside this region, liquid and gas can both exist at the same temperature and pressure. But there is nothing that tells us how much gas there should be and how much liquid: atoms can happily move from the liquid state to the gas state. This means that while the density of gas and liquid is fixed, the *average* density of the system is not. It can vary between the gas density and the liquid density simply by changing the amount of liquid. The upshot of this argument is that inside the co-existence curves, the isotherms simply become flat lines, reflecting the fact that the density can take any value. This is shown in graph on the right of Figure 37.

Figure 37: The co-existence curve in red, resulting in constant pressure regions consisting of a harmonious mixture of vapour and liquid.

To illustrate the physics of this situation, suppose that we sit at some fixed density $\rho = 1/v$ and cool the system down from a high temperature to $T < T_c$ at a point inside the co-existence curve so that we're now sitting on one of the flat lines. Here, the system is neither entirely liquid, nor entirely gas. Instead it will split into gas, with density $1/v_{\text{gas}}$, and liquid, with density $1/v_{\text{liquid}}$ so that the average density remains $1/v$. The system undergoes *phase separation*. The minimum energy configuration will typically be a single phase of liquid and one of gas because the interface between the two costs energy. (We will derive an expression for this energy in Section 5.5). The end result is shown on the right. In the presence of gravity, the higher density liquid will indeed sink to the bottom.

Figure 38:

Meta-Stable States

We've understood what replaces the unstable region of the van der Waals phase diagram. But we seem to have removed more states than anticipated: parts of the Van der Waals isotherm that had $dp/dv < 0$ are contained in the co-existence region and replaced by the flat pressure lines. This is the region of the p-V phase diagram that is contained between the two dotted lines in the figure to the right. The outer dotted line is the co-existence curve. The inner dotted curve is constructed to pass through the stationary points of the van der Waals isotherms. It is called the *spinodial curve*.

Figure 39:

The van der Waals states which lie between the spinodial curve and the co-existence curve are good states. But they are meta-stable. One can show that their Gibbs free energy is higher than that of the liquid-gas equilibrium at the same p and T. However, if we compress the gas very slowly we can coax the system into this state. It is known as a supercooled vapour. It is delicate. Any small disturbance will cause some amount of the gas to condense into the liquid. Similarly, expanding a liquid beyond the co-existence curve results in an meta-stable, superheated liquid.

5.1.2 The Clausius-Clapeyron Equation

We can also choose to plot the liquid-gas phase diagram on the $p - T$ plane. Here the co-existence region is squeezed into a line: if we're sitting in the gas phase and increase the pressure just a little bit at at fixed $T < T_c$ then we jump immediately to the liquid phase. This appears as a discontinuity in the volume. Such discontinuities are the sign of a phase transition. The end result is sketched in the figure to the right; the thick solid line denotes the presence of a phase transition.

Figure 40:

Either side of the line, all particles are either in the gas or liquid phase. We know from (5.4) that the Gibbs free energies of these two states are equal,

$$G_{\text{liquid}} = G_{\text{gas}}$$

So G is continuous as we move across the line of phase transitions. Suppose that we sit on the line itself and move along it. How does G change? We can easily compute this from (4.21),

$$\begin{aligned} dG_{\text{liquid}} &= -S_{\text{liquid}}dT + V_{\text{liquid}}dp \\ = dG_{\text{gas}} &= -S_{\text{gas}}dT + V_{\text{gas}}dp \end{aligned}$$

But this gives us a nice expression for the slope of the line of phase transitions in the p-T plane. It is

$$\frac{dp}{dT} = \frac{S_{\text{gas}} - S_{\text{liquid}}}{V_{\text{gas}} - V_{\text{liquid}}}$$

We usually define the *latent heat*

$$L = T(S_{\text{gas}} - S_{\text{liquid}})$$

This is the energy released as we pass through the phase transition. We see that the slope of the line in the p-T plane is determined by the ratio of latent heat released in the phase transition and the discontinuity in volume. The result is known as the *Clausius-Clapeyron* equation,

$$\frac{dp}{dT} = \frac{L}{T(V_{\text{gas}} - V_{\text{liquid}})} \tag{5.6}$$

There is a classification of phase transitions, due originally to Ehrenfest. When the nth derivative of a thermodynamic potential (either F or G usually) is discontinuous, we say we have an nth order phase transition. In practice, we nearly always deal with first, second and (very rarely) third order transitions. The liquid-gas transition releases latent heat, which means that $S = -\partial F/\partial T$ is discontinuous. Alternatively, we can say that $V = \partial G/\partial p$ is discontinuous. Either way, it is a *first order phase transition*. The Clausius-Clapeyron equation (5.6) applies to any first order transition.

As we approach $T \to T_c$, the discontinuity diminishes and $S_{\text{liquid}} \to S_{\text{gas}}$. At the critical point $T = T_c$ we have a second order phase transition. Above the critical point, there is no sharp distinction between the gas phase and liquid phase.

For most simple materials, the phase diagram above is part of a larger phase diagram which includes solids at smaller temperatures or higher pressures. A generic version of such a phase diagram is shown to the right. The van der Waals equation is missing the physics of solidification and includes only the liquid-gas line.

Figure 41:

An Approximate Solution to the Clausius-Clapeyron Equation

We can solve the Clausius-Clapeyron solution if we make the following assumptions:

- The latent heat L is constant.
- $V_{\text{gas}} \gg V_{\text{liquid}}$, so $V_{\text{gas}} - V_{\text{liquid}} \approx V_{\text{gas}}$. For water, this is an error of less than 0.1%
- Although we derived the phase transition using the van der Waals equation, now we've got equation (5.6) we'll pretend the gas obeys the ideal gas law $pV = Nk_BT$.

With these assumptions, it is simple to solve (5.6). It reduces to

$$\frac{dp}{dT} = \frac{Lp}{Nk_BT^2} \quad \Rightarrow \quad p = p_0 e^{-L/Nk_BT}$$

5.1.3 The Critical Point

Let's now return to discuss some aspects of life at the critical point. We previously worked out the critical temperature (5.2) by looking for solutions to simultaneous equations $\partial p/\partial v = \partial^2 p/\partial v^2 = 0$. There's a slightly more elegant way to find the critical point which also quickly gives us p_c and v_c as well. We rearrange the van der Waals equation (5.1) to get a cubic,

$$pv^3 - (pb + k_B T)v^2 + av - ab = 0$$

For $T < T_c$, this equation has three real roots. For $T > T_c$ there is just one. Precisely at $T = T_c$, the three roots must therefore coincide (before two move off onto the complex plane). At the critical point, this curve can be written as

$$p_c(v - v_c)^3 = 0$$

Comparing the coefficients tells us the values at the critical point,

$$k_B T_c = \frac{8a}{27b} \quad , \quad v_c = 3b \quad , \quad p_c = \frac{a}{27b^2} \tag{5.7}$$

The Law of Corresponding States

We can invert the relations (5.7) to express the parameters a and b in terms of the critical values, which we then substitute back into the van der Waals equation. To this end, we define the *reduced* variables,

$$\bar{T} = \frac{T}{T_c} \quad , \quad \bar{v} = \frac{v}{v_c} \qquad \bar{p} = \frac{p}{p_c}$$

The advantage of working with $\bar{T}$, $\bar{v}$ and $\bar{p}$ is that it allows us to write the van der Waals equation (5.1) in a form that is universal to all gases, usually referred to as the law of corresponding states

$$\bar{p} = \frac{8}{3}\frac{\bar{T}}{\bar{v} - 1/3} - \frac{3}{\bar{v}^2}$$

Moreover, because the three variables T_c, p_c and v_c at the critical point are expressed in terms of just two variables, a and b (5.7), we can construct a combination of them which is independent of a and b and therefore supposedly the same for all gases. This is the universal compressibility ratio,

$$\frac{p_c v_c}{k_B T_c} = \frac{3}{8} = 0.375 \tag{5.8}$$

Figure 42: The co-existence curve for gases. Data is plotted for Ne, Ar, Kr,Xe, N_2, O_2, CO and CH_4.

Comparing to real gases, this number is a little high. Values range from around 0.28 to 0.3. We shouldn't be too discouraged by this; after all, we knew from the beginning that the van der Waals equation is unlikely to be accurate in the liquid regime. Moreover, the fact that gases have a critical point (defined by three variables T_c, p_c and v_c) guarantees that a similar relationship would hold for any equation of state which includes just two parameters (such as a and b) but would most likely fail to hold for equations of state that included more than two parameters.

Dubious as its theoretical foundation is, the law of corresponding states is the first suggestion that something remarkable happens if we describe a gas in terms of its reduced variables. More importantly, there is striking experimental evidence to back this up! Figure 42 shows the Guggenheim plot, constructed in 1945. The co-existence curve for 8 different gases in plotted in reduced variables: $\bar{T}$ along the vertical axis; $\bar{\rho} = 1/\bar{v}$ along the horizontal. The gases vary in complexity from the simple monatomic gas Ne to the molecule CH_4. As you can see, the co-existence curve for all gases is essentially the same, with the chemical make-up largely forgotten. There is clearly something interesting going on. How to understand it?

Critical Exponents

We will focus attention on physics close to the critical point. It is not immediately obvious what are the right questions to ask. It turns out that the questions which have the most interesting answer are concerned with how various quantities change as we approach the critical point. There are lots of ways to ask questions of this type since there are many quantities of interest and, for each of them, we could approach

the critical point from different directions. Here we'll look at the behaviour of three quantities to get a feel for what happens.

First, we can ask what happens to the difference in (inverse) densities $v_{\text{gas}} - v_{\text{liquid}}$ as we approach the critical point along the co-existence curve. For $T < T_c$, or equivalently $\bar{T} < 1$, the reduced van der Waals equation (5.8) has two stable solutions,

$$\bar{p} = \frac{8\bar{T}}{3\bar{v}_{\text{liquid}} - 1} - \frac{3}{\bar{v}_{\text{liquid}}^2} = \frac{8\bar{T}}{3\bar{v}_{\text{gas}} - 1} - \frac{3}{\bar{v}_{\text{gas}}^2}$$

If we solve this for $\bar{T}$, we have

$$\bar{T} = \frac{(3\bar{v}_{\text{liquid}} - 1)(3\bar{v}_{\text{gas}} - 1)(\bar{v}_{\text{liquid}} + \bar{v}_{\text{gas}})}{8\bar{v}_{\text{gas}}^2\bar{v}_{\text{liquid}}^2}$$

Notice that as we approach the critical point, $\bar{v}_{\text{gas}}, \bar{v}_{\text{liquid}} \to 1$ and the equation above tells us that $\bar{T} \to 1$ as expected. We can see exactly how we approach $\bar{T} = 1$ by expanding the right right-hand side for small $\epsilon \equiv \bar{v}_{\text{gas}} - \bar{v}_{\text{liquid}}$. To do this quickly, it's best to notice that the equation is symmetric in $\bar{v}_{\text{gas}}$ and $\bar{v}_{\text{liquid}}$, so close to the critical point we can write $\bar{v}_{\text{gas}} = 1 + \epsilon/2$ and $\bar{v}_{\text{liquid}} = 1 - \epsilon/2$. Substituting this into the equation above and keeping just the leading order term, we find

$$\bar{T} \approx 1 - \frac{1}{16}(\bar{v}_{\text{gas}} - \bar{v}_{\text{liquid}})^2$$

Or, re-arranging, as we approach T_c along the co-existence curve,

$$v_{\text{gas}} - v_{\text{liquid}} \sim (T_c - T)^{1/2} \tag{5.9}$$

This is the answer to our first question.

Our second variant of the question is: how does the volume change with pressure as we move along the critical isotherm. It turns out that we can answer this question without doing any work. Notice that at $T = T_c$, there is a unique pressure for a given volume $p(v, T_c)$. But we know that $\partial p/\partial v = \partial^2 p/\partial v^2 = 0$ at the critical point. So a Taylor expansion around the critical point must start with the cubic term,

$$p - p_c \sim (v - v_c)^3 \tag{5.10}$$

This is the answer to our second question.

Our third and final variant of the question concerns the compressibility, defined as

$$\kappa = -\frac{1}{v}\left.\frac{\partial v}{\partial p}\right|_T \tag{5.11}$$

We want to understand how κ changes as we approach $T \to T_c$ from above. In fact, we met the compressibility before: it was the feature that first made us nervous about the van der Waals equation since κ is negative in the unstable region. We already know that at the critical point $\partial p/\partial v|_{T_c} = 0$. So expanding for temperatures close to T_c, we expect

$$\left.\frac{\partial p}{\partial v}\right|_{T;v=v_c} = -a(T - T_c) + \ldots$$

This tells us that the compressibility should diverge at the critical point, scaling as

$$\kappa \sim (T - T_c)^{-1} \tag{5.12}$$

We now have three answers to three questions: (5.9), (5.10) and (5.12). Are they right?! By which I mean: do they agree with experiment? Remember that we're not sure that we can trust the van der Waals equation at the critical point so we should be nervous. However, there is also reason for some confidence. Notice, in particular, that in order to compute (5.10) and (5.12), we didn't actually need any details of the van der Waals equation. We simply needed to assume the existence of the critical point and an analytic Taylor expansion of various quantities in the neighbourhood. Given that the answers follow from such general grounds, one may hope that they provide the correct answers for a gas in the neighbourhood of the critical point even though we know that the approximations that went into the van der Waals equation aren't valid there. Fortunately, that isn't the case: physics is much more interesting than that!

The experimental results for a gas in the neighbourhood of the critical point do share one feature in common with the discussion above: they are completely independent of the atomic make-up of the gas. However, the scaling that we computed using the van der Waals equation is not fully accurate. The correct results are as follows. As we approach the critical point along the co-existence curve, the densities scale as

$$v_{\text{gas}} - v_{\text{liquid}} \sim (T_c - T)^\beta \quad \text{with} \quad \beta \approx 0.32$$

(Note that the exponent β has nothing to do with inverse temperature. We're just near the end of the course and running out of letters and β is the canonical name for this exponent). As we approach along an isotherm,

$$p - p_c \sim (v - v_c)^\delta \quad \text{with} \quad \delta \approx 4.8$$

Finally, as we approach T_c from above, the compressibility scales as

$$\kappa \sim (T - T_c)^{-\gamma} \quad \text{with} \quad \gamma \approx 1.2$$

The quantities β, γ and δ are examples of *critical exponents*. We will see more of them shortly. The van der Waals equation provides only a crude first approximation to the critical exponents.

Fluctuations

We see that the van der Waals equation didn't do too badly in capturing the dynamics of an interacting gas. It gets the qualitative behaviour right, but fails on precise quantitative tests. So what went wrong? We mentioned during the derivation of the van der Waals equation that we made certain approximations that are valid only at low density. So perhaps it is not surprising that it fails to get the numbers right near the critical point $v = 3b$. But there's actually a deeper reason that the van der Waals equation fails: fluctuations.

This is simplest to see in the grand canonical ensemble. Recall that back in Section 1 that we argued that $\Delta N/N \sim 1/\sqrt{N}$, which allowed us to happily work in the grand canonical ensemble even when we actually had fixed particle number. In the context of the liquid-gas transition, fluctuating particle number is the same thing as fluctuating density $\rho = N/V$. Let's revisit the calculation of ΔN near the critical point. Using (1.45) and (1.48), the grand canonical partition function can be written as $\log \mathcal{Z} = \beta V p(T, \mu)$, so the average particle number (1.42) is

$$\langle N \rangle = V \left.\frac{\partial p}{\partial \mu}\right|_{T,V}$$

We already have an expression for the variance in the particle number in (1.43),

$$\Delta N^2 = \frac{1}{\beta} \left.\frac{\partial \langle N \rangle}{\partial \mu}\right|_{T,V}$$

Dividing these two expressions, we have

$$\frac{\Delta N^2}{N} = \frac{1}{V\beta} \left.\frac{\partial \langle N \rangle}{\partial \mu}\right|_{T,V} \left.\frac{\partial \mu}{\partial p}\right|_{T,V} = \frac{1}{V\beta} \left.\frac{\partial \langle N \rangle}{\partial p}\right|_{T,V}$$

But we can re-write this expression using the general relationship between partial derivatives $\partial x/\partial y|_z \, \partial y/\partial z|_x \, \partial z/\partial x|_y = -1$. We then have

$$\frac{\Delta N^2}{N} = -\frac{1}{\beta} \left.\frac{\partial \langle N \rangle}{\partial V}\right|_{p,T} \frac{1}{V} \left.\frac{\partial V}{\partial p}\right|_{N,T}$$

This final expression relates the fluctuations in the particle number to the compressibility (5.11). But the compressibility is diverging at the critical point and this means that there are large fluctuations in the density of the fluid at this point. The result is that any simple equation of state, like the van der Waals equation, which works only with the average volume, pressure and density will miss this key aspect of the physics.

Understanding how to correctly account for these fluctuations is the subject of *critical phenomena*. It has close links with the *renormalization group* and *conformal field theory* which also arise in particle physics and string theory. You will meet some of these ideas in next year's Statistical Field Theory course. Here we will turn to a different phase transition which will allow us to highlight some of the key ideas.

5.2 The Ising Model

The Ising model is one of the touchstones of modern physics; a simple system that exhibits non-trivial and interesting behaviour.

The Ising model consists of N sites in a d-dimensional lattice. On each lattice site lives a quantum spin that can sit in one of two states: spin up or spin down. We'll call the eigenvalue of the spin on the ith lattice site s_i. If the spin is up, $s_i = +1$; if the spin is down, $s_i = -1$.

The spins sit in a magnetic field that endows an energy advantage to those which point up,

$$E_B = -B \sum_{i=1}^{N} s_i$$

(A comment on notation: B should be properly denoted H. We're sticking with B to avoid confusion with the Hamiltonian. There is also a factor of the magnetic moment which has been absorbed into the definition of B). The lattice system with energy E_B is equivalent to the two-state system that we first met when learning the techniques of statistical mechanics back in Section 1.2.3. However, the Ising model contains an additional complication that makes the sysem much more interesting: this is an interaction between neighbouring spins. The full energy of the system is therefore,

$$E = -J \sum_{\langle ij \rangle} s_i s_j - B \sum_i s_i \tag{5.13}$$

The notation $\langle ij \rangle$ means that we sum over all "nearest neighbour" pairs in the lattice. The number of such pairs depends both on the dimension d and the type of lattice.

We'll denote the number of nearest neighbours as q. For example, in $d = 1$ a lattice has $q = 2$; in $d = 2$, a square lattice has $q = 4$. A square lattice in d dimensions has $q = 2d$.

If $J > 0$, neighbouring spins prefer to be aligned ($\uparrow\uparrow$ or $\downarrow\downarrow$). In the context of magnetism, such a system is called a *ferromagnet*. If $J < 0$, the spins want to anti-align ($\uparrow\downarrow$). This is an *anti-ferromagnet*. In the following, we'll choose $J > 0$ although for the level of discussion needed for this course, the differences are minor.

We work in the canonical ensemble and introduce the partition function

$$Z = \sum_{\{s_i\}} e^{-\beta E[s_i]} \tag{5.14}$$

While the effect of both $J > 0$ and $B \neq 0$ is to make it energetically preferable for the spins to align, the effect of temperature will be to randomize the spins, with entropy winnning out over energy. Our interest is in the average spin, or average *magnetization*

$$m = \frac{1}{N} \sum_i \langle s_i \rangle = \frac{1}{N\beta} \frac{\partial \log Z}{\partial B} \tag{5.15}$$

The Ising Model as a Lattice Gas

Before we develop techniques to compute the partition function (5.14), it's worth pointing out that we can drape slightly different words around the mathematics of the Ising model. It need not be interpreted as a system of spins; it can also be thought of as a lattice description of a gas.

To see this, consider the same d-dimensional lattice as before, but now with particles hopping between lattice sites. These particles have hard cores, so no more than one can sit on a single lattice site. We introduce the variable $n_i \in \{0, 1\}$ to specify whether a given lattice site, labelled by i, is empty ($n_i = 0$) or filled ($n_i = 1$). We can also introduce an attractive force between atoms by offering them an energetic reward if they sit on neighbouring sites. The Hamiltonian of such a lattice gas is given by

$$E = -4J \sum_{\langle ij \rangle} n_i n_j - \mu \sum_i n_i$$

where μ is the chemical potential which determines the overall particle number. But this Hamiltonian is trivially the same as the Ising model (5.13) if we make the identification

$$s_i = 2n_i - 1 \in \{-1, 1\}$$

The chemical potenial μ in the lattice gas plays the role of magnetic field in the spin system while the magnetization of the system (5.15) measures the average density of particles away from half-filling.

5.2.1 Mean Field Theory

For general lattices, in arbitrary dimension d, the sum (5.14) cannot be performed. An exact solution exists in $d = 1$ and, when $B = 0$, in $d = 2$. (The $d = 2$ solution is originally due to Onsager and is famously complicated! Simpler solutions using more modern techniques have since been discovered).

Here we'll develop an approximate method to evaluate Z known as *mean field theory*. We write the interactions between neighbouring spins in term of their deviation from the average spin m,

$$\begin{aligned} s_i s_j &= [(s_i - m) + m][(s_j - m) + m] \\ &= (s_i - m)(s_j - m) + m(s_j - m) + m(s_i - m) + m^2 \end{aligned}$$

The mean field approximation means that we assume that the fluctuations of spins away from the average are small which allows us to neglect the first term above. Notice that this isn't the statement that the variance of an individual spin is small; that can never be true because s_i takes values $+1$ or -1 so $\langle s_i^2 \rangle = 1$ and the variance $\langle (s_i - m)^2 \rangle$ is always large. Instead, the mean field approximation is a statement about fluctuations between spins on neighbouring sites, so the first term above can be neglected when summing over $\sum_{\langle ij \rangle}$. We can then write the energy (5.13) as

$$\begin{aligned} E_{\text{mf}} &= -J \sum_{\langle ij \rangle} [m(s_i + s_j) - m^2] - B \sum_i s_i \\ &= \frac{1}{2} JNqm^2 - (Jqm + B) \sum_i s_i \end{aligned} \tag{5.16}$$

where the factor of $Nq/2$ in the first term is simply the number of nearest neighbour pairs $\sum_{\langle ij \rangle}$. The factor or $1/2$ is there because $\sum_{\langle ij \rangle}$ is a sum over pairs rather than a sum of individual sites. (If you're worried about this formula, you should check it for a simple square lattice in $d = 1$ and $d = 2$ dimensions). A similar factor in the second term cancelled the factor of 2 due to $(s_i + s_j)$.

We see that the mean field approximation has removed the interactions. The Ising model reduces to the two state system that we saw way back in Section 1. The result of the interactions is that a given spin feels the average effect of its neighbour's spins through a contribution to the effective magnetic field,

$$B_{\text{eff}} = B + Jqm$$

Once we've taken into account this extra contribution to B_{eff}, each spin acts indepen-

Figure 43: $\tanh(Jqm\beta)$ for $Jq\beta < 1$

Figure 44: $\tanh(Jqm\beta)$ for $Jq\beta > 1$

dently and it is easy to write the partition function. It is

$$\begin{aligned} Z &= e^{-\frac{1}{2}\beta JNqm^2}\left(e^{-\beta B_{\text{eff}}} + e^{\beta B_{\text{eff}}}\right)^N \\ &= e^{-\frac{1}{2}\beta JNqm^2} 2^N \cosh^N \beta B_{\text{eff}} \end{aligned} \tag{5.17}$$

However, we're not quite done. Our result for the partition function Z depends on B_{eff} which depends on m which we don't yet know. However, we can use our expression for Z to self-consistently determine the magnetization (5.15). We find,

$$m = \tanh(\beta B + \beta Jqm) \tag{5.18}$$

We can now solve this equation to find the magnetization for various values of T and B: $m = m(T, B)$. It is simple to see the nature of the solutions using graphical methods.

B=0

Let's first consider the situation with vanishing magnetic field, $B = 0$. The figures above show the graph linear in m compared with the tanh function. Since $\tanh x \approx x - \frac{1}{3}x^3 + \ldots$, the slope of the graph near the origin is given by βJq. This then determines the nature of the solution.

- The first graph depicts the situation for $\beta Jq < 1$. The only solution is $m = 0$. This means that at high temperatures $k_B T > Jq$, there is no average magnetization of the system. The entropy associated to the random temperature fluctuations wins over the energetically preferred ordered state in which the spins align.

- The second graph depicts the situation for $\beta Jq > 1$. Now there are three solutions: $m = \pm m_0$ and $m = 0$. It will turn out that the middle solution, $m = 0$, is unstable. (This solution is entirely analogous to the unstable solution of the van der Waals equation. We will see this below when we compute the free energy).

For the other two possible solutions, $m = \pm m_0$, the magnetization is non-zero. Here we see the effects of the interactions begin to win over temperature. Notice that in the limit of vanishing temperature, $\beta \to \infty$, $m_0 \to 1$. This means that all the spins are pointing in the same direction (either up or down) as expected.

- The critical temperature separating these two cases is

$$k_B T_c = Jq \tag{5.19}$$

The results described above are perhaps rather surprising. Based on the intuition that things in physics always happen smoothly, one might have thought that the magnetization would drop slowly to zero as $T \to \infty$. But that doesn't happen. Instead the magnetization turns off abruptly at some finite value of the temperature $T = T_c$, with no magnetization at all for higher temperatures. This is the characteristic behaviour of a phase transition.

$\mathbf{B \neq 0}$

For $B \neq 0$, we can solve the consistency equation (5.18) in a similar fashion. There are a couple of key differences to the $B \neq 0$ case. Firstly, there is now no phase transition at fixed B as we vary temperature T. Instead, for very large temperatures $k_B T \gg Jq$, the magnetization goes smoothly to zero as

$$m \to \frac{B}{k_B T} \qquad \text{as } T \to \infty$$

At low temperatures, the magnetization again asymptotes to the state $m \to \pm 1$ which minimizes the energy. Except this time, there is no ambiguity as to whether the system chooses $m = +1$ or $m = -1$. This is entirely determined by the sign of the magnetic field B.

In fact the low temperature behaviour requires slightly more explanation. For small values of B and T, there are again three solutions to (5.18). This follows simply from continuity: there are three solutions for $T < T_c$ and $B = 0$ shown in Figure 44 and these must survive in some neighbourhood of $B = 0$. One of these solutions is again unstable. However, of the remaining two only one is now stable: that with $\text{sign}(m) = \text{sign}(B)$. The other is meta-stable. We will see why this is the case shortly when we come to discuss the free energy.

Figure 45: Magnetization with $B = 0$ and the phase transtion

Figure 46: Magnetization at $B \neq 0$.

The net result of our discussion is depicted in the figures above. When $B = 0$ there is a phase transition at $T = T_c$. For $T < T_c$, the system can sit in one of two magnetized states with $m = \pm m_0$. In contrast, for $B \neq 0$, there is no phase transition as we vary temperature and the system has at all times a preferred magnetization whose sign is determined by that of B. Notice however, we do have a phase transition if we fix temperature at $T < T_c$ and vary B from negative to positive. Then the magnetization jumps discontinuously from a negative value to a positive value. Since the magnetization is a first derivative of the free energy (5.15), this is a first order phase transition. In contrast, moving along the temperature axis at $B = 0$ results in a second order phase transition at $T = T_c$.

5.2.2 Critical Exponents

It is interesting to compare the phase transition of the Ising model with that of the liquid-gas phase transition. The two are sketched in the Figure 47 above. In both cases, we have a first order phase transition and a quantity jumps discontinuously at $T < T_c$. In the case of the liquid-gas, it is the density $\rho = 1/v$ that jumps as we vary pressure; in the case of the Ising model it is the magnetization m that jumps as we vary the magnetic field. Moreover, in both cases the discontinuity disappears as we approach $T = T_c$.

We can calculate critical exponents for the Ising model. To compare with our discussion for the liquid-gas critical point, we will compute three quantities. First, consider the magnetization at $B = 0$. We can ask how this magnetization decreases as we tend towards the critical point. Just below $T = T_c$, m is small and we can Taylor expand (5.18) to get

$$m \approx \beta Jqm - \frac{1}{3}(\beta Jqm)^3 + \ldots$$

Figure 47: A comparison of the phase diagram for the liquid-gas system and the Ising model.

The magnetization therefore scales as

$$m_0 \sim \pm(T_c - T)^{1/2} \tag{5.20}$$

This is to be compared with the analogous result (5.9) from the van der Waals equation. We see that the values of the exponents are the same in both cases. Notice that the derivative dm/dT becomes infinite as we approach the critical point. In fact, we had already anticipated this when we drew the plot of the magnetization in Figure 45.

Secondly, we can sit at $T = T_c$ and ask how the magnetization changes as we approach $B = 0$. We can read this off from (5.18). At $T = T_c$ we have $\beta Jq = 1$ and the consistency condition becomes $m = \tanh(B/Jq + m)$. Expanding for small B gives

$$m \approx \frac{B}{Jq} + m - \frac{1}{3}\left(\frac{B}{Jq} + m\right)^3 + \ldots \approx \frac{B}{Jq} + m - \frac{1}{3}m^3 + \mathcal{O}(B^2)$$

So we find that the magnetization scales as

$$m \sim B^{1/3} \tag{5.21}$$

Notice that this power of 1/3 is again familiar from the liquid-gas transition (5.10) where the van der Waals equation gave $v_{\text{gas}} - v_{\text{liquid}} \sim (p - p_c)^{1/3}$.

Finally, we can look at the magnetic susceptibility χ, defined as

$$\chi = N \left.\frac{\partial m}{\partial B}\right|_T$$

This is analogous to the compressibility κ of the gas. We will ask how χ changes as we approach $T \to T_c$ from above at $B = 0$. We differentiate (5.18) with respect to B to get

$$\chi = \frac{N\beta}{\cosh^2 \beta Jqm}\left(1 + \frac{Jq}{N}\chi\right)$$

We now evaluate this at $B = 0$. Since we want to approach $T \to T_c$ from above, we can also set $m = 0$ in the above expression. Evaluating this at $B = 0$ gives us the scaling

$$\chi = \frac{N\beta}{1 - Jq\beta} \sim (T - T_c)^{-1} \tag{5.22}$$

Once again, we see that same critical exponent that the van der Waals equation gave us for the gas (5.12).

5.2.3 Validity of Mean Field Theory

The phase diagram and critical exponents above were all derived using the mean field approximation. But this was an unjustified approximation. Just as for the van der Waals equation, we can ask the all-important question: are our results right?

There is actually a version of the Ising model for which the mean field theory is exact: it is the $d = \infty$ dimensional lattice. This is unphysical (even for a string theorist). Roughly speaking, mean field theory works for large d because each spin has a large number of neighbours and so indeed sees something close to the average spin.

But what about dimensions of interest? Mean field theory gets things most dramatically wrong in $d = 1$. In that case, no phase transition occurs. We will derive this result below where we briefly describe the exact solution to the $d = 1$ Ising model. There is a general lesson here: in low dimensions, both thermal and quantum fluctuations are more important and invariably stop systems forming ordered phases.

In higher dimensions, $d \geq 2$, the crude features of the phase diagram, including the existence of a phase transition, given by mean field theory are essentially correct. In fact, the very existence of a phase transition is already worthy of comment. The defining feature of a phase transition is behaviour that jumps discontinuously as we vary β or B. Mathematically, the functions must be non-analytic. Yet all properties of the theory can be extracted from the partition function Z which is a sum of smooth, analytic functions (5.14). How can we get a phase transition? The loophole is that Z is only necessarily analytic if the sum is finite. But there is no such guarantee when the number of lattice sites $N \to \infty$. We reach a similar conclusion to that of Bose-Einstein condensation: phase transitions only strictly happen in the thermodynamic limit. There are no phase transitions in finite systems.

What about the critical exponents that we computed in (5.20), (5.21) and (5.22)? It turns out that these are correct for the Ising model defined in $d \geq 4$. (We will briefly sketch why this is true at the end of this Chapter). But for $d = 2$ and $d = 3$, the critical exponents predicted by mean field theory are only first approximations to the true answers.

For $d = 2$, the exact solution (which goes quite substantially past this course) gives the critical exponents to be,

$$\begin{aligned} m_0 \sim (T_c - T)^\beta \quad &\text{with} \quad \beta = \frac{1}{8} \\ m \sim B^{1/\delta} \quad &\text{with} \quad \delta = 15 \\ \chi \sim (T - T_c)^{-\gamma} \quad &\text{with} \quad \gamma = \frac{7}{4} \end{aligned}$$

The biggest surprise is in $d = 3$ dimensions. Here the critical exponents are not known exactly. However, there has been a great deal of numerical work to determine them. They are given by

$$\beta \approx 0.32 \quad , \quad \delta \approx 4.8 \quad , \quad \gamma \approx 1.2$$

But these are exactly the same critical exponents that are seen in the liquid-gas phase transition. That's remarkable! We saw above that the mean field approach to the Ising model gave the same critical exponents as the van der Waals equation. But they are both wrong. And they are both wrong in the same, complicated, way! Why on earth would a system of spins on a lattice have anything to do with the phase transition between a liquid and gas? It is as if all memory of the microscopic physics — the type of particles, the nature of the interactions — has been lost at the critical point. And that's exactly what happens.

What we're seeing here is evidence for *universality*. There is a single theory which describes the physics at the critical point of the liquid gas transition, the 3d Ising model and many other systems. This is a theoretical physicist's dream! We spend a great deal of time trying to throw away the messy details of a system to focus on the elegant essentials. But, at a critical point, Nature does this for us! Although critical points in two dimensions are well understood, there is still much that we don't know about critical points in three dimensions. This, however, is a story that will have to wait for another day.

5.3 Some Exact Results for the Ising Model

This subsection is something of a diversion from our main interest. In later subsections, we will develop the idea of mean field theory. But first we pause to describe some exact results for the Ising model using techniques that do not rely on the mean field approximation. Many of the results that we derive have broader implications for systems beyond the Ising model.

As we mentioned above, there is an exact solution for the Ising model in $d = 1$ dimension and, when $B = 0$, in $d = 2$ dimensions. Here we will describe the $d = 1$ solution but not the full $d = 2$ solution. We will, however, derive a number of results for the $d = 2$ Ising model which, while falling short of the full solution, nonetheless provide important insights into the physics.

5.3.1 The Ising Model in $d = 1$ Dimensions

We start with the *Ising chain*, the Ising model on a one dimensional line. Here we will see that the mean field approximation fails miserably, giving qualitatively incorrect results: the exact results shows that there are no phase transitions in the Ising chain.

The energy of the system (5.13) can be trivially rewritten as

$$E = -J \sum_{i=1}^{N} s_i s_{i+1} - \frac{B}{2} \sum_{i=1}^{N} (s_i + s_{i+1})$$

We will impose periodic boundary conditions, so the spins live on a circular lattice with $s_{N+1} \equiv s_1$. The partition function is then

$$Z = \sum_{s_1=\pm 1} \dots \sum_{s_N=\pm 1} \prod_{i=1}^{N} \exp\left(\beta J s_i s_{i+1} + \frac{\beta B}{2}(s_i + s_{i+1})\right) \tag{5.23}$$

The crucial observation that allows us to solve the problem is that this partition function can be written as a product of matrices. We adopt notation from quantum mechanics and define the 2×2 matrix,

$$\langle s_i | T | s_{i+1} \rangle \equiv \exp\left(\beta J s_i s_{i+1} + \frac{\beta B}{2}(s_i + s_{i+1})\right) \tag{5.24}$$

The row of the matrix is specified by the value of $s_i = \pm 1$ and the column by $s_{i+1} = \pm 1$. T is known as the *transfer matrix* and, in more conventional notation, is given by

$$T = \begin{pmatrix} e^{\beta J + \beta B} & e^{-\beta J} \\ e^{-\beta J} & e^{\beta J - \beta B} \end{pmatrix}$$

The sums over the spins $\sum_{s_i}$ and product over lattice sites $\prod_i$ in (5.23) simply tell us to multiply the matrices defined in (5.24) and the partition function becomes

$$Z = \mathrm{Tr}\ (\langle s_1 | T | s_2 \rangle \langle s_2 | T | s_3 \rangle \dots \langle s_N | T | s_1 \rangle) = \mathrm{Tr}\, T^N \tag{5.25}$$

where the trace arises because we have imposed periodic boundary conditions. To complete the story, we need only compute the eigenvalues of T to determine the partition function. A quick calculation shows that the two eigenvalues of T are

$$\lambda_\pm = e^{\beta J}\cosh\beta B \pm \sqrt{e^{2\beta J}\cosh^2\beta B - 2\sinh 2\beta J} \tag{5.26}$$

where, clearly, $\lambda_- < \lambda_+$. The partition function is then

$$Z = \lambda_+^N + \lambda_-^N = \lambda_+^N\left(1 + \frac{\lambda_-^N}{\lambda_+^N}\right) \approx \lambda_+^N \tag{5.27}$$

where, in the last step, we've used the simple fact that if λ_+ is the largest eigenvalue then $\lambda_-^N/\lambda_+^N \approx 0$ for very large N.

The partition function Z contains many quantities of interest. In particular, we can use it to compute the magnetisation as a function of temperature when $B = 0$. This, recall, is the quantity which is predicted to undergo a phase transition in the mean field approximation, going abruptly to zero at some critical temperature. In the $d = 1$ Ising model, the magnetisation is given by

$$m = \frac{1}{N\beta}\left.\frac{\partial \log Z}{\partial B}\right|_{B=0} = \frac{1}{\lambda_+\beta}\left.\frac{\partial \lambda_+}{\partial B}\right|_{B=0} = 0$$

We see that the true physics for $d = 1$ is very different than that suggested by the mean field approximation. When $B = 0$, there is no magnetisation! While the J term in the energy encourages the spins to align, this is completely overwhelmed by thermal fluctuations for any value of the temperature.

There is a general lesson in this calculation: thermal fluctuations always win in one dimensional systems. They never exhibit ordered phases and, for this reason, never exhibit phase transitions. The mean field approximation is bad in one dimension.

5.3.2 2d Ising Model: Low Temperatures and Peierls Droplets

Let's now turn to the Ising model in $d = 2$ dimensions. We'll work on a square lattice and set $B = 0$. Rather than trying to solve the model exactly, we'll have more modest goals. We will compute the partition function in two different limits: high temperature and low temperature. We start here with the low temperature expansion.

The partition function is given by the sum over all states, weighted by $e^{-\beta E}$. At low temperatures, this is always dominated by the lowest lying states. For the Ising model,

we have

$$Z = \sum_{\{s_i\}} \exp\left(\beta J \sum_{\langle ij \rangle} s_i s_j\right)$$

The low temperature limit is $\beta J \to \infty$, where the partition function can be approximated by the sum over the first few lowest energy states. All we need to do is list these states.

The ground states are easy. There are two of them: spins all up or spins all down. For example, the ground state with spins all up looks like

Each of these ground states has energy $E = E_0 = -2NJ$.

The first excited states arise by flipping a single spin. Each spin has $q = 4$ nearest neighbours – denoted by red lines in the example below – each of which leads to an energy cost of $2J$. The energy of each first excited state is therefore $E_1 = E_0 + 8J$.

There are, of course, N different spins that we we can flip and, correspondingly, the first energy level has a degeneracy of N.

To proceed, we introduce a diagrammatic method to list the different states. We draw only the "broken" bonds which connect two spins with opposite orientation and, as in the diagram above, denote these by red lines. We further draw the flipped spins as red dots, the unflipped spins as blue dots. The energy of the state is determined simply by the number of red lines in the diagram. Pictorially, we write the first excited state as

$E_1 = E_0 + 8J$

Degeneracy $= N$

The next lowest state has six broken bonds. It takes the form

$$E_2 = E_0 + 12J$$
$$\text{Degeneracy} = 2N$$

where the extra factor of 2 in the degeneracy comes from the two possible orientations (vertical and horizontal) of the graph.

Things are more interesting for the states which sit at the third excited level. These have 8 broken bonds. The simplest configuration consists of two, disconnected, flipped spins

$$E_3 = E_0 + 16J$$
$$\text{Degeneracy} = \tfrac{1}{2}N(N-5) \tag{5.28}$$

The factor of N in the degeneracy comes from placing the first graph; the factor of $N-5$ arises because the flipped spin in the second graph can sit anywhere apart from on the five vertices used in the first graph. Finally, the factor of $1/2$ arises from the interchange of the two graphs.

There are also three further graphs with the same energy E_3. These are

$$E_3 = E_0 + 16J$$
$$\text{Degeneracy} = N$$

and

$$E_3 = E_0 + 16J$$
$$\text{Degeneracy} = 2N$$

where the degeneracy comes from the two orientations (vertical and horizontal). And, finally,

$$E_3 = E_0 + 16J$$
$$\text{Degeneracy} = 4N$$

where the degeneracy comes from the four orientations (rotating the graph by 90°).

Adding all the graphs above together gives us an expansion of the partition function in power of $e^{-\beta J} \ll 1$. This is

$$Z = 2e^{2N\beta J}\left(1 + Ne^{-8\beta J} + 2Ne^{-12\beta J} + \frac{1}{2}(N^2 + 9N)e^{-16\beta J} + \ldots\right) \tag{5.29}$$

where the overall factor of 2 originates from the two ground states of the system. We'll make use of the specific coefficients in this expansion in Section 5.3.4. Before we focus on the physics hiding in the low temperature expansion, it's worth making a quick comment that something quite nice happens if we take the log of the partition function,

$$\log Z = \log 2 + 2N\beta J + Ne^{-8\beta J} + 2Ne^{-12\beta J} + \frac{9}{2}Ne^{-16\beta J} + \ldots$$

The thing to notice is that the N^2 term in the partition function (5.29) has cancelled out and $\log Z$ is proportional to N, which is to be expected since the free energy of the system is extensive. Looking back, we see that the N^2 term was associated to the disconnected diagrams in (5.28). There is actually a general lesson hiding here: the partition function can be written as the exponential of the sum of connected diagrams. We saw exactly the same issue arise in the cluster expansion in (2.37).

Peierls Droplets

Continuing the low temperature expansion provides a heuristic, but physically intuitive, explanation for why phase transitions happen in $d \geq 2$ dimensions but not in $d = 1$. As we flip more and more spins, the low energy states become *droplets*, consisting of a region of space in which all the spins are flipped, surrounded by a larger sea in which the spins have their original alignment. The energy cost of such a droplet is roughly

$$E \sim 2JL$$

where L is the perimeter of the droplet. Notice that the energy does not scale as the area of the droplet since all spins inside are aligned with their neighbours. It is only those on the edge which are misaligned and this is the reason for the perimeter scaling. To understand how these droplets contribute to the partition function, we also need to know their degeneracy. We will now argue that the degeneracy of droplets scales as

$$\text{Degeneracy} \sim e^{\alpha L}$$

for some value of α. To see this, consider firstly the problem of a random walk on a 2d square lattice. At each step, we can move in one of four directions. So the number of paths of length L is

$$\#\text{paths} \sim 4^L = e^{L \log 4}$$

Of course, the perimeter of a droplet is more constrained that a random walk. Firstly, the perimeter can't go back on itself, so it really only has three directions that it can move in at each step. Secondly, the perimeter must return to its starting point after L steps. And, finally, the perimeter cannot self-intersect. One can show that the number of paths that obey these conditions is

$$\#\text{paths} \sim e^{\alpha L}$$

where $\log 2 < \alpha < \log 3$. Since the degeneracy scales as $e^{\alpha L}$, the entropy of the droplets is proportional to L.

The fact that both energy and entropy scale with L means that there is an interesting competition between them. At temperatures where the droplets are important, the partition function is schematically of the form

$$Z \sim \sum_L e^{\alpha L} e^{-2\beta J L}$$

For large β (i.e. low temperature) the partition function converges. However, as the temperature increases, one reaches the critical temperature

$$k_B T_c \approx \frac{2J}{\alpha} \tag{5.30}$$

where the partition function no longer converges. At this point, the entropy wins over the energy cost and it is favourable to populate the system with droplets of arbitrary sizes. This is the how one sees the phase transition in the partition function. For temperature above T_c, the low-temperature expansion breaks down and the ordered magnetic phase is destroyed.

We can also use the droplet argument to see why phase transitions don't occur in $d = 1$ dimension. On a line, the boundary of any droplet always consists of just two points. This means that the energy cost to forming a droplet is always $E = 2J$, regardless of the size of the droplet. But, since the droplet can exist anywhere along the line, its degeneracy is N. The net result is that the free energy associated to creating a droplet scales as

$$F \sim 2J - k_B T \log N$$

and, as $N \to \infty$, the free energy is negative for any $T > 0$. This means that the system will prefer to create droplets of arbitrary length, randomizing the spins. This is the intuitive reason why there is no magnetic ordered phase in the $d = 1$ Ising model.

5.3.3 2d Ising Model: High Temperatures

We now turn to the 2d Ising model in the opposite limit of high temperature. Here we expect the partition function to be dominated by the completely random, disordered configurations of maximum entropy. Our goal is to find a way to expand the partition function in $\beta J \ll 1$.

We again work with zero magnetic field, $B = 0$ and write the partition function as

$$Z = \sum_{\{s_i\}} \exp\left(\beta J \sum_{\langle ij \rangle} s_i s_j\right) = \sum_{\{s_i\}} \prod_{\langle ij \rangle} e^{\beta J s_i s_j}$$

There is a useful way to rewrite $e^{\beta J s_i s_j}$ which relies on the fact that the product $s_i s_j$ only takes ± 1. It doesn't take long to check the following identity:

$$\begin{aligned} e^{\beta J s_i s_j} &= \cosh \beta J + s_i s_j \sinh \beta J \\ &= \cosh \beta J \left(1 + s_i s_j \tanh \beta J\right) \end{aligned}$$

Using this, the partition function becomes

$$\begin{aligned} Z &= \sum_{\{s_i\}} \prod_{\langle ij \rangle} \cosh \beta J \left(1 + s_i s_j \tanh \beta J\right) \\ &= (\cosh \beta J)^{qN/2} \sum_{\{s_i\}} \prod_{\langle ij \rangle} \left(1 + s_i s_j \tanh \beta J\right) \end{aligned} \tag{5.31}$$

where the number of nearest neighbours is $q = 4$ for the 2d square lattice.

With the partition function in this form, there is a natural expansion which suggests itself. At high temperatures $\beta J \ll 1$ which, of course, means that $\tanh \beta J \ll 1$. But the partition function is now naturally a product of powers of $\tanh \beta J$. This is somewhat analogous to the cluster expansion for the interacting gas that we met in Section 2.5.3. As in the cluster expansion, we will represent the expansion graphically.

We need no graphics for the leading order term. It has no factors of $\tanh \beta J$ and is simply

$$Z \approx (\cosh \beta J)^{2N} \sum_{\{s_i\}} 1 = 2^N (\cosh \beta J)^{2N}$$

That's simple.

Let's now turn to the leading correction. Expanding the partition function (5.31), each power of $\tanh \beta J$ is associated to a nearest neighbour pair $\langle ij \rangle$. We'll represent this by drawing a line on the lattice:

$$= \; s_i s_j \tanh \beta J$$

But there's a problem: each factor of $\tanh \beta J$ in (5.31) also comes with a sum over all spins s_i and s_j. And these are $+1$ and -1 which means that they simply sum to zero,

$$\sum_{s_i, s_j} s_i s_j = +1 - 1 - 1 + 1 = 0$$

How can we avoid this? The only way is to make sure that we're summing over an even number of spins on each site, since then we get factors of $s_i^2 = 1$ and no cancellations. Graphically, this means that every site must have an even number of lines attached to it. The first correction is then of the form

$$= \; (\tanh \beta J)^4 \sum_{\{s_i\}} s_1 s_2 \, s_2 s_3 \, s_3 s_4 \, s_4 s_1 = 2^4 (\tanh \beta J)^4$$

There are N such terms since the upper left corner of the square can be on any one of the N lattice sites. (Assuming periodic boundary conditions for the lattice). So including the leading term and first correction, we have

$$Z = 2^N (\cosh \beta J)^{2N} \left(1 + N (\tanh \beta J)^4 + \ldots\right)$$

We can go further. The next terms arise from graphs of length 6 and the only possibilities are rectangles, oriented as either landscape or portrait. Each of them can sit on

one of N sites, giving a contribution

$$= \; 2N(\tanh \beta J)^4$$

Things get more interesting when we look at graphs of length 8. We have four different types of graphs. Firstly, there are the trivial, disconnected pair of squares

$$= \; \frac{1}{2}N(N-5)(\tanh \beta J)^8$$

Here the first factor of N is the possible positions of the first square; the factor of $N-5$ arises because the possible location of the upper corner of the second square can't be on any of the vertices of the first, but nor can it be on the square one to the left of the upper corner of the first since that would give a graph that looks like which has three lines coming off the middle site and therefore vanishes when we sum over spins. Finally, the factor of 1/2 comes because the two squares are identical.

The other graphs of length 8 are a large square, a rectangle and a corner. The large square gives a contribution

$$= \; N(\tanh \beta J)^8$$

There are two orientations for the rectangle. Including these gives a factor of 2,

$$= \; 2N(\tanh \beta J)^8$$

Finally, the corner graph has four orientations, giving

$$= \; 4N(\tanh \beta J)^8$$

Adding all contributions together gives us the first few terms in high temperature expansion of the partition function

$$Z = 2^N(\cosh \beta J)^{2N}\Big(1 + N(\tanh \beta J)^4 + 2N(\tanh \beta J)^6 + \frac{1}{2}(N^2+9N)(\tanh \beta J)^8 + \ldots\Big) \tag{5.32}$$

There's some magic hiding in this expansion which we'll turn to in Section 5.3.4. First, let's just see how the high energy expansion plays out in the $d = 1$ dimensional Ising model.

The Ising Chain Revisited

Let's do the high temperature expansion for the $d = 1$ Ising chain with periodic boundary conditions and $B = 0$. We have the same partition function (5.31) and the same issue that only graphs with an even number of lines attached to each vertex contribute. But, for the Ising chain, there is only one such term: it is the closed loop. This means that the partition function is

Figure 48:

$$Z = 2^N (\cosh \beta J)^N \left(1 + (\tanh \beta J)^N\right)$$

In the limit $N \to \infty$, $(\tanh \beta J)^N \to 0$ at high temperatures and even the contribution from the closed loop vanishes. We're left with

$$Z = (2 \cosh \beta J)^N$$

This agrees with our exact result for the Ising chain given in (5.27), which can be seen by setting $B = 0$ in (5.26) so that $\lambda_+ = 2 \cosh \beta J$.

5.3.4 Kramers-Wannier Duality

In the previous sections we computed the partition function perturbatively in two extreme regimes of low temperature and high temperature. The physics in the two cases is, of course, very different. At low temperatures, the partition function is dominated by the lowest energy states; at high temperatures it is dominated by maximally disordered states. Yet comparing the partition functions at low temperature (5.29) and high temperature (5.32) reveals an extraordinary fact: the expansions are the same! More concretely, the two series agree if we exchange

$$e^{-2\beta J} \longleftrightarrow \tanh \beta J \tag{5.33}$$

Of course, we've only checked the agreement to the first few orders in perturbation theory. Below we shall prove that this miracle continues to all orders in perturbation theory. The symmetry of the partition function under the interchange (5.33) is known as *Kramers-Wannier duality*. Before we prove this duality, we will first just assume that it is true and extract some consequences.

We can express the statement of the duality more clearly. The Ising model at temperature β is related to the same model at temperature $\tilde{\beta}$, defined as

$$e^{-2\tilde{\beta}J} = \tanh \beta J \tag{5.34}$$

This way of writing things hides the symmetry of the transformation. A little algebra shows that this is equivalent to

$$\sinh 2\tilde{\beta}J = \frac{1}{\sinh 2\beta J}$$

Notice that this is a hot/cold duality. When βJ is large, $\tilde{\beta}J$ is small. Kramers-Wannier duality is the statement that, when $B = 0$, the partition functions of the Ising model at two temperatures are related by

$$\begin{aligned} Z[\beta] &= \frac{2^N(\cosh \beta J)^{2N}}{2e^{2N\tilde{\beta}J}} Z[\tilde{\beta}] \\ &= 2^{N-1}(\cosh \beta J \sinh \beta J)^N Z[\tilde{\beta}] \end{aligned} \tag{5.35}$$

This means that if you know the thermodynamics of the Ising model at one temperature, then you also know the thermodynamics at the other temperature. Notice however, that it does not say that all the physics of the two models is equivalent. In particular, when one system is in the ordered phase, the other typically lies in the disordered phase.

One immediate consequence of the duality is that we can use it to compute the exact critical temperature T_c. This is the temperature at which the partition function in singular in the $N \to \infty$ limit. (We'll discuss a more refined criterion in Section 5.4.3). If we further assume that there is just a single phase transition as we vary the temperature, then it must happen at the special self-dual point $\beta = \tilde{\beta}$. This is

$$k_B T = \frac{2J}{\log(\sqrt{2}+1)} \approx 2.269\, J$$

The exact solution of Onsager confirms that this is indeed the transition temperature. It's also worth noting that it's fully consistent with the more heuristic Peierls droplet argument (5.30) since $\log 2 < \log(\sqrt{2}+1) < \log 3$.

Proving the Duality

So far our evidence for the duality (5.35) lies in the agreement of the first few terms in the low and high temperature expansions (5.29) and (5.32). Of course, we could keep computing further and further terms and checking that they agree, but it would be nicer to simply prove the equality between the partition functions. We shall do so here.

The key idea that we need can actually be found by staring hard at the various graphs that arise in the two expansions. Eventually, you will realise that they are the same, albeit drawn differently. For example, consider the two "corner" diagrams

The two graphs are *dual*. The red lines in the first graph intersect the black lines in the second as can be seen by placing them on top of each other:

The same pattern occurs more generally: the graphs appearing in the low temperature expansion are in one-to-one correspondence with the dual graphs of the high temperature expansion. Here we will show how this occurs and how one can map the partition functions onto each other.

Let's start by writing the partition function in the form (5.31) that we met in the high temperature expansion and presenting it in a slightly different way,

$$\begin{aligned} Z[\beta] &= \sum_{\{s_i\}} \prod_{\langle ij \rangle} (\cosh \beta J + s_i s_j \sinh \beta J) \\ &= \sum_{\{s_i\}} \prod_{\langle ij \rangle} \sum_{k_{ij}=0,1} C_{k_{ij}}[\beta J] \, (s_i s_j)^{k_{ij}} \end{aligned}$$

where we have introduced the rather strange variable k_{ij} associated to each nearest neighbour pair that takes values 0 and 1, together with the functions.

$$C_0[\beta J] = \cosh \beta J \quad \text{and} \quad C_1[\beta J] = \sinh \beta J$$

The variables in the original Ising model were spins on the lattice sites. The observation that the graphs which appear in the two expansions are dual suggests that it might be

profitable to focus attention on the links between lattice sites. Clearly, we have one link for every nearest neighbour pair. If we label these links by l, we can trivially rewrite the partition function as

$$Z = \sum_{k_l=0,1} \prod_l \sum_{\{s_i\}} C_{k_l}[\beta J]\,(s_i s_j)^{k_l}$$

Notice that the strange label k_{ij} has now become a variable that lives on the links l rather than the original lattice sites i.

At this stage, we do the sum over the spins s_i. We've already seen that if a given spin, say s_i, appears in a term an odd number of times, then that term will vanish when we sum over the spin. Alternatively, if the spin s_i appears an even number of times, then the sum will give 2. We'll say that a given link l is turned on in configurations with $k_l = 1$ and turned off when $k_l = 0$. In this language, a term in the sum over spin s_i contributes only if an even number of links attached to site i are turned on. The partition function then becomes

$$Z = 2^N \sum_{k_l} \prod_l C_{k_l}[\beta J]\Big|_{\text{Constrained}} \tag{5.36}$$

Now we have something interesting. Rather than summing over spins on lattice sites, we're now summing over the new variables k_l living on links. This looks like the partition function of a totally different physical system, where the degrees of freedom live on the links of the original lattice. But there's a catch – that big "Constrained" label on the sum. This is there to remind us that we don't sum over all k_l configurations; only those for which an even number of links are turned on for every lattice site. And that's annoying. It's telling us that the k_l aren't really independent variables. There are some constraints that must be imposed.

Fortunately, for the 2d square lattice, there is a simple way to solve the constraint. We introduce yet more variables, $\tilde{s}_i$ which, like the original spin variables, take values ± 1. However, the $\tilde{s}_i$ do not live on the original lattice sites. Instead, they live on the vertices of the dual lattice. For the 2d square lattice, the dual vertices are drawn in the figure. The original lattice sites are in white; the dual lattice sites in black.

Figure 49:

The link variables k_l are related to the two nearest spin variables $\tilde{s}_i$ as follows:

$$k_{12} = \frac{1}{2}(1 - \tilde{s}_1 \tilde{s}_2)$$

$$
\begin{aligned}
k_{13} &= \frac{1}{2}(1-\tilde{s}_2\tilde{s}_3) \\
k_{14} &= \frac{1}{2}(1-\tilde{s}_3\tilde{s}_4) \\
k_{15} &= \frac{1}{2}(1-\tilde{s}_1\tilde{s}_4)
\end{aligned}
$$

Notice that we've replaced four variables k_l taking values $0, 1$ with four variables $\tilde{s}_i$ taking values ± 1. Each set of variables gives 2^4 possibilities. However, the map is not one-to-one. It is not possible to construct for all values of k_l using the parameterization in terms of $\tilde{s}_i$. To see this, we need only look at

$$
\begin{aligned}
k_{12}+k_{13}+k_{14}+k_{15} &= 2-\frac{1}{2}(\tilde{s}_1\tilde{s}_2+\tilde{s}_2\tilde{s}_3+\tilde{s}_3\tilde{s}_4+\tilde{s}_1\tilde{s}_4) \\
&= 2-\frac{1}{2}(\tilde{s}_1+\tilde{s}_3)(\tilde{s}_2+\tilde{s}_4) \\
&= 0, 2, \text{or } 4
\end{aligned}
$$

In other words, the number of links that are turned on must be even. But that's exactly what we want! Writing the k_l in terms of the auxiliary spins $\tilde{s}_i$ automatically solves the constraint that is imposed on the sum in (5.36). Moreover, it is simple to check that for every configuration $\{k_l\}$ obeying the constraint, there are two configurations of $\{\tilde{s}_i\}$. This means that we can replace the constrained sum over $\{k_l\}$ with an unconstrained sum over $\{\tilde{s}_i\}$. The only price we pay is an additional factor of $1/2$.

$$
Z[\beta] = \frac{1}{2}\,2^N \sum_{\{\tilde{s}_i\}} \prod_{\langle ij\rangle} C_{\frac{1}{2}(1-\tilde{s}_i\tilde{s}_j)}[\beta j]
$$

Finally, we'd like to find a simple expression for C_0 and C_1 in terms of $\tilde{s}_i$. That's easy enough. We can write

$$
\begin{aligned}
C_k[\beta J] &= \cosh\beta J\,\exp\left(k\log\tanh\beta J\right) \\
&= (\sinh\beta J\cosh\beta J)^{1/2}\exp\left(-\frac{1}{2}\tilde{s}_i\tilde{s}_j\log\tanh\beta J\right)
\end{aligned}
$$

Substituting this into our newly re-written partition function gives

$$
\begin{aligned}
Z[\beta] &= 2^{N-1}\sum_{\{\tilde{s}_i\}}\prod_{\langle ij\rangle}(\sinh\beta J\cosh\beta J)^{1/2}\exp\left(-\frac{1}{2}\tilde{s}_i\tilde{s}_j\log\tanh\beta J\right) \\
&= 2^{N-1}(\sinh\beta J\cosh\beta J)^N\sum_{\{\tilde{s}_i\}}\exp\left(-\frac{1}{2}\log\tanh\beta J\sum_{\langle ij\rangle}\tilde{s}_i\tilde{s}_j\right)
\end{aligned}
$$

But this final form of the partition function in terms of the dual spins $\tilde{s}_i$ has exactly the same functional form as the original partition function in terms of the spins s_i. More precisely, we can write

$$Z[\beta] = 2^{N-1}(\sinh 2\beta J)^N Z[\tilde{\beta}]$$

where

$$e^{-2\tilde{\beta}J} = \tanh \beta J$$

as advertised previously in (5.34). This completes the proof of Kramers-Wannier duality in the 2d Ising model on a square lattice.

The concept of *duality* of this kind is a major feature in much of modern theoretical physics. The key idea is that when the temperature gets large there may be a different set of variables in which a theory can be written where it appears to live at low temperature. The same idea often holds in quantum theories, where duality maps strong coupling problems to weak coupling problems.

The duality in the Ising model is special for two reasons: firstly, the new variables $\tilde{s}_i$ are governed by the same Hamiltonian as the original variables s_i. We say that the Ising model is self-dual. In general, this need not be the case — the high temperature limit of one system could look like the low-temperature limit of a very different system. Secondly, the duality in the Ising model can be proven explicitly. For most systems, we have no such luck. Nonetheless, the idea that there may be dual variables in other, more difficult theories, is compelling. Commonly studied examples include the exchange particles and vortices in two dimensions, and electrons and magnetic monopoles in three dimensions.

5.4 Landau Theory

We saw in Sections 5.1 and 5.2 that the van der Waals equation and mean field Ising model gave the same (sometimes wrong!) answers for the critical exponents. This suggests that there should be a unified way to look at phase transitions. Such a method was developed by Landau. It is worth stressing that, as we saw above, the Landau approach to phase transitions often only gives qualitatively correct results. However, its advantage is that it is extremely straightforward and easy. (Certainly much easier than the more elaborate methods needed to compute critical exponents more accurately).

The Landau theory of phase transitions is based around the free energy. We will illustrate the theory using the Ising model and then explain how to extend it to different systems. The free energy of the Ising model in the mean field approximation is readily attainable from the partition function (5.17),

$$F = -\frac{1}{\beta}\log Z = \frac{1}{2}JNqm^2 - \frac{N}{\beta}\log\left(2\cosh\beta B_{\text{eff}}\right) \tag{5.37}$$

So far in this course, we've considered only systems in equilibrium. The free energy, like all other thermodynamic potentials, has only been defined on equilibrium states. Yet the equation above can be thought of as an expression for F as a function of m. Of course, we could substitute in the equilibrium value of m given by solving (5.18), but it seems a shame to throw out $F(m)$ when it is such a nice function. Surely we can put it to some use!

The key step in Landau theory is to treat the function $F = F(T, V; m)$ seriously. This means that we are extending our viewpoint away from equilibrium states to a whole class of states which have a constant average value of m. If you want some words to drape around this, you could imagine some external magical power that holds m fixed. The free energy $F(T, V; m)$ is then telling us the equilibrium properties in the presence of this magical power. Perhaps more convincing is what we do with the free energy in the absence of any magical constraint. We saw in Section 4 that equilibrium is guaranteed if we sit at the minimum of F. Looking at extrema of F, we have the condition

$$\frac{\partial F}{\partial m} = 0 \quad \Rightarrow \quad m = \tanh\beta B_{\text{eff}}$$

But that's precisely the condition (5.18) that we saw previously. Isn't that nice!

In the context of Landau theory, m is called an *order parameter*. When it takes non-zero values, the system has some degree of order (the spins have a preferred direction in which they point) while when $m = 0$ the spins are randomised and happily point in any direction.

For any system of interest, Landau theory starts by identifying a suitable order parameter. This should be taken to be a quantity which vanishes above the critical temperature at which the phase transition occurs, but is non-zero below the critical temperature. Sometimes it is obvious what to take as the order parameter; other times less so. For the liquid-gas transition, the relevant order parameter is the difference in densities between the two phases, $v_{\text{gas}} - v_{\text{liquid}}$. For magnetic or electric systems, the order parameter is typically some form of magnetization (as for the Ising model) or

the polarization. For the Bose-Einstein condensate, superfluids and superconductors, the order parameter is more subtle and is related to off-diagonal long-range order in the one-particle density matrix[11], although this is usually rather lazily simplified to say that the order parameter can be thought of as the macroscopic wavefunction $|\psi|^2$.

Starting from the existence of a suitable order parameter, the next step in the Landau programme is to write down the free energy. But that looks tricky. The free energy for the Ising model (5.37) is a rather complicated function and clearly contains some detailed information about the physics of the spins. How do we just write down the free energy in the general case? The trick is to assume that we can expand the free energy in an analytic power series in the order parameter. For this to be true, the order parameter must be small which is guaranteed if we are close to a critical point (since $m = 0$ for $T > T_c$). The nature of the phase transition is determined by the kind of terms that appear in the expansion of the free energy. Let's look at a couple of simple examples.

5.4.1 Second Order Phase Transitions

We'll consider a general system (Ising model; liquid-gas; BEC; whatever) and denote the order parameter as m. Suppose that the expansion of the free energy takes the general form

$$F(T;m) = F_0(T) + a(T)m^2 + b(T)m^4 + \ldots \tag{5.38}$$

One common reason why the free energy has this form is because the theory has a symmetry under $m \to -m$, forbidding terms with odd powers of m in the expansion. For example, this is the situation in the Ising model when $B = 0$. Indeed, if we expand out the free energy (5.37) for the Ising model for small m using $\cosh x \approx 1 + \frac{1}{2}x^2 + \frac{1}{4!}x^4 + \ldots$ and $\log(1+y) \approx y - \frac{1}{2}y^2 + \ldots$ we get the general form above with explicit expressions for $F_0(T)$, $a(T)$ and $b(T)$,

$$F_{Ising}(T;m) = -Nk_BT\log 2 + \left(\frac{NJq}{2}(1 - Jq\beta)\right)m^2 + \left(\frac{N\beta^3J^4q^4}{24}\right)m^4 + \ldots$$

The leading term $F_0(T)$ is unimportant for our story. We are interested in how the free energy changes with m. The condition for equilibrium is given by

$$\frac{\partial F}{\partial m} = 0 \tag{5.39}$$

But the solutions to this equation depend on the sign of the coefficients $a(T)$ and

[11] See, for example, the book "*Quantum Liquids*" by Anthony Leggett

Figure 50: Free energy when $a(T) > 0$

Figure 51: Free energy when $a(T) < 0$

$b(T)$. Moreover, this sign can change with temperature. This is the essence of the phase transitions. In the following discussion, we will assume that $b(T) > 0$ for all T. (If we relax this condition, we have to also consider the m^6 term in the free energy which leads to interesting results concerning so-called tri-critical points).

The two figures above show sketches of the free energy in the case where $a(T) > 0$ and $a(T) < 0$. Comparing to the explicit free energy of the Ising model, $a(T) < 0$ when $T > T_c = Jq/k_B$ and $a(T) < 0$ when $T < T_c$. When $a(T) > 0$, we have just a single equilibrium solution to (5.39) at $m = 0$. This is typically the situation at high temperatures. In contrast, at $a(T) < 0$, there are three solutions to (5.39). The solution $m = 0$ clearly has higher free energy: this is now the unstable solution. The two stable solutions sit at $m = \pm m_0$. For example, if we choose to truncate the free energy (5.38) at quartic order, we have

$$m_0 = \sqrt{\frac{-a}{2b}} \qquad T < T_c$$

If $a(T)$ is a smooth function then the equilibrium value of m changes continuously from $m = 0$ when $a(T) > 0$ to $m \neq 0$ at $a(T) < 0$. This describes a second order phase transition occurring at T_c, defined by $a(T_c) = 0$.

Once we know the equilibrium value of m, we can then substitute this back into the free energy $F(T; m)$ in (5.38). This gives the thermodynamic free energy $F(T)$ of the system in equilibrium that we have been studying throughout this course. For the quartic free energy, we have

$$F(T) = \begin{cases} F_0(T) & T > T_c \\ F_0(T) - a^2/4b & T < T_c \end{cases} \tag{5.40}$$

Because $a(T_c) = 0$, the equilibrium free energy $F(T)$ is continuous at $T = T_c$. Moreover, the entropy $S = -\partial F/\partial T$ is also continuous at $T = T_c$. However, if you differentiate the

equilibrium free energy twice, you will get a term a'^2/b which is generically not vanishing at $T = T_c$. This means that the heat capacity $C = T\partial S/\partial T$ changes discontinuously at $T = T_c$, as befits a second order phase transition. A word of warning: if you want to compute equilibrium quantities such as the heat capacity, it's important that you first substitution in the equilibrium value of m and work with (5.40) rather than i (5.38). If you don't, you miss the fact that the magnetization also changes with T.

We can easily compute critical exponents within the context of Landau theory. We need only make further assumptions about the behaviour of $a(T)$ and $b(T)$ in the vicinity of T_c. If we assume that near $T = T_c$, we can write

$$b(T) \approx b_0 \quad , \quad a(T) \approx a_0(T - T_c) \tag{5.41}$$

then we have

$$m_0 \approx \pm\sqrt{\frac{a_0}{2b_0}}(T_c - T)^{1/2} \qquad T < T_c$$

which reproduces the critical exponent (5.9) and (5.20) that we derived for the van der Waals equation and Ising model respectively.

Notice that we didn't put any discontinuities into the free energy. Everything in $F(T; m)$ was nice and smooth. When Taylor expanded, it has only integer powers of m and T as shown in (5.38) and (5.41). But the minima of F behave in a non-analytic fashion as seen in the expression for m_0 above.

Landau's theory of phase transitions predicts this same critical exponent for all values of the dimension d of the system. But we've already mentioned in previous contexts that the critical exponent is in fact only correct for $d \geq 4$. We will understand how to derive this criterion from Landau theory in the next section.

Spontaneous Symmetry Breaking

As we approach the end of the course, we're touching upon a number of ideas that become increasingly important in subsequent developments in physics. We already briefly met the idea of universality and critical phenomena. Here I would like to point out another very important idea: *spontaneous symmetry breaking*.

The free energy (5.38) is invariant under the $\mathbf{Z}_2$ symmetry $m \to -m$. Indeed, we said that one common reason that we can expand the free energy only in even powers of m is that the underlying theory also enjoys this symmetry. But below T_c, the system must pick one of the two ground states $m = +m_0$ or $m = -m_0$. Whichever choice it makes breaks the $\mathbf{Z}_2$ symmetry. We say that the symmetry is spontaneously broken by the choice of ground state of the theory.

Spontaneous symmetry breaking has particularly dramatic consequences when the symmetry in question is continuous rather than discrete. For example, consider a situation where the order parameter is a complex number ψ and the free energy is given by (5.38) with $m = |\psi|^2$. (This is effectively what happens for BECs, superfluids and superconductors). Then we should only look at the $m > 0$ solutions so that the ground state has $|\psi|^2 = +m_0$. But this leaves the phase of ψ completely undetermined. So there is now a continuous choice of ground states: we get to sit anywhere on the circle parameterised by the phase of ψ. Any choice that the system makes spontaneously breaks the $U(1)$ rotational symmetry which acts on the phase of ψ. Some beautiful results due to Nambu and Goldstone show that the much of the physics of these systems can be understood simply as a consequence of this symmetry breaking. The ideas of spontaneous symmetry breaking are crucial in both condensed matter physics and particle physics. In the latter context, it is intimately tied with the Higgs mechanism.

5.4.2 First Order Phase Transitions

Let us now consider a situation where the expansion of the free energy also includes odd powers of the order parameter

$$F(T;m) = F_0(T) + \alpha(T)m + a(T)m^2 + \gamma(T)m^3 + b(T)m^4 + \ldots$$

For example, this is the kind of expansion that we get for the Ising model free energy (5.37) when $B \neq 0$, which reads

$$F_{\text{Ising}}(T;m) = -Nk_BT\log 2 + \frac{JNq}{2}m^2 - \frac{N}{2k_BT}(B+Jqm)^2 + \frac{N}{24(k_BT)^3}(B+Jqm)^4 + \ldots$$

Notice that there is no longer a symmetry relating $m \to -m$: the B field has a preference for one sign over the other.

If we again assume that $b(T) > 0$ for all temperatures, the crude shape of the free energy graph again has two choices: there is a single minimum, or two minima and a local maximum.

Let's start at suitably low temperatures for which the situation is depicted in Figure 52. The free energy once again has a double well, except now slightly skewed. The local maximum is still an unstable point. But this time around, the minima with the lower free energy is preferred over the other one. This is the true ground state of the system. In contrast, the point which is locally, but not globally, a minimum corresponds to a meta-stable state of the system. In order for the system to leave this state, it must first fluctuate up and over the energy barrier separating the two.

Figure 52: The free energy of the Ising model for $B < 0$, $B = 0$ and $B > 0$.

In this set-up, we can initiate a first order phase transition. This occurs when the coefficient of the odd terms, $\alpha(T)$ and $\gamma(T)$ change sign and the true ground state changes discontinuously from $m < 0$ to $m > 0$. In some systems this behaviour occurs when changing temperature; in others it could occur by changing some external parameter. For example, in the Ising model the first order phase transition is induced by changing B.

At very high temperature, the double well potential is lost in favour of a single minimum as depicted in the figure to the right. There is a unique ground state, albeit shifted from $m = 0$ by the presence of the $\alpha(T)$ term above (which translates into the magnetic field B in the Ising model). The temperature at which the meta-stable ground state of the system is lost corresponds to the spinodial point in our discussion of the liquid-gas transition.

Figure 53:

One can play further games in Landau theory, looking at how the shape of the free energy can change as we vary temperature or other parameters. One can also use this framework to give a simple explanation of the concept of hysteresis. You can learn more about these from the links on the course webpage.

5.4.3 Lee-Yang Zeros

You may have noticed that the flavour of our discussion of phase transitions is a little different from the rest of the course. Until now, our philosophy was to derive everything from the partition function. But in this section, we dumped the partition function as soon as we could, preferring instead to work with the macroscopic variables such as the free energy. Why didn't we just stick with the partition function and examine phase transitions directly?

The reason, of course, is that the approach using the partition function is hard! In this short section, which is somewhat tangential to our main discussion, we will describe how phase transitions manifest themselves in the partition function.

For concreteness, let's go back to the classical interacting gas of Section 2.5, although the results we derive will be more general. We'll work in the grand canonical ensemble, with the partition function

$$\mathcal{Z}(z,V,T) = \sum_N z^N Z(N,V,T) = \sum_N \frac{z^N}{N!\lambda^{3N}} \int \prod_i d^3r_i \; e^{-\beta \sum_{j<k} U(r_{jk})} \tag{5.42}$$

To regulate any potential difficulties with short distances, it is useful to assume that the particles have hard-cores so that they cannot approach to a distance less than r_0. We model this by requiring that the potential satisfies

$$U(r_{jk}) = 0 \quad \text{for} \quad r_{jk} < r_0$$

But this has an obvious consequence: if the particles have finite size, then there is a maximum number of particles, N_V, that we can fit into a finite volume V. (Roughly this number is $N_V \sim V/r_0^3$). But that, in turn, means that the canonical partition function $Z(N,V,T) = 0$ for $N > N_V$, and the grand partition function $\mathcal{Z}$ is therefore a finite polynomial in the fugacity z, of order N_V. But if the partition function is a finite polynomial, there can't be any discontinuous behaviour associated with a phase transition. In particular, we can calculate

$$pV = k_B T \log \mathcal{Z} \tag{5.43}$$

which gives us pV as a smooth function of z. We can also calculate

$$N = z\frac{\partial}{\partial z} \log \mathcal{Z} \tag{5.44}$$

which gives us N as a function of z. Eliminating z between these two functions (as we did for both bosons and fermions in Section 3) tells us that pressure p is a smooth function of density N/V. We're never going to get the behaviour that we derived from the Maxwell construction in which the plot of pressure vs density shown in Figure 37 exhibits a discontinous derivative.

The discussion above is just re-iterating a statement that we've alluded to several times already: there are no phase transitions in a finite system. To see the discontinuous behaviour, we need to take the limit $V \to \infty$. A theorem due to Lee and Yang[12] gives us a handle on the analytic properties of the partition function in this limit.

[12]This theorem was first proven for the Ising model in 1952. Soon afterwards, the same Lee and Yang proposed a model of parity violation in the weak interaction for which they won the 1957 Nobel prize.

The surprising insight of Lee and Yang is that if you're interested in phase transitions, you should look at the zeros of $\mathcal{Z}$ in the complex z-plane. Let's firstly look at these when V is finite. Importantly, at finite V there can be no zeros on the positive real axis, $z > 0$. This follows follows from the defintion of $\mathcal{Z}$ given in (5.42) where it is a sum of positive quantities. Moreover, from (5.44), we can see that $\mathcal{Z}$ is a monotonically increasing function of z because we necessarily have $N > 0$. Nonetheless, $\mathcal{Z}$ is a polynomial in z of order N_V so it certainly has N_V zeros somewhere in the complex z-plane. Since $\mathcal{Z}^\star(z) = \mathcal{Z}(z^\star)$, these zeros must either sit on the real negative axis or come in complex pairs.

However, the statements above rely on the fact that $\mathcal{Z}$ is a finite polynomial. As we take the limit $V \to \infty$, the maximum number of particles that we can fit in the system diverges, $N_V \to \infty$, and $\mathcal{Z}$ is now defined as an infinite series. But infinite series can do things that finite ones can't. The Lee-Yang theorem says that as long as the zeros of $\mathcal{Z}$ continue to stay away from the positive real axis as $V \to \infty$, then no phase transitions can happen. But if one or more zeros happen to touch the positive real axis, life gets more interesting.

More concretely, the Lee-Yang theorem states:

- **Lee-Yang Theorem:** The quantity

$$\Theta = \lim_{V\to\infty} \left(\frac{1}{V} \log \mathcal{Z}(z, V, T) \right)$$

  exists for all $z > 0$. The result is a continuous, non-decreasing function of z which is independent of the shape of the box (up to some sensible assumptions such as Surface Area/$V \sim V^{-1/3}$ which ensures that the box isn't some stupid fractal shape).

  Moreover, let R be a fixed, volume independent, region in the complex z plane which contains part of the real, positive axis. If R contains no zero of $\mathcal{Z}(z, V, T)$ for all $z \in R$ then Θ is a an analytic function of z for all $z \in R$. In particular, all derivatives of Θ are continuous.

In other words, there can be no phase transitions in the region R even in the $V \to \infty$ limit. The last result means that, as long as we are safely in a region R, taking derivatives of with respect to z commutes with the limit $V \to \infty$. In other words, we are allowed to use (5.44) to write the particle density $n = N/V$ as

$$\lim_{V\to\infty} n = \lim_{V\to\infty} z\frac{\partial}{\partial z}\left(\frac{p}{k_B T}\right) = z\frac{\partial\Theta}{\partial z}$$

However, if we look at points z where zeros appear on the positive real axis, then Θ will generally not be analytic. If $d\Theta/dz$ is discontinuous, then the system is said to undergo a first order phase transition. More generally, if $d^m\Theta/dz^m$ is discontinuous for $m = n$, but continuous for all $m < n$, then the system undergoes an nth order phase transition. We won't offer a proof of the Lee-Yang theorem. Instead illustrate the general idea with an example.

A Made-Up Example

Ideally, we would like to start with a Hamiltonian which exhibits a first order phase transition, compute the associated grand partition function $\mathcal{Z}$ and then follow its zeros as $V \to \infty$. However, as we mentioned above, that's hard! Instead we will simply make up a partition function $\mathcal{Z}$ which has the appropriate properties. Our choice is somewhat artificial,

$$\mathcal{Z}(z, V) = (1 + z)^{[\alpha V]}(1 + z^{[\alpha V]})$$

Here α is a constant which will typically depend on temperature, although we'll suppress this dependence in what follows. Also,

$$[x] = \text{Integer part of } x$$

Although we just made up the form of $\mathcal{Z}$, it does have the behaviour that one would expect of a partition function. In particular, for finite V, the zeros sit at

$$z = -1 \quad \text{and} \quad z = e^{\pi i(2n+1)/[\alpha V]} \quad n = 0, 1, \ldots, [\alpha V] - 1$$

As promised, none of the zeros sit on the positive real axis.However, as we increase V, the zeros become denser and denser on the unit circle. From the Lee-Yang theorem, we expect that no phase transition will occur for $z \neq 1$ but that something interesting could happen at $z = 1$.

Let's look at what happens as we send $V \to \infty$. We have

$$\begin{aligned}
\Theta &= \lim_{V\to\infty} \frac{1}{V} \log \mathcal{Z}(z, V) \\
&= \lim_{V\to\infty} \frac{1}{V} \left([\alpha V] \log(1 + z) + \log(1 + z^{[\alpha V]})\right) \\
&= \begin{cases} \alpha \log(1 + z) & |z| < 1 \\ \alpha \log(1 + z) + \alpha \log z & |z| > 1 \end{cases}
\end{aligned}$$

We see that Θ is continuous for all z as promised. But it is only analytic for $|z| \neq 1$.

We can extract the physics by using (5.43) and (5.44) to eliminate the dependence on z. This gives us the equation of state, with pressure p as a function of $n = V/N$. For $|z| < 1$, we have

$$p = \alpha k_B T \log\left(\frac{\alpha}{\alpha - n}\right) \qquad n \in [0, \alpha/2) \quad , \quad p < k_B T \log 2$$

While for $|z| > 1$, we have

$$p = \alpha k_B T \log\left(\frac{2\alpha n}{(2\alpha - n)^2}\right) \qquad n \in (3\alpha/2, 2\alpha) \quad , \quad p > k_B T \log 2$$

They key point is that there is a jump in particle density of $\Delta n = \alpha$ at $p = \alpha k_B T \log 2$. Plotting this as a function of p vs $v = 1/n$, we find that we have a curve that is qualitatively identical to the pressure-volume plot of the liquid-gas phase diagram under the co-existence curve. (See, for example, figure 37). This is a first order phase transition.

5.5 Landau-Ginzburg Theory

Landau's theory of phase transition focusses only on the average quantity, the order parameter. It ignores the fluctuations of the system, assuming that they are negligible. Here we sketch a generalisation which attempts to account for these fluctuations. It is known as Landau-Ginzburg theory.

The idea is to stick with the concept of the order parameter, m. But now we allow the order parameter to vary in space so it becomes a function $m(\vec{r})$. Let's restrict ourselves to the situation where there is a symmetry of the theory $m \to -m$ so we need only consider even powers in the expansion of the free energy. We add to these a gradient term whose role is to captures the fact that there is some stiffness in the system, so it costs energy to vary the order parameter from one point to another. (For the example of the Ising model, this is simply the statement that nearby spins want to be aligned). The free energy is then given by

$$F[m(\vec{r})] = \int d^d r \ \left[a(T)m^2 + b(T)m^4 + c(T)(\nabla m)^2\right] \tag{5.45}$$

where we have dropped the constant $F_0(T)$ piece which doesn't depend on the order parameter and hence plays no role in the story. Notice that we start with terms quadratic in the gradient: a term linear in the gradient would violate the rotational symmetry of the system.

We again require that the free energy is minimised. But now F is a functional – it is a function of the function $m(\vec{r})$. To find the stationary points of such objects we need to use the same kind of variational methods that we use in Lagrangian mechanics. We write the variation of the free energy as

$$\begin{aligned}\delta F &= \int d^d r \ \left[2am\,\delta m + 4bm^3\,\delta m + 2c\nabla m\cdot\nabla\delta m\right] \\ &= \int d^d r \ \left[2am + 4bm^3 - 2c\nabla^2 m\right]\delta m\end{aligned}$$

where to go from the first line to the second we have integrated by parts. (We need to remember that $c(T)$ is a function of temperature but does not vary in space so that ∇ doesn't act on it). The minimum of the free energy is then determined by setting $\delta F = 0$ which means that we have to solve the Euler-Lagrange equations for the function $m(\vec{r})$,

$$c\nabla^2 m = am + 2bm^3 \tag{5.46}$$

The simplest solutions to this equation have m constant, reducing us back to Landau theory. We'll assume once again that $a(T) > 0$ for $T > T_c$ and $a(T) < 0$ for $T < T_c$). Then the constant solutions are $m = 0$ for $T < T_c$ and $m = \pm m_0 = \pm\sqrt{-a/2b}$ for $T < T_c$. However, allowing for the possibility of spatial variation in the order parameter also opens up the possibility for us to search for more interesting solutions.

Domain Walls

Suppose that we have $T < T_c$ so there exist two degenerate ground states, $m = \pm m_0$. We could cook up a situation in which one half of space, say $x < 0$, lives in the ground state $m = -m_0$ while the other half of space, $x > 0$ lives in $m = +m_0$. This is exactly the situation that we already met in the liquid-gas transition and is depicted in Figure 38. It is also easy to cook up the analogous configuration in the Ising model. The two regions in which the spins point up or down are called domains. The place where these regions meet is called the *domain wall*.

We would like to understand the structure of the domain wall. How does the system interpolate between these two states? The transition can't happen instantaneously because that would result in the gradient term $(\nabla m)^2$ giving an infinite contribution to the free energy. But neither can the transition linger too much because any point at which $m(\vec{r})$ differs significantly from the value m_0 costs free energy from the m^2 and m^4 terms. There must be a happy medium between these two.

To describe the system with two domains, $m(\vec{r})$ must vary but it need only change in one direction: $m = m(x)$. Equation (5.46) then becomes an ordinary differential equation,

$$\frac{d^2m}{dx^2} = \frac{am}{c} + \frac{2bm^3}{c}$$

This equation is easily solved. We should remember that in order to have two vacua, $T < T_c$ which means that $a < 0$. We then have

$$m = m_0 \tanh\left(\sqrt{\frac{-a}{2c}}x\right)$$

where $m_0 = \sqrt{-a/2b}$ is the constant ground state solution for the spin. As $x \to \pm\infty$, the tanh function tends towards ± 1 which means that $m \to \pm m_0$. So this solution indeed interpolates between the two domains as required. We learn that the width of the domain wall is given by $\sqrt{-2c/a}$. Outside of this region, the magnetisation relaxes exponentially quickly back to the ground state values.

We can also compute the cost in free energy due to the presence of the domain wall. To do this, we substitute the solution back into the expression for the free energy (5.45). The cost is not proportional to the volume of the system, but instead proportional to the area of the domain wall. This means that if the system has linear size L then the free energy of the ground state scales as L^d while the free energy required by the wall scales only as L^{d-1}. It is simple to find the parametric dependence of this domain wall energy without doing any integrals; the energy per unit area scales as $\sqrt{-ca^3/b}$. Notice that as we approach the critical point, and $a \to 0$, the two vacua are closer, the width of the domain wall increases and its energy decreases.

5.5.1 Correlations

One of the most important applications of Landau-Ginzburg theory is to understand the correlations between fluctuations of the system at different points in space. Suppose that we know that the system has an unusually high fluctuation away from the average at some point in space, let's say the origin $\vec{r} = 0$. What is the effect of this on nearby points?

There is a simple way to answer this question that requires us only to solve the differential equation (5.46). However, there is also a more complicated way to derive the same result which has the advantage of stressing the underlying physics and the role played by fluctuations. Below we'll start by deriving the correlations in the simple manner. We'll then see how it can also be derived using more technical machinery.

We assume that the system sits in a given ground state, say $m = +m_0$, and imagine small perturbations around this. We write the magnetisation as

$$m(\vec{r}) = m_0 + \delta m(\vec{r}) \tag{5.47}$$

If we substitute this into equation (5.46) and keep only terms linear in δm, we find

$$c\nabla^2 \delta m + \frac{2a}{c}\delta m = 0$$

where we have substituted $m_0^2 = -a/2b$ to get this result. (Recall that $a < 0$ in the ordered phase). We now perturb the system. This can be modelled by putting a delta-function source at the origin, so that the above equation becomes

$$c\nabla^2 \delta m + \frac{2a}{c}\delta m = \frac{1}{2c}\,\delta^d(0)$$

where the strength of the delta function has been chosen merely to make the equation somewhat nicer. It is straightforward to solve the asymptotic behaviour of this equation. Indeed, it is the same kind of equation that we already solved when discussing the Debye-Hückel model of screening. Neglecting constant factors, it is

$$\delta m(\vec{r}) \sim \frac{e^{-r/\xi}}{r^{(d-1)/2}} \tag{5.48}$$

This tells us how the perturbation decays as we move away from the origin. This equation has several names, reflecting the fact that it arises in many contexts. In liquids, it is usually called the *Ornstein-Zernicke* correlation. It also arises in particle physics as the Yukawa potential. The length scale ξ is called the *correlation length*

$$\xi = \sqrt{\frac{-c}{2a}} \tag{5.49}$$

The correlation length provides a measure of the distance it takes correlations to decay. Notice that as we approach a critical point, $a \to 0$ and the correlation length diverges. This provides yet another hint that we need more powerful tools to understand the physics at the critical point. We will now take the first baby step towards developing these tools.

5.5.2 Fluctuations

The main motivation to allow the order parameter to depend on space is to take into the account the effect of fluctuations. To see how we can do this, we first need to think a little more about the meaning of the quantity $F[m(\vec{r})]$ and what we can use it for.

To understand this point, it's best if we go back to basics. We know that the true free energy of the system can be equated with the log of the partition function (1.36). We'd like to call the true free energy of the system F because that's the notation that we've been using throughout the course. But we've now called the Landau-Ginzburg functional $F[m(\vec{r})]$ and, while it's closely related to the true free energy, it's not quite the same thing as we shall shortly see. So to save some confusion, we're going to change notation at this late stage and call the true free energy A. Equation (1.36) then reads $A = -k_B T \log Z$, which we write this as

$$e^{-\beta A} = Z = \sum_n e^{-\beta E_n}$$

We would like to understand the right way to view the functional $F[m(\vec{r})]$ in this framework. Here we give a heuristic and fairly handwaving argument. A fuller treatment involves the ideas of the renormalisation group.

The idea is that each microstate $|n\rangle$ of the system can be associated to some specific function of the spatially varying order parameter $m(\vec{r})$. To illustrate this, we'll talk in the language of the Ising model although the discussion generalises to any system. There we could consider associate a magnetisation $m(\vec{r})$ to each lattice site by simply averaging over all the spins within some distance of that point. Clearly, this will only lead to functions that take values on lattice sites rather than in the continuum. But if the functions are suitably well behaved it should be possible to smooth them out into continuous functions $m(\vec{r})$ which are essentially constant on distance scales smaller than the lattice spacing. In this way, we get a map from the space of microstates to the magnetisation, $|n\rangle \mapsto m(\vec{r})$. But this map is not one-to-one. For example, if the averaging procedure is performed over enough sites, flipping the spin on just a single site is unlikely to have much effect on the average. In this way, many microstates map onto the same average magnetisation. Summing over just these microstates provides a first principles construction of the $F[m(\vec{r})]$,

$$e^{-\beta F[m(\vec{r})]} = \sum_{n|m(\vec{r})} e^{-\beta E_n} \tag{5.50}$$

Of course, we didn't actually perform this procedure to get to (5.45): we simply wrote it down the most general form in the vicinity of a critical point with a bunch of unknown coefficients $a(T)$, $b(T)$ and $c(T)$. But if we were up for a challenge, the above procedure tells us how we could go about figuring out those functions from first principles. More importantly, it also tells us what we should do with the Landau-Ginzburg free energy. Because in (5.50) we have only summed over those states that correspond to a particular

value of $m(\vec{r})$. To compute the full partition function, we need to sum over all states. But we can do that by summing over all possible values of $m(\vec{r})$. In other words,

$$Z = \int Dm(\vec{r})\ e^{-\beta F[m(\vec{r})]} \tag{5.51}$$

This is a tricky beast: it is a *functional integral*. We are integrating over all possible function $m(\vec{r})$, which is the same thing as performing an infinite number of integrations. (Actually, because the order parameters $m(\vec{r})$ arose from an underlying lattice and are suitably smooth on short distance scales, the problem is somewhat mitigated).

The result (5.51) is physically very nice, albeit mathematically somewhat daunting. It means that we should view the Landau-Ginzburg free energy as a new effective Hamiltonian for a continuous variable $m(\vec{r})$. It arises from performing the partition function sum over much of the microscopic information, but still leaves us with a final sum, or integral, over fluctuations in an averaged quantity, namely the order parameter.

To complete the problem, we need to perform the function integral (5.51). This is *hard*. Here "hard" means that the majority of unsolved problems in theoretical physics can be boiled down to performing integrals of this type. Yet the fact it's hard shouldn't dissuade us, since there is a wealth of rich and beautiful physics hiding in the path integral, including the deep reason behind the magic of universality. We will start to explore some of these ideas in next year's course on Statistical Field Theory.

Statistical Mechanics

Daniel F. Styer

May 2018

Statistical Mechanics

Dan Styer
Schiffer Professor of Physics
Oberlin College
Oberlin, Ohio 44074-1088

Dan.Styer@oberlin.edu
http://www.oberlin.edu/physics/dstyer

© 16 May 2018 Daniel F. Styer

The copyright holder grants the freedom to copy, modify, convey, adapt, and/or redistribute this work under the terms of the Creative Commons Attribution Share Alike 4.0 International License. A copy of that license is available at https://creativecommons.org/licenses/by-sa/4.0/legalcode.

Although, as a matter of history, statistical mechanics owes its origin to investigations in thermodynamics, it seems eminently worthy of an independent development, both on account of the elegance and simplicity of its principles, and because it yields new results and places old truths in a new light.

— J. Willard Gibbs
Elementary Principles in Statistical Mechanics

Contents

0 Preface **1**

1 The Properties of Matter in Bulk **4**

1.1 What is Statistical Mechanics About? 4

1.2 Outline of Book 5

1.3 Fluid Statics 6

1.4 Phase Diagrams 7

1.5 Additional Problems 8

2 Principles of Statistical Mechanics **11**

2.1 Microscopic Description of a Classical System 11

2.2 Macroscopic Description of a Large Equilibrium System 15

2.3 Fundamental Assumption 16

2.4 Statistical Definition of Entropy 19

2.5 Entropy of a Monatomic Ideal Gas 21

2.6 Qualitative Features of Entropy 27

2.7 Using Entropy to Find (Define) Temperature and Pressure 37

2.8 Additional Problems 47

3 Thermodynamics **49**

3.1 Heat and Work 49

3.2 Entropy 51

3.3 Heat Engines . . . 55
3.4 Multivariate Calculus . . . 57
3.5 Thermodynamic Quantities . . . 64
3.6 The Thermodynamic Dance . . . 67
3.7 Non-fluid Systems . . . 74
3.8 Thermodynamics Applied to Fluids . . . 75
3.9 Thermodynamics Applied to Phase Transitions . . . 83
3.10 Thermodynamics Applied to Chemical Reactions . . . 86
3.11 Thermodynamics Applied to Light . . . 90
3.12 Additional Problems . . . 95

4 Ensembles **99**
4.1 The Canonical Ensemble . . . 99
4.2 Meaning of the Term "Ensemble" . . . 101
4.3 Classical Monatomic Ideal Gas . . . 101
4.4 Energy Dispersion in the Canonical Ensemble . . . 104
4.5 Temperature as a Control Variable for Energy (Canonical Ensemble) . . . 107
4.6 The Equivalence of Canonical and Microcanonical Ensembles . . . 108
4.7 The Grand Canonical Ensemble . . . 108
4.8 The Grand Canonical Ensemble in the Thermodynamic Limit . . . 109
4.9 Summary of Major Ensembles . . . 111
4.10 Quantal Statistical Mechanics . . . 112
4.11 Ensemble Problems I . . . 114
4.12 Ensemble Problems II . . . 121

5 Classical Ideal Gases **123**
5.1 Classical Monatomic Ideal Gases . . . 123
5.2 Classical Diatomic Ideal Gases . . . 126
5.3 Heat Capacity of an Ideal Gas . . . 126
5.4 Specific Heat of a Hetero-nuclear Diatomic Ideal Gas . . . 131
5.5 Chemical Reactions Between Gases . . . 134
5.6 Problems . . . 134

6 Quantal Ideal Gases **138**

6.1 Introduction . . . 138

6.2 The Interchange Rule . . . 138

6.3 Quantum Mechanics of Independent Identical Particles . . . 139

6.4 Statistical Mechanics of Independent Identical Particles . . . 146

6.5 Quantum Mechanics of Free Particles . . . 151

6.6 Fermi-Dirac Statistics . . . 152

6.7 Bose-Einstein Statistics . . . 154

6.8 Specific Heat of the Ideal Fermion Gas . . . 159

6.9 Additional Problems . . . 167

7 Harmonic Lattice Vibrations **170**

7.1 The Problem . . . 170

7.2 Statistical Mechanics of the Problem . . . 170

7.3 Normal Modes for a One-dimensional Chain . . . 171

7.4 Normal Modes in Three Dimensions . . . 171

7.5 Low-temperature Heat Capacity . . . 171

7.6 More Realistic Models . . . 173

7.7 What is a Phonon? . . . 173

7.8 Additional Problems . . . 173

8 Interacting Classical Fluids **175**

8.1 Introduction . . . 175

8.2 Perturbation Theory . . . 177

8.3 Variational Methods . . . 181

8.4 Distribution Functions . . . 182

8.5 Correlations and Scattering . . . 185

8.6 The Hard Sphere Fluid . . . 185

9 Strongly Interacting Systems and Phase Transitions **188**

9.1 Introduction to Magnetic Systems and Models . . . 188

9.2 Free Energy of the One-Dimensional Ising Model . . . 188

9.3 The Mean-Field Approximation . . . 193

9.4 Correlation Functions in the Ising Model . . . 193

9.5 Computer Simulation . . . 199

9.6 Additional Problems . . . 208

A Series and Integrals **214**

B Evaluating the Gaussian Integral **215**

C Clinic on the Gamma Function **216**

D Volume of a Sphere in d Dimensions **218**

E Stirling's Approximation **221**

F The Euler-MacLaurin Formula and Asymptotic Series **223**

G Ramblings on the Riemann Zeta Function **224**

H Tutorial on Matrix Diagonalization **230**

H.1 What's in a name? . . . 230

H.2 Vectors in two dimensions . . . 231

H.3 Tensors in two dimensions . . . 233

H.4 Tensors in three dimensions . . . 236

H.5 Tensors in d dimensions . . . 237

H.6 Linear transformations in two dimensions . . . 238

H.7 What does "eigen" mean? . . . 239

H.8 How to diagonalize a symmetric matrix . . . 240

H.9 A glance at computer algorithms . . . 246

H.10 A glance at non-symmetric matrices and the Jordan form . . . 247

I Catalog of Misconceptions **251**

J Thermodynamic Master Equations **253**

K Useful Formulas **254**

Chapter 0

Preface

This is a book about statistical mechanics at the advanced undergraduate level. It assumes a background in classical mechanics through the concept of phase space, in quantum mechanics through the Pauli exclusion principle, and in mathematics through multivariate calculus. (Section 9.2 also assumes that you can can diagonalize a 2×2 matrix.)

The book is in draft form and I would appreciate your comments and suggestions for improvement. In particular, if you have an idea for a new problem, or for a new class of problems, I would appreciate hearing about it. If you are studying statistical mechanics and find that you are asking yourself "Why is... true" or "What would happen if... were changed?" or "I wish I were more familiar with...", then you have the germ of a new problem... so please tell me!

The specific goals of this treatment include:

- To demonstrate the extraordinary range of applicability of the ideas of statistical mechanics. These ideas are applicable to crystals and magnets, superconductors and solutions, surfaces and even bottles of light. I am always irritated by books that apply statistical mechanics only to fluids, or worse, only to the ideal gas.
- To introduce some modern concepts and tools such as correlation functions and computer simulation. While a full treatment awaits the final two chapters, the need for these tools is presented throughout the book.
- To develop qualitative insight as well as analytic and technical skills. Serving this goal are particular sections of text (e.g. section 2.6) and particular problems (e.g. problem 2.4) as well as overall attention to conceptual issues. Also in support of this goal is an attempt to eliminate misconceptions as well as present correct ideas (see appendix I, "Catalog of Misconceptions"). I particularly aim to foster a qualitative understanding of those central concepts, entropy and chemical potential, emphasizing that the latter is no more difficult to understand (and no less!) than temperature.

- To review classical and quantum mechanics, and mathematics, with a special emphasis on difficult points. It is my experience that some topics are so subtle that they are never learned on first exposure, no matter how well they are taught. A good example is the interchange rule for identical particles in quantum mechanics. The rule is so simple and easy to state, yet its consequences are so dramatic and far reaching, that any student will profit from seeing it treated independently in both a quantum mechanics course and a statistical mechanics course.

- To develop problem solving skills. In the sample problems I attempt to teach both strategy and tactics—rather than just show *how* to solve a problem I point out *why* the steps I use are profitable. (See the index entry on "problem solving tips".) Throughout I emphasize that doing a problem involves more than simply reaching an algebraic result: it also involves trying to understand that result and appreciate its significance.

The problems are a central part of this book. (Indeed, I wrote the problems first and designed the book around them.) In any physics course, the problems play a number of roles: they check to make sure you understand the material, they force you to be an active (and thus more effective) participant in your own learning, they expose you to new topics and applications. For this reason, the problems spread over a wide range of difficulty, and I have identified them as such.

- Easy problems (marked with an E) might better be called "questions". You should be able to answer them after just a few moment's thought, usually without the aid of pencil and paper. Their main purpose is to check your understanding or to reinforce conceptual points already made in the text. You should do all the easy problems as you come to them, but they are not usually appropriate for assigned problem sets. The level of these problems is similar to that of problems found on the Graduate Record Examination in physics. . . some of them are even set up in multiple-choice format.

- Intermediate problems (marked with an I) are straightforward applications of the text material. They might better be called "exercises" because they exercise the skills you learned in the text without calling them into stress. They are useful both for checking your understanding and for forcing your participation. Sometimes the detailed derivation of results described in the text is left for such an exercise, with appropriate direction and clues.

- Difficult problems (marked with one to three D's, to indicate the level of difficulty) are more ambitious and usually require the student to pull together knowledge and skills from several different pieces of this book, and from other physics courses as well. Sometimes these problems introduce and develop topics that are not mentioned in the text at all.

- In addition, some easy or intermediate problems are marked with a star (e.g. I*) to indicate that they are "essential problems". These problems are so important to the development of the course material that you *must* do them to have any hope of being able to understand the course material. Essential problems are either relied upon in later text material (e.g. problem 1.2) or else cover important topics that are not covered in the text itself (e.g. problem 3.6). In the latter case, I covered that important

topic through a problem rather than trough text material because I thought it was easier to understand in that form.

I have attempted to make all of the problems at all of the levels interesting in their own right, rather than burdens to complete out of necessity. For that reason, I recommend that you read all of the problems, even the ones that you don't do.

The bibliographies at the end of each chapter are titled "Resources" rather than "Additional Reading" because they include references to computer programs and world wide web pages as well as to books and articles. The former will no doubt go out of date quickly, but to ignore them would be criminal.

Ideas. Perhaps book should come with disk of some public domain programs. . . or maybe I should set up a web page for the book.

Appendices J and K should appear as rear and front endpapers, respectively.

Chapter 1

The Properties of Matter in Bulk

1.1 What is Statistical Mechanics About?

Statistical mechanics treats matter in bulk. While most branches of physics... classical mechanics, atomic physics, quantum mechanics, nuclear physics... deal with one or two or a few dozen particles, statistical mechanics deals with, typically, about a mole of particles at one time. A mole is 6.02×10^{23}, considerably larger than a few dozen. Let's compare this to a number often considered large, namely the U.S. national debt. This debt is (2018) about 21 trillion dollars, so the national debt is less than 40 trillionth of a mole of dollars.[1] Even so, a mole of water molecules occupies only 18 ml or about half a fluid ounce... it's just a sip.

The huge number of particles present in the systems studied by statistical mechanics means that the traditional questions of physics are impossible to answer. For example, the traditional question of classical mechanics is the time-development problem: Given the positions and velocities of all the particles now, find out what they will be at some future time. This problem has has not been completely solved for three gravitating bodies... clearly we will get nowhere asking the same question for 6.02×10^{23} bodies! But in fact, a solution of the time-development problem for a mole of water molecules would be useless even if it could be obtained. Who cares where each molecule is located? No experiment will ever be able to find out. To make progress, we have to ask different questions, question like "How does the pressure change with volume?", "How does the temperature change upon adding particles?", "What is the mean distance between atoms?", or "What is the probability for finding two atoms separated by a given distance?". Thus the challenge of statistical mechanics is two-fold: first find the questions, and only then find the answers.

[1] In contrast, the Milky Way galaxy contains about 0.3 or 0.6 trillionth of a mole of stars. The entire universe probably contains fewer than a mole of stars.

1.2 Outline of Book

This book begins with a chapter, *the properties of matter in bulk*, that introduces statistical mechanics and shows why it is so fascinating.

It proceeds to discuss the *principles of statistical mechanics*. The goal of this chapter is to motivate and then produce a conceptual definition for that quantity of central importance: entropy. In contrast to, say, quantum mechanics, it is not useful to cast the foundations of statistical mechanics into a mathematically rigorous "postulate, theorem, proof" mold. Our arguments in this chapter are often heuristic and suggestive; "plausibility arguments" rather than proofs.

Once we have defined entropy and know a few of its properties, what can we do with it? The subject of thermodynamics asks what can be discovered about substance by just knowing that entropy exists, without knowing a formula for it. It is one of the most fascinating fields in all of science, because it produces a large number of dramatic and unexpected results based on this single modest assumption. This book's chapter on *thermodynamics* begins begins by developing a concrete operational definition for entropy, in terms of heat and work, to complement the conceptual definition produced in the previous chapter. It goes on to apply entropy to situations as diverse as fluids, phase transitions, and light.

The chapter on *ensembles* returns to issues of principle, and it produces formulas for the entropy that are considerably easier to apply than the one produced in chapter 2. Armed with these easier formulas, the rest of the book uses them in various applications.

The first three applications are to the classic topics of *classical ideal gases*, *quantal ideal gases*, including Fermi-Dirac and Bose-Einstein statistics, and *harmonic lattice vibrations* or phonons.

The subject of ideal gases (i.e. gases of non-interacting particles) is interesting and often useful, but it clearly does not tell the full story... for example, the classical ideal gas can never condense into a liquid, so it cannot show any of the fascinating and practical phenomena of phase transitions. The next chapter treats *weakly interacting fluids*, using the tools of perturbation theory and the variational method. The correlation function is introduced as a valuable tool. This is the first time in the book that we ask questions more detailed than the questions of thermodynamics.

Finally we treat *strongly interacting systems and phase transitions*. Here our emphasis is on magnetic systems. Tools include mean field theory, transfer matrices, correlation functions, and computer simulations. Under this heading fall some of the most interesting questions in all of science... some answered, many still open.

The first five chapters (up to and including the chapter on *classical ideal gases*) are essential background to the rest of the book, and they must be treated in the sequence presented. The last four chapters are independent and can be treated in any order.

1.3 Fluid Statics

I mentioned above that statistical mechanics asks questions like "How does the pressure change with volume?". But what is pressure? Most people will answer by saying that pressure is force per area:

$$\text{pressure} = \frac{\text{force}}{\text{area}}. \tag{1.1}$$

But force is a vector and pressure is a scalar, so how can this formula be correct? The aim of this section is to investigate what this formula means and find out when it is correct.[2]

1.3.1 Problems

1.1 (I) **The rotating water glass**
A cylinder containing a fluid of mass density ρ is placed on the center of a phonograph turntable and rotated with constant angular velocity ω. After some initial sloshing of the fluid, everything calms down to a steady state.

a. The pressure is a function of height h and distance from the axis r. Show that the variation of pressure with radial distance is

$$\frac{\partial p(r,h)}{\partial r} = \rho\omega^2 r, \tag{1.2}$$

while the variation with vertical distance is

$$\frac{\partial p(r,h)}{\partial h} = -\rho g. \tag{1.3}$$

(Where g is the acceleration of gravity.)

[2] As such, the aim of this section is quite modest. If you want to learn more about the interesting subject of fluid flow, see the "Resources" section of this chapter.

b. The pressure at the surface of the fluid at the center of the cylinder ($r = 0$, $h = h_c$) is of course atmospheric pressure p_a. Integrate the differential equations of part (a.) to show that, at any point in the fluid,

$$p(r, h) = p_a + \tfrac{1}{2}\rho\omega^2 r^2 - \rho g(h - h_c). \tag{1.4}$$

c. Show that the profile of the fluid surface is given by

$$y(r) = \frac{\omega^2}{2g} r^2. \tag{1.5}$$

1.4 Phase Diagrams

Too often, books such as this one degenerate into a study of gases... or even into a study of the ideal gas! Statistical mechanics in fact applies to all sorts of materials: fluids, crystals, magnets, metals, polymers, starstuff, even light. I want to show you some of the enormous variety of behaviors exhibited by matter in bulk, and that can (at least in principle) be explained through statistical mechanics.

Because the axes of a phase diagram are pressure and temperature, the misconception arises that phase diagrams plot pressure as a function of temperature. No. Pressure and temperature are independent variables. For example, volume is a function of pressure and temperature, $V(T, p)$. Instead, the lines on a phase diagram mark the places where there are cliffs in the function $V(T, p)$.

End with the high Tc phase diagram of Amnon Aharony discussed by MEF at Gibbs Symposium. Birgeneau.

Resources

The problems of fluid flow are neglected in the typical American undergraduate physics curriculum. An introduction to these fascinating problems can be found in the chapters on elasticity and fluids in any introductory physics book, such as

F.W. Sears, M.W. Zemansky, and H.D. Young, *University Physics*, fifth edition (Addison-Wesley, Reading, Massachusetts, 1976), chapters 10, 12, and 13, or

D. Halliday, R. Resnick, and J. Walker, *Fundamentals of Physics*, fourth edition (John Wiley, New York, 1993), sections 16–1 to 16–7.

More idiosyncratic treatments are given by

R.P. Feynman, R.B. Leighton, and M. Sands, *The Feynman Lectures on Physics* (Addison-Wesley, Reading, Massachusetts, 1964), chapters II-40 and II-41, and

Jearl Walker *The Flying Circus of Physics* (John Wiley, New York, 1975), chapter 4.

Hansen and McDonald

An excellent description of various states of matter (including liquid crystals, antiferromagnets, superfluids, spatially modulated phases, and more) extending our section on "Phase Diagrams" is

> Michael E. Fisher, "The States of Matter—A Theoretical Perspective" in W.O. Milligan, ed., *Modern Structural Methods* (The Robert A. Welch Foundation, Houston, Texas, 1980) pp. 74–175.

1.5 Additional Problems

1.2 (I*) **Compressibility, expansion coefficient**

The "isothermal compressibility" of a substance is defined as

$$\kappa_T(p,T) = -\frac{1}{V(p,T)}\frac{\partial V(p,T)}{\partial p}, \tag{1.6}$$

where the volume $V(p,T)$ is treated as a function of pressure and temperature.

a. Justify the name "compressibility". If a substance has a large κ_T is it hard or soft? Since "squeeze" is a synonym for "compress", is "squeezeabilty" a synonym for "compressibility"? Why were the negative sign and the factor of $1/V$ included in the definition?

b. The "expansion coefficient" is

$$\beta(p,T) = \frac{1}{V(p,T)}\frac{\partial V(p,T)}{\partial T}. \tag{1.7}$$

In most situations β is positive, but it can be negative. Give one such circumstance.

c. What are κ_T and β in a region of two-phase coexistence (for example, a liquid in equilibrium with its vapor)?

d. Find κ_T and β for an ideal gas, $V(p,T) = Nk_BT/p$.

e. Show that

$$\frac{\partial \kappa_T(p,T)}{\partial T} = -\frac{\partial \beta(p,T)}{\partial p} \tag{1.8}$$

for all substances.

f. Verify this relation for the ideal gas. (Clue: The two expressions are *not* both equal to $-Nk_B/p^2V$.)

1.3 (I*) **Heat capacity as a susceptibility**

Later in this book we will find that the "heat capacity" of a fluid[3] is

$$C_V(T,V,N) = \frac{\partial E(T,V,N)}{\partial T}, \tag{1.9}$$

[3]Technically, the "heat capacity at constant volume".

where the energy $E(T, V, N)$ is considered as a function of temperature, volume, and number of particles. The heat capacity is easy to measure experimentally and is often the first quantity observed when new regimes of temperature or pressure are explored. (For example, the first sign of superfluid He^3 was an anomalous dip in the measured heat capacity of that substance.)

a. Explain how to measure the heat capacity of a gas given a strong, insulated bottle, a thermometer, a resistor, a voltmeter, an ammeter, and a clock.

b. Near the superfluid transition temperature T_c, the heat capacity of Helium is given by

$$C_V(T) = -A \ln(|T - T_c|/T_c). \tag{1.10}$$

Sketch the heat capacity and the energy as a function of temperature in this region.

c. The heat capacity is one member of a class of thermodynamic quantities called "susceptibilities". Why does it have that name? (Clues: A change in temperature causes a change in energy, but how much of a change? If the heat capacity is relatively high, is the system relatively sensitive or insensitive (i.e. susceptible or insusceptible) to such temperature changes?)

d. Interpret the isothermal compressibility (1.6) as a susceptibility. (Clue: A change in pressure causes a change in volume.)

1.4 (I) **The meaning of "never"**

(This problem is modified from Kittel and Kroemer, *Thermal Physics*, second edition, page 53.)

It has been said[4] that "six monkeys, set to strum unintelligently on typewriters for millions of years, would be bound in time to write all the books in the British Museum". This assertion gives a misleading impression concerning very large numbers.[5] Consider the following situation:

- The quote considers six monkeys. Let's be generous and allow 10^{10} monkeys to work. (About twice the present human population.)
- The quote vaguely mentions "millions of years". The age of the universe is about 14 billion years, so let's allow all of that time, about 10^{18} seconds.
- The quote wants to write out all the books in a very large library. Let's be modest and demand only the production of Shakespeare's *Hamlet*, a work of about 10^5 characters.
- Finally, assume that a monkey can type ten characters per second, and for definiteness, assume a keyboard of 29 characters (letters, comma, period, space... ignore caPitALIzaTion).

a. Show that the probability that a given sequence of 10^5 characters comes out through a random striking of 10^5 keys is

$$\frac{1}{29^{100\,000}} \approx 10^{-146\,240}. \tag{1.11}$$

How did you perform the arithmetic?

[4] J. Jeans, *Mysterious Universe* (Cambridge University Press, Cambridge, 1930) p. 4.

[5] An insightful discussion of the "monkeys at typewriters" problem, and its implications for biological evolution, is given by Richard Dawkins in his book *The Blind Watchmaker* (Norton, New York, 1987) pp. 43–49.

b. Show that the probability of producing *Hamlet* through the "unintelligent strumming" of 10^{10} monkeys over 10^{18} seconds is about $10^{-146\,211}$, which is small enough to be considered zero for most purposes.

1.5 **Human genetics**

There are about 21,000 genes in the human genome. Suppose that each gene could have any of three possible states (called "alleles"). (For example, if there were a single gene for hair color, the three alleles might be black, brown, and blond.) Then how many genetically distinct possible people would there be? Compare to the current worldwide human population. Estimate how long it would take for every possible genetically distinct individual to be realized. Compare to the age of the universe.

Chapter 2

Principles of Statistical Mechanics

In a book on classical or quantum mechanics, the chapter corresponding to this one would be titled "Foundations of Classical (or Quantum) Mechanics". Here, I am careful to use the term "principles" rather than "foundations". The term "foundations" suggests rock solid, logically rigorous, hard and fast rules, such as the experimental evidence that undergirds quantum theory. Statistical mechanics lacks such a rigorous undergirding. Remember that our first job in statistical mechanics is to find the questions. Of course, we can ask any question we wish, but we need to find profitable questions. Such a task is not and cannot be one of rigor.

Our treatment here is based on classical mechanics, not quantum mechanics. This approach is easier and more straightforward than the approach through quantum mechanics, and—as you will soon see—there are quite enough difficulties and subtleties in the classical approach! After we investigate the principles of statistical mechanics from a classical perspective, we will outline (in section 4.10) the generalization to quantum mechanics.

2.1 Microscopic Description of a Classical System

This section deals with classical mechanics, not statistical mechanics. But any work that attempts to build macroscopic knowledge from microscopic knowledge—as statistical mechanics does—must begin with a clear and precise statement of what that microscopic knowledge is.

The microscopic description of a physical system has two components: First, "What are the parts of the system? How do they affect each other?", second, "How are those parts arranged?". The first question is answered by giving the *mechanical parameters*[1] of the system. The second is answered by giving its *dynamical variables*. Rather than give formal definitions of these terms, we give two examples.

[1] The mechanical parameters are sometimes called "parameters in the Hamiltonian", or "external parameters", or "fields".

The earth-moon system. In this system, the mechanical parameters are the mass of the earth, the mass of the moon, and (because the earth and moon interact gravitationally), the gravitational constant G. The dynamical variables are the position and velocity (or momentum) of each body. (Alternative dynamical variables are the position of the center of mass and the separation between the bodies, plus the total momentum of the system and the angular momentum of the two bodies about the center of mass.) You can see from this example that the mechanical parameters give you the knowledge to write down the Hamiltonian for the system, while the dynamical variables are the quantities that change according to the laws governed through that Hamiltonian. The mechanical parameters do not depend upon the initial condition of the system—the dynamical variables do. Often (although not always) the mechanical parameters are time-constant while the dynamical variables are time-varying.

Helium atoms in a box. It is natural to begin the description by saying that there are N atoms, each of mass m. But this innocent beginning is neither obvious nor precisely correct. By saying that the only thing we need to know about each atom is its mass, we are modeling the atoms as point particles. A more precise model would describe the system as N nuclei and $2N$ electrons, but then our treatment would necessarily involve quantum mechanics rather than classical mechanics. Furthermore we would not gain anything by this more precise and more difficult description... we know from experience that under ordinary conditions the nuclei and electrons *do* bind themselves together as atoms. Even the more precise description would not result in unassailable rigor, because the nuclei themselves are made up of nucleons and the nucleons of quarks. In fact, a model-building process similar to this one went on unmentioned even in our treatment of the earth-moon system: When we said that we needed to know only the masses of the two bodies, we were assuming (a good but not perfect assumption) that the distribution of matter through the earth and moon was irrelevant to their motion. We will adopt the model that replaces helium atoms by point particles, but you should keep in mind that it *is* a model.

To continue in our microscopic description, we need to know how the atoms interact with each other. A common model is that atoms interact in a pairwise fashion through some sort of atom-atom interaction potential such as the "Lennard-Jones 6–12 potential":

$$\text{atom-atom potential energy } = -\frac{a}{r^6} + \frac{b}{r^{12}}. \tag{2.1}$$

Here the quantities a and b are mechanical parameters.

An important part of the microscopic description of helium atoms in a box is a description of the box. We must know where the walls are located and how the walls interact with the atoms. There are many different models for the wall-atom interaction: common models include the "hard wall" and "soft wall" potential energy functions sketched in figure 2.1. But even by assuming that the walls and atoms interact through a potential at all we are making a dramatic and easily overlooked assumption: the assumption of a smooth, non-atomic wall. Real box walls are themselves made up of atoms, which can of course move, so there is no fixed "zero-distance" point as implied by the graphs in the figure. The assumption of fixed, smooth walls implies that when an atom collides with the wall its energy is unchanged, while it is quite possible for the atom to gain or loose some energy while colliding with a real, atomic wall. The assumption of a smooth wall is particularly suspicious, and we will find in chapter 4 that it can be relaxed very profitably.

Figure 2.1:

The final step is to recognize that there might be externally electric, magnetic, or gravitational fields that affect the system. If these external fields are present and relevant they will have to be added to the list of mechanical parameters.

At last we are ready to turn to the dynamical variables for the helium-atoms-in-a-box system. These are (mercifully) easy to describe: they are just the position and momentum of each particle, a total of $2N$ 3-dimensional vectors. A shorthand name for this information is "point in phase space". Phase space is an abstract space with one coordinate corresponding to each dynamical variable. Thus in our example the coordinates are the positions and momenta for particle number 1, number 2,... number 9,... and number N:

$$(x_1, y_1, z_1, p_{x,1}, p_{y,1}, p_{z,1}, x_2, y_2, z_2, p_{x,2}, p_{y,2}, p_{z,2}, \ldots x_9, y_9, z_9, p_{x,9}, p_{y,9}, p_{z,9}, \ldots x_N, y_N, z_N, p_{x,N}, p_{y,N}, p_{z,N}). \tag{2.2}$$

For a system of N particles, phase space is $6N$-dimensional. A single point in phase space gives the position *and* momentum of *every* particle in the system.

2.1.1 Time development of a classical system

Given a microscopic description such as either of the two above, what can we do with it? The time development of the system is represented by the motion of a point in phase space. That point will snake through many points in the phase space but all of the points visited will have the same energy.[2] Because *many* points of the given energy will be visited, it is natural to ask whether, in fact, *all* phase space points corresponding to a given energy will eventually be visited by a system started at any one of those points.

It is easy to find systems for which this statement is false, but all such examples seem to be in one way or another atypical. For example, consider two or three or even many millions of non-interacting particles in a

[2]For the earth-moon model, all the points visited will also have the same total momentum and angular momentum, but this is *not* the case for the helium-in-a-smooth-box model.

hard-walled box, and start them all traveling straight up and down. They will travel straight up and down forever. The points in phase space with identical energy, but with the particles traveling left and right, will never be visited. This example is atypical because if the particles interacted, even slightly, then they would fall out of the "straight up and down" regions of phase space.

Problems

2.1 (Q) **Three-body interactions: microscopic**
Make up a problem involving two charged point particles and a polarizable atom.

2.2 **Mechanical parameters and dynamical variables**
Here is a classical mechanics problem: "A pendulum bob of mass m swings at the end of a cord which runs through a small hole in the ceiling. The cord is being pulled up through the hole so that its length is $\ell(t) = \ell_0 - \alpha t$. At time $t = 0$ the bob is at rest and at angle θ_0. Find the subsequent motion of the bob." *Do not* solve this problem. Instead, list the mechanical parameters and the dynamical variables that appear in it.

2.3 **A Hamiltonian**
Write down the total Hamiltonian of a neutral plasma of N protons and N electrons moving in a rectangular box with interior dimensions of $L_x \times L_y \times L_z$, assuming that i) any proton or electron interacts with the wall material through a potential energy function

$$W(d) = \begin{cases} W_0 \left(\dfrac{1}{d^2} - \dfrac{1}{a^2} \right) & \text{for } d < a \\ 0 & \text{otherwise,} \end{cases} \tag{2.3}$$

where d is the (perpendicular) distance of the particle in question from the wall, and ii) the system is subject to a uniform electric field in the $\hat{\mathbf{x}}$ direction of magnitude E. List the mechanical parameters that appear in this Hamiltonian, and distinguish them from the dynamical variables.

2.4 (Q) **For discussion: Mechanical parameters, dynamical variables, and modeling**
List the mechanical parameters and dynamical variables of these systems:

a. Hydrogen molecules enclosed in a sphere.

b. Water molecules in a box.

c. A mixture of hydrogen molecules and helium atoms in a box.

To what extent are you making models as you generate descriptions? To what extent are you making assumptions? (For example, by using non-relativistic classical mechanics.)

2.2 Macroscopic Description of a Large Equilibrium System

The title of this section introduces two new terms... large and equilibrium. A system is large if it contains many particles, but just how large is "large enough"? A few decades ago, physicists dealt with systems of one or two or a dozen particles, or else they dealt with systems with about 10^{23} particles, and the distinction was clear enough. Now that computer chips have such small lines, systems of intermediate size are coming under investigation. (This field carries the romantic name of "mesoscopic physics".) In practical terms, it is usually easy enough to see when the term "large" applies, but there is no rigorous criterion and there have been some surprises in this regard.

A similar situation holds for the term equilibrium. A system is said to be in equilibrium if its macroscopic properties are not changing with time. Thus a cup of tea, recently stirred and with churning, swirling flow, is not at equilibrium, whereas some time later the same cup, sedate and calm, is at equilibrium. But some of the properties of the latter cup *are* changing with time: for example, the height of water molecule number 173 changes rapidly. Of course this is not a macroscopic property, but then there is no rigorous definition of macroscopic *vs.* microscopic properties. Once again there is little difficulty in practice, but a rigorous criterion is wanting and some excellent physicists have been fooled. (For example, a mixture of hydrogen gas and oxygen gas can behave like a gas in equilibrium. But if a spark is introduced, these chemicals will react to form water. The gas mixture is in equilibrium as far as its physical properties are concerned, but not as far as its chemical properties are concerned.)

With these warnings past, we move on to the macroscopic description. As with the microscopic description, it has two parts: the first saying what the system is and the second saying what condition the system is in. For definiteness, we consider the helium-in-a-smooth-box model already introduced.

To say what this system is, we again list mechanical parameters. At first these are the same as for the microscopic description: the number of particles N, the mass of each particle m, and a description of the atom-atom interaction potential (e.g. the Lennard-Jones parameters a and b). But we usually don't need to specify the atom-wall interaction or even the location of the walls... instead we specify only the volume of the container V. The reason for this is not hard to see. If we deal with a large system with typical wall-atom interactions (short range) and typical walls (without intricate projections and recesses) then very few particles will be interacting with the walls at any one instant, so we expect that the details of wall-atom interaction and container shape will be affect only a tiny minority of atoms and hence be irrelevant to macroscopic properties. There are of course exceptions: for example, in a layered substance such as graphite or mica, or one of the high-temperature superconductors, the shape of the container might well be relevant. But in this book we will not often deal with such materials.

Finally, what condition is the system in? (In other words, what corresponds to the dynamical variables in a microscopic description?) Clearly the "point in phase space", giving the positions and momenta of each and every particle in the system, is far too precise to be an acceptable *macroscopic* description. But at the same time it is not acceptable to say we don't care anything about microscopic quantities, because the energy of this system is conserved, so energy will be a feature of both the microscopic and macroscopic

descriptions. In fact, for the helium-in-a-smooth-box model, energy is the *only* quantity that is conserved,[3] so it is the only item in the list of macroscopic descriptors. (Other systems, such as the earth-moon system, will have additional conserved quantities, and hence additional items in that list.)

What equilibrium is not

There's a common misconception that at equilibrium, the sample is uniform. This might not be true: A mixture of ice and water at atmospheric pressure and temperature 0 C is at equilibrium but is not uniform. Or, consider a container of air 50 kilometers tall with its base on Earth at sea level: most of the molecules huddle at the bottom of the container, and the top is near-vacuum.

2.5 (Q) **Three-body interactions: macroscopic**
Even though three-body interactions do exist, they can usually be ignored in a macroscopic description. (Just as the wall-atom interaction does exist, but it can usually be ignored in a macroscopic description.) Why?

2.6 (Q,E) **Lost in space**
A collection of N asteroids floats in space far from other gravitating bodies. Model each asteroid as a hard sphere of radius R and mass m. What quantities are required for a microscopic description of this system? For a macroscopic description?

2.3 Fundamental Assumption

Define microstate and macrostate.

There are many microstates corresponding to any given macrostate. The collection of all such microstates is called an "ensemble". (Just as a musical ensemble is a collection of performers.)

Note: An ensemble is a (conceptual) collection of macroscopic systems. It is *not* the collection of atoms that makes up a macroscopic system.

(Terminology: A microstate is also called a "configuration" or a "complexion"... both poor terms. A macrostate is also called a "thermodynamic state". A "corresponding" microstate is sometimes called an "accessible" or a "consistent" microstate.)

A system is said to be "isolated" if no energy goes in or out, and if the mechanical parameters (N, V, etc.) are also fixed. Most of the systems you deal with in classical mechanics classes, for example, are isolated.

The fundamental assumption of statistical mechanics is:

[3] The momentum and angular momentum of the helium atoms is not conserved, because there are external forces due to the box.

An isolated system in an equilibrium macrostate has equal probability of being in any of the microstates corresponding to that macrostate.

Conceptual difficulties:

1. What is equilibrium?
2. What is probability? (Any experiment is done on *one* system.)
3. May be false! (Can be motivated by the ergodic hypothesis, but this is just suggestive, and the ergodic hypothesis itself has not been proven and may be false. Cellular automata may violate this assumption... but do physical/biological systems? See for example Andrew I. Adamatzky, *Identification of Cellular Automata* (Taylor and Francis, 1995) and the essay review by Normand Mousseau, in *Contemporary Physics*, **37** (1996) 321–323; Stephen Wolfram, *A New Kind of Science* (Wolfram Media, 2002).)

Practical difficulties.

1. Gives way to find average values of microscopic quantities, such as

$$\langle \text{kinetic energy} \rangle, \langle \text{potential energy} \rangle, \langle \text{height} \rangle,$$

   but not things like temperature and pressure.
2. Based on model. (In particular, infinitely hard smooth walls.)
3. For point particle model, requires infinitely thin sheet in phase space. Work instead with the volume of phase space corresponding to energies from E to $E + \Delta E$, and at the end of the calculation take the limit $\Delta E \to 0$.

To examine these practical difficulties in more detail, consider again the helium-in-a-smooth-box model. Suppose we want to find the number of microstates with energy ranging from E to $E + \Delta E$, in a system with N identical particles in a box of volume V.

Your first answer might be that there are an infinite number of points in phase space satisfying this criterion. This is true but it misses the point: A thimble and a Mack truck both contain an infinite number of points, but the Mack truck carries more because it has more volume. Clearly the microstate count we desire is some sort of measure of phase-space volume. I'll call the region of phase space with energy ranging from E to $E + \Delta E$ by the name $\sigma(E, \Delta E)$, and call its volume

$$\int_{\sigma(E,\Delta E)} d\Gamma. \qquad \text{(Volume in phase space.)} \tag{2.4}$$

However, this volume isn't precisely what we want. Permutation argument for N identical particles. (For a system with N identical particles, there are $N!$ points in phase space corresponding to the same microstate.) A better measure of the microstate count is thus

$$\frac{1}{N!}\int_{\sigma(E,\Delta E)} d\Gamma. \qquad \text{(Delabeled volume in phase space.)} \tag{2.5}$$

There remains one more problem. We desire a count, which is dimensionless, and the above expression gives a phase-space volume, which has the dimensions of

$$(\text{angular momentum})^{3N}.$$

The solution to this problem is straightforward if clunky. Pick a quantity with the dimensions of angular momentum. Any quantity will do, so make an arbitrary choice. Call the quantity h_0. (Clearly, by the time we produce a measurable physical result at the end of a calculation, it had better be independent of our choice of h_0, just as the value of any measurable result in a classical mechanics problem has to be independent of choice of the potential energy zero.) Then define the microstate count as the dimensionless quantity

$$\Omega(E,\Delta E,V,N) = \frac{1}{h_0^{3N}}\frac{1}{N!}\int_{\sigma(E,\Delta E)} d\Gamma. \qquad \text{(Dimensionless, delabeled volume in phase space.)} \tag{2.6}$$

In poker, there are 52 cards in a deck, so the probability of drawing any given card from a shuffled deck is 1/52. The number 52 plays a fundamental role in answering any question concerning probability in poker.

In statistical mechanics, there are Ω microstates corresponding to a macrostate, so the probability of encountering any given microstate is (if the fundamental assumption is correct) $1/\Omega$. The number Ω plays the same fundamental role in statistical mechanics that 52 does in poker.

2.7 Microstate count for a mixture

What expression corresponds to (2.6) for a collection of N_{He} Helium atoms and N_{Ar} Argon atoms?

2.4 Statistical Definition of Entropy

Logarithms and dimensions
You cannot take the logarithm of a number with dimensions. Perhaps you have heard this rule phrased as "you can't take the logarithm of 3.5 meters" or "you can't take the logarithm of five oranges". Why not? A simple argument is "Well, what would be the units of ln(3.5 meters)?" A more elaborate argument follows. The logarithm function is the inverse of the exponential function:

$$y = \ln(x) \quad \text{means the same as} \quad x = e^y.$$

But remember that

$$x = e^y = 1 + y + \frac{1}{2!}y^2 + \frac{1}{3!}y^3 + \cdots.$$

If y had the dimensions of length, then the expression above would be a meaningless sum of 1 plus a length plus an area plus a volume plus so forth.
You cannot exponentiate a number with dimensions, and you cannot take the logarithm of a number with dimensions.

Additivity (see problem 2.10) ... don't physically mix.

$$S(E, \Delta E, V, N) = k_B \ln \Omega(E, \Delta E, V, N) \tag{2.7}$$

The constant k_B in this definition is called the "Boltzmann constant". It is clear from the argument that the Boltzmann constant could have been chosen to have any value: 1, π, whatever. For historical reasons, it was chosen to have the value

$$k_B = 1.38 \times 10^{-23} \text{ joule/kelvin}. \tag{2.8}$$

There is no particular physical significance to this number: its logical role is analogous to 2.54 cm/inch or 4.186 joule/calorie. In other words, the origin of this number is not to be sought in nature, but in the history of the definition of the joule and the kelvin. It is a conversion factor.

Problems

2.8 (E) **Accessible regions of phase space**
Suppose that N non-interacting particles, each of mass m, move freely in a one-dimensional box (i.e. an infinite square well). Denote the position coordinates by $x_1, x_2, \ldots, x_N$ and the momentum coordinates by $p_1, p_2, \ldots, p_N$. The box restricts all the positions to fall between $x_i = 0$ and $x_i = L$. The energy of the system lies between E and $E + \Delta E$.

a. If only one particle is present, draw the system's phase space and shade the regions of phase space that are accessible.

b. If two particles are present then phase space is four dimensional, which makes it difficult to draw. Draw separately the part of phase space involving positions and the part involving momenta. Shade the accessible regions of phase space.

c. Suppose two particles are present, and consider the slice of phase space for which $x_1 = (2/3)L$ and p_2 equals some constant called $\tilde{p}_2$. Draw a (carefully labeled) sketch of this slice with the accessible regions shaded.

d. Describe the accessible regions of phase space if N particles are present.

2.9 **Accessible configurations of a spin system**

Consider an isolated system of N spin-$\frac{1}{2}$ atoms in a magnetic field H. The atoms are fixed at their lattice sites and the spins do not interact. Each atom has a magnetic moment m that can point either "up" (parallel to the field H) or "down" (antiparallel to H). A microstate (or configuration) of this system is specified by giving the direction of every spin. An up spin has energy $-mH$, a down spin has energy $+mH$, so a configuration with $n_\uparrow$ up spins and $n_\downarrow$ down spins has energy

$$E = -(n_\uparrow - n_\downarrow)mH. \tag{2.9}$$

This system is called the "ideal paramagnet".

a. Not every energy is possible for this model. What is the maximum possible energy? The minimum? What is the minimum possible non-zero energy difference between configurations?

b. Suppose we know that the system has $n_\uparrow$ up spins and $n_\downarrow$ down spins, but we do not know how these spins are arranged. How many microstates are consistent with this knowledge?

c. The variables $n_\uparrow$ and $n_\downarrow$ cannot be determined directly from macroscopic measurements. Find expressions for $n_\uparrow$ and $n_\downarrow$ in terms of N, E, and H. (Hand a paramagnet sample to an experimentalist and ask her to find the number of up spins. She will just look at you quizzically. But ask her to find the number, the energy, and the magnetic field and she'll be happy to.)

d. Consider the energy range from E to $E + \Delta E$ where ΔE is small compared to NmH but large compared to mH. What is the approximate number of states $\Omega(E, \Delta E, H, N)$ lying in this energy range? Express your answer in a form that does not include the quantities $n_\uparrow$ or $n_\downarrow$.

2.10 **Microstates for a combined system**

System #1 is in a macrostate with three corresponding microstates, labeled A, B, and C. System #2 is in a macrostate with four corresponding microstates, labeled α, β, γ, and δ. How many microstates are accessible to the combined system consisting of system #1 and system #2? List all such microstates.

2.11 (E) **The logarithm**

Suppose that a differentiable function satisfies

$$f(xy) = f(x) + f(y) \tag{2.10}$$

for all positive x and y. Show that

$$f(x) = k \ln(x). \tag{2.11}$$

[Clues: 1) Take derivative with respect to x, then set $x = 1$. 2) Set $y = 1$ in equation (2.10).]

2.5 Entropy of a Monatomic Ideal Gas

So far in this chapter, we have been dealing very abstractly with a very general class of physical systems. We have made a number of assumptions that are reasonable but that we have not tested in practice. It is time to put some flesh on these formal bones. We do so by using our statistical definition of entropy to calculate the entropy of a monatomic ideal gas. (Here "monatomic" means that we approximate the atoms by point particles, and "ideal" means that those particles do not interact with each other. In addition, we assume that the gas contains only one chemical species and that classical mechanics provides an adequate description. Thus a more precise name for our system would be the "pure classical monatomic idea gas", but in this case we wisely prefer brevity to precision.) Working with this concrete example will show us that what we have said is sensible (at least for this system), and guide us in further general developments.

The previous pages have been remarkably free of equations for a physics book. Now is the time to remedy that situation. Before studying this section, you need to know that the volume of a d-dimensional sphere is

$$V_d(r) = \frac{\pi^{d/2}}{(d/2)!} r^d. \tag{2.12}$$

If you don't already know this, then read appendix D, "Volume of a Sphere in d Dimensions", before reading this section. And if you don't know the meaning of $x!$, where x is a half-integer, then you should read appendix C, "Clinic on the Gamma Function", before reading appendix D. Finally, if you don't know Stirling's approximation for the factorial function, namely

$$\ln n! \approx n \ln n - n \qquad \text{for } n \gg 1, \tag{2.13}$$

then you should also read appendix E, "Stirling's Approximation", before reading further. (Do not be discouraged by this long list of prerequisites. This mathematical material is quite interesting in its own right and will be valuable throughout this book.)

We consider a system of N identical, classical, non-interacting point particles, each of mass m. The kinetic energy of this system is

$$\frac{1}{2m}(p_1^2 + p_2^2 + \cdots + p_N^2), \tag{2.14}$$

and the potential energy is

$$\begin{cases} 0 & \text{if all particles are inside container} \\ \infty & \text{otherwise.} \end{cases} \tag{2.15}$$

(One sometimes hears that the ideal gas has "no potential energy". It is true that there is no potential energy due to atom-atom interaction, but, as the above expression makes clear, there is indeed a potential

energy term due to atom-wall interaction. Because of the character we assume for that term, however, the numerical value of the potential energy is always zero. Note also that the ideal gas is not the same as the "hard-sphere gas". In the hard-sphere model two atoms have infinite potential energy if they are separated by a distance of twice the hard-sphere radius or less. In the ideal gas model two atoms do not interact. It is permissible even for two atoms to occupy the same location... only in the model, of course!)

Now that the system is completely specified, it is time to begin the problem. We wish to calculate the entropy

$$S(E, \Delta E, V, N) = k_B \ln \frac{W(E, \Delta E, V, N)}{N! h_0^{3N}}, \tag{2.16}$$

where the function W represents the volume in phase space corresponding to energies from E to $E + \Delta E$ (i.e. the volume of the region $\sigma(E, \Delta E)$). Before jumping into this (or any other) problem, it is a good idea to list a few properties that we expect the solution will have... this list might guide us in performing the calculation; it will certainly allow us to check the answer against the list to see if either our mathematics or our expectations need revision. We expect that:

- We will be able to take the limit as $\Delta E \to 0$ and get sensible results.
- The entropy S will depend on only the volume of the container and not on its shape.
- If we double the size of the system, by doubling E, V, and N, then we will double S. (Additivity.)
- S will depend on h_0 in a trivial, "sea-level" fashion.

The formal expression for the volume of the accessible region of phase space is

$$\begin{aligned} W(E, \Delta E, V, N) &= \text{accessible volume in phase space} && (2.17) \\ &= \int_{\sigma(E,\Delta E)} d\Gamma && (2.18) \\ &= \int dx_1 \int dy_1 \int dz_1 \cdots \int dx_N \int dy_N \int dz_N \cdots \\ &\qquad \int dp_{x,1} \int dp_{y,1} \int dp_{z,1} \cdots \int dp_{x,N} \int dp_{y,N} \int dp_{z,N}. \end{aligned}$$

The complexity of this integral rests entirely in the complexity of the shape of $\sigma(E, \Delta E)$ rather than in the complexity of the integrand, which is just 1. Fortunately the integral factorizes easily into a position part and a momentum part, and the position part factorizes into a product of integrals for each particle (see problem 2.8). For, say, particle number 5, if the particle is inside the container it contributes 0 to the energy, so the total energy might fall between E and $E + \Delta E$ (depending on other factors). But if it is outside the container, then it contributes ∞ to the energy, which *always* exceeds the limit $E + \Delta E$. Thus the integral is just the volume of the container:

$$\int dx_5 \int dy_5 \int dz_5 = V. \tag{2.19}$$

This integral depends on the volume V but is independent of the shape of the container. We will soon see that, as a consequence, the entropy depends on volume but not shape (which is in accord with our expectations).

The integrals over momentum space do not factorize, so we must consider the entire $3N$-dimensional momentum space rather than N separate 3-dimensional spaces. We know that the total potential energy is zero (unless it is infinite), so the energy restriction is taken up entirely by the kinetic energy. Equation (2.14) tells us that the momentum space points with energy E fall on the surface of a sphere of radius $\sqrt{2mE}$. Thus the accessible region in momentum space is a shell with inner radius $\sqrt{2mE}$ and with outer radius $\sqrt{2m(E+\Delta E)}$. (Notice that we are counting *all* the microstates within the accessible region of phase space, not just "typical" microstates there. For example, one microstate to be counted has all of the particles at rest, except for one particle that has all the energy of the system and is heading due west. This is to be counted just as seriously as is the microstate in which the energy is divided up with precise equality among the several particles, and they are traveling in diverse specified directions. Indeed, the system has exactly the same probability of being in either of these two microstates.) Using equation (2.12) for the volume of a $3N$-dimensional sphere, the volume of that shell is

$$\frac{\pi^{3N/2}}{(3N/2)!}\left[(2m(E+\Delta E))^{3N/2}-(2mE)^{3N/2}\right]. \tag{2.20}$$

I prefer to write this result in a form with all the dimensionfull quantities lumped together, namely as

$$\frac{\pi^{3N/2}}{(3N/2)!}(2mE)^{3N/2}\left[\left(1+\frac{\Delta E}{E}\right)^{3N/2}-1\right]. \tag{2.21}$$

The quantity in square brackets is dimensionless.

To find the accessible volume of the entire phase space, we multiply the above result by V^N, the result of performing N separate position integrals. Thus

$$W(E,\Delta E,V,N)=\frac{(2\pi mEV^{2/3})^{3N/2}}{(3N/2)!}\left[\left(1+\frac{\Delta E}{E}\right)^{3N/2}-1\right]. \tag{2.22}$$

As promised, W depends upon the variables E, ΔE, V, and N. It also depends upon the "unmentioned" mechanical parameter m. The arguments on page 15 have been vindicated... the phase space volume W depends only upon V and not upon the detailed shape of the container.

At last we can find the entropy! It is

$$S=k_B\ln\frac{W}{N!h_0^{3N}}=k_B\ln\left\{\left(\frac{2\pi mEV^{2/3}}{h_0^2}\right)^{3N/2}\frac{1}{N!(3N/2)!}\left[\left(1+\frac{\Delta E}{E}\right)^{3N/2}-1\right]\right\} \tag{2.23}$$

or

$$\frac{S}{k_B}=\tfrac{3}{2}N\ln\left(\frac{2\pi mEV^{2/3}}{h_0^2}\right)-\ln N!-\ln(\tfrac{3}{2}N)!+\ln\left[\left(1+\frac{\Delta E}{E}\right)^{3N/2}-1\right]. \tag{2.24}$$

How does this expression compare to our list of expectations on page 22?

- If we take the limit $\Delta E\to 0$, the entropy approaches $-\infty$, contrary to expectations.
- The entropy S depends on the volume of the container but not on its shape, in accord with expectations.

- If we double E, V, and N, then S will not exactly double, contrary to expectations.
- The entropy S does depend on h_0 in a "sea-level" fashion, in accord with expectations.

Only two of our four expectations have been satisfied. (And it was obvious even from equation (2.16) that the fourth expectation would be correct.) How could we have gone so far wrong?

The trouble with expression (2.24) for the entropy of a monatomic ideal gas is that it attempts to hold for systems of any size. In justifying the definition of entropy (2.7) (and in writing the list of expectations on page 22) we relied upon the assumption of a "large" system, but in deriving expression (2.24) we never made use of that assumption. On the strength of this revised analysis we realize that our expectations will hold only approximately for finite systems: they will hold to higher and higher accuracy for larger and larger systems, but they will hold exactly only for infinite systems.

There is, of course, a real problem in examining an infinite system. The number of particles is infinite, as is the volume, the energy, and of course the entropy too. Why do we need an equation for the entropy when we already know that it's infinite? Once the problem is stated, the solution is clear: We need an expression not for the total entropy S, but for the entropy per particle $s = S/N$. More formally, we want to examine the system in the "thermodynamic limit", in which

$$N \to \infty \quad \text{in such a way that} \quad \frac{E}{N} \to e, \frac{V}{N} \to v, \text{ and } \frac{\Delta E}{N} \to \delta e. \tag{2.25}$$

In this limit we expect that the entropy will grow linearly with system size, i.e. that

$$S(E, \Delta E, V, N) \to N s(e, v, \delta e). \tag{2.26}$$

The quantities written in lower case, such as e, the energy per particle, and v, the volume per particle, play the same role in statistical mechanics as "per capita" quantities do in demographics. (The gross national product of the United States is much larger than the gross national product of Kuwait, but that is just because the United States is much larger than Kuwait. The GNP per capita is higher in Kuwait than in the United States.)

Let's take the thermodynamic limit of expression (2.24) (the entropy of a *finite* system) to find the entropy per particle of an *infinite* monatomic ideal gas. The first thing to do, in preparing to take the thermodynamic limit, is to write V as vN, E as eN, and ΔE as $\delta e\, N$ so that the only size-dependent variable is N. This results in

$$\frac{S}{k_B} = \tfrac{3}{2} N \ln\left(\frac{2\pi m e v^{2/3} N^{5/3}}{h_0^2}\right) - \ln N! - \ln(\tfrac{3}{2}N)! + \ln\left[\left(1 + \frac{\delta e}{e}\right)^{3N/2} - 1\right]. \tag{2.27}$$

Next we use Stirling's approximation,

$$\ln n! \approx n \ln n - n \quad \text{for } n \gg 1, \tag{2.28}$$

to simplify the expressions like $\ln(\frac{3}{2}N)!$ above. Thus for large values of N we have approximately (an approximation that becomes exact as $N \to \infty$)

$$\begin{aligned}\frac{S}{k_B} &\approx \tfrac{3}{2}N\ln\left(\frac{2\pi mev^{2/3}N^{5/3}}{h_0^2}\right) - N\ln N + N - (\tfrac{3}{2}N)\ln(\tfrac{3}{2}N) + \tfrac{3}{2}N + \ln\left[\left(1+\frac{\delta e}{e}\right)^{3N/2} - 1\right] \\ &= \tfrac{3}{2}N\ln\left(\frac{2\pi mev^{2/3}}{h_0^2}\right) + \tfrac{3}{2}N\ln N^{5/3} - N\ln N + N - \tfrac{3}{2}N\ln(\tfrac{3}{2}N) + \tfrac{3}{2}N + \ln\left[\left(1+\frac{\delta e}{e}\right)^{3N/2} - 1\right].\end{aligned}$$

The first term on the right increases linearly with N. The next bunch of terms is

$$\begin{aligned}& \tfrac{3}{2}N\ln N^{5/3} - N\ln N + N - \tfrac{3}{2}N\ln(\tfrac{3}{2}N) + \tfrac{3}{2}N \\ =\ & \tfrac{5}{2}N\ln N - N\ln N + N - \tfrac{3}{2}N\ln(\tfrac{3}{2}) - \tfrac{3}{2}N\ln(N) + \tfrac{3}{2}N \\ =\ & N\left[\tfrac{5}{2} - \tfrac{3}{2}\ln(\tfrac{3}{2})\right],\end{aligned}$$

which again increases linearly with N. The final term is, as N grows,

$$\ln\left[\left(1+\frac{\delta e}{e}\right)^{3N/2} - 1\right] \approx \ln\left(1+\frac{\delta e}{e}\right)^{3N/2} \tag{2.29}$$

$$= \tfrac{3}{2}N\ln\left(1+\frac{\delta e}{e}\right). \tag{2.30}$$

This term not only increases linearly with N in the thermodynamic limit, it also vanishes as $\delta e \to 0$! (This is a general principle: One must first take the thermodynamic limit $N \to \infty$, and only then take the "thin phase space limit" $\delta e \equiv \Delta E/N \to 0$.)

Our expectation (2.26) that in the thermodynamic limit the entropy would be proportional to the system size N has been fully vindicated. So has the expectation that we could let $\Delta E \to 0$, although we have seen that we must do so carefully. The end result is that the entropy per particle of the pure classical monatomic ideal gas is

$$s(e,v) = k_B\left[\tfrac{3}{2}\ln\left(\frac{4\pi mev^{2/3}}{3h_0^2}\right) + \tfrac{5}{2}\right]. \tag{2.31}$$

This is called the "Sackur-Tetrode formula".[4] It is often written as

$$S(E,V,N) = k_B N\left[\tfrac{3}{2}\ln\left(\frac{4\pi mEV^{2/3}}{3h_0^2N^{5/3}}\right) + \tfrac{5}{2}\right], \tag{2.32}$$

with the understanding that it should be applied only to very large systems, i.e. to systems effectively at the thermodynamic limit.

Problems

2.12 Entropy of a spin system

Consider again the ideal paramagnet of problem 2.9.

[4] Otto Sackur (1880–1914) was a German physical chemist. Hugo Tetrode (1895–1931) was a Dutch theoretical physicist. Each independently uncovered this equation in 1912.

a. Write down an expression for $\ln \Omega(E, \Delta E, H, N)$ as a function of E. Simplify it using Stirling's approximation for large values of N. (Clue: Be careful to never take the logarithm of a number with dimensions.)

b. Find an expression for the entropy per spin $s(e, H)$ as a function of the energy per spin e and magnetic field H in the thermodynamic limit.

c. Sketch the resulting entropy as a function of the dimensionless quantity $u \equiv e/mH$. Does it take on the proper limits as $e \to \pm mH$? (In your sketch pay special attention to endpoints of the domain, and to any places where the function or its derivative suffers a discontinuity, or a kink, or goes to zero or infinity. In general, it is important that the sketch convey a correct impression of the qualitative character of the function, and less important that it be quantitatively accurate.)

2.13 **The approach to the thermodynamic limit**

For the classical monatomic ideal gas, plot entropy as a function of particle number using both the "finite size" form (2.24) and the Sackur-Tetrode form (2.32). We will see in problem 4.11 that for a gas at room temperature and atmospheric pressure, it is appropriate to use

$$EV^{2/3}/h_0^2 = (1.66 \times 10^{29}\ \mathrm{kg}^{-1})N^{5/3}. \tag{2.33}$$

Use the masses of argon and krypton. All other things being equal, is the thermodynamic limit approached more rapidly for atoms of high mass or for atoms of low mass?

2.14 **Other energy conventions**

In the text we found the entropy of a monatomic ideal gas by assuming that the potential energy of an atom was zero if the atom were inside the box and infinite if it were outside the box. What happens if we choose a different conventional zero of potential energy so that the potential energy is U for an atom inside the box and infinite for an atom outside the box?

2.15 **Other worlds**

Find the entropy as a function of E, V, and N in the thermodynamic limit for a monatomic ideal gas in a world with arbitrary[5] spatial dimensionality d.

2.16 **Ideal gas mixtures**

Consider a large sample of classical monatomic ideal gas that is a mixture of two components: N_A particles of mass m_A and N_B particles of mass m_B. If $N \equiv N_A + N_B$, show that the entropy is

$$S(E, V, N_A, N_B) \quad = \quad +k_B N_A \left[\tfrac{3}{2} \ln \left(\frac{4\pi m_A E V^{2/3}}{3h_0^2 N^{5/3}} \right) + \tfrac{5}{2} \right]$$

[5] Why, you wonder, should anyone care about a world that is not three-dimensional? For three reasons: (1) There are important physical approximations of two-dimensional worlds (namely surfaces) and of one-dimensional worlds (namely polymers). (2) The more general formulation might help you in unexpected ways. For example, Ken Wilson and Michael Fisher were trying to understand an important problem concerning critical points. They found that their technique could not solve the problem in three dimensions, but it *could* solve the problem in four dimensions. Then they figured out how to use perturbation theory to slide carefully from four dimensions to three dimensions, thus making their solution relevant to real physical problems. Wilson was awarded the Nobel Prize for this work. This illustrates the third reason, namely you never can tell what will be important and hence: (3) Knowledge is better than ignorance.

$$+k_B N_B \left[\tfrac{3}{2}\ln\left(\frac{4\pi m_B E V^{2/3}}{3h_0^2 N^{5/3}}\right)+\tfrac{5}{2}\right] \tag{2.34}$$
$$-k_B N\left[\left(\frac{N_A}{N}\right)\ln\left(\frac{N_A}{N}\right)+\left(\frac{N_B}{N}\right)\ln\left(\frac{N_B}{N}\right)\right].$$

(Clue: Use the result of problem D.2.)

2.6 Qualitative Features of Entropy

The concept of entropy is notoriously difficult to grasp. Even the consummate mathematician and physicist Johnny von Neumann claimed that "nobody really knows what entropy is anyway." Although we have an exact and remarkably simple formula for the entropy of a macrostate in terms of the number of corresponding microstates, this simplicity merely hides the subtle characterization needed for a real understanding of the entropy concept. To gain that understanding, we must examine the truly wonderful (in the original meaning of that overused word) surprises that this simple formula presents when applied to real physical systems.

2.6.1 Surprises

The monatomic ideal gas

Let us examine the Sackur-Tetrode formula (2.32) qualitatively to see whether it agrees with our understanding of entropy as proportional to the number of microstates corresponding to a given macrostate. If the volume V is increased, then the formula states that the entropy S increases, which certainly seems reasonable: If the volume goes up, then each particle has more places where it can be, so the entropy ought to increase. If the energy E is increased, then S increases, which again seems reasonable: If there is more energy around, then there will be more different ways to split it up and share it among the particles, so we expect the entropy to increase. (Just as there are many more ways to distribute a large booty among a certain number of pirates than there are to distribute a small booty.) But what if the mass m of each particle increases? (Experimentally, one could compare the entropy of, say, argon and krypton under identical conditions. See problem 2.27.) Our formula shows that entropy increases with mass, but is there any way to understand this qualitatively?

In fact, I can produce not just one but two qualitative arguments concerning the dependence of S on m. Unfortunately the two arguments give opposite results! The first relies upon the fact that

$$E=\frac{1}{2m}\sum_i p_i^2, \tag{2.35}$$

so for a given energy E, any individual particle may have a momentum ranging from 0 to $\sqrt{2mE}$. A larger mass implies a wider range of possible momenta, which suggests more microstates and a greater entropy. The second argument relies upon the fact that

$$E=\frac{m}{2}\sum_i v_i^2, \tag{2.36}$$

so for a given energy E, any individual particle may have a speed ranging from 0 to $\sqrt{2E/m}$. A larger mass implies a narrowed range of possible speeds, which suggests fewer microstates and a smaller entropy. The moral is simple: Qualitative arguments can backfire!

> That's the *moral* of the paradox. The *resolution* of the paradox is both deeper and more subtle: It hinges on the fact that the proper home of statistical mechanics is phase space, not configuration space, because Liouville's theorem implies conservation of volume in phase space, not in configuration space. This issue deeply worried the founders of statistical mechanics. See Ludwig Boltzmann, *Vorlesungen über Gastheorie* (J.A. Barth, Leipzig, 1896–98), part II, chapters III and VII [translated into English by Stephen G. Brush: *Lectures on Gas Theory* (University of California Press, Berkeley, 1964)]; J. Willard Gibbs, *Elementary Principles in Statistical Mechanics* (C. Scribner's Sons, New York, 1902), page 3; and Richard C. Tolman, *The Principles of Statistical Mechanics* (Oxford University Press, Oxford, U.K., 1938), pages 45, 51–52.

Freezing water

It is common to hear entropy associated with "disorder," "smoothness," or "homogeneity." How do these associations stand up to the simple situation of a bowl of liquid water placed into a freezer? Initially the water is smooth and homogeneous. As its temperature falls, the sample remains homogeneous until the freezing point is reached. At the freezing temperature the sample is an inhomogeneous mixture of ice and liquid water until all the liquid freezes. Then the sample is homogeneous again as the temperature continues to fall. Thus the sample has passed from homogeneous to inhomogeneous to homogeneous, yet all the while its entropy has decreased. (We will see later that the entropy of a sample always decreases as its temperature falls.)

Suppose the ice is then cracked out of its bowl to make slivers, which are placed back into the bowl and allowed to rest at room temperature until they melt. The jumble of irregular ice slivers certainly seems disordered relative to the homogeneous bowl of meltwater, yet it is the ice slivers that have the lower entropy. The moral here is that the huge number of microscopic degrees of freedom in the meltwater completely overshadow the minute number of macroscopic degrees of freedom in the jumbled ice slivers. But the analogies of entropy to "disorder" or "smoothness" invite us to ignore this moral and concentrate on the system's gross appearance and nearly irrelevant macroscopic features.

Reentrant phases

When the temperature falls at constant pressure, most pure materials pass from gas to liquid to solid. But the unusual materials called "liquid crystals," which consist of rod-like molecules, display a larger number of phases. For typical liquid crystals, the high-temperature liquid phase is isotropic, meaning that the positions and the orientations of the molecules are scattered about nearly at random. At lower temperatures, the substance undergoes a transition into the so-called "nematic" phase, in which the molecules tend to orient

Figure 2.2: Phase diagram of a liquid crystal mixture. The variable "composition" refers to the molecular weight ratio of 6OCB to 8OCB.

in the same direction but in which positions are still scattered. At still lower temperatures it passes into the "smectic" phase, in which the molecules orient in the same direction and their positions tend to fall into planes. Finally, at even lower temperatures, the molecules freeze into a conventional solid. The story told so far reinforces the picture of "entropy as disorder," with lower-temperature (hence lower entropy) phases showing more and more qualitative order.

But not all liquid crystals behave in exactly this fashion. One material called "hexyloxy-cyanobiphenyl" or "6OCB" passes from isotropic liquid to nematic to smectic *and then back to nematic again* as the temperature is lowered. The first transition suggests that the nematic phase is "less orderly" than the smectic phase, while the second transition suggests the opposite!

One might argue that the lower-temperature nematic phase — the so-called "reentrant nematic" — is somehow qualitatively different in character from the higher-temperature nematic, but the experiments summarized in figure 2.2 demonstrate that this is not the case. These experiments involve a similar liquid crystal material called "octyloxy-cyanobiphenyl" or "8OCB" which has no smectic phase at all. Adding a bit of 8OCB into a sample of 6OCB reduces the temperature range over which the smectic phase exists. Adding a bit more reduces that range further. Finally, addition of enough 8OCB makes the smectic phase disappear altogether. The implication of figure 2.2 is clear: there is no qualitative difference between the usual nematic and the reentrant nematic phases — you can move continuously from one to the other in the temperature–composition phase diagram.

Figure 2.3: Phase diagram for a mixture of water and nicotine.

The implication of reentrant phases for entropy is profound: Under some conditions the nematic phase has more entropy than the smectic phase and under other conditions less, while in all cases the nematic is qualitatively less ordered.

Another example of reentrant behavior appears in the phase diagram of the mixture of water and nicotine (see figure 2.3). For a wide range of mixing ratios, this mixture is a homogeneous solution at high temperatures, segregates into water-rich and nicotine-rich phases at moderate temperatures, yet becomes homogeneous again at low temperatures. At mixing ratios closer to pure water or pure nicotine, the mixture is homogeneous at all temperatures. Thus the high-temperature and reentrant homogeneous phases are in fact the same phase. (Reentrant phases are also encountered in type-II superconductors.)

Rust

According to a common misconception, "The entropy law describes the tendency for all objects to rust, break, fall apart, wear out, and otherwise move to a less ordered state."[6] Or "Entropy tends to make our eyes grow weaker as we age . . . entropy causes iron and other metals to rust."[7] Or "Entropy imposes itself in the form of limitation and in the natural tendency toward confusion and chaos, toward decay and destruction. Entropy is why ice cream melts and people die. It is why cars rust and forests burn. Preponderantly, the second law describes effects that we don't want to see."[8]

At least one error is held in common by these three quotes. Does rusting lead to increased entropy?

[6] Cutler J. Cleveland and Robert Kaufmann "Fundamental principles of energy," in *Encyclopedia of Earth*, last updated 29 August 2008, http://www.eoearth.org/article/Fundamental_principles_of_energy.

[7] Louis M. Savary, *Teilhard de Chardin: The Divine Milieu Explained* (Paulist Press, Mahwah, NJ, 2007) page 163.

[8] Gilbert L. Wedekind, *Spiritual Entropy* (Xulon Press, Fairfax, VA, 2003) page 68.

The rust reaction is

$$4\,\mathrm{Fe} + 3\,\mathrm{O_2} \to 2\,\mathrm{Fe_2O_3}.$$

According to standard tables[9] the entropy (at room temperature 298.15 K and at pressure 10^5 Pa) of one mole of Fe is 27.280 J/K, of one mole of O_2 is 205.147 J/K, and of one mole of Fe_2O_3 is 87.404 J/K. The entropy of the reactants is 724.561 J/K, the entropy of the products is 174.808 J/K, so the reaction results in an entropy *decrease* of 549.753 J/K.

It is easy to understand why this should be so: gases typically have much greater entropies that solids. And of course this doesn't mean that during rusting, the entropy of the universe decreases: although the entropy of the iron plus oxygen decreases, the entropy of the surroundings increases by even more. Nevertheless, it is clear that rusting itself involves a decrease in entropy, not an increase.

2.6.2 Entropy and the lattice gas

The four examples above should caution us about relying on qualitative arguments concerning entropy. Here is another situation[10] to challenge your intuition: The next page shows two configurations of $13^2 = 169$ squares tossed down on an area that has $35 \times 35 = 1225$ empty spaces, each of which could hold a square. (This system is called the "lattice gas model.") The two configurations were produced by two different computer programs (*Toss1* and *Toss2*) that used different rules to position the squares. (The rules will be presented in due course; for the moment I shall reveal only that both rules employ random numbers.) Which configuration do you think has the greater entropy? Be sure look at the configurations, ponder, and make a guess (no matter how ill-informed) before reading on.

[9] Ihsan Barin, *Thermochemical Data of Pure Substances* (VCH Publishers, Weinheim, Germany, 1995) pages 675, 1239, and 702.

[10] The argument of this section was invented by Edward M. Purcell and is summarized in Stephen Jay Gould, *Bully for Brontosaurus* (W.W. Norton, New York, 1991), pages 265–268, 260–261. The computer programs mentioned, which work under MS-DOS, are available for free downloading through http://www.oberlin.edu/physics/dstyer/.

Figure 2.4: A lattice gas configuration generated by the program *Toss1*.

Figure 2.5: A lattice gas configuration generated by the program *Toss2*.

Before analyzing these pictures, I must first confess that my question was very misleading. I asked "Which configuration has the greater entropy?", but entropy is not defined in terms of a single configuration (a single microstate). Instead, entropy is defined for a macrostate, and is related to the number of microstates that the system can take on and still be classified in that same macrostate. Instead of asking the question I did, I should have pointed out that I had two classes, two pools, of microstates, and that I used my two computer programs to select one member from each of those two pools. The selection was done at random, so the selected configuration should be considered typical. Thus my question should have been "Which configuration was drawn from the larger pool?", or "Which configuration is typical of the larger class?". Given this corrected question, you might want to go back and look at the configurations again.

I have asked these questions of a number of people (both students and professionals) and most of them guess that the lower configuration (the output of program *Toss2*), is typical of the class with larger entropy. The lower configuration is smoother, less clumpy. They look at the upper configuration and see patterns, which suggests some orderly process for producing that configuration, while the smoothness of the lower configuration suggests a random, disorderly construction process.

Let me end the suspense and tell you how my computer programs produced the two configurations. The top configuration was produced by tossing $169 = (13)^2$ squares down at random onto an area with $35 \times 35 = 1225$ locations, subject only to the rule that two squares could not fall on the same location. The bottom configuration was produced in exactly the same manner except that there was an additional rule, namely that two squares could not fall on adjacent locations either. Thus the top configuration was drawn from the pool of *all* patterns with 169 squares, while the bottom configuration was drawn from the much smaller pool of patterns with 169 squares *and with no two squares adjacent*. The top configuration is typical of the class with more configurations and hence greater entropy.

Look again at the bottom configuration. You will notice that there are no squares immediately adjacent to any given square. This "nearest neighbor exclusion" rule acts to spread out the squares, giving rise to the smooth appearance that tricks so many into guessing that the bottom configuration is typical of a class with high entropy.

Now look again at the top configuration. You will notice holes and clumps of squares, the inhomogeneities that lead many to guess that it is typical of a small class. But in fact one should *expect* a random configuration to have holes — only a very exceptional configuration is perfectly smooth.[11] This involves the distinction between a *typical* configuration and an *average* configuration. Typical configurations have holes: some have holes in the upper right, some in the middle left, some in the very center. Because the holes fall in various locations, the average configuration — the one produced by adding all the configurations and dividing by the number of configurations — is smooth. The average configuration is actually atypical. (Analogy: A typical person is not of average height. A typical person is somewhat taller or somewhat shorter than average, and very few people are exactly of average height. Any clothing manufacturer that produced only shirts of average size would quickly go bankrupt.) The presence of holes or clumps, therefore, need not be an indication of a pattern or of a design. However, we humans tend to find patterns wherever we look, even when no design is present.[12] In just this way the ancient Greeks looked into the nighttime sky, with stars

[11] In his book *The Second Law* (Scientific American Books, New York, 1984), P.W. Atkins promotes the idea that entropy is a measure of homogeneity. (This despite the everyday observation of two-phase coexistence.) To buttress this argument, the book presents six illustrations (on pages 54, 72, 74, 75, and 77) of "equilibrium lattice gas configurations." Each configuration has 100 occupied sites on a 40×40 grid. If the occupied sites had been selected at random, then the probability of any site being occupied would be 100/1600, and the probability of any given pair of sites both being occupied would be $1/(16)^2$. The array contains $2 \times 39 \times 39$ adjacent site pairs, so the expected number of occupied adjacent pairs would be $2(39/16)^2 = 11.88$. The actual numbers of occupied nearest-neighbor pairs in the six illustrations are 0, 7, 3, 7, 4, and 3. A similar calculation shows that the expected number of empty rows or columns in a randomly occupied array is $(15/16)^{10} \times 2 \times 40 = 41.96$. The actual numbers for the six illustrations are 28, 5, 5, 4, 4, and 0. I am confident that the sites in these illustrations were not occupied at random, but rather to give the impression of uniformity.

[12] Look again at figure 2.4. Do you see the dog catching a ball in the middle left? Do you see the starship *Enterprise* in the bottom right?

sprinkled about at random, and saw the animals, gods, and heroes that became our constellations.

2.6.3 Entropy and poker

An excellent illustration of the nature of entropy is given by the card game poker. There are many possible hands in poker, some valuable and most less so. For example, the hand

$$A\heartsuit, K\heartsuit, Q\heartsuit, J\heartsuit, 10\heartsuit \tag{2.37}$$

is an example of a royal flush, the most powerful hand in poker. There are only four royal flushes (the royal flush of hearts, of diamonds, of spades, and of clubs) and any poker player who has ever been dealt a royal flush will remember it for the rest of his life.

By contrast, no one can remember whether he has been dealt the hand

$$4\diamondsuit, 3\diamondsuit, J\heartsuit, 2\spadesuit, 7\diamondsuit \tag{2.38}$$

because this hand is a member of an enormous class of not-particularly-valuable poker hands. But the probability of being dealt this hand is *exactly* the same as the probability of being dealt the royal flush of hearts. The reason that one hand is memorable and the other is not has nothing to do with the rarity of that particular hand, it has everything to do with the size of the class of which the hand is a member.

This illustration of the importance of class rather than individual configuration is so powerful and so graphic that I shall call that distinction "the poker paradox".

2.6.4 Conclusion

It is often said that entropy is a measure of the disorder of a system. This qualitative concept has at least three failings: First, it is vague. There is no precise definition of disorder. Some find the abstract paintings of Jackson Pollock to be disorderly; others find them pregnant with structure. Second, it uses an emotionally charged word. Most of us have feelings about disorder (either for it or against it), and the analogy encourages us to transfer that like or dislike from disorder, where our feelings are appropriate, to entropy, where they are not. The most important failing, however, is that the analogy between entropy and disorder invites us to think about a single configuration rather than a class of configurations. In the lattice gas model there are many "orderly" configurations (such as the checkerboard pattern of figure 2.6) that are members of both classes. There are many other "orderly" configurations (such as the solid block pattern of figure 2.7) that are members only of the larger (higher entropy!) class.[13] The poker hand

$$2\clubsuit, 4\diamondsuit, 6\heartsuit, 8\spadesuit, 10\clubsuit$$

is very orderly, but a member of a very large class of nearly worthless poker hands.

[13] Someone might raise the objection: "Yes, but how many configurations would you have to draw from the pool, on average, before you obtained exactly the special configuration of figure 2.7?" The answer is, "Precisely the same number that you would need to draw, on average, before you obtained exactly the special configuration of figure 2.4." These two configurations are equally special and equally rare.

Figure 2.6: An orderly lattice gas configuration that is a member of both the large class and the small class of configurations.

Figure 2.7: An orderly lattice gas configuration that is a member of only the large (high entropy!) class of configurations.

Given the clear need for an intuition concerning entropy, and the appealing but unsatisfactory character of the simile "entropy as disorder," what is to be done? I suggest an additional simile, namely "entropy as freedom," which should be used not by itself but in conjunction with "entropy as disorder."

"Freedom" means a range of possible actions, while "entropy" means a range of possible microstates. If only one microstate corresponds to a certain macrostate, then the system has no freedom to choose its microstate — and it has zero entropy. If you are free to manage your own bedroom, then you may keep it

either neat or messy, just as high entropy macrostates encompass both orderly and disorderly microstates. The entropy gives the number of ways that the constituents of a system can be arranged and still be a member of the club (or class). If the class entropy is high, then there are a number of different ways to satisfy the class membership criteria. If the class entropy is low, then that class is very demanding — very restrictive — about which microstates it will admit as members. In short, the advantage of the "entropy as freedom" analogy is that it focuses attention on the variety of microstates corresponding to a macrostate whereas the "entropy as disorder" analogy invites focus on a single microstate.

While "entropy as freedom" has these benefits, it also has two of the drawbacks of "entropy as disorder." First, the term "freedom" is laden with even more emotional baggage than the term "disorder." Second, it is even more vague: political movements from the far right through the center to the extreme left all characterize themselves as "freedom fighters." Is there any way to reap the benefits of this analogy without sinking into the mire of drawbacks?

For maximum advantage, I suggest using *both* of these analogies. The emotions and vaguenesses attached to "freedom" are very different from those attached to "disorder," so using them together tends to cancel out the emotion. A simple sentence like "For macrostates of high entropy, the system has the freedom to chose one of a large number of microstates, and the bulk of such microstates are microscopically disordered" directs attention *away* from glib emotional baggage and *toward* the perspective of "more entropy means more microstates."

But perhaps it is even better to avoid analogies all together. Just think of entropy as describing the number of microstates consistent with the prescribed macrostate. And then you will remember always that entropy applies to a macrostate, i.e. a class of microstates, rather than to an individual microstate.

Problems

2.17 Entropy of the classical monatomic ideal gas: limits

a. Every macrostate has at least one corresponding microstate. Use this to show that in all cases $S \geq 0$.

b. Use the Sackur-Tetrode formula (2.31) to find the entropy of the classical monatomic ideal gas in the limit of high density (i.e. volume per particle approaches zero). [This absurd result indicates a breakdown in the approximation of "ideal" non-interacting particles. At high densities, as the particles crowd together, it is no longer possible to ignore the repulsive interactions which are responsible for the "size" of each atom.]

c. What is the entropy of the classical monatomic ideal gas when the energy equals the ground-state energy (i.e. $E \to 0$)? [This absurd result indicates a breakdown in the "classical" approximation. As we will see in chapter 6, quantum mechanics becomes a significant effect at low energies. Historically, this absurdity was one of the first indications that classical mechanics could not be universally true.]

2.18 (Q,E) **The coin toss**

If you flip a coin ten times, you expect on average to get five heads and five tails.

a. The pattern `HHHHHHHHHH` violates this expectation dramatically. What is the probability of obtaining this pattern?

b. The pattern `HTHTHTHTHT` matches this expectation exactly. What is the probability of obtaining this pattern?

c. What is the probability of obtaining the pattern `HTTTHHTTHT`?

d. What is the probability of obtaining a pattern with one tail and nine heads?

2.19 (Q) **Random walks**

You are presented with two pictures that supposedly sketch the progress of a random walker as it moves through space. One was drawn by a computer using a random number generator, the other by a human being. One picture is more or less uniform, the other has some spaces dark with lines and other spaces hardly touched. Which picture was drawn in which way?

2.20 **Poker**

In the game of poker, a hand consists of five cards drawn from a deck of 52 cards. The cards are evenly divided into four suits (hearts, diamonds, clubs, spades).

a. How many hands are there in poker? Explain your reasoning.

b. How many hands are flushes? (That is, all five cards of the same suit.)

c. Harry plays poker every evening, and each evening he is dealt 100 hands. How long must Harry play, on average, before being dealt the hand

$$4\diamondsuit, 3\diamondsuit, J\heartsuit, 2\spadesuit, 7\diamondsuit \tag{2.39}$$

2.7 Using Entropy to Find (Define) Temperature and Pressure

References: Schroeder sections 3.1, 3.4, and 3.5. Reif section 3.3. Kittel and Kroemer pages 30–41.

You will remember that one of the problems with our ensemble approach to statistical mechanics is that, while one readily sees how to calculate ensemble average values of microscopic (mechanical) quantities like the kinetic energy, the potential energy, or the mean particle height, it is hard to see how to calculate the sort of macroscopic (thermodynamic) quantity that we're really more interested in anyway, such as the temperature or pressure. In fact, it is not even clear how we should define these important quantities. Our task in this lecture is to motivate such definitions. Because our arguments are motivations rather than deductions, they will be suggestive rather than definitive. The arguments can be made substantially more solid, but only at the cost of dramatic increase in mathematical complexity. (See David Ruelle's *Statistical Mechanics: Rigorous Results* [Physics QC174.8.R84 1989].) At this stage in your education, you should just accept the arguments and the definitions, realizing that they are not definitive, and go on to learn how to

use them. If you're still worried about them a year from now, then that would be a good time to read the more complex arguments in Ruelle's book.

An analogy will help explain the mathematical level of this lecture. Why is the period of the simple harmonic oscillator independent of amplitude? I can give two answers: 1) At larger amplitudes, the particle has more distance to move, but it also moves faster, and these two effects exactly cancel. 2) Solve the problem mathematically, and you'll see that $\omega = \sqrt{k/m}$ which is clearly unaffected by amplitude. The first answer gives more insight into what is going on physically, but it is not definitive. (It does not explain why the two effects cancel exactly rather that have one dominate the other, which is the case for non-parabolic oscillators.) The second argument is bulletproof, but it gives little insight. It is best to use both types of argument in tandem. Unfortunately, the bulletproof arguments for this section are really too elaborate and formal to present here. (In a few weeks I will give you a sample of a rigorous statistical mechanical argument: "The Grand Canonical Ensemble in the Thermodynamic Limit", section 4.8. It is the easiest such argument that I know of, and you will see that it's not very easy at all!)

2.7.1 Rapidly increasing character of the $\Omega(E)$ function

For a monatomic ideal gas ("in the thermodynamic limit") the entropy is

$$S(E,V,N) \equiv k_B \ln \Omega(E,V,N) = k_B N \left[\frac{3}{2} \ln \left(\frac{4\pi m E V^{2/3}}{3 h_0^2 N^{5/3}} \right) + \frac{5}{2} \right]. \tag{2.40}$$

Thus $\Omega(E,V,N)$, the number of microstates consistent with the macrostate (or thermodynamic state) specified by E, V, and N, is

$$\Omega(E,V,N) = e^{(5/2)N} \left(\frac{4\pi m E V^{2/3}}{3 h_0^2 N^{5/3}} \right)^{(3/2)N}. \tag{2.41}$$

In particular, as a function of energy,

$$\Omega(E) = (\text{constant}) E^{(3/2)N}. \tag{2.42}$$

This is a very rapidly increasing function of E. (The function $f(x) = x^2$ increases more rapidly than $f(x) = x$. The function $f(x) = x^3$ increases more rapidly than $f(x) = x^2$. The function $f(x) = x^{10^{23}}$ increases very rapidly indeed!) Although we have proven this result only for the ideal gas, it is plausible[14] for interacting systems as well: If you throw more energy into a system, there are more ways for the energy to be arranged. (There is an argument to this effect in Reif's section 2.5. The result is not, however, *always* true... check out the ideal paramagnet.)

As the particle number N grows larger and larger (approaching the "thermodynamic limit") this increase of Ω with E becomes more and more rapid.

[14] Through the booty argument of page 27.

2.7.2 Two isolated systems

Consider two systems, A and B, enclosed in our perfectly reflecting no-skid walls. System A has energy E_A, volume V_A, and particle number N_A; similarly for system B.

I have drawn the two systems adjacent, but they might as well be miles apart because they can't affect each other through the no-skid walls. No energy can flow from one system to the other because no energy can move from the atoms in the system into the walls. The formal name for such walls is "insulating" or "adiabatic". As we have discussed previously, such perfectly insulating walls do not exist in nature (if nothing else, system A affects system B through the gravitational interaction), but good experimental realizations exist and they are certainly handy conceptually.

The total system consisting of A plus B is characterized by the macroscopic quantities E_A, E_B, V_A, V_B, N_A, and N_B. The number of microstates corresponding to this macrostate is

$$\Omega_T(E_A, E_B, V_A, V_B, N_A, N_B) = \Omega_A(E_A, V_A, N_A)\Omega_B(E_B, V_B, N_B). \tag{2.43}$$

2.7.3 Two systems in thermal contact

Consider the same two systems, A and B, as before, but now allow the wall between them to be an ordinary, energy-permeable wall. Such a wall is called "diathermal". (The walls between the systems and the outside world are still insulating.)

Now the mechanical parameters V_A, V_B, N_A, and N_B still remain constant when the two systems are brought together, but the energies E_A and E_B change. (Of course the total energy $E_T \equiv E_A + E_B$ remains constant as individual energies change.) Such contact between two systems, where energy can flow between the systems

but where the mechanical parameters don't change, is called "thermal contact". Once they are in contact, I don't *know* what the distribution of energy between A and B is: It could be that A is in its ground state and all the excess energy is in B, or vice versa, and it could change rapidly from one such extreme to the other. But while I no longer have *knowledge* of the distribution, I do have an *expectation*. My expectation, drawn from daily experience, is that energy will flow from one system to another until E_A and E_B take on some pretty well-defined final equilibrium values. The exact energies will then fluctuate about those equilibrium values, but those fluctuations will be small. And what characterizes this final equilibrium situation? Your first guess might be that the energy on each side should be equal. But no, if a big system were in contact with a small system, then you'd expect more energy to end up in the big system. Then you might think that the energy per particle on each side should be equal, but that's not right either. (Ice and liquid water are in equilibrium at the freezing point, despite the fact that the energy per particle of liquid water is much greater than than of ice.) The correct expectation (known to any cook) is that:

> When two systems A and B are brought into thermal contact, E_A and E_B will change until they reach equilibrium values $E_A^{(e)}$ and $E_B^{(e)}$, with negligible fluctuations, and at equilibrium both systems A and B will have the same "temperature".

This is the qualitative property of temperature that will allow us, in a few moments, to produce a rigorous, quantitative definition. To do so we now look at the process microscopically instead of through our macroscopic expectations.

Once the two systems are in thermal contact the energy of system A could be anything from 0 to the total energy E_T.[15] For any given E_A, the number of accessible microstates of the combined system, A plus B, is $\Omega_A(E_A)\Omega_B(E_T - E_A)$. Thus the total number of microstates of the combined system is

$$\Omega_T(E_T) = \sum_{E_A=0}^{E_T} \Omega_A(E_A)\Omega_B(E_T - E_A). \tag{2.44}$$

(Please don't tell me the obvious... that E_A is a continuous quantity so you have to integrate rather than sum. This is a seat-of-the-pants argument. If you want attention to fine points, look them up in Ruelle's book.)

Let us examine the character of the summand, $\Omega_A(E_A)\Omega_B(E_T - E_A)$, as a function of E_A. The functions $\Omega_A(E_A)$ and $\Omega_B(E_B)$ are both rapidly increasing functions of their arguments, so $\Omega_B(E_T - E_A)$ is a rapidly *decreasing* function of E_A. Thus the product $\Omega_A(E_A)\Omega_B(E_T - E_A)$ will start out near zero, then jump up for a narrow range of E_A values and then fall quickly to near zero again.

[15] Here and in the rest of this section, we assume that the ground state energy of both system A and system B has been set by convention to zero.

In conclusion, for most values of E_A the summand nearly vanishes, so most of the contribution to the sum comes from its largest term, which is located at an energy we will call $\hat{E}_A$:

$$\Omega_T(E_T) = \Omega_A(\hat{E}_A)\Omega_B(E_T - \hat{E}_A) + \text{ small change.} \tag{2.45}$$

The sharply peaked character of the product $\Omega_A(E_A)\Omega_B(E_T - E_A)$ becomes more and more pronounced as the thermodynamic limit is approached, so in this limit the "small change" can be ignored. You can see that for macroscopic systems the "microscopically most probable value" $\hat{E}_A$ should be interpreted as the thermodynamic equilibrium value $E_A^{(e)}$.

We locate this value by maximizing

$$\Omega_A(E_A)\Omega_B(E_T - E_A). \tag{2.46}$$

To make life somewhat easier, we will not maximize this function directly but, what is the same thing, maximize its logarithm

$$f(E_A) \equiv \ln[\Omega_A(E_A)\Omega_B(E_T - E_A)] \tag{2.47}$$

$$= \ln \Omega_A(E_A) + \ln \Omega_B(E_T - E_A). \tag{2.48}$$

Recognizing the entropy terms we have

$$k_B f(E_A) = S_A(E_A) + S_B(E_T - E_A), \tag{2.49}$$

and differentiation produces

$$\frac{\partial(k_B f(E_A))}{\partial E_A} = \frac{\partial S_A}{\partial E_A} + \frac{\partial S_B}{\partial E_B}\underbrace{\frac{\partial E_B}{\partial E_A}}_{-1}. \tag{2.50}$$

We locate the maximum by setting the above derivative to zero, which tells us that at equilibrium

$$\frac{\partial S_A}{\partial E_A} = \frac{\partial S_B}{\partial E_B}. \tag{2.51}$$

Our expectation is that the temperature will be the same for system A and system B at equilibrium, while our microscopic argument shows that $\partial S/\partial E$ will be the same for system A and system B at equilibrium. May we conclude from this that

$$\frac{\partial S}{\partial E} = \text{Temperature ?} \tag{2.52}$$

No we cannot! All we can conclude is that

$$\frac{\partial S}{\partial E} = \text{invertible function (Temperature).} \tag{2.53}$$

To find out which invertible function to choose, let us test these ideas for the ideal gas. Using

$$S(E) = k_B \ln C + \tfrac{3}{2} k_B N \ln E + \tfrac{5}{2} k_B N \tag{2.54}$$

we find

$$\frac{\partial S}{\partial E} = \frac{3}{2} k_B \frac{N}{E}, \tag{2.55}$$

so a large value of E results in a small value of $\partial S/\partial E$. Because we are used to thinking of high temperature as related to high energy, it makes sense to *define (absolute) temperature* through

$$\frac{1}{T(E,V,N)} \equiv \frac{\partial S(E,V,N)}{\partial E}. \tag{2.56}$$

Applying this definition to the ideal gas gives the famous result

$$\frac{E}{N} = \frac{3}{2} k_B T. \tag{2.57}$$

In fact, this result is *too* famous, because people often forget that it applies only to a classical monatomic ideal gas. It is a common misconception that temperature is defined through equation (2.57) rather than through equation (2.56), so that temperature is always proportional to the average energy per particle or to the average kinetic energy per particle. If this misconception were correct, then when ice and water were in equilibrium, the molecules in the ice and the molecules in the water would have the same average energy per particle! In truth, these proportionalities hold exactly only for noninteracting classical point particles.

2.7.4 Two systems in thermal and mechanical contact

Consider the same two systems, A and B, as before, but now allow them to exchange both energy and volume. (The wall between A and B is a sliding, diathermal wall. The walls between the systems and the outside world remains insulating and rigid.)

Now the numbers N_A and N_B remain constant, but the energies E_A and E_B, and the volumes V_A and V_B, do change. Both the total energy $E_T \equiv E_A + E_B$ and the total volume $V_T \equiv V_A + V_B$ remain constant as the individual energies and volumes change. Because of the type of contact, I don't *know* the distribution of energy between A and B, nor do I *know* the distribution of volume between A and B. But just as with energy, I do have an *expectation*.

> When two systems A and B are brought into thermal and mechanical contact, E_A, E_B, V_A, and V_B will change until they reach equilibrium values $E_A^{(e)}$, $E_B^{(e)}$, $V_A^{(e)}$, and $V_B^{(e)}$, with negligible fluctuations, and at equilibrium systems A and B will have the same temperature and the same "pressure".

Just as with energy and temperature, this qualitative expectation will lead us, after a microscopic argument, to a rigorous, quantitative definition of pressure.

> In most cases the system will first reach a preliminary mechanical equilibrium as the wall slides quickly to make the pressure equal on both sides. Then more slowly, as energy seeps from one side to the other, the two temperatures will equilibrate. As the temperatures change the pressure on each side changes, and the wall again slides, more slowly now, to reflect these changes. The argument that follows could be carried out using a sliding insulated wall, instead of a sliding diathermal wall, but in this case energy is still transferred from A to B through mechanical work, so the argument is in fact more complicated that way.

The total number of microstates of the combined system is

$$\Omega_T(E_T, V_T) = \sum_{E_A} \sum_{V_A} \Omega_A(E_A, V_A)\Omega_B(E_T - E_A, V_T - V_A). \tag{2.58}$$

For an ideal gas, $\Omega(V) = CV^N$, but in general we expect $\Omega(V)$ to be a rapidly increasing function of V. (In analogy to the "pirate booty argument", we might call this the "sardine argument.") Just as before, this leads us to expect that the product $\Omega_A(E_A, V_A)\Omega_B(E_T - E_A, V_T - V_A)$ will nearly vanish except in the vicinity of the equilibrium values $E_A = E_A^{(e)}$ and $V_A = V_A^{(e)}$, and that the sum can be approximated by its largest value

$$\Omega_T(E_T, V_T) \approx \Omega_A(E_A^{(e)}, V_A^{(e)})\Omega_B(E_T - E_A^{(e)}, V_T - V_A^{(e)}). \tag{2.59}$$

We locate the equilibrium values by maximizing the product $\Omega_A(E_A, V_A)\Omega_B(E_T - E_A, V_T - V_A)$ with respect to both E_A and V_A. This leads directly to the requirements

$$\frac{\partial S_A}{\partial E_A} = \frac{\partial S_B}{\partial E_B}, \tag{2.60}$$

as before, and

$$\frac{\partial S_A}{\partial V_A} = \frac{\partial S_B}{\partial V_B}. \tag{2.61}$$

We conclude that

$$\frac{\partial S}{\partial V} = \text{invertible function (Pressure, Temperature)}. \tag{2.62}$$

The dimensions of $\partial S/\partial V$ are

$$\left[\frac{\partial S}{\partial V}\right] = \frac{\mathrm{J/K}}{\mathrm{m}^3} \tag{2.63}$$

while those for pressure are

$$[p] = \frac{\mathrm{J}}{\mathrm{m}^3}. \tag{2.64}$$

This motivates us to *define pressure* through

$$\frac{p(E,V,N)}{T(E,V,N)} \equiv \frac{\partial S(E,V,N)}{\partial V}. \tag{2.65}$$

For an ideal gas,

$$S(V) = k_B N[C + \ln V]. \tag{2.66}$$

so

$$\frac{\partial S}{\partial V} = \frac{k_B N}{V} = \frac{p}{T}, \tag{2.67}$$

leading to

$$pV = Nk_B T, \tag{2.68}$$

the ideal gas equation of state! (Which is again too famous for its own good. Too many statistical mechanics courses degenerate into "an intensive study of the ideal gas", and it is a shame to waste such a beautiful, powerful, and diverse subject on the study of only one substance, especially since that substance doesn't even exist in nature!)

2.7.5 Two systems that can exchange energy, volume, and particles

We return to the same situation as before, but now the wall between A and B is not only diathermal and sliding, but also allows particles to flow. (Imagine piercing a few tiny holes in the wall.)

The reasoning in this case exactly parallels the reasoning already given two above, so I will not belabor it. I will merely point out that we can now define a new quantity, the *chemical potential* μ, which governs the flow of particles just as temperature governs the flow of energy or pressure governs the flow of volume. It is

$$-\frac{\mu(E,V,N)}{T(E,V,N)} \equiv \frac{\partial S(E,V,N)}{\partial N}. \tag{2.69}$$

2.7.6 The meaning of chemical potential

The upshot of all our definitions is that

$$dS = \frac{1}{T}\,dE + \frac{p}{T}\,dV - \frac{\mu}{T}\,dN. \tag{2.70}$$

What do these definitions mean physically? The first term says that if two systems are brought together so that they can exchange energy but not volume or particle number, then in the system with high temperature the energy will decrease while in the system with low temperature the energy will increase. The second term says that if two systems with the same temperature are brought together so that they can exchange volume but not particle number, then in the system with high pressure the volume will increase while in the system with low pressure the volume will decrease. The third term says that if two systems with the same temperature and pressure are brought together so that they can exchange particles, then in the system with high chemical potential the number will decrease while in the system with low chemical potential the number will increase. In summary:

Exchange of energy?	Exchange of volume?	Exchange of particles?	Result:
Yes.	Prohibited.	Prohibited.	System with high T decreases in E.
Permitted, but $\Delta T = 0$.	Yes.	Prohibited.	System with high p increases in V.
Permitted, but $\Delta T = 0$.	Permitted, but $\Delta p = 0$.	Yes.	System with high μ decreases in N.

Chemists have a great name for chemical potential... they call it "escaping tendency", because particles tend to move from a region of high μ to a region of low μ just as energy tend to move from a region of high T to a region of low T. (G.W. Castellan, *Physical Chemistry*, 2nd edition, Addison-Wesley, Reading, Mass., 1971, section 11–2.) You can see that chemical potential is closely related to density, and indeed for an ideal gas

$$\mu = k_B T \ln(\lambda^3(T)\rho), \tag{2.71}$$

where $\rho = N/V$ is the number density and $\lambda(T)$ is a certain length, dependent upon the temperature, that we will see again (in equation (5.4)).

> Once again, remember that this relation holds *only* for the ideal gas. At the freezing point of water, when ice and liquid water are in equilibrium, the chemical potential is the same for both the solid and the liquid, but the densities clearly are not!

Problems

2.21 (Q) **The location of $E_A^{(e)}$**
In the figure on page 41, does the equilibrium value $E_A^{(e)}$ fall where the two curves $\Omega_A(E_A)$ and $\Omega_B(E_T - E_A)$ cross? If not, is there a geometrical interpretation, in terms of that figure, that does characterize the crossing?

2.22 (Q) **Meaning of temperature and chemical potential**
The energy of a microscopic system can be dissected into its components: the energy due to motion,

the energy due to gravitational interaction, the energy due to electromagnetic interaction. Can the temperature be dissected into components in this way? The chemical potential? (Clue: Because the temperature and chemical potential are derivatives of the entropy, ask yourself first whether the entropy can be dissected into components.)

2.23 (E) **Chemical potential of an ideal gas**

a. Show that the chemical potential of a pure classical monatomic ideal gas is

$$\mu = -\frac{3}{2}k_BT\ln\left(\frac{4\pi mEV^{2/3}}{3h_0^2N^{5/3}}\right) \tag{2.72}$$

$$= -k_BT\ln\left[\left(\frac{2\pi mk_BT}{h_0^2}\right)^{3/2}\frac{V}{N}\right]. \tag{2.73}$$

b. Show that when the temperature is sufficiently high and the density sufficiently low (and these are, after all, the conditions under which the ideal gas approximation is valid) the chemical potential is negative.

c. Show that

$$\frac{\partial\mu(T,V,N)}{\partial T} = \frac{\mu(T,V,N)}{T} - \frac{3}{2}k_B. \tag{2.74}$$

At a fixed, low density, does μ increase or decrease with temperature?

2.24 **Ideal paramagnet, take three**

Find the chemical potential $\mu(E,H,N)$ of the ideal paramagnet. (Your answer must *not* be a function of T.) (To work this problem you must have already worked problems 2.9 and 2.12.)

Resources

Percival as CM for chaos and SM. Hilborn on chaos.

PAS chaos programs, public domain stadium and wedge. CUPS stadium.

Tolman, Ruelle.

2.8 Additional Problems

2.25 **Accessible configurations in another spin system**

The *Ising model* for a ferromagnetic material such as nickel is different from the "ideal paramagnet" discussed in problem 2.9. In this model the spins reside at lattice sites and may point either up or down, but in contrast to the ideal paramagnet model, two adjacent spins *do* interact. Specifically, if two adjacent spins point in the same direction, the pair has energy $-J$. If they point in different directions, the pair has energy $+J$. (In the ferromagnetic situation considered here, $J > 0$.) We consider here a one-dimensional Ising model of N sites with zero external magnetic field.

a. Not every energy is possible for this model. What is the maximum possible energy? The minimum? What is the smallest possible energy spacing between configurations?

b. Consider the energy range from E to $E + \Delta E$ where ΔE is small compared to NJ but large compared to J. What is the approximate number of states $\Omega(E, \Delta E, J, N)$ lying within this energy range?

c. Write down an expression for the entropy $S(E, \Delta E, J, N)$ as a function of E. (Clue: Be careful to never take the logarithm of a number with dimensions.)

d. Take the thermodynamic limit to find the entropy per atom as a function of the energy per atom: $s(e, J)$. (Clue: Use Stirling's approximation.)

e. Check your work by comparing the Ising entropy per spin $s(e, J)$ when the interactions vanish ($J = 0$) to the ideal paramagnet entropy per spin $s(e, H)$ in zero field. (See problem 2.12. Clue: If $J = 0$, then $e = 0$ also. Why?)

General clues: It helps to concentrate not on the status of the *spins* (up or down) but on the status of the *pairs* (alike or different). How many nearest neighbor pairs are there? For each given configuration of pairs, how many corresponding configurations of spins are there?

2.26 **Shake and spin**

Make up a problem involving an ideal paramagnet with atoms free to move around.

2.27 **Entropy as a function of mass**

Use the Sackur-Tetrode formula to find the difference between the entropy of a sample of argon gas and a sample krypton gas under identical conditions. Use the data in Ihsan Barin, *Thermochemical Data of Pure Substances*, third edition (VCH Publishers, New York, 1995) to test this prediction at the temperatures 300 K and 2000 K.

2.28 (Q,D) **For discussion: "The icy cold of outer space"**
Is outer space at a high temperature or a low temperature? Before answering, consider these points:

a. The environment of outer space consists of protons and photons streaming away from the sun. The two systems are virtually uncoupled (do not interact with each other). I'm asking for the temperature of the protons.

b. It was implicitly (perhaps illicitly) assumed in section 2.7.3 that the centers of mass of the two systems in thermal contact were in the same reference frame. This is a general principle: When referring to the temperature of a moving body, we mean the temperature as measured by a thermometer moving along with that body.

c. Remember that in the arguments of section 2.7.3 the two systems could exchange energy *but not particles*.

d. What if we approximated the protons by an ideal gas?

e. Look at the *Handbook of Chemistry and Physics* under index entry "Temperature, Earth's atmosphere".

2.29 (Q,D) **Errors in Reif**
Scan Reif pages 47–61, but don't read them in detail as they are rife with errors. Point out the major error in quantum mechanics made in each of the following passages. (Use a maximum of three sentences per passage.)

a. The examples on pages 48 and 49.

b. The second paragraph on page 57.

c. The first paragraph on page 60.

2.30 **Polymers**
I'd like to think up some problem concerning polymers of fixed length *vs.* polymers of fixed "activity for length". The activity is analogous to chemical potential in that it is a control knob for polymer length. It might render the whole concept more concrete.

Chapter 3

Thermodynamics

3.1 Heat and Work

We've seen how to calculate—at least in principle, although our techniques were primitive and difficult to use—the entropy $S(E, V, N)$. We've also argued ("the booty argument", section 2.7.1) that for most systems entropy will increase monotonically with energy when V and N are fixed. Thus we (can)/(will always be able to) invert the function to find energy as a function of entropy, volume, and number:

$$E(S, V, N). \tag{3.1}$$

Recall that V and N are just stand-ins as representative mechanical parameters.

But how can we measure entropy? Walk into any laboratory and the experimentalist there will say "I have thermometers for measuring temperature. I have resistors, ammeters, and voltmeters, and a I have a procedure for using these instruments to measure the heat flowing into a sample. I have calibers for measuring volume and manometers for measuring pressure. But I have no instrument to measure the number of microstates consistent with a given macrostate." Although we have produced a *conceptual definition* of entropy, we need also an *operational definition* of entropy.

An "operational definition" defines the target word, not in terms of other words, but in terms of a procedure that produces a decision or a numerical value concerning that target word. For example, an operational definition for the word *tuberculosis* is "a disease marked by wasting, cough, fever, and formation of small cheesy lumps in the lungs". In contrast, a conceptual definition is "a disease caused by the bacillus bacterium *Mycobacterium tuberculosis*". The operational definition tells you how to diagnose the disease; the conceptual definition tells you what's going on and suggests a cure. We need to produce an operational definition to go along with our conceptual definition of entropy, and that's the starting point of this chapter.

Heat:

Change in energy with no change in mechanical parameters =
Change in energy due to temperature difference between system and surroundings =
Heat (absorbed by system) = Q

Configuration work.

Change in energy due to change in mechanical parameters =
Change in energy when system is wrapped in insulating ("no-skid") walls =
Configuration work (done by system) = W_{conf}

Dissipative work.

Change in energy due to non-equilibrium effects =
Change in energy due to, e.g., shaking, stirring, or running current through the system =
Dissipative work (done by system) = W_{diss}

It is an empirical fact (sometimes called "the second law of thermodynamics"), that dissipative work always increases the energy of the system. Joule the brewer; story about honeymoon at Cascade de Sallanches.

Caution: Heat and work refer to mechanisms for energy *change*, not to types of energy. Do not think that heat goes into a change of kinetic energy (or into "mixed up energy") whereas work goes into a change of potential energy (or into "ordered energy").

Any change due to heating could also be done via dissipative work, and there's no way you could tell the difference. Joule called this "the conversion of work to heat", although that's not correct using the modern definitions of "work" and "heat".

Analogy: You can put money into your bank account through depositing cash or checks, or through an electronic funds transfer. You can take money out of your bank account through these three mechanisms too. But once the money is inside your bank account, you cannot say "This dollar was deposited through cash, while that dollar was deposited through electronic funds transfer." Once they're in your account, you can't distinguish the dollars through the mechanism of entrance. Similarly with energy. You can increase the temperature and energy of a cup of water by putting it near a flame, or by putting it near a high-temperature resistor, or by stirring it rigorously. Given a high-temperature cup of water, there's no way to tell which of these mechanisms was used.

$$\Delta E = Q + W_{\text{conf}} + W_{\text{diss}} \tag{3.2}$$

Problems

3.1 **What is heat?**

Below is an excerpt from a sixth-grade science textbook. Find at least six errors in the excerpt.

> *What is heat?*
>
> You have learned that all matter is made up of atoms. Most of these atoms combine to form molecules. These molecules are always moving—they have kinetic energy. *Heat* is the energy of motion (kinetic energy) of the particles that make up any piece of matter.
>
> The amount of heat a material has depends on how many molecules it has and how fast the molecules are moving. The greater the number of molecules and the faster they move, the greater the number of collisions between them. These collisions produce a large amount of heat.
>
> How is heat measured? Scientists measure heat by using a unit called a *calorie*. A calorie is the amount of heat needed to raise the temperature of 1 gram of 1 water 1 degree centigrade (Celsius).
>
> A gram is a unit used for measuring mass. There are about 454 grams in 1 pound.
>
> *What is temperature?*
>
> The amount of hotness in an object is called its temperature. A thermometer is used to measure temperature in units called degrees. Most thermometers contain a liquid.

3.2 Entropy

$$\begin{aligned} dE &= T\,dS - p\,dV + \mu\,dN \\ &= đQ + đW_{\text{conf}} + đW_{\text{diss}} \\ &\geq đQ + đW_{\text{conf}} \end{aligned}$$

but

$$đW_{\text{conf}} = -p\,dV + \mu\,dN$$

so

$$T\,dS \geq đQ$$

where the equality holds for quasistatic changes.

The operational definition of entropy is that, for a quasistatic change,

$$dS = \frac{đQ}{T}. \tag{3.3}$$

If we walk into a laboratory with *this* definition of entropy, the experimentalist will say "Sure I can find this quantity. I've got thermometers; and I've got resistors, ammeters, and voltmeters, plus a procedure for using them to calculate heat flow."

For any change (perhaps quasistatic, perhaps not)

$$dS \geq \frac{đQ}{T}. \tag{3.4}$$

(The entropy change might be due to heat, might be due to dissipative work, might be due to a combination of the two.)

Configurational entropy vs. thermal entropy

The operational, thermodynamic definition for entropy, equation (3.3), was in fact discovered before the more fundamental, conceptual, statistical definition (2.7) was. Because of this dual definition some people get the misconception that there are two kinds of entropy: the statistical kind ("configurational entropy") and the operational kind ("thermal entropy"). That's false. These are two different ways of looking at the same thing.

3.2 (I) **Ideal gas with variables S and V.**

a. Use equation (2.32) to show that for a sample of monatomic ideal gas with a fixed number of particles, the energy function is

$$E(S,V) = E_0 \left(\frac{V_0}{V}\right)^{2/3} e^{S/S_0}, \tag{3.5}$$

where E_0, V_0, and S_0 are constants with the dimensions of energy, volume, and entropy.

b. Use the formulas

$$T(S,V) = \frac{\partial E(S,V)}{\partial S} \quad \text{and} \quad p(S,V) = -\frac{\partial E(S,V)}{\partial V} \tag{3.6}$$

to find the temperature and pressure functions for this sample.

3.3 (I) **Dependence of heat and work on path**

Suppose the ideal gas sample mentioned in the previous problem undergoes a change from an initial state with volume V_0 and entropy S_0 to a final state with volume $8V_0$ and entropy $(1+5\ln 2)S_0$. There are an infinite number of quasistatic paths that execute this change. For example:

A First increase the entropy from $S = S_0$ to $S = (1+5\ln 2)S_0$, keeping the volume fixed at $V = V_0$, then expand from $V = V_0$ to $V = 8V_0$, keeping the entropy fixed at $S = (1+5\ln 2)S_0$.

B Increase the volume and entropy simultaneously along the path

$$S(V) = S_0 \left[\frac{5}{3} \ln \left(\frac{V}{V_0} \right) + 1 \right]. \tag{3.7}$$

C First expand from $V = V_0$ to $V = 8V_0$, keeping the entropy fixed at $S = S_0$, then increase the entropy from $S = S_0$ to $S = (1+5\ln 2)S_0$, keeping the volume fixed at $V = 8V_0$.

Each path has the same change in volume, entropy, and energy, but each will have a different heat and work. (Remember that each path is quasistatic.)

a. Calculate the heat absorbed by the sample and the work done on the sample for each of these three paths.

b. Find the difference $Q + W$ for each of these three paths.

c. A fourth path from the initial to the final state involves a non-quasistatic change and the absorption of heat $Q = 3E_0$. What work is done on the sample during this change?

(In making up this problem I choose somewhat peculiar initial and final states, and connecting paths, so that the above quantities could be calculated analytically. Despite these peculiarities the principle illustrated by the problem should be clear: heat and work depend upon path, but their difference the energy change does not.)

3.4 (I*) **Work in terms of p and V**

We have so far considered work and heat in a fluid system specified by the variables S, V, and N, and have often assumed that N is constant. Another way of specifying such a system is through the variables p (pressure) and V. This problem considers quasistatic processes described as paths in the (p, V) plane.

a. Show that the work done on the system in changing quasistatically from the initial state (V_i, p_i) to the final state (V_f, p_f) along the path shown in the figure below is the negative of the area under the curve. What is the work done if the path is traversed in the opposite direction?

b. What is the work done on the system through the same change along the path below?

c. Finally, what is the work done on the system if it comes back to its starting point as in the cycle shown below? What if the cycle is instead executed in a counterclockwise fashion?

3.5 (I) **Fluid work**

In a quasistatic, adiabatic process, a sample of gas expands from $V_i = 1$ m^3 to $V_f = 8$ m^3, while the pressure drops from $p_i = 32$ Pa to $p_f = 1$ Pa. It is observed that during this process the pressure and volume are related by

$$pV^\gamma = \text{ constant}, \tag{3.8}$$

where the constant γ is $\frac{5}{3}$. Of course, this is not the only process leading from this particular initial state to that particular final state. Find the total work done on the system and heat absorbed by the system in each of the following quasistatic processes, all of which connect the same two thermodynamic states.

a. The adiabatic process described above.

b. At constant pressure, the gas expands from V_i to V_f. (Heat must be added to the system during this expansion in order to keep the pressure constant.) Then at constant volume, the the pressure is reduced from p_i to p_f. (Heat must be extracted from the system during this stage of the process.)

c. The volume is increased and heat is supplied at just the right rate to make the pressure decrease linearly with volume.

d. As in part (b.) but with the two steps performed in opposite order.

3.3 Heat Engines

The emphasis of this course is on the properties of matter. This section is independent of the properties of matter! It is included because:

1. historically important
2. expected coverage (e.g. for GRE)
3. important from an applications and engineering standpoint (See, for example, "Hurricane heat engines" by H.E. Willoughby, *Nature* **401** (14 October 1999) 649–650. Or "A thermoacoustic Stirling heat engine" by S. Backhaus and G.W. Swift, *Nature* **399** (27 May 1999) 335–338, and "Traveling-wave thermoacoustic electric generator," by S. Backhaus, E. Tward, and M. Petach, *Applied Physics Letters* **85** (9 August 2004) 1085–1087, concerning an engine that might replace nuclear thermal power sources in deep space missions.)
4. fascinating

This section follows Reif sections 3.1 and 5.11.

Heat engines do not include electrical and chemical engines (such as muscles). The Carnot theorems don't apply to them. The name Carnot is French, so it is pronounced "Car-no" rather than "Car-not".

Problems

3.6 (I*) **The Carnot cycle**
Describe the Carnot cycle. Always means quasistatic. Take this order:

1. Expand at constant $T = T_1$ from A to B, decreasing pressure. Absorb heat q_1 from high-temperature heat bath.
2. Expand even more, adibatically, from B to C, decreasing pressure still more.
3. Contract at constant $T = T_2$ from C to D, increasing pressure. Expel heat q_2 into low-temperature heat bath.
4. Contract even more, adibatically, increasing pressure still more, until system comes back to original state A.

3.7 (E) **The Carnot cycle**
(This problem is stolen from a GRE Physics test.)
The p-V diagram for a quasistatic Carnot cycle is shown in Reif page 189. Legs bc and da represent isotherms, while ab and cd represent adiabats. A system is carried through the cycle abcd, taking in heat q_1 from the hot reservoir at temperature T_1 and releasing heat q_2 to the cold reservoir at temperature T_2. Which of the following statements is false?

a. $q_1/T_1 = q_2/T_2$.
b. The entropy of the hot reservoir decreases.
c. The entropy of the system increases.
d. The work w is equal to the net heat absorbed, $q_1 - q_2$.
e. The efficiency of the cycle is independent of the working substance.

3.8 (D) **The liquifying Carnot cycle**
Suppose a quasistatic Carnot cycle is executed with a working substance of steam rather than ideal gas. Furthermore, suppose that the cycle straddles the liquid-vapor coexistence curve, so that when the working substance is at a high volume and low pressure it is steam, but when it is at a low volume and high pressure it is liquid water. Then the (p, V) diagram of the cycle resembles the following:

a. Explain the significance of the flat portions of the isotherms.

b. Of the four points A, B, C, and D, which represent liquid and which represent gas?

c. Below is a list of properties of the Carnot cycle executed with an ideal gas. Which of these properties remain true for our liquifying Carnot cycle?

   i. The efficiency is $1 - T_1/T_2$.

   ii. The adiabats BC and DA are described by $pV^\gamma =$ constant.

   iii. The work done is the area enclosed by the cycle.

   iv. More.

3.4 Multivariate Calculus

What is a section on multivariate calculus doing in a physics book... particularly a physics book which assumes that you already know multivariate calculus? The answer is that you went through your multivariate calculus course learning one topic after another, and there are some subtle topics that you covered early in the course that really couldn't be properly understood until you had covered other topics that came later in the course. (This is *not* the fault of your teacher in multivariate calculus, because the later topics could not be understood *at all* without an exposure to the earlier topics.) This section goes back and investigates five subtle points from multivariate calculus to make sure they don't trip you when they are applied to thermodynamics.

3.4.1 What is a partial derivative?

Given a function $f(x, y, z)$, what is the meaning of $\partial f/\partial y$? Many will answer that

$$\frac{\partial f}{\partial y} \text{ is the change of } f \text{ with } y \text{ while everything else is held constant.} \tag{3.9}$$

This answer is WRONG! If f changes, then f^2 changes, and $\sin(f)$ changes, and so forth, so it can't be that "everything else" is held constant. The proper answer is that

$$\frac{\partial f}{\partial y} \text{ is the change of } f \text{ with } y \text{ while all other variables are held constant.} \tag{3.10}$$

Thus it becomes essential to keep clear which quantities are variables and which are functions. This is not usually hard in the context of mathematics: the functions are f, g, and h while the variables are x, y, and z. But in the context of physics we use symbols like E, V, p, T, and N which suggest the quantities they represent, and it is easy to mix up the functions and the variables.

An illustration from geometry makes this point very well. Consider the set of all right circular cylinders. Each cylinder can be specified uniquely by the variables r, radius, and h, height. If you know r and h for a cylinder you can readily calculate any quantity of interest—such as the area of the top, $T(r, h)$, the area

of the side $S(r,h)$, and the volume $V(r,h)$—as shown on the left side of the table below. But this is not the only way to specify each cylinder uniquely. For example, if you know the height and the side area of the cylinder, you may readily calculate the radius and hence find our previous specification. Indeed, the specification through the variables S and h is just as good as the one through the variables r and h, as is shown on the right side of the table below. [There are many other possible specifications (e.g. r and S, or T and V) but these two sets will be sufficient to make our point.]

Describing a Cylinder

variables:		**variables:**	
radius	r	side area	S
height	h	height	h
functions:		**functions:**	
top area	$T(r,h) = \pi r^2$	radius	$r(S,h) = S/2\pi h$
side area	$S(r,h) = 2\pi r h$	top area	$T(S,h) = S^2/4\pi h^2$
volume	$V(r,h) = \pi r^2 h$	volume	$V(S,h) = S^2/4\pi h$

All of this is quite straightforward and ordinary. But now we ask one more question concerning the geometry of cylinders, namely "How does the volume change with height?" The last line of the table presents two formulas for volume, so taking appropriate derivative gives us either

$$\frac{\partial V}{\partial h} = \pi r^2 = T \quad \text{or} \quad \frac{\partial V}{\partial h} = -S^2/4\pi h^2 = -T. \tag{3.11}$$

What? Is $\partial V/\partial h$ equal to T or to $-T$? It can't be equal to both!

The problem with equation (3.11) is that we were careless about specifying the variables. The two expressions for volume,

$$V(r,h) = \pi r^2 h \quad \text{and} \quad V(S,h) = S^2/4\pi h, \tag{3.12}$$

are in fact two completely different functions, with completely different variables, so we should *expect* completely different derivatives. If we increase the height h keeping the radius r fixed, the figure on the left below makes it clear that the volume increases. But if we increase the height and keep the side area S fixed, the radius will have to decrease as shown on the right below. The change on the right adds to the volume at the top of the cylinder but subtracts from the volume all along the sides. It would be most remarkable if the two volume changes were equal and, as our derivatives have shown, they are not.

A mathematician would say that we got ourselves in trouble in equation (3.11) because we gave two different functions the same name. A physicist would reply that they both represent the volume, so they deserve the same name. Rather than get into an argument, it is best to write out the variables of all functions explicitly, thus rewriting (3.11) as

$$\frac{\partial V(r,h)}{\partial h} = \pi r^2 = T(r,h) \quad \text{or} \quad \frac{\partial V(S,h)}{\partial h} = -S^2/4\pi h^2 = -T(S,h). \tag{3.13}$$

(Physicists often neglect to write out the full list of variables, which saves some time and some ink but which invites error. R.H. Price and J.D. Romano (*Am. J. Phys.* **66** (1998) 114) expose a situation in which a physicist published a deep error, which he made by neglecting to write out an explicit variable list.)

It becomes tiresome to write out the entire argument list for every function, so a shorthand notation has been developed. A right hand parenthesis is written after the partial derivative, and the functional arguments that are not being differentiated are listed as subscripts to that parenthesis. Thus the derivatives above are written as

$$\frac{\partial V(r,h)}{\partial h} = \left.\frac{\partial V}{\partial h}\right)_r \quad \text{and} \quad \frac{\partial V(S,h)}{\partial h} = \left.\frac{\partial V}{\partial h}\right)_S . \tag{3.14}$$

The expression on the left is read "the partial derivative of V with respect to h while r is held constant".

3.4.2 The Legendre transformation

Let us return to a description of cylinders in terms of the variables r and h. Clearly, one of the functions of interest is the volume

$$V(r,h). \tag{3.15}$$

A glance at the table on page 58 (or a moment's thought about geometry) shows that the total differential of V as a function of r and h is

$$dV = S(r,h)\,dr + T(r,h)\,dh, \tag{3.16}$$

whence

$$S(r,h) = \left.\frac{\partial V}{\partial r}\right)_h \quad \text{and} \quad T(r,h) = \left.\frac{\partial V}{\partial h}\right)_r . \tag{3.17}$$

Thus knowledge of the function $V(r,h)$ gives us a bonus... if we know $V(r,h)$, then we can take simple derivatives to find the other quantities of interest concerning cylinders, namely $S(r,h)$ and $T(r,h)$. Because

of the central importance of $V(r,h)$, it is called a "master function" and the total differential (3.16) is called a "master equation".

Is there any way to find a similarly convenient "master description" in terms of the variables S and h? Indeed there is, and it is given by the "Legendre transformation". In the Legendre transformation from the variables r and h to the variables S and h, we change the focus of our attention from the master function $V(r,h)$ to the function

$$\Phi(S,h) = V(r(S,h),h) - Sr(S,h). \tag{3.18}$$

(The above equation is written out in full with all arguments shown. It is more usually seen as

$$\Phi = V - Sr, \tag{3.19}$$

although this form raises the possibility that variables and functions will become mixed up.) The total differential of Φ is

$$\begin{aligned} d\Phi &= dV - S\,dr - r\,dS && (3.20)\\ &= S\,dr + T\,dh - S\,dr - r\,dS && (3.21)\\ &= -r\,dS + T\,dh. && (3.22) \end{aligned}$$

We have found a new master function! It is

$$\Phi(S,h), \tag{3.23}$$

and the new master equation is

$$d\Phi = -r(S,h)\,dS + T(S,h)\,dh, \tag{3.24}$$

giving rise immediately to

$$r(S,h) = -\left.\frac{\partial\Phi}{\partial S}\right)_h \quad \text{and} \quad T(S,h) = \left.\frac{\partial\Phi}{\partial h}\right)_S. \tag{3.25}$$

This description has all the characteristics of a master description: once the master function is known, all the other interesting functions can be found by taking straightforward derivatives.

3.4.3 Maxwell relations

Suppose

$$df = A(x,y)\,dx + B(x,y)\,dy. \tag{3.26}$$

Then

$$A(x,y) = \left.\frac{\partial f}{\partial x}\right)_y \quad \text{and} \quad B(x,y) = \left.\frac{\partial f}{\partial y}\right)_x.$$

But because

$$\frac{\partial^2 f(x,y)}{\partial x \partial y} = \frac{\partial^2 f(x,y)}{\partial y \partial x}$$

it follows that

$$\left.\frac{\partial A}{\partial y}\right)_x = \left.\frac{\partial B}{\partial x}\right)_y . \tag{3.27}$$

This is called a "Maxwell relation".

Applied to equation (3.16), this tells us at a glance that

$$\left.\frac{\partial S}{\partial h}\right)_r = \left.\frac{\partial T}{\partial r}\right)_h . \tag{3.28}$$

We know that these two derivatives are equal without needing to find either one of them! Applied to equation (3.24), it tells us with equal ease that

$$\left.\frac{\partial r}{\partial h}\right)_S = -\left.\frac{\partial T}{\partial S}\right)_h . \tag{3.29}$$

3.4.4 Implicit function theorem

Suppose $f(x, y)$ is a function of the variables x and y. What is

$$\left.\frac{\partial y}{\partial x}\right)_f , \tag{3.30}$$

the slope of a contour of constant f?

Start with

$$df = \left.\frac{\partial f}{\partial x}\right)_y dx + \left.\frac{\partial f}{\partial y}\right)_x dy. \tag{3.31}$$

which holds for any differential change dx and dy. But we're not interested in *any* differential change: to evaluate the slope (3.30), we need a change in which $df = 0$ so

$$0 = \left.\frac{\partial f}{\partial x}\right)_y dx + \left.\frac{\partial f}{\partial y}\right)_x dy \qquad \text{with } dx,\, dy \text{ on contour of } f.$$

Thus

$$\frac{dy}{dx} = -\frac{\left.\frac{\partial f}{\partial x}\right)_y}{\left.\frac{\partial f}{\partial y}\right)_x} \qquad \text{with } dx,\, dy \text{ on contour of } f$$

and, writing the restriction "with dx, dy on contour of f" into the symbols of the equation,

$$\left.\frac{\partial y}{\partial x}\right)_f = -\frac{\left.\frac{\partial f}{\partial x}\right)_y}{\left.\frac{\partial f}{\partial y}\right)_x}. \tag{3.32}$$

Note that you get the wrong answer if you "cancel the small quantity ∂f from numerator and denominator of the ratio." That's because "the small quantity ∂f with constant y" is *different* from "the small quantity ∂f with constant x". I need to write a few words and a figure concerning why.

3.4.5 Multivariate chain rule

Suppose $f(x, y)$ and $g(x, y)$ are two function of the variables x and y. Then again we have

$$df = \left.\frac{\partial f}{\partial x}\right)_y dx + \left.\frac{\partial f}{\partial y}\right)_x dy \tag{3.33}$$

for any differential change dx and dy.

What if we are interested, not in any change, but in a change along a contour of constant g? Specifically, what if we need to find the change of f with x while moving on a contour of constant g? Then just take the exact differential above and apply it to a change dx, dy along the contour with constant g. Divide by the differential quantity dx:

$$\frac{df}{dx} = \left.\frac{\partial f}{\partial x}\right)_y \frac{dx}{dx} + \left.\frac{\partial f}{\partial y}\right)_x \frac{dy}{dx} = \left.\frac{\partial f}{\partial x}\right)_y + \left.\frac{\partial f}{\partial y}\right)_x \frac{dy}{dx} \qquad \text{with } dx,\ dy \text{ on contour of } g.$$

Now write the restriction "with dx, dy on contour of g" into the symbols of the equation to find

$$\left.\frac{\partial f}{\partial x}\right)_g = \left.\frac{\partial f}{\partial x}\right)_y + \left.\frac{\partial f}{\partial y}\right)_x \left.\frac{\partial y}{\partial x}\right)_g, \tag{3.34}$$

the multivariate chain rule.

Problems

3.9 **Partial derivatives in space**

A point on the plane can be specified either by the variables (x, y) or by the variables (x', y') where

$$\begin{aligned} x' &= +\cos(\theta)\, x + \sin(\theta)\, y, \\ y' &= -\sin(\theta)\, x + \cos(\theta)\, y. \end{aligned} \tag{3.35}$$

If $f(x, y)$ is some function of location on the plane, then write expressions for

$$\left.\frac{\partial f}{\partial x'}\right)_{y'}, \quad \left.\frac{\partial f}{\partial y'}\right)_{x'}, \quad \text{and} \quad \left.\frac{\partial f}{\partial x}\right)_{y'} \tag{3.36}$$

in terms of

$$\left.\frac{\partial f}{\partial x}\right)_{y} \quad \text{and} \quad \left.\frac{\partial f}{\partial y}\right)_{x}. \tag{3.37}$$

Interpret $\partial f/\partial x)_{y'}$ geometrically as a directional derivative. (That is, $\partial f/\partial x)_{y'}$ is the slope of f along which curve in the plane?) Given this interpretation, does it have the expected limits as $\theta \to 0$ and as $\theta \to \pi/2$?

3.10 Maxwell relations for a three-variable system

Suppose $\Phi(x, y, z)$ satisfies

$$d\Phi = A(x, y, z)\, dx + B(x, y, z)\, dy + C(x, y, z)\, dz. \tag{3.38}$$

State three Maxwell relations relating various first derivatives of $A(x, y, z)$, $B(x, y, z)$, and $C(x, y, z)$, and a fourth Maxwell relation relating various second derivatives of these functions.

3.11 The cylinder model with three variables

In the "three-variable cylinder model" the cylinders are described by height h, radius r, and density ρ. The master function is mass $M(h, r, \rho)$, and the master equation is

$$dM(h, r, \rho) = \rho S(h, r)\, dr + \rho T(r)\, dh + V(h, r)\, d\rho, \tag{3.39}$$

where $S(h, r)$ is the side area, $T(r)$ is the top area, and $V(h, r)$ is the volume. Perform a Legendre transformation to a description in terms of the variables h, S, and ρ using the new master function

$$\Phi(h, S, \rho) = M - \rho S r. \tag{3.40}$$

a. Write down the new master equation.

b. Write down the three first-order Maxwell relations and confirm their correctness using explicit formulas such as $M(h, S, \rho) = \rho S^2/(4\pi h)$.

c. Interpret $\Phi(h, S, \rho)$ physically.

3.12 Multivariate chain rule

Invent a problem going through the chain rule argument of section 3.8.1 with the cylinder model.

3.13 Contours of constant side area

Find contours of constant side area preparing for/using the technique of section 3.8.3.

3.5 Thermodynamic Quantities

By a thermodynamic "quantity" I mean either a variable or a function. We have already seen that for fluids we may regard the entropy as a function of energy, volume, and number, $S(E, V, N)$, or the energy as a function of entropy, volume, and number, $E(S, V, N)$: using the term "quantity" avoids prejudging the issue of whether entropy, for example, is to be treated as a variable or a function.

It is traditional to divide thermodynamic quantities into two classes: "extensive" and "intensive". Extensive quantities increase linearly with system size while intensive quantities are independent of system size. (You might ask, "Why can't a quantity depend on system size in some other fashion, such as quadratically?" The answer is that this is not impossible—for example, the radius of a spherical sample increses as the cube root of the system size—but it rarely happens in practice for quantities that concern us—see problem 3.18.) Thus extensive quantities are characteristic of the *sample* while intensive quantities are characteristic of the *substance*.

Examples of extensive quantities are energy, volume, number, entropy, and the magnetization $\mathbf{M}$ (i.e. the magnetic dipole moment of the entire sample):

$$E, \quad V, \quad N, \quad S, \quad \mathbf{M}. \tag{3.41}$$

Examples of intensive quantities are temperature, pressure, chemical potential, and the applied magnetic field $\mathbf{H}$:

$$T, \quad p, \quad \mu, \quad \mathbf{H}. \tag{3.42}$$

If one extensive quantities is divided by another extensive quantity, the quotent is an intensive quantity. The most frequently used quantities of this type are the *number densities*, such as

$$e = \frac{E}{N}, \quad s = \frac{S}{N}, \quad \text{and} \quad v = \frac{V}{N}. \tag{3.43}$$

There are also *volume densities* like

$$u = \frac{E}{V}, \quad \mathcal{S} = \frac{S}{V}, \quad \text{and} \quad \rho = \frac{N}{V}. \tag{3.44}$$

Quantities such as E/S are also intensive, but they are rarely used and don't have special names.

Note that in defining densities such as above, we are not making any assumptions about how the matter in the sample is arranged. For liquid water in equilibrium with ice (e.g. at atmospheric pressure and a temperature of 273 K) the water has a greater number density (and a greater entropy density) than the ice, but the number density of the system is still just the total number divided by the total volume.

Here is a valuable problem solving tip. You know from previous courses that it is important to keep track of the dimensions of physical quantities: if you derive an equation for which the left side has the dimensions of meters and the right side has the dimensions of meters/second, your derivation must be wrong. In thermodynamics there is the equally important check of extensivity: if the left side is extensive and the right side is intensive, then the equation must be wrong. You should keep a running check of these properties

as you solve problems. The first time you produce an equation that is incorrect in its dimensions or its extensivity, go back immediately and correct your error.

Problem 1.2 has already introduced the "isothermal compressibility"

$$\kappa_T(p,T) = -\frac{1}{V}\frac{\partial V(p,T)}{\partial p} \tag{3.45}$$

and the "expansion coefficient"

$$\beta(p,T) = \frac{1}{V}\frac{\partial V(p,T)}{\partial T}. \tag{3.46}$$

Both of these quantities are defined so as to be intensive.

Another quantity of interest is the "heat capacity", defined informally as "the amount of heat required to quasistatically raise the temperature of a sample by one kelvin". (The heat capacity was introduced briefly in problem 1.3.) Because this definition refers to a *sample*, heat capacity is an extensive quantity. The definition is incomplete because it doesn't say what path is taken while the heat is added, and we know that heat is a path-dependent quantity.[1] For fluids, the two most frequently used heat capacities are the heat capacity at constant volume and at constant pressure, denoted by

$$C_V \quad \text{and} \quad C_p. \tag{3.47}$$

Thus C_p (which will in general be a function of temperature, pressure, and number) is the amount of heat required to increase the temperature of a sample by one kelvin while the sample is under a constant pressure (often, in earth-bound laboratories, a pressure of one atmosphere). In contrast C_V (which will in general be a function of temperature, volume, and number) is the amount of heat required to increase the temperature of a sample by one kelvin while holding it in a strong box to prevent it from expanding. Because liquids expand little upon heating, C_V and C_p are nearly equal for liquids. But the two quantities differ dramatically for gases.

For small quasistatic changes, we know that

$$\Delta S = \frac{Q}{T} \tag{3.48}$$

whence

$$C = \lim_{\Delta T \to 0} \frac{T\Delta S}{\Delta T}. \tag{3.49}$$

This gives us the formal definitions of heat capacity,

$$C_V = T\frac{\partial S(T,V,N)}{\partial T} \quad \text{and} \quad C_p = T\frac{\partial S(T,p,N)}{\partial T}, \tag{3.50}$$

and from these equations it is again clear that heat capacity is an extensive quantity.

The intensive quantities analogous to heat capacities are called "specific heats":

$$c_V = \frac{C_V}{N} \quad \text{and} \quad c_p = \frac{C_p}{N}. \tag{3.51}$$

The specific heat is readily measured experimentally and we will come back to it again and again.

[1] In truth, it's worse than this, because it doesn't even specify the end point of the process. If we are describing the fluid sample by V and T, for example, and the starting point were (V_i, T_i), then the end point must have $T_f = T_i + 1$ K, but as far as our informal definition goes, the final volume could be anything.

Problems

3.14 (E) **Which one was it?**

For a system with a fixed number of particles, the reciprocal of the absolute (kelvin) temperature T is given by which of the following derivatives? Explain your reasoning. Do not look up the answer or, if you have memorized it, don't use that piece of your memory. (Clue: Dimensional analysis will work, but there's an even easier way.)

$$\frac{\partial p(S,V)}{\partial V} \quad \frac{\partial p(S,V)}{\partial S} \quad \frac{\partial S(E,p)}{\partial p} \quad \frac{\partial V(E,p)}{\partial p} \quad \frac{\partial S(E,V)}{\partial E} \tag{3.52}$$

3.15 (I) **Acceptable forms for the entropy**

Below are seven equations purporting to be equations for $S(E,V,N)$ for various thermodynamic systems. However, four of them are not physically acceptable. Find the four inpermissible equations and for each indicate why it cannot be correct. The quantities T_0 and v_0 are in all cases positive, intensive quantities with the dimensions of temperature and volume, respectively.

$$S(E,V,N) = \left(\frac{k_B^2}{v_0 T_0}\right)^{1/3} (EVN)^{1/3} \tag{3.53}$$

$$S(E,V,N) = \left(\frac{k_B v_0^2}{T_0^2}\right)^{1/3} \left(\frac{EN}{V}\right)^{2/3} \tag{3.54}$$

$$S(E,V,N) = \left(\frac{k_B v_0^2}{T_0^2}\right)^{1/3} \frac{EN}{V} \tag{3.55}$$

$$S(E,V,N) = \left(\frac{k_B}{T_0}\right)^{1/2} \left(NE + \frac{k_B T_0 V^2}{v_0^2}\right)^{1/2} \tag{3.56}$$

$$S(E,V,N) = k_B N \ln(EV/N^2 k_B T_0 v_0) \tag{3.57}$$

$$S(E,V,N) = k_B N \exp(-EV/N^2 k_B v_0) \tag{3.58}$$

$$S(E,V,N) = k_B N \exp(-EV^2/N^2 k_B T_0 v_0^2) \tag{3.59}$$

3.16 (I) **Heat capacities for the ideal gas**

Use equation (2.32) to find the heat capacities

$$C_V(T,V,N) = T \left.\frac{\partial S}{\partial T}\right)_{V,N} \qquad \text{and} \qquad C_p(T,p,N) = T \left.\frac{\partial S}{\partial T}\right)_{p,N}$$

for the pure classical monatomic ideal gas.

3.17 (I) **Heat capacity at a phase transition**

What is C_p for a fluid when the liquid and gas coexist?

3.18 (D) **Gravitational potential energy**

Consider a uniform sphere of mass M and radius R.

a. Use dimensional analysis to show that the gravitational potential energy of the sphere is of the form

$$U_G = -cG\frac{M^2}{R}, \tag{3.60}$$

where c is a dimensionless constant independent of M and R.

b. Is U_G intensive, extensive, or neither?

c. Is the total energy E (which is equal to U_G plus contributions from other forms of energy) intensive, extensive, or neither?

d. Estimate the value of U_G for a glass of water.

You should conclude from this problem that the total energy E is in principle non-extensive, but that the deviations from being extensive are negligible for everyday thermodynamic systems.

e. (Optional and quite time consuming.) Use ideas from classical mechanics to show that the dimensionless constant in equation (3.60) is $c = 3/5$.

3.6 The Thermodynamic Dance

A more conventional name for this section would be "Changes of Variable in Thermodynamics", but the changes of variable involved always remind me of an elaborate dance where partners are exchanged and where patterns seem to dissolve and then reappear, but in fact are always present, even if hidden.

3.6.1 Description in terms of variables (S, V, N)

We know that in many circumstances (e.g. pure, non-magnetic fluids) the thermodynamic state of a system is uniquely specified by giving the entropy S, the volume V, and the particle number N. The master function for this description is the energy

$$E(S, V, N) \tag{3.61}$$

whose total differential is

$$dE = T\,dS - p\,dV + \mu\,dN. \tag{3.62}$$

Remember that a "master function" is one from which all functions of interest can be obtained by simple differentiation... the equation above, for example, shows directly that

$$p(S, V, N) = -\left.\frac{\partial E}{\partial V}\right)_{S,N}. \tag{3.63}$$

Remember also that the variables V and N are just representative mechanical parameters. Other mechanical parameters such as particle mass m, particle radius r_{HS}, or the numbers of two different species, N_{H_2} and N_{He}, will be relevant in other circumstances.

3.6.2 Description in terms of variables (T, V, N)

For many purposes the description in terms of S, V, and N is awkward. (Entropy, for example, can be difficult to measure and to understand.) In these situations a more natural set of variables might be temperature T, volume V, and number N. We can change[2] to this set through a Legendre transformation (a process sometimes called "trading in an S for a T") obtaining the new master function

$$F(T, V, N) = E - TS, \tag{3.64}$$

which is called the "Helmholtz potential" (or the "Helmholtz free energy"[3]). and which has the associated master equation

$$dF = -S\,dT - p\,dV + \mu\,dN. \tag{3.65}$$

(In some books, the Helmholtz potential is denoted by A rather than F.)

Lots of information can be read directly from the master equation. For example, the entropy can be found through

$$S(T, V, N) = -\left.\frac{\partial F}{\partial T}\right)_{V,N}. \tag{3.66}$$

It is less obvious how to find the energy, but a little thought shows that

$$E(T, V, N) = F + TS = F - T\left.\frac{\partial F}{\partial T}\right)_{V,N} = \left.\frac{\partial (F/T)}{\partial (1/T)}\right)_{V,N}. \tag{3.67}$$

This equation is frequently used and carries the name "Gibbs-Helmholtz equation".

> One sometimes hears that "energy must be regarded as a function of (S, V, N) while Helmholtz potential must be regarded as a function of (T, V, N)". This is false. The energy $E(T, V, N)$ above is a perfectly good function but it is *not* a master function. The energy is the master function for the description in terms of (S, V, N) but *not* for the description in terms of (T, V, N). Similarly, one could find the Helmholtz potential as a function of (S, V, N), but it would not be a master function.

Another immediate consequence of the master equation is that

$$p(T, V, N) = -\left.\frac{\partial F}{\partial V}\right)_{T,N}. \tag{3.68}$$

This equation is *extremely* well know... it is just the "equation of state", relating pressure, temperature, volume, and number. Examples are the famous ideal gas equation of state

$$p(T, V, N) = \frac{N k_B T}{V} \tag{3.69}$$

[2]When we changed variable from (S, V, N) to (E, V, N) (equation 3.1), I argued that this inversion was mathematically legitimate. Here I will not be so formal and just assume that everything will come out all right. (See, however, problem 3.27.) Clifford Truesdell has called this "the principle of thoughtless invertibility".

[3]The term "free" in this name is of only historical significance. Its origin lies in the fact that F is related to the maximum amount of energy that is available ("free") for conversion into work through an isothermal process.

and the only slightly less famous van der Waals equation of state for non-ideal gases

$$p(T,V,N) = \frac{k_B T}{V/N - v_0} - \frac{e_0 v_0}{(V/N)^2}, \tag{3.70}$$

where v_0 and e_0 are positive empirical constants with the dimensions of volume and energy respectively.

Notice that $S(T,V,N)$, $E(T,V,N)$, and $p(T,V,N)$ are all interesting quantities, but none of them are master functions, that is, none of them contain *all* the thermodynamic information. It is a common misconception that the equation of state contains everything one could want to know about a substance. In fact, for example, helium and nitrogen are both nearly ideal gases at room temperature (so they obey nearly the same equation of state) but the specific heat of nitrogen is about 67% greater than that of helium (so they differ dramatically in some thermodynamic properties).

Yet another immediate consequence of the master equation is the Maxwell relation

$$\left.\frac{\partial S}{\partial V}\right)_{T,N} = \left.\frac{\partial p}{\partial T}\right)_{V,N}. \tag{3.71}$$

Each of the derivatives above is not only a mathematical expression, but also an invitation to perform an experiment. The derivative on the left is measured through an experiment like the following: A sample in a container of variable volume (such as a piston) is placed within a thermostatically controlled bath (so that the temperature doesn't change) and is heated in a slow and carefully monitored way (so that the heat absorbed quasistatically can be divided by the temperature to find the entropy change). As the substance is heated at constant temperature, the volume of the piston must change. Dividing the heat absorbed by the temperature and the volume change gives (for small volume changes) the derivative on the left. This experiment is not impossible, but clearly it is difficult and expensive.

Consider in turn the experiment on the right. The sample is in a "strong box" container of fixed volume and its pressure and temperature are measured as the temperature is changed. The change doesn't need to be controlled carefully and the heat absorbed doesn't need to be monitored: you can just blast your sample with a propane torch. Dividing the measured change in pressure by the measured change in temperature gives (for small temperature changes) the derivative on the right. It is remarkable that the results of these two very different experiments is always the same... and it shows how thermodynamics can save a lot of experimental labor! But it is still more remarkable—nearly unbelievable—that we know these two experiments give the same results because we know that the entropy function $S(E,V,N)$ exists, even though we might not know what that function is.[4] (This Maxwell relation is discussed further in problem 3.23.)

3.6.3 Description in terms of variables (S,p,N)

I don't need to stop with this second description. Both of the descriptions above used volume as a variable. Most experiments executed here on Earth's surface are actually performed with a constant pressure (namely

[4] If a material existed for which these two experiments *did not* give identical results, then we could use that substance to build a perpetual motion machine.

one atmosphere) rather than with a constant volume, suggesting that pressure should be one of our variables. We may start with the description in terms of (S, V, N) and then trade in a V for a p, obtaining a master function

$$H(S, p, N) = E + pV, \tag{3.72}$$

called the "enthalpy", and the related master equation

$$dH = T\,dS + V\,dp + \mu\,dN. \tag{3.73}$$

We could pause here to write down derivatives of $H(S, p, N)$, Maxwell relations, and so forth, but this would merely be repetitive of what we did in the last section. Instead, we ask for yet another description with the advantages of using both temperature instead of entropy and pressure instead of volume.

3.6.4 Description in terms of variables (T, p, N)

In this description, the master function

$$G(T, p, N) = F + pV \tag{3.74}$$

is called the "Gibbs potential" (or the "Gibbs free energy"[5]) and the master equation is

$$dG = -S\,dT + V\,dp + \mu\,dN. \tag{3.75}$$

(It is most unfortunate that in some books the Gibbs potential is denoted F, the symbol that most books reserve for the Helmholtz potential.)

It is immediately obvious that

$$\mu(T, p, N) = \left.\frac{\partial G}{\partial N}\right)_{T,p}. \tag{3.76}$$

It is also true that G is an extensive quantity, and that it is a function of *only one* extensive quantity, namely N. It follows that G must increase linearly with N, whence

$$\mu(T, p) = \frac{G(T, p, N)}{N}. \tag{3.77}$$

Notice that μ, which appeared from equation (3.76) to depend upon N, is actually independent of N.

The chemical potential has so far been a wallflower in the thermodynamic dance. But there is no reason why we cannot trade in an N for a μ in the same way that we traded in a V for a p.

[5] I prefer the name "Gibbs potential" because it is like electrostatic potential — the quantity itself is not of intrinsic interest, but one can find many interesting things by taking its derivatives. The name "free energy" is particularly inappropriate: What do you buy from a power company? You've been told "Turn off the lamps to conserve energy", but this is silly: energy is conserved whether the lamps are on or off. There's plenty of thermal energy in any room, but you can't get work out of that energy because the room has uniform temperature. So you don't buy energy from a power company, you in fact buy Gibbs potential. For this reason it should be called "Gibbs expensive energy". I am trying to get the federal government to change the name of the Department of Energy to the Department of Gibbs Potential, but so far I've had no success.

3.6.5 Description in terms of variables (T, V, μ)

Here the master function is

$$\Pi(T, V, \mu) = F - \mu N \tag{3.78}$$

with master equation

$$d\Pi = -S\,dT - p\,dV - N\,d\mu. \tag{3.79}$$

You might wonder why Π doesn't have a name. This is because

$$\Pi = F - \mu N = F - G = -pV \tag{3.80}$$

or, to put it more formally,

$$\Pi(T, V, \mu) = -p(T, \mu)V, \tag{3.81}$$

so Π already has the name of "negative pressure times volume".

3.6.6 Intensive description in terms of variables (T, μ)

Putting equations (3.79) and (3.81) together gives

$$-p\,dV - V\,dp = -S\,dT - p\,dV - N\,d\mu \tag{3.82}$$

whence

$$dp = \frac{S}{V}\,dT + \frac{N}{V}\,d\mu. \tag{3.83}$$

In other words, we have produced a new master function, the pressure

$$p(T, \mu) \tag{3.84}$$

with master equation

$$dp = \mathcal{S}\,dT + \rho\,d\mu \tag{3.85}$$

where we have used the volume densities

$$\mathcal{S} = S/V \quad \text{and} \quad \rho = N/V. \tag{3.86}$$

The master function $p(T, \mu)$ differs in important ways from the others that we have seen. First, it is intensive rather than extensive. Second, it is a function of two rather than three variables, and the two variables are both intensive. It is clear that we cannot get any information about the system size out of $p(T, \mu)$... thus it fails the usual test of a master function, namely that it must provide *all* thermodynamic information. On the other hand, $p(T, \mu)$ provides all the information about intensive quantities. The problems in this chapter demonstrate that $p(T, \mu)$ is a surprisingly useful master function.

This entire section is summarized in appendix J.

Problems

3.19 Stumbling in the thermodynamic dance

a. From the thermodynamic assembly with variables T, p, and N, with master equation

$$dG = -S\,dT + V\,dp + \mu\,dN \tag{3.87}$$

one is tempted to produce an assembly with variables T, p, and μ, using master function

$$\Phi = G - \mu N. \tag{3.88}$$

Why does this fail?

b. Show that what is really desired is an intensive-only description in terms of the variables T and p, for which the master equation is

$$d\mu = -s\,dT + v\,dp, \tag{3.89}$$

where $s = S/N$ and $v = V/N$. (This result is called the "Gibbs-Duhem equation".)

3.20 Not a master function

The function $E(S, V, N)$ is a master function: all thermodynamic quantities of interest can be obtained by taking suitable derivatives of $E(S, V, N)$. For the variables (T, V, N) the master function is the Helmholtz potential $F(T, V, N)$. While the energy $E(T, V, N)$ of course remains a function, it is no longer a master function. Prove this to yourself by considering two different substances, A and B, with Helmholtz potentials $F_A(T, V, N)$ and $F_B(T, V, N) = F_A(T, V, N) + aTV^2/N$. Show that these two substances have identical energies $E(T, V, N)$ but different equations of state $p(T, V, N)$.

3.21 Thermodynamics of a new substance

The entropy of a newly discovered gas is determined to be

$$S(E, V, N) = \left(\frac{k_B}{T_0}\right)^{1/2} \left(NE + \frac{k_B T_0 V^2}{v_0^2}\right)^{1/2},$$

where the constants T_0 and v_0 are positive, intensive quantities with the dimensions of temperature and volume, respectively.

a. Verify that this equation is acceptable in its dimensions and in its extensivity.

b. Find $T(S, V, N)$ and $p(S, V, N)$ for this gas.

c. Find $C_V(T, V, N)$ for this gas.

3.22 **Energy from the Gibbs potential**

The text claims that "any thermodynamic function can be obtained from a master function by taking derivatives." To back up this claim, show that

$$E(T,p,N) = G(T,p,N) - T\left.\frac{\partial G}{\partial T}\right)_{p,N} - p\left.\frac{\partial G}{\partial p}\right)_{T,N} \tag{3.90}$$

$$= \left.\frac{\partial(G/p)}{\partial(1/p)}\right)_{T,N} - T\left.\frac{\partial G}{\partial T}\right)_{p,N} \tag{3.91}$$

$$= \left.\frac{\partial(G/T)}{\partial(1/T)}\right)_{p,N} - p\left.\frac{\partial G}{\partial p}\right)_{T,N}. \tag{3.92}$$

Find an expression for $F(T,p,N)$ in terms of $G(T,p,N)$.

3.23 **A Maxwell relation: special cases**

a. Apply the Maxwell relation (3.71) to the the special case of an ideal gas (equation of state $pV = Nk_BT$) to show that

$$S(T,V,N) = Nk_B\ \ln(V/V_0(T)), \tag{3.93}$$

where $V_0(T)$ is an undetermined function of integration that differs from one ideal gas to another. (The remarkable character of our Maxwell relation comes into sharp focus when applied to this special case: The "mechanical-type" experiments which uncover the equation of state enable us to determine much about the entropy function even in the absence of any "heat-type" experiments.)

b. Another special case worth examination is two-phase coexistence of liquid and gas. Verify the Maxwell relation both for "typical" two-phase coexistence—in which the low-temperature phase has higher density than the high-temperature phase—and for the unusual cases—such as the coexistence of water and ice—where the low-temperature phase has a higher density.

3.24 **Chemical potential for mixtures**

If a system contains a mixture of two chemical substances, say N_A molecules of substance A and N_B molecules of substance B, then the list of mechanical parameters must be expanded, and the entropy is a function

$$S(E,V,N_A,N_B). \tag{3.94}$$

In this case the chemical potential of substance A is

$$\mu_A(E,V,N_A,N_B) = -T(E,V,N_A,N_B)\left.\frac{\partial S}{\partial N_A}\right)_{E,V,N_B} \tag{3.95}$$

and similarly for B. Notice that the chemical potential of substance A depends upon the number of molecules of B present.

a. Perform the thermodynamic dance down to the function $F(T,V,N_A,N_B)$ to show that

$$\mu_A(T,V,N_A,N_B) = \left.\frac{\partial F}{\partial N_A}\right)_{T,V,N_B} \tag{3.96}$$

and similarly for B.

b. Argue that, because μ_A and μ_B are intensive, their functional dependence on V, N_A, and N_B, must be through the number densities $\rho_A = N_A/V$ and $\rho_B = N_B/V$, i.e. that

$$\mu_A(T, V, N_A, N_B) = \mu_A(T, \rho_A, \rho_B). \tag{3.97}$$

c. Consider the Gibbs potential $G(T, p, N_A, N_B)$ and modify the extensivity argument that produced equation (3.77) to show that

$$G(T, p, N_A, N_B) = \mu_A(T, p, N_A/N_B)N_A + \mu_B(T, p, N_A/N_B)N_B, \tag{3.98}$$

where the chemical potentials are independent of the overall numbers N_A and N_B but may depend on their ratio.

3.25 **More parameters**
What is the generalization of

$$dE = T\,dS - p\,dV + \mu\,dN \tag{3.99}$$

for systems that can be lifted and tossed? (That is, for systems in which height h and velocity $\mathbf{v}$ are mechanical parameters.)

3.26 **Another mechanical parameter**
Recall expression (2.32) for the entropy of a monatomic ideal gas, and recall that $E = \frac{3}{2}Nk_BT$. Find expressions for the change in entropy and in volume if the masses of the particles are varied at constant p, T, and N. Suggest experiments that could verify these relations.

3.27 **Which variables to exchange?**
In carrying out the thermodynamic dance, we have exchanged the variable S for the function $T(S, V, N)$, and we have exchanged the variable V for the function $p(T, V, N)$, and more. Why would it never be appropriate to exchange the volume for the temperature? (Clue: Consider the volume of water at temperatures just above and below freezing.)

3.7 Non-fluid Systems

All the results of the previous section relied on the underlying assumption that the system was a pure fluid and thus could be described by specifying, for example, the temperature T, the volume V, and the particle number N. Recall that the latter two were simply examples of many possible mechanical parameters that could be listed, such as the molecular mass, the molecular moment of inertia, or pair interaction parameters such as the hard-sphere radius or the Lennard-Jones parameters.

But of course nature presents us with many materials that are not pure fluids! One obvious example is a fluid mixture, the specification of which requires the number of molecules of each constituent. Another example is a crystal of a layered material such as graphite. To find the energy, for example, it is not sufficient

to specify only the volume. You must know the area of the layers and the height to which these layers are stacked.

In this book we will focus instead on a third example, namely magnets. The fundamental thermodynamic relation for magnetic systems is

$$dE = T\,dS - M\,dH, \tag{3.100}$$

where H, the applied magnetic field, is an intensive mechanical parameter and M, the magnetization (total magnetic dipole moment of the sample), is extensive. Just as the thermodynamic equations for fluids presented in the previous section implicitly assume that the magnetic properties of the sample can be ignored (either because the substance is non-magnetic or because the magnetic field does not change), so the equation above implicitly assumes that the volume and number specification of the sample can be ignored.

In another course, you may have learned a mnemonic for remembering the thermodynamic differentials and Maxwell relations of a pure fluid systems with a constant particle number. Such mnemonics encourage the very worst problem solving strategy, namely "poke around until you discover an equation that fits". Anyone who uses this strategy finds it impossible to investigate mixtures, crystals, magnets, or any other member of the rich array of materials that nature has so generously spread before us. Instead of memorizing equations and hoping that the right one will present itself, you should think about what sort of equation you will need to solve a problem and then derive it. Appendix J will remind you of the strategy of the thermodynamic dance and will help you keep your signs straight.

3.28 **Magnetic systems**

Show that for magnetic systems (see equation (3.100)),

$$\left.\frac{\partial M}{\partial T}\right)_H = \left.\frac{\partial S}{\partial H}\right)_T \tag{3.101}$$

and

$$\left.\frac{\partial H}{\partial T}\right)_M = -\left.\frac{\partial S}{\partial M}\right)_T. \tag{3.102}$$

3.8 Thermodynamics Applied to Fluids

In the rest of this chapter we apply the general results of section 3.6, "The Thermodynamic Dance", to particular concrete situations. We begin with fluids, that is, systems whose thermodynamic states are adequately described by giving the variables temperature T, volume V, and particle number N. Furthermore, in this section we will not allow particles to enter or leave our system, so only two variables, T and V, are needed.

3.8.1 Heat capacities

Is there any relation between

$$C_p(T,p) = T\left.\frac{\partial S}{\partial T}\right)_p, \tag{3.103}$$

the heat capacity at constant pressure, and

$$C_V(T,V) = T\left.\frac{\partial S}{\partial T}\right)_V, \tag{3.104}$$

the heat capacity at constant volume? Remembering that entirely different experiments are used to measure the two quantities, you might guess that there is not. But mathematically, the difference between C_p and C_V is related to a change in variable from (T,V) to (T,p), so you might begin to suspect a relation. [You might want to prepare for this section by working problem 3.12.]

Begin with the mathematical definition of the total differential of entropy,

$$dS = \left.\frac{\partial S}{\partial T}\right)_V dT + \left.\frac{\partial S}{\partial V}\right)_T dV. \tag{3.105}$$

The above holds for any infinitesimal change. We restrict ourselves to infinitesimal changes at constant pressure, and divide by dT, to find

$$\left.\frac{\partial S}{\partial T}\right)_p = \left.\frac{\partial S}{\partial T}\right)_V + \left.\frac{\partial S}{\partial V}\right)_T \left.\frac{\partial V}{\partial T}\right)_p. \tag{3.106}$$

Multiplying both sides by T gives a heat capacity relation

$$C_p = C_V + T\left.\frac{\partial S}{\partial V}\right)_T \left.\frac{\partial V}{\partial T}\right)_p. \tag{3.107}$$

This relationship is correct but not yet written in its most convenient form. For example, reference books provide tabulations of C_p and C_V, but not of $\partial S/\partial V)_T$ or $\partial V/\partial T)_p$.

The first step in changing the heat capacity relation into "standard form" is an easy one. Recall from problem 1.2 that the expansion coefficient (an intensive tabulated quantity[6]) is defined by

$$\beta(T,p) = \frac{1}{V}\left.\frac{\partial V}{\partial T}\right)_p \tag{3.108}$$

whence

$$C_p = C_V + TV\beta\left.\frac{\partial S}{\partial V}\right)_T. \tag{3.109}$$

The second step is less obvious. At equation (3.71) I discussed the Maxwell relation

$$\left.\frac{\partial S}{\partial V}\right)_T = \left.\frac{\partial p}{\partial T}\right)_V, \tag{3.110}$$

and mentioned that the right hand side was much easier to measure. Thus we have the preferable expression

$$C_p = C_V + TV\beta\left.\frac{\partial p}{\partial T}\right)_V. \tag{3.111}$$

This expression is still not very convenient, however, because the quantity $\partial p/\partial T)_V$ is neither named nor tabulated. Although easier to measure than $\partial S/\partial V)_T$ is, its measurement still requires a constant-volume

[6] It is easy to see why $\partial V/\partial T)_p$ itself is not tabulated. Instead of requiring a tabulation for iron, you would need a tabulation for a five gram sample of iron, for a six gram sample of iron, for a seven gram sample of iron, etc.

strong box. Measurements at constant pressure are easier to perform and hence more frequently tabulated. We now write $\partial p/\partial T)_V$ in terms of such quantities.

The total differential of $V(T,p)$ is

$$dV = \left.\frac{\partial V}{\partial T}\right)_p dT + \left.\frac{\partial V}{\partial p}\right)_T dp \tag{3.112}$$

or, in terms of named quantities,

$$dV = V\beta\, dT - V\kappa_T\, dp. \tag{3.113}$$

(The isothermal compressibility

$$\kappa_T(T,V) \equiv -\frac{1}{V}\left.\frac{\partial V}{\partial p}\right)_T \tag{3.114}$$

was defined in problem 1.2.) Restricting the above total differential to changes at constant V gives

$$V\beta\, dT = V\kappa_T\, dp \tag{3.115}$$

or

$$\frac{\beta}{\kappa_T} = \left.\frac{\partial p}{\partial T}\right)_V. \tag{3.116}$$

This immediately gives us the final form for our relationship between heat capacities,

$$C_p = C_V + TV\frac{\beta^2}{\kappa_T}. \tag{3.117}$$

This is a far from obvious result that is true for all fluids and, I emphasize, was derived assuming only that entropy exists and without the use of any explicit formula for the entropy.

Note that T, V, and κ_T are all positive quantities. The expansion coefficient β may be either positive or negative, but it enters the relationship as β^2. Thus

$$C_p \geq C_V. \tag{3.118}$$

Can we understand these results physically? Heat capacity is the amount of heat required to raise the temperature of a sample by one kelvin. If the sample is heated at constant volume, all of the heat absorbed goes into increasing the temperature. But if the sample is heated at constant pressure, then generally the substance will expand as well as increase its temperature, so some of the heat absorbed goes into increasing the temperature and some is converted into expansion work. Thus we expect that more heat will be needed for a given change of temperature at constant pressure than at constant volume, i.e. that $C_p \geq C_V$. We also expect that the relation between C_p and C_V will depend upon the expansion coefficient β. It is hard, however, to turn these qualitative observations into the precise formula (3.117) without recourse to the formal mathematics that we have used. It is surprising, for example, that β enters into the formula as β^2, so that materials which contract with increasing temperature still have $C_p \geq C_V$.

For solids and liquids, there is little expansion and thus little expansion work, whence $C_p \approx C_V$. But for gases, $C_p \gg C_V$. What happens at two-phase coexistence?

So far we have discussed only the difference between the two heat capacities. It is conventional to also define the "heat capacity ratio"

$$\gamma \equiv \frac{C_p}{C_V}, \tag{3.119}$$

which can be regarded as a function of T and V or as a function of T and p. But the result above shows that for all values of T and V,

$$\gamma(T, V) \geq 1. \tag{3.120}$$

We can apply these results to the ideal gas with equation of state

$$pV = Nk_BT. \tag{3.121}$$

Problem 1.2 used this equation to show that for an ideal gas,

$$\beta = \frac{1}{T} \quad \text{and} \quad \kappa_T = \frac{1}{p}. \tag{3.122}$$

Thus the heat capacity relation (3.117) becomes

$$C_p = C_V + TV\frac{p}{T^2} = C_V + \frac{pV}{T} = C_V + k_BN, \tag{3.123}$$

and we have

$$\gamma = \frac{C_p}{C_V} = 1 + \frac{k_BN}{C_V}. \tag{3.124}$$

3.8.2 Energy as a function of temperature and volume

You know that $E(S, V)$ is a master function with the famous master equation

$$dE = T\,dS - p\,dV. \tag{3.125}$$

For the variables T and V the master function is the Helmholtz potential $F(T, V)$ with master equation

$$dF = -S\,dT - p\,dV. \tag{3.126}$$

Thus in terms of the variables T and V the energy in no longer a master function, but that doesn't mean that it's not a function at all. It is possible to consider the energy as a function of T and V, and in this subsection we will find the total differential of $E(T, V)$.

We begin by finding the total differential of $S(E, V)$ and finish off by substituting that expression for dS into the master equation (3.125). The mathematical expression for that total differential is

$$dS = \left.\frac{\partial S}{\partial T}\right)_V dT + \left.\frac{\partial S}{\partial V}\right)_T dV, \tag{3.127}$$

but we showed in the previous subsection that this is more conveniently written as

$$dS = \frac{C_V}{T}\,dT + \frac{\beta}{\kappa_T}\,dV. \tag{3.128}$$

Thus

$$dE = T\,dS - p\,dV = T\left[\frac{C_V}{T}\,dT + \frac{\beta}{\kappa_T}\,dV\right] - p\,dV \tag{3.129}$$

and, finally,

$$dE = C_V\, dT + \left[T\frac{\beta}{\kappa_T} - p\right] dV. \tag{3.130}$$

The most interesting consequence of this exercise is that the heat capacity C_V, which was defined in terms of an entropy derivative, is also equal to an energy derivative:

$$C_V(T,V) \equiv T\left.\frac{\partial S}{\partial T}\right)_V = \left.\frac{\partial E}{\partial T}\right)_V. \tag{3.131}$$

For an ideal gas

$$\left[T\frac{\beta}{\kappa_T} - p\right] = 0, \tag{3.132}$$

whence

$$dE = C_V\, dT + 0\, dV. \tag{3.133}$$

Thus the energy, which for most fluids depends upon both temperature and volume, depends in the ideal gas only upon temperature:

$$E(T,V) = E(T). \tag{3.134}$$

The same is true of the heat capacity

$$C_V(T,V) = \left.\frac{\partial E}{\partial T}\right)_V = C_V(T), \tag{3.135}$$

and the heat capacity ratio

$$\gamma(T) = 1 + \frac{k_B N}{C_V(T)}. \tag{3.136}$$

Thermodynamics proves that for an ideal gas C_V depends only upon T and not V, but experiment shows that in fact C_V is almost independent of T as well. From the point of view of thermodynamics this is just an unexpected experimental result and nothing deeper. From the point of view of statistical mechanics this is not just a coincidence but is something that can be proved... we will do so in chapter 5.

3.8.3 Quasistatic adiabatic processes

This topic, titled by a string of polysyllabic words, sounds like an arcane and abstract one. In fact it has direct applications to the speed of sound, the manufacture of refrigerators, to the inflation of soccer balls, and to the temperature of mountain tops. [You might want to prepare for this section by working problem 3.13.]

If a fluid is changed in a quasistatic, adiabatic manner, its entropy will remain constant during the change. So, for example, if the state of the system is described by the two variables temperature and volume, then during a quasistatic adiabatic process the system will move along contour lines[7] of constant entropy in the (T,V) plane, such as those below.

[7] Such contours are sometimes called isoentropic curves or, even worse, adiabats. The latter name is not just a linguistic horror, it also omits the essential qualifier that only *quasistatic* adiabatic processes move along curves of constant entropy.

Our aim in this section is to find an equation V(T) for such constant-entropy contours.

We begin with the now-familiar total differential for the entropy as a function of temperature and volume,

$$dS = \frac{C_V}{T}\,dT + \frac{\beta}{\kappa_T}\,dV. \tag{3.137}$$

This equation holds for any infinitesimal change, but we will apply it now to changes along an entropy contour, i.e. changes for which $dS = 0$. For such changes

$$\frac{\beta}{\kappa_T}\,dV = -\frac{C_V}{T}\,dT \tag{3.138}$$

whence the differential equation for an entropy contour is

$$\frac{dV(T)}{dT} = -\frac{C_V(T,V)\kappa_T(T,V)}{T\beta(T,V)}. \tag{3.139}$$

The quantities on the right are usually (not always) positive, so the entropy contours usually (not always) slope downward. This is as much as we can say for a general fluid.

But for an ideal gas we can fill in the functions on the right to find

$$\frac{dV(T)}{dT} = -\frac{C_V(T)}{p(T,V)} = -\frac{C_V(T)V}{Nk_BT}. \tag{3.140}$$

Using equation (3.136), this result is often written in the form

$$\frac{dV(T)}{dT} = \frac{1}{1-\gamma(T)}\frac{V}{T}. \tag{3.141}$$

This is as far as we can go for an arbitrary ideal gas.

But for an ideal gas with constant heat capacity, and hence with constant γ, this differential equation is readily solved using separation of variables. We write

$$(1-\gamma)\frac{dV}{V} = \frac{dT}{T} \tag{3.142}$$

with the immediate solution

$$(1-\gamma)\ln V = \ln T + \text{ constant}, \tag{3.143}$$

giving

$$V^{1-\gamma} = \text{constant } T \quad \text{or} \quad TV^{\gamma-1} = \text{constant}. \tag{3.144}$$

Remembering that $\gamma \geq 1$, this confirms the downward slope of the entropy contours on the (V, T) plane.

This result is most frequently used with the variables p and T, rather than V and T:

$$pV^{\gamma} = \text{constant}. \tag{3.145}$$

The above equation holds only for quasistatic, adiabatic processes in ideal gases that have heat capacities independent of temperature. You might think that it is such a restricted result that it holds little interest. This equation is used frequently in applications, in the graduate record examination, and in physics oral examinations! You should memorize it and know how to work with it.

Problems

3.29 **Intensive vs. extensive variables**

Equation (3.134), $E(T, V) = E(T)$, states that for an ideal gas the energy is a function of temperature alone. How is it possible for E, an extensive quantity, to be a function of only T, an intensive quantity?

3.30 **Heat capacity at constant pressure**

Equation (3.131) shows that the heat capacity at constant volume, which is defined in terms of an entropy derivative, is also equal to an energy derivative. You might suspect a similar relation between $C_p(T, p, N)$ and

$$\left.\frac{\partial E(T, p, N)}{\partial T}\right)_{p,N}.$$

Show that such a suspicion is *not* correct, and instead find an expression for C_p in terms of a derivative of enthalpy.

3.31 **Qualitative heat capacity relations**

(This problem is stolen from a GRE Physics test.)

For an ideal gas, C_p is greater than C_V because:

a. The gas does work on its surroundings when its pressure remains constant while its temperature is increased.

b. The heat input per unit increase in temperature is the same for processes at constant pressure and at constant volume.

c. The pressure of the gas remains constant when its temperature remains constant.

d. The increase in the gas's internal energy is greater when the pressure remains constant than when the volume remains constant.

e. The heat needed is greater when the volume remains constant than when the pressure remains constant.

3.32 **Heat capacities in a magnetic system**

For a magnetic system (see equation (3.100)), show that

$$C_H = T\left.\frac{\partial S}{\partial T}\right)_H, \quad C_M = T\left.\frac{\partial S}{\partial T}\right)_M, \quad \beta = \left.\frac{\partial M}{\partial T}\right)_H, \quad \text{and} \quad \chi_T = \left.\frac{\partial M}{\partial H}\right)_T \tag{3.146}$$

are related through

$$C_M = C_H - T\beta^2/\chi_T. \tag{3.147}$$

3.33 **Isothermal compressibility**

a. Show that the isothermal compressibility, defined in problem 1.2 as

$$\kappa_T = -\frac{1}{V}\left.\frac{\partial V}{\partial p}\right)_{T,N}, \tag{3.148}$$

is also given by

$$\kappa_T = \frac{1}{\rho}\left.\frac{\partial \rho}{\partial p}\right)_T = \frac{1}{\rho^2}\left.\frac{\partial \rho}{\partial \mu}\right)_T = \frac{1}{\rho^2}\left.\frac{\partial^2 p}{\partial \mu^2}\right)_T, \tag{3.149}$$

where ρ is the number density N/V. Clue:

$$\left.\frac{\partial \rho}{\partial p}\right)_T = \left.\frac{\partial \rho}{\partial \mu}\right)_T \left.\frac{\partial \mu}{\partial p}\right)_T. \tag{3.150}$$

b. What does this result tell you about the relation between density and chemical potential?

c. In part (a.) we began with a description in terms of the three variables T, p, N and then reduced it to an intensive-only description, which requires just two variables, such as μ and T. Reverse this process to show that

$$\kappa_T = \frac{V}{N^2}\left.\frac{\partial N}{\partial \mu}\right)_{T,V} = -\frac{1}{N}\left.\frac{\partial V}{\partial \mu}\right)_{T,N}. \tag{3.151}$$

3.34 **Pressure differential**

By regarding pressure as a function of temperature T and number density ρ, show that

$$dp = \frac{\beta}{\kappa_T}\,dT + \frac{1}{\rho\kappa_T}\,d\rho. \tag{3.152}$$

3.35 **Isothermal *vs.* adiabatic compressibility**

In class we derived a remarkable relation between the heat capacities C_p and C_V. This problem uncovers a similar relation between the isothermal and adiabatic compressibilities,

$$\kappa_T = -\frac{1}{V}\left.\frac{\partial V}{\partial p}\right)_T \quad \text{and} \quad \kappa_S = -\frac{1}{V}\left.\frac{\partial V}{\partial p}\right)_S. \tag{3.153}$$

The adiabatic compressibility κ_S is the compressibility measured when the fluid is thermally insulated. (It is related to the speed of sound: see problem 3.44.)

a. Use

$$dS = \left.\frac{\partial S}{\partial T}\right)_p dT + \left.\frac{\partial S}{\partial p}\right)_T dp \tag{3.154}$$

to show that

$$\left.\frac{\partial T}{\partial p}\right)_S = \frac{\beta T}{C_p/V}. \tag{3.155}$$

Sketch an experiment to measure this quantity directly.

b. From the mathematical relation

$$dV = \left.\frac{\partial V}{\partial p}\right)_T dp + \left.\frac{\partial V}{\partial T}\right)_p dT \tag{3.156}$$

derive the multivariate chain rule

$$\left.\frac{\partial V}{\partial p}\right)_S = \left.\frac{\partial V}{\partial p}\right)_T + \left.\frac{\partial V}{\partial T}\right)_p \left.\frac{\partial T}{\partial p}\right)_S, \tag{3.157}$$

whence

$$\kappa_S = \kappa_T - \frac{\beta^2 T}{C_p/V}. \tag{3.158}$$

c. Finally, show that

$$\gamma \equiv \frac{C_p}{C_V} = \frac{\kappa_T}{\kappa_S}. \tag{3.159}$$

3.36 **Change of chemical potential with temperature**

Prove that

$$\left.\frac{\partial \mu}{\partial T}\right)_{p,N} = -\frac{S}{N}, \tag{3.160}$$

and that

$$\left.\frac{\partial \mu}{\partial T}\right)_{V,N} = -\left.\frac{\partial S}{\partial N}\right)_{T,V} = -\frac{S}{N} + \frac{\beta}{\rho\kappa_T}. \tag{3.161}$$

How's that for weird?

3.9 Thermodynamics Applied to Phase Transitions

At phase coexistence between, say, gas and liquid

$$\mu_G(p,T) = \mu_L(p,T). \tag{3.162}$$

The Clausius-Clapeyron equation relates the slope of a phase coexistence line $p_x(T)$ with the volume and entropy changes when the sample crosses that line:

$$\frac{dp_x}{dT} = \frac{\Delta S}{\Delta V}. \tag{3.163}$$

Problems

3.37 Thermodynamics applied to the triple point

A triple point has the usual arrangement of a rising gas-solid coexistence line splitting into two rising coexistence lines: a liquid-solid line and a gas-liquid line.

a. Relate the slopes of the three coexistence lines that meet at the triple point to the entropy changes $\Delta S_{gl} = S_{\text{gas}} - S_{\text{liquid}}$ and $\Delta S_{ls} = S_{\text{liquid}} - S_{\text{solid}}$.

b. Show that the gas-solid coexistence line, when extended beyond the triple point, must extend into the liquid phase as show below left. It must *not* extend into the gas phase as shown below right.

[This result is called "the Schreinemakers 180° rule" after Franciscus A. H. Schreinemakers, Dutch chemist, 1864–1945.]

c. Show that in most cases, at the triple point the gas-solid coexistence line has nearly the same slope as the gas-liquid coexistence line.

d. The diagram below illustrates the Wikipedia article "Triple point" (15 December 2014). The caption is "A typical phase diagram. The solid green line applies to most substances; the dotted green line gives the anomalous behaviour of water." Critique this diagram.

3.38 **Magnetic phase transitions**

The phase diagrams for magnetic systems are usually plotted with temperature on the vertical axis and magnetic field H on the horizontal axis. Show that if such a diagram displays two-phase coexistence along the line $T_x(H)$, then that phase boundary has slope

$$\frac{dT_x}{dH} = -\frac{\Delta M}{\Delta S} \tag{3.164}$$

where ΔM is the magnetization change accompanying the phase transition.

3.39 **Curvature of phase coexistence lines**

a. Define

$$\beta' = \left.\frac{\partial v}{\partial T}\right)_p = v\beta \quad \text{and} \quad \kappa'_T = -\left.\frac{\partial v}{\partial p}\right)_T = v\kappa_T$$

and show that, for small changes of temperature δT and pressure δp, the resulting change in chemical potential will be

$$\delta\mu = -s\,\delta T + v\,\delta p + \tfrac{1}{2}\left[-\frac{c_p}{T}(\delta T)^2 + 2\beta'(\delta T)(\delta p) - \kappa'_T(\delta p)^2\right] + \cdots.$$

b. Show that for a change along a phase coexistence boundary $p_x(T)$,

$$\delta p = \frac{dp_x}{dT}\delta T + \tfrac{1}{2}\frac{d^2p_x}{dT^2}(\delta T)^2 + \cdots.$$

c. If $\Delta s = s_A - s_B$, and similarly for Δv, Δc_p, etc., show that for any point on a phase coexistence boundary

$$0 = -\Delta s\,\delta T + \Delta v\,\delta p + \tfrac{1}{2}\left[-\frac{\Delta c_p}{T}(\delta T)^2 + 2\Delta\beta'(\delta T)(\delta p) - \Delta\kappa'_T(\delta p)^2\right] + \cdots.$$

d. Conclude that the slope of the phase coexistence curve is

$$\frac{dp_x}{dT} = \frac{\Delta s}{\Delta v},$$

the usual Clausius-Clapeyron equation, while the second derivative is

$$\frac{d^2 p_x}{dT^2} = \frac{\Delta c_p}{T\Delta v} - 2\frac{\Delta\beta'\Delta s}{(\Delta v)^2} + \frac{\Delta\kappa'_T(\Delta s)^2}{(\Delta v)^3}, \tag{3.165}$$

the "second-order Clausius-Clapeyron equation".

3.10 Thermodynamics Applied to Chemical Reactions

3.10.1 Thermodynamics of mixtures

So far in this book we have considered mostly pure substances. What happens if we have a mixture of several chemical species, say the four substances A, B, C, and D? In this case the mechanical parameters will include the numbers N_A, N_B, N_C, and N_D, and the entropy

$$S(E, V, N_A, N_B, N_C, N_D) \tag{3.166}$$

will have the total differential

$$dS = \frac{1}{T}\,dE + \frac{p}{T}\,dV - \sum_{i=A}^{D} \frac{\mu_i}{T}\,dN_i. \tag{3.167}$$

The differential above defines the quantity

$$\mu_A(E, V, N_A, N_B, N_C, N_D) = -T(E, V, N_A, N_B, N_C, N_D)\left.\frac{\partial S}{\partial N_A}\right)_{E,V,N_B,N_C,N_D}. \tag{3.168}$$

Notice that the chemical potential of substance A can depend upon the number of molecules of B, C, and D present, so it is not necessarily the same as the chemical potential of pure A. (For example, the chemical potential of sugar dissolved in a cup of water is not equal to the chemical potential of pure sugar, unless the solution is saturated and there is undissolved sugar at the bottom of the cup.)

It is convenient to define the total number of molecules

$$N = N_A + N_B + N_C + N_D \tag{3.169}$$

and the (intensive) fractional amounts

$$f_A = \frac{N_A}{N}, \text{ etc.} \tag{3.170}$$

The thermodynamic dance can be followed for mixtures just as well as it can be for pure substances. One noteworthy result is

$$\begin{aligned} G(T, p, N_A, N_B, N_C, N_D) &\equiv E - TS + pV \\ &= \sum_{i=A}^{D} \mu_i(T, p, f_A, f_B, f_C, f_D)N_i \end{aligned} \tag{3.171}$$

3.10.2 Equilibrium condition for a chemical reaction

Suppose that α molecules of substance A can combine with β molecules of substance B to produce γ molecules of substance C and δ molecules of substance D. If this can happen, then of course the reverse can also occur: γ molecules of substance C may combine with δ molecules of substance D to produce α molecules of substance A and β molecules of substance B. The first action is called a "forward step" of the chemical reaction, and the second is called a "backward step". This chemical reaction is symbolized by

$$\alpha \text{ A} + \beta \text{ B} \Longleftrightarrow \gamma \text{ C} + \delta \text{ D}. \tag{3.172}$$

Thus if we mix together a certain number of molecules each of A, B, C, and D, there can be a reaction and we might not end up with the same distribution of molecules that we started off with. It is possible that there will be more forward than backwards steps, in which case we will end up with more C's and D's and fewer A's and B's than we had at the start, or it could go predominantly the other way. It is allowable for the system to exist in a wide range of conditions, varying from "mostly reactants" to "mostly products", but experience teaches us that the overwhelmingly likely condition is one with some of each are present. In this condition reactions still proceed, in both the forward and backward directions, but steps in both directions proceed at the same rate so there is no net change in the number of molecules of any species. This condition is called "chemical equilibrium".

The above paragraph precisely parallels the discussion in section 2.7, "Using entropy to find (define) temperature and pressure". In that section we saw that if, for example, two systems could exchange volume, it was possible for all the volume to go to one system or for all of it to go to the other system, but that the overwhelmingly likely possibility was the "equilibrium condition" in which the two systems shared volume in such a way that each had the same pressure. The equilibrium condition was the one with a maximum number of microstates (or, equivalently, a maximum entropy) with respect to the distribution of volume.

Precisely the same effect is at work in chemical equilibrium. The equilibrium state is characterized by a maximum in the entropy with respect to the number of number of reaction steps taken. At this equilibrium point the entropy doesn't change when the chemical reaction proceeds by one step in either direction. The change in entropy through a forward step is of course

$$-\alpha \frac{\partial S}{\partial N_A} - \beta \frac{\partial S}{\partial N_B} + \gamma \frac{\partial S}{\partial N_C} + \delta \frac{\partial S}{\partial N_D}, \tag{3.173}$$

and the change in entropy through a backward step is the negative of this. In either case, the equilibrium condition is that this change in entropy vanish, i.e. that

$$\alpha \mu_A + \beta \mu_B = \gamma \mu_C + \delta \mu_D. \tag{3.174}$$

Interpretation in terms of "escaping tendency".

If we specialize to the case of liquid-gas phase equilibrium, this equation is just equation (3.162).

We cannot go further without formulas for the chemical potentials μ. Complete formulas are not available within the realm of thermodynamics, but the next section shows that thermodynamics can put severe constraints on the permissible functional forms for μ.

3.10.3 Chemical potential of an ideal gas

I emphasized on page 69 that the equation of state did not contain full thermodynamic information. Thus, for example, knowledge of the equation of state $V(T,p,N)$ is not sufficient to uncover the master function $G(T,p,N)$. On the other hand, that knowledge is sufficient to restrict the functional form of the Gibbs potential, which in turn restricts the possible functional forms of the chemical potential. In this subsection we uncover that restriction.

Our strategy is to recognize that volume is the derivative of $G(T,p,N)$ with respect to p, and chemical potential is the derivative of $G(T,p,N)$ with respect to N. Given $V(T,p,N)$, we can integrate with respect to p to find $G(T,p,N)$, and then differentiate with respect to N to find $\mu(T,p)$. This process does not completely determine the Gibbs potential or the chemical potential, because a constant of integration is involved. But it does restrict the permissible functional forms of $G(T,p,N)$ and $\mu(T,p)$.

The master thermodynamic equation for variables (T,p,N) is

$$dG = -S\,dT + V\,dp + \mu\,dN, \tag{3.175}$$

whence

$$V(T,p,N) = \left.\frac{\partial G}{\partial p}\right)_{T,N}. \tag{3.176}$$

Thus for a substance obeying the ideal gas equation of state, we have

$$\frac{Nk_BT}{p} = \left.\frac{\partial G}{\partial p}\right)_{T,N}. \tag{3.177}$$

Integrating both sides with respect to p gives

$$Nk_BT\int\frac{dp}{p} = \int\left.\frac{\partial G}{\partial p}\right)_{T,N}dp \tag{3.178}$$

or

$$Nk_BT\ln\left(\frac{p}{\tilde{p}(T,N)}\right) = G(T,p,N). \tag{3.179}$$

The role of "constant of integration" is played by $\tilde{p}(T,N)$, which must be constant with respect to p but which may vary with T and N.

In fact, careful examination of the above equation shows what the dependence of $\tilde{p}(T,N)$ on N must be. The quantities G and N above are extensive, while T and p are intensive, so $\tilde{p}$ must be intensive so that both sides will depend linearly on system size. Thus $\tilde{p}$ is independent of N, and we have

$$G(T,p,N) = Nk_BT\ln\left(\frac{p}{\tilde{p}(T)}\right), \tag{3.180}$$

whence

$$\mu(T,p) = \frac{G(T,p,N)}{N} = k_BT\ln\left(\frac{p}{\tilde{p}(T)}\right). \tag{3.181}$$

This is the form of the Gibbs potential and of the chemical potential for any ideal gas. The function $\tilde{p}(T)$ is undetermined, and it is through differences in $\tilde{p}(T)$ that one ideal gas differs from another. But the dependence on p and on N are completely pinned down by the above results.

This result is frequently written in terms of the number density $\rho = N/V$. For an ideal gas $p = k_B T\rho$, and we can define the function $\tilde{\rho}(T)$ through $\tilde{p}(T) = k_B T\tilde{\rho}(T)$. Then we can write

$$\mu(T,\rho) = k_B T \ln\left(\frac{\rho}{\tilde{\rho}(T)}\right). \tag{3.182}$$

What if the ideal gas is a mixture of, say, the four chemical species A, B, C, and D? Exactly the same result applies for each species:

$$\mu_A(T,\rho_A) = k_B T \ln\left(\frac{\rho_A}{\tilde{\rho}_A(T)}\right) \tag{3.183}$$

where the function $\tilde{\rho}_A(T)$ is the same for a pure ideal gas of A and for an ideal gas mixture. The chemical potential μ_A is independent of the densities ρ_B, ρ_C, and ρ_D. I am looking for a purely thermodynamic argument to proved this result, but until I find it you'll have to accept this statistical argument: In an ideal gas the molecules of A don't interact with the molecules of B, C, and D, so they don't know whether or not such molecules are even present. The behavior of A thus can't depend upon the number of B, C, and D molecules present, and it will be the same even if there are 0 molecules of B, C, and D, that is if the gas is pure.

3.10.4 The equilibrium condition for idea gas reactions

[Before plunging in, I want to point out that there's a contradiction embodied in the term "ideal gas chemical reaction". In an ideal gas, molecules do not interact... when two molecules approach, they just pass right through each other. But a chemical reaction is a form of interaction! Clearly, in this section we are thinking of molecules that interact weakly unless they are very close to each other, but once they approach they interact strongly and react quickly.]

Putting together the general equilibrium condition for chemical reactions

$$\alpha\mu_A + \beta\mu_B = \gamma\mu_C + \delta\mu_D \tag{3.184}$$

with expression (3.183) for the chemical potential of an ideal gas gives the equilibrium condition

$$\alpha k_B T \ln\left(\frac{\rho_A}{\tilde{\rho}_A(T)}\right) + \beta k_B T \ln\left(\frac{\rho_B}{\tilde{\rho}_B(T)}\right) = \gamma k_B T \ln\left(\frac{\rho_C}{\tilde{\rho}_C(T)}\right) + \delta k_B T \ln\left(\frac{\rho_D}{\tilde{\rho}_D(T)}\right). \tag{3.185}$$

Using the properties of logarithms, this can be rewritten as

$$\left(\frac{\rho_A}{\tilde{\rho}_A(T)}\right)^{\alpha}\left(\frac{\rho_B}{\tilde{\rho}_B(T)}\right)^{\beta} = \left(\frac{\rho_C}{\tilde{\rho}_C(T)}\right)^{\gamma}\left(\frac{\rho_D}{\tilde{\rho}_D(T)}\right)^{\delta},$$

or, after minor rearrangement, as

$$\frac{\rho_C^{\gamma}\rho_D^{\delta}}{\rho_A^{\alpha}\rho_B^{\beta}} = \frac{\tilde{\rho}_C^{\gamma}(T)\tilde{\rho}_D^{\delta}(T)}{\tilde{\rho}_A^{\alpha}(T)\tilde{\rho}_B^{\beta}(T)}. \tag{3.186}$$

Chemists like to write this equation in terms of "the concentration of A", written as [A], namely as

$$\frac{[\mathrm{C}]^\gamma[\mathrm{D}]^\delta}{[\mathrm{A}]^\alpha[\mathrm{B}]^\beta} = K(T), \tag{3.187}$$

where $K(T)$ is called the "equilibrium constant" (despite the fact that it is a function of temperature). This result is called the "law of mass action" despite the fact that it is not a fundamental law (we have just derived it) and even despite the fact that (because it holds only for ideal gases) it's not precisely true for any real substance! (And also, I might add, despite the fact that the "law" has nothing to do with "mass" or with "action".) Even though the result is poorly named, it shows just how valuable thermodynamics is in connecting disparate areas of research. Who would have guessed that the law of mass action follows from only thermodynamics and the ideal gas equation of state? We have proven the result without once doing a chemical experiment!

Problems

3.40 **Chemical potential for ideal gas mixtures**
The entropy $S(E, V, N_A, N_B)$ of a classical monatomic ideal gas mixture was uncovered using statistical mechanics in problem 2.16. Invert the result of that problem to find $E(S, V, N_A, N_B)$, then show that the temperature and pressure of an ideal gas mixture obey familiar results for pure ideal gases. Show that

$$\mu_A(T, p, f_A, f_B) = k_B T \ln\left[\left(\frac{h_0^2}{2\pi m_A (k_B T)^{5/3}}\right)^{3/2} p f_A\right], \tag{3.188}$$

and relate this expression to the form (3.183). Verify relation (3.172).

3.41 **Relations between equilibrium constants**
You measure the equilibrium constants for these three chemical reactions:

$$\begin{aligned} 2\ \mathrm{H_2} + \mathrm{O_2} &\iff 2\ \mathrm{H_2O} \\ \mathrm{H_2} + \mathrm{Cl_2} &\iff 2\ \mathrm{HCl} \\ \mathrm{Cl_2} + 2\ \mathrm{O_2} &\iff 2\ \mathrm{ClO_2} \end{aligned}$$

(Call those constants K_A, K_B, and K_C.) Find an expression for the equilibrium constant for the reaction

$$4\ \mathrm{ClO_2} + 2\ \mathrm{H_2O} \iff 4\ \mathrm{HCl} + 5\ \mathrm{O_2}$$

3.11 Thermodynamics Applied to Light

Another non-fluid system (see section 3.7).

See Robert E. Kelly, "Thermodynamics of blackbody radiation," *Am. J. Phys.* **49** (1981) 714–719, and Max Planck, *The Theory of Heat Radiation*, part II.

How do you get a "box of light?" This is blackbody (cavity) radiation: meaning radiation in thermal equilibrium. (Usually it's also in equilibrium with the cavity walls, but you could imagine a container with mirror walls...)

3.11.1 Fundamentals

The fundamental thermodynamic equation for this system involves the master function $E(S, V)$. It is

$$dE = T\,dS - p\,dV. \tag{3.189}$$

This equation differs from previously encountered master equations in that there is no term for $\mu\,dN$. From the classical perspective, this is because radiation is made up of fields, not particles. From the quantal perspective, this is because photon number is not conserved.

Thermodynamics binds one quantity to another, but it must use experiment (or statistical mechanics) to find the values being bound. For example, thermodynamics tells us that

$$C_p = C_V + TV\frac{\beta^2}{\kappa_T},$$

but it cannot calculate either C_p, or C_V, or β, or κ_T: these quantities must be found by experiment (or through a statistical mechanical calculation). The empirical result that we will employ is that for blackbody radiation,

$$p = \frac{1}{3}\frac{E}{V} = \frac{1}{3}u. \tag{3.190}$$

This equation of state is the parallel for blackbody radiation to $pV = Nk_BT$ for ideal gases. It was discovered by experiment, but it can be derived from electrodynamics as well.

3.11.2 Energy density as a function of temperature

Consider the energy as a function of volume and temperature. Because $E(V, T)$ is extensive, but in the argument list only V is extensive (recall that N doesn't appear in the argument list), we must have

$$E(V, T) = Vu(T). \tag{3.191}$$

That is, the energy density u depends *only* upon temperature. What is this dependence?

We seek a differential equation for $u(T)$. Compare

$$dE = d(Vu) = V\,du + u\,dV = V\frac{du}{dT}\,dT + u\,dV \tag{3.192}$$

with

$$dE = T\,dS - p\,dV = T\,dS - \tfrac{1}{3}u\,dV. \tag{3.193}$$

Together these give us a formula for dS:

$$dS = \left(\frac{V}{T}\frac{du}{dT}\right) dT + \left[\frac{4}{3}\frac{u}{T}\right] dV. \tag{3.194}$$

The Maxwell relation associated with this differential is

$$\left.\frac{\partial(\quad)}{\partial V}\right)_T = \left.\frac{\partial[\quad]}{\partial T}\right)_V \tag{3.195}$$

or

$$\frac{1}{T}\frac{du}{dT} = \frac{4}{3}\frac{d[u/T]}{dT} = \frac{4}{3}\left[\frac{1}{T}\frac{du}{dT} - \frac{1}{T^2}u\right]. \tag{3.196}$$

A rearrangement gives

$$\frac{du}{dT} = 4\frac{u}{T} \tag{3.197}$$

or

$$4\frac{dT}{T} = \frac{du}{u}. \tag{3.198}$$

The solution is

$$4\ln T = \ln u + \text{ const} \tag{3.199}$$

whence

$$u(T) = \sigma T^4. \tag{3.200}$$

The Stefan-Boltzmann law!

3.11.3 Quasistatic adiabatic expansion of radiation

Consider a sample of radiation undergoing quasistatic adiabatic change of volume. The entropy is a constant during this process, although the value of that constant will of course depend on the particular sample that's expanding.

$$dE = T\,dS - p\,dV \tag{3.201}$$

but $dS = 0$, $E = uV$, and $p = u/3$ so

$$\begin{aligned} u\,dV + V\,du &= -\tfrac{1}{3}u\,dV \\ V\,du &= -\tfrac{4}{3}u\,dV \\ \ln u &= -\tfrac{4}{3}\ln V + \text{ const} \\ u &= KV^{-4/3} \end{aligned}$$

Recognizing that this constant will depend upon which adiabat is taken, i.e. that it will depend on the entropy, we write

$$u(S,V) = K(S)V^{-4/3}. \tag{3.202}$$

Using the Stefan-Boltzmann result $u = \sigma T^4$ we find that

$$T(S,V) = \frac{C(S)}{V^{1/3}}. \tag{3.203}$$

This explains the cooling of the universe as it expands from the initial "hot big bang" to the current "3° K microwave background."

3.11.4 Thermodynamics of the energy spectrum

What if we consider not just the energy density per volume, but the energy density per volume and wavelength? Let

$$\bar{u}(T,\lambda)\,d\lambda \tag{3.204}$$

represent the energy per volume due to that radiation with wavelength between λ and $\lambda + d\lambda$. As you know, quantum mechanics was discovered through Planck's efforts to find a theoretical explanation for the measured function $\bar{u}(T,\lambda)$. This is not the place to describe Planck's work. Instead I want to focus on a purely thermodynamic result that was known long before Planck started his investigations.

This is Wien's law[8], which states that the function $\bar{u}(T,\lambda)$, which you'd think could have any old form, must be of the form

$$\bar{u}(T,\lambda) = T^5 f(\lambda T). \tag{3.205}$$

An immediate consequence of Wien's law is the Wien displacement theorem: The wavelength $\hat{\lambda}$ which maximizes $\bar{u}(T,\lambda)$ is inversely proportional to temperature:

$$\hat{\lambda}(T) = \frac{\text{constant}}{T}, \tag{3.206}$$

where the constant is the value of x that maximizes $f(x)$. The consequences of the Wien displacement theorem are familiar from daily life: low temperature objects (such as people) radiate largely in the infrared, moderate temperature objects (such as horseshoes in the forge) radiate largely in the red, while high temperature objects (such as the star Sirius) radiate largely in the blue.

I stated earlier that Wien's law is a purely thermodynamic result. That's almost true, but it also relies on one more fact from electrodynamics, a result called "no mode hoping":

> If the volume makes a quasistatic, adiabatic change from V_1 to V_2, then the light of wavelength in the range λ_1 to $\lambda_1 + d\lambda_1$ shifts into the range λ_2 to $\lambda_2 + d\lambda_2$ where
>
> $$\frac{\lambda_1}{V_1^{1/3}} = \frac{\lambda_2}{V_2^{1/3}}. \tag{3.207}$$

(This result may be derived rigorously from Maxwell's equations, but it's reasonable through this analogy: A string of length L vibrating in, say, its third mode, has wavelength $\lambda = \frac{3}{2}L$. If the length of the string is slowly changed, then the wave remains in its third mode, so λ/L is constant.)

Now we're ready to begin the derivation. Consider the quasistatic adiabatic expansion of light, and while that expansion is going on focus your attention on the light in wavelength range λ to $\lambda + d\lambda$. According to the "no mode hopping" result, during this expansion the quantity

$$\lambda/V^{1/3} \tag{3.208}$$

[8] Wien is pronounced like the English "veen".

remains constant during the expansion. Furthermore, according to equation (3.203) the quantity

$$TV^{1/3} \tag{3.209}$$

also remains constant. Multiplying these two equations, we find that the volume-independent quantity

$$T\lambda \tag{3.210}$$

remains constant during the expansion: this number characterizes the expansion.

Another such volume-independent constant can be found by repeating the reasoning of subsection 3.11.3, Quasistatic adiabatic expansion of radiation, but considering not the energy of all the radiation, but the energy of the radiation with wavelengths from λ to $\lambda+\Delta\lambda$. This energy is $E = \bar{u}V\Delta\lambda$, and the pressure due to this segment of the radiation is $p = \frac{1}{3}\bar{u}\Delta\lambda$. During a quasistatic adiabatic expansion, $dE = -p\,dV$, so

$$(V\Delta\lambda)\,d\bar{u} + (\bar{u}\Delta\lambda)\,dV + (\bar{u}V)\,d[\Delta\lambda] = -\tfrac{1}{3}\bar{u}\Delta\lambda\,dV. \tag{3.211}$$

During the expansion the volume and wavelengths are changing through (see equation 3.207)

$$\begin{aligned} \lambda &= cV^{1/3} \\ d\lambda &= c\tfrac{1}{3}V^{-2/3}\,dV = \frac{1}{3}\frac{\lambda}{V}\,dV \\ d[\Delta\lambda] &= \frac{1}{3}\frac{\Delta\lambda}{V}\,dV \end{aligned} \tag{3.212}$$

so we have

$$(V\Delta\lambda)\,d\bar{u} = -\tfrac{5}{3}\bar{u}\Delta\lambda\,dV. \tag{3.213}$$

Thus

$$\begin{aligned} V\,d\bar{u} &= -\tfrac{5}{3}\bar{u}\,dV \\ \ln\bar{u} &= -\tfrac{5}{3}\ln V + \text{ const} \\ \bar{u} &= KV^{-5/3} \end{aligned} \tag{3.214}$$

But equation (3.203) shows how a larger volume is related to a lower temperature, so the quantity

$$\frac{\bar{u}(T,\lambda)}{T^5} \tag{3.215}$$

remains constant during the expansion.

Thus we have two volume-independent ways to characterize the particular curve taken by this expansion. In the thermodynamics of light, a state is specified by two variables, so a curve is specified by only one parameter. Hence these two characterizations cannot be independent: one must be a function of the other. Thus

$$\frac{\bar{u}(T,\lambda)}{T^5} = f(\lambda T)$$

or

$$\bar{u}(T,\lambda) = T^5 f(\lambda T). \tag{3.216}$$

Wien's law.

Problems

3.42 **Heat capacity of light**

Show that, for blackbody radiation, $C_V = 4E/T$.

Resources

Thermodynamic tables. (G.N. Lewis and M. Randall) Zemansky. Practical heat engines. Callen. Fermi.

Math book. e.g. Taylor and Mann?

Picture of Smithsonian crystal on www?

Thermodynamic data (e.g. steam tables) on www?

3.12 Additional Problems

3.43 **Cool mountain air**

Model the earth's atmosphere as an ideal gas (nitrogen) in a uniform gravitational field. Ignore all winds. Let m denote the mass of a gas molecule, g the acceleration of gravity, and z the height above sea level.

a. Use ideas from Newtonian mechanics to show that the change of atmospheric pressure p with height z is

$$\frac{dp}{dz} = -\frac{mg}{k_B T(z)} p(z). \tag{3.217}$$

b. If the atmosphere is a poor conductor of heat, then the decrease in pressure with height is due to an adiabatic expansion. (*Clue:* In other words: At the bottom of the mountain, fill an insulated balloon with air at the local density, pressure, and temperature. Transport that balloon to the top of the mountain. During the journey the balloon will expand, so the density, pressure, and temperature will change. According to our assumption, during this journey the balloon's density, pressure, and temperature will match those of the atmosphere outside.) Show that under this assumption

$$\frac{dp}{dT} = \frac{\gamma}{\gamma - 1} \frac{p(T)}{T} \tag{3.218}$$

and hence that

$$\frac{dT}{dz} = -\frac{\gamma - 1}{\gamma} \frac{mg}{k_B}. \tag{3.219}$$

Evaluate this expression in kelvin per kilometer for nitrogen, which has $\gamma = 1.4$.

c. In contrast, if the atmosphere were a good conductor of heat, then temperature would be uniform. Find $p(z)$ under such circumstances. Denote the sea-level pressure and temperature by p_0 and T_0.

d. Similarly find $p(z)$ for an adiabatic atmosphere.

3.44 **The speed of sound**

When a sound wave passes through a fluid (liquid or gas), the period of vibration is short compared to the time necessary for significant heat flow, so the compressions may be considered adiabatic.

Analyze the compressions and rarefactions of fluid in a tube. The equilibrium mass density is ρ_0. Apply $\mathbf{F} = m\mathbf{a}$ to a slug of fluid of thickness Δx, and show that if the variations in pressure $p(x,t)$ are small then pressure satisfies the wave equation

$$\frac{\partial^2 p}{\partial t^2} = c^2 \frac{\partial^2 p}{\partial x^2} \tag{3.220}$$

where c, the speed of sound, is given by

$$c = \frac{1}{\sqrt{\rho_0 \kappa_S}}. \tag{3.221}$$

Optional: Use the results of problems 1.2 and 3.35 to show that, for an ideal gas,

$$c = \sqrt{\gamma \frac{k_B T}{m}}. \tag{3.222}$$

3.45 **Thermodynamics of a plastic rod**

(This problem is based on Reif problem 5.14.)

For a restricted range of lengths L and temperatures T, the tension force in a stretched plastic rod is

$$F(T, L) = aT^2(L - L_0), \tag{3.223}$$

where a is a positive constant and L_0 is the relaxed (unstretched) length of the rod. When $L = L_0$, the heat capacity C_L of the rod (measured at constant length) is given by $C_L(T, L_0) = bT$, where b is independent of temperature.

a. Write down the fundamental thermodynamic relation for this rod, expressing dE in terms of dS and dL.

b. Compute $(\partial S/\partial L)_T$. (Clue: Derive an appropriate Maxwell relation for the assembly with variables T and L.)

c. Knowing $S(T_0, L_0)$, integrate along an appropriate path to find $S(T, L)$ at any temperature and length within the range of applicability of the equation for $F(T, L)$.

d. If you start at $T = T_i$ and $L = L_i$ and then quasi-statically stretch a thermally insulated rod until it reaches length L_f, what is the final temperature T_f? Show than when $L_0 \leq L_i < L_f$, the rod is cooled by this process.

e. Find the heat capacity $C_L(L, T)$ of the rod when its length is not necessarily L_0.

f. Find $(\partial T/\partial L)_S$ for arbitrary T and L. Can insulated stretches warm as well as cool the rod?

3.46 **Magnetic cooling**

At low temperatures, paramagnetic salts obey the Curie law

$$M = c\frac{H}{T}, \tag{3.224}$$

where c is a positive constant (see equation (3.100)). Assume that the heat capacity C_H is a constant independent of temperature and field. Suppose a sample at magnetic field H_i and temperature T_i is wrapped in insulation, and then the magnetic field is slowly reduced to zero. Find the final temperature, and show that it is less than T_i. This technique, known as "adiabatic demagnetization" is the refrigeration method used to produce the temperatures from about 1 kelvin to 1 microkelvin.

3.47 **Thermodynamics of an electrochemical cell**

Reif 5.16.

3.48 **Thermodynamics and evolution**

Read the essay "Thermodynamics and Evolution" by John W. Patterson, in *Scientists Confront Creationism*, Laurie R. Godfrey, ed. (Norton, New York, 1983), pages 99–116, on reserve in the science library.

a. When a snowflake forms, its surroundings increase in entropy ("become more disordered"). What is the name of the heat flow associated with this entropy change?

b. Patterson argues that $\Delta S < 0$ on Earth, due to biological evolution, and that $\Delta S > 0$ somewhere else in the universe in order to make up for it. Where is that entropy increase taking place?

c. Patterson feels the need to invoke "self-organization" and Prigogine (pages 110–111) to explain *how* his ram pumps could be made. Is this necessary? List two or more situations from nature in which water *does* flow uphill.

3.49 **Entropy and evolution**

Creationists sometimes claim that the second law of thermodynamics prohibits biological evolution.

a. The surface of the Sun (mean temperature 5778 K) heats the surface of the Earth (mean temperature 288 K) through visible and near-infrared radiation. The solar energy absorbed by the Earth each second is 1.732×10^{17} J. What is the entropy change per second (due to this process) of the Sun? The Earth? Does the entropy of "Sun plus Earth" increase or decrease?

b. Yet the mean temperature of the Earth changes slowly, if at all. This is because almost all of the solar energy absorbed by the Earth is then emitted through far-infrared radiation which in turn heats "outer space" — the cosmic microwave background (CMB; temperature 2.728 K). What is the entropy change per second (due to this process) of the Earth? The CMB? Does the entropy of "Earth plus CMB" increase or decrease?

c. Now refine the model by supposing that, due to evolution, the entropy of the Earth is not exactly constant, but is decreasing. (In this case the entropy of the CMB would have to be increasing faster than rate predicted in part (b).) Suppose that, due to evolution, each individual organism is 1000 times "more improbable" than the corresponding individual was 100 years ago. In other words, if Ω_i is the number of microstates consistent with the specification of an organism 100 years ago, and if Ω_f is the number of microstates consistent with the specification of today's "improved and less probable" organism, then $\Omega_f = 10^{-3}\Omega_i$. What is the corresponding change in entropy per organism?

d. The population of Earth is about 10^{18} eukaryotic individuals and 10^{32} prokaryotic individuals. If the estimate of part (c) holds for each one of them, what is the change in entropy due to evolution each second?

e. How accurately would you have to measure the entropy flux of part (b) in order to notice the diversion of entropy flux calculated in part (d)? Has any scientific quantity ever been measured to this accuracy?

f. It is generally agreed that the greatest rate of evolution fell during the Cambrian period, from 542 million years ago to 488 million years ago. During this so-called "Cambrian explosion" multicellular organisms first formed and then radiated into remarkable variety. Suppose that during the Cambrian period entropy was diverted into the evolution of living things at the rate calculated in part (d). And suppose that at the end of the Cambrian there were 10^{18} multicellular individuals. How much "improved and less probable" would each organism be, relative to its single-celled ancestor at the beginning of the Cambrian period?

The moral of the story? There's plenty of entropy to go around.

Chapter 4

Ensembles

After the last chapter's dip into thermodynamics, we return to the discussion of the principles of statistical mechanics begun in chapter 2.

4.1 The Canonical Ensemble

Examples of "systems" and "baths":

- Bottle of wine/a swimming pool.
- Sample of gas/laboratory temperature control apparatus.
- One ideal gas molecule/sample of gas.
- But *not* one interacting gas molecule/sample of gas. (The Boltzmann distribution is for systems within ensembles, not for molecules within systems.)

We denote a microstate by x. Exactly what is meant by a microstate will depend upon circumstances. For a system of point particles it means specifying all the positions and momenta. For a spin system it means specifying the "up" or "down" state of each spin. We'll see other examples later.

I'll try to use different letters for microscopic and macroscopic (thermodynamic) quantities. For example $H(\mathsf{x})$ versus E (the thermodynamic energy is $E = \langle H(\mathsf{x})\rangle$; $\mathcal{M}(\mathsf{x})$ versus $M = \langle \mathcal{M}(\mathsf{x})\rangle$.

Summary

The canonical ensemble in general:

The probability that the system is in the microstate x is proportional to the "Boltzmann factor"

$$e^{-H(\mathsf{x})/k_BT}. \tag{4.1}$$

The normalization factor is called the "partition function" or "sum over all states" (German "Zustandsumme"):

$$Z(T,V,N) = \sum_{\text{microstates } \mathsf{x}} e^{-H(\mathsf{x})/k_BT}. \tag{4.2}$$

(Note that Z is independent of x.) Thus the probability that the system is in microstate x is

$$\frac{e^{-H(\mathsf{x})/k_BT}}{Z(T,V,N)}. \tag{4.3}$$

The connection to thermodynamics is that the Helmholtz free energy is

$$F(T,V,N) = -k_BT \ln Z(T,V,N). \tag{4.4}$$

Note that in finding Z we sum over *all* microstates: the low-energy ones, the high-energy ones, the "orderly" ones (e.g. all atoms heading west, or all atoms heading east), the "disorderly" ones (e.g. atoms heading in scattered directions).

The canonical ensemble for a pure classical monatomic fluid:

The probability that the system is in the microstate Γ is proportional to the "Boltzmann factor"

$$e^{-H(\Gamma)/k_BT}. \tag{4.5}$$

Writing out all the normalizations correctly gives: the probability that the system is in some microstate within the phase space volume element $d\Gamma$ about Γ is

$$\frac{e^{-H(\Gamma)/k_BT}}{N!h_0^{3N}Z(T,V,N)}d\Gamma, \tag{4.6}$$

where the partition function is

$$Z(T,V,N) = \frac{1}{N!h_0^{3N}} \int e^{-H(\Gamma)/k_BT}\, d\Gamma. \tag{4.7}$$

(The integral runs over all of phase space.) This is an example of "partition function", namely the partition function for a pure classical monatomic fluid. It does not apply to mixtures, to crystals, to the ideal paramagnet. In contrast, the definition of "partition function" is equation (4.2), the "sum over all states" of the Boltzmann factor.

4.2 Meaning of the Term "Ensemble"

An ensemble is just a collection. For example, a musical group is an ensemble. Think of a bunch of systems filling up a huge gymnasium.

- Microcanonical ensemble: A collection of many systems, all with the same volume, particle number, particle mass, chemical composition, etc., and each with energy $E < H(x) \leq E + \Delta E$.
- Canonical ensemble: A collection of many systems, all with the same volume, particle number, particle mass, chemical composition, etc., and each connected to the same heat bath (whence each with the same temperature T).

The names (selected by J. Willard Gibbs) are pretty poor.

4.3 Classical Monatomic Ideal Gas

I've told you that it's easier to do calculations in the canonical ensemble than in the microcanonical ensemble. Today I'm going to demonstrate the truth of this assertion.

Remember how we found the entropy $S(E, V, N)$ for a classical monatomic ideal gas? The hard part involved finding the volume of a shell in $3N$-dimensional space. It took us three or four hours to derive the Sackur-Tetrode formula (equation (2.32)), namely

$$S(E, V, N) = k_B N \left[\tfrac{3}{2} \ln \left(\frac{4\pi m E V^{2/3}}{3 h_0^2 N^{5/3}} \right) + \tfrac{5}{2} \right]. \tag{4.8}$$

It took us another hour or so to show that the energy of this system is (equation (2.57))

$$E = \tfrac{3}{2} N k_B T \tag{4.9}$$

whence

$$S(T, V, N) = k_B N \left[\tfrac{3}{2} \ln \left(\frac{2\pi m k_B T V^{2/3}}{h_0^2 N^{2/3}} \right) + \tfrac{5}{2} \right]. \tag{4.10}$$

In this section, we'll derive this same equation using the canonical ensemble. While the derivation is no stroll in the park, most people find it considerably easier than the microcanonical derivation.

Our strategy will be: (1) Integrate the Boltzmann factor over all phase space to find the partition function $Z(T, V, N)$. (2) Find the Helmholtz free energy using

$$F(T, V, N) = -k_B T \ln(Z(T, V, N)). \tag{4.11}$$

And finally (3) use the thermodynamic result

$$S(T, V, N) = -\left. \frac{\partial F}{\partial T} \right)_{V,N} \tag{4.12}$$

to find the entropy.

The Hamiltonian for this model system is

$$H(\Gamma) = \frac{1}{2m}(p_{x,1}^2 + p_{y,1}^2 + p_{z,1}^2 + p_{x,2}^2 + \cdots + p_{z,N}^2) + U(\mathbf{x}_1) + U(\mathbf{x}_2) + \cdots + U(\mathbf{x}_N), \tag{4.13}$$

where the potential energy function for each individual atom as a function of its location $\mathbf{x}$ is

$$U(\mathbf{x}) = \begin{cases} 0 & \text{if } \mathbf{x} \text{ is inside the container} \\ \infty & \text{if outside.} \end{cases} \tag{4.14}$$

This has been background. Now we're ready to roll up our sleeves and plunge into the work. Any questions? Okay, let's go.

The canonical partition function is

$$\begin{aligned} Z(T,V,N) &= \frac{1}{N!h_0^{3N}} \int d\Gamma\, e^{-\beta H(\Gamma)} \qquad (4.15) \\ &= \frac{1}{N!h_0^{3N}} \int_{-\infty}^{+\infty} dx_1 \int_{-\infty}^{+\infty} dy_1 \int_{-\infty}^{+\infty} dz_1 \cdots \int_{-\infty}^{+\infty} dx_N \int_{-\infty}^{+\infty} dy_N \int_{-\infty}^{+\infty} dz_N \\ &\quad \times \int_{-\infty}^{+\infty} dp_{x,1} \int_{-\infty}^{+\infty} dp_{y,1} \int_{-\infty}^{+\infty} dp_{z,1} \cdots \int_{-\infty}^{+\infty} dp_{x,N} \int_{-\infty}^{+\infty} dp_{y,N} \int_{-\infty}^{+\infty} dp_{z,N}\, e^{-\beta H(\Gamma)}. \end{aligned}$$

There are N triplets of integrals over positions and N triplets of integrals over momenta... in total, a $6N$-dimensional integral. Because the Hamiltonian consists of a sum of terms, the Boltzmann factor consists of a product of quantities:

$$e^{-\beta H(\Gamma)} = e^{-\beta(p_{x,1}^2/2m)} e^{-\beta(p_{y,1}^2/2m)} \times \cdots \times e^{-\beta(p_{z,N}^2/2m)} e^{-\beta U(\mathbf{x}_1)} e^{-\beta U(\mathbf{x}_2)} \times \cdots \times e^{-\beta U(\mathbf{x}_N)}. \tag{4.16}$$

More importantly, however, each of these quantities depends on a different set of variables. Therefore the partition function also breaks up into a product of quantities:

$$\begin{aligned} Z(T,V,N) &= \frac{1}{N!h_0^{3N}} \left[\int_{-\infty}^{+\infty} dx_1 \int_{-\infty}^{+\infty} dy_1 \int_{-\infty}^{+\infty} dz_1\, e^{-\beta U(\mathbf{x}_1)} \right] \times \cdots \times \left[\int_{-\infty}^{+\infty} dx_N \int_{-\infty}^{+\infty} dy_N \int_{-\infty}^{+\infty} dz_N\, e^{-\beta U(\mathbf{x}_N)} \right] \\ &\quad \times \left[\int_{-\infty}^{+\infty} dp_{x,1}\, e^{-\beta(p_{x,1}^2/2m)} \right] \times \cdots \times \left[\int_{-\infty}^{+\infty} dp_{z,N}\, e^{-\beta(p_{z,N}^2/2m)} \right]. \end{aligned}$$

The partition function, which in general is a $6N$-dimensional integral, has been reduced in this case to a product of N three-dimensional integrals over position and $3N$ one-dimensional integrals over momentum. (This near-miraculous simplification comes about because all of the limits of integration are constants... if the limits of integration of $p_{y,1}$ were functions of $p_{x,1}$—as they are in the microcanonical ensemble—then it wouldn't happen.)

Moreover, the N position integrals differ only in the dummy variable of integration: they all have the same value. The same holds for the $3N$ integrals over momentum. The partition function is nothing more than

$$\begin{aligned} Z(T,V,N) &= \frac{1}{N!h_0^{3N}} \left[\int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} dy \int_{-\infty}^{+\infty} dz\, e^{-\beta U(\mathbf{x})} \right]^N \\ &\quad \times \left[\int_{-\infty}^{+\infty} dp\, e^{-\beta(p^2/2m)} \right]^{3N}. \end{aligned} \tag{4.17}$$

It is not difficult to evaluate the two integrals remaining. The position integral ranges over all space, but for any point in space outside of the container, its integrand is

$$e^{-\beta U(\mathbf{x})} = e^{-\beta(\infty)} = 0. \tag{4.18}$$

Meanwhile, for any point in space within the container, the integrand is

$$e^{-\beta U(\mathbf{x})} = e^{-\beta(0)} = 1. \tag{4.19}$$

Thus

$$\int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} dy \int_{-\infty}^{+\infty} dz\, e^{-\beta U(\mathbf{x})} = \int\int\int_{\text{container}} dx_1 dy_1 dz_1\, 1 = V. \tag{4.20}$$

Meanwhile, the momentum integral is a clear cousin to the Gaussian integral (equation B.5)

$$\int_{-\infty}^{+\infty} e^{-u^2}\, du = \sqrt{\pi}. \tag{4.21}$$

Use of the substitution $u = \sqrt{\beta/2m}\, p$ in the momentum integral gives

$$\int_{-\infty}^{+\infty} dp\, e^{-\beta(p^2/2m)} = \sqrt{2m/\beta} \int_{-\infty}^{+\infty} e^{-u^2}\, du = \sqrt{2\pi m k_B T}. \tag{4.22}$$

Putting all this together gives us

$$Z(T,V,N) = \frac{V^N}{N! h_0^{3N}} \left(\sqrt{2\pi m k_B T}\right)^{3N} = \frac{1}{N!} \left(\frac{2\pi m k_B T V^{2/3}}{h_0^2}\right)^{3N/2}. \tag{4.23}$$

We've completed the first step of our three-step plan. Now we must find the free energy

$$F(T,V,N) = -k_B T \ln Z = -k_B T \ln \left[\frac{1}{N!} \left(\frac{2\pi m k_B T V^{2/3}}{h_0^2}\right)^{3N/2}\right]. \tag{4.24}$$

Something's a bit funny here. If we double V and N, we should double F, but this doesn't exactly happen: that factor of $N!$ is sure to mess things up. What's gone wrong? We ran into the same problem in when we used the microcanonical ensemble (at equation (2.24)) and it has the same solution: we must take the thermodynamic limit. For large values of N,

$$\begin{aligned} F(T,V,N) &= -k_B T \ln \left[\frac{1}{N!} \left(\frac{2\pi m k_B T V^{2/3}}{h_0^2}\right)^{3N/2}\right] \\ &= -k_B T \left[\ln \left(\frac{2\pi m k_B T V^{2/3}}{h_0^2}\right)^{3N/2} - \ln N!\right] \\ &\approx -k_B T \left[\tfrac{3}{2} N \ln \left(\frac{2\pi m k_B T V^{2/3}}{h_0^2}\right) - N \ln N + N\right]. \end{aligned}$$

But

$$-N \ln N = \tfrac{3}{2} N (-\tfrac{2}{3} \ln N) = \tfrac{3}{2} N \ln(N^{-2/3}),$$

so

$$F(T,V,N) = -k_BTN\left[\tfrac{3}{2}\ln\left(\frac{2\pi mk_BTV^{2/3}}{h_0^2N^{2/3}}\right)+1\right]. \tag{4.25}$$

This feels a lot healthier: If we double both V and N, then the ratio $(V^{2/3}/N^{2/3})$ is unchanged, so the free energy exactly doubles.

Now the final step: Take the derivative with respect to temperature to find the entropy. To do this, recognize that within the square brackets we have

$$\left[\tfrac{3}{2}\ln(T) + \text{ a lot of things independent of } T\right].$$

With this realization the derivative becomes straightforward:

$$S(T,V,N) = -\left.\frac{\partial F}{\partial T}\right)_{V,N} = k_BN\left[\tfrac{3}{2}\ln\left(\frac{2\pi mk_BTV^{2/3}}{h_0^2N^{2/3}}\right)+1\right] + k_BTN\left[\tfrac{3}{2}\frac{1}{T}\right] \tag{4.26}$$

or

$$S(T,V,N) = k_BN\left[\tfrac{3}{2}\ln\left(\frac{2\pi mk_BTV^{2/3}}{h_0^2N^{2/3}}\right)+\tfrac{5}{2}\right]. \tag{4.27}$$

Wow. This is exactly the same as expression (4.10), which came from a (considerably longer) microcanonical derivation.

We've come a long way and it is certainly appropriate to take a moment here to bask in our triumph. But while doing so, a small but troubling question might arise: Why *should* we get the same answer from a canonical and a microcanonical calculation? After all, the ensembles in question are quite distinct: In a microcanonical ensemble each system has the same energy. In a canonical ensemble the individual systems can have any energy they'd like, as reflected by states that range from crystalline to plasma. Why should we get identical entropies from such very different collections?

Loose remark: One often hears that for a gas at temperature T, the probability of an atom having energy E is proportional to the "Boltzman factor" e^{-E/k_BT}. This is true only for non-interacting atoms. If the atoms interact then the kinetic energy belongs to atoms but the potential energy belongs to atom pairs, so it's impossible to define "the energy of an atom". What is always true is that in the canonical ensemble at temperature T, the probability of a system having energy E is proportional to e^{-E/k_BT}.

4.4 Energy Dispersion in the Canonical Ensemble

The systems in the canonical ensemble are not restricted to having just one particular energy or falling within a given range of energies. Instead, systems with any energy from the ground state energy to infinity are present in the ensemble, but systems with higher energies are less probable. In this circumstance, it is important to ask not only for the mean energy, but also for the dispersion (uncertainty, fluctuation, spread, standard deviation) in energy.

Terminology: "Uncertainty" suggests that there's one correct value, but measurement errors prevent your knowing it. (For example: "You are 183.3 ± 0.3 cm tall.") "Dispersion" suggests that there are several

values, each one correct. (For example: "The mean height of people in this room is 172 cm, with a dispersion (as measured by the standard deviation) of 8 cm.") "Fluctuation" is similar to "dispersion," but suggests that the value changes with time. (For example: My height fluctuates between when I slouch and when I stretch.) This book will use the term "dispersion" or, when tradition dictates, "fluctuation." Other books use the term "uncertainty."

The energy of an individual member of the ensemble we call $H(\mathsf{x})$, whereas the average energy of the ensemble members we call E:

$$H(\mathsf{x}) = \text{microscopic energy of an individual system} \tag{4.28}$$

$$E = \langle H(\mathsf{x})\rangle = \text{thermodynamic energy for the ensemble.} \tag{4.29}$$

The dispersion in energy ΔE is given through

$$\begin{aligned}
\Delta E^2 &= \langle (H(\mathsf{x}) - E)^2\rangle && (4.30)\\
&= \langle H^2(\mathsf{x}) - 2H(\mathsf{x})E + E^2\rangle \\
&= \langle H^2(\mathsf{x})\rangle - 2E^2 + E^2 \\
&= \langle H^2(\mathsf{x})\rangle - E^2 \\
&= \langle H^2(\mathsf{x})\rangle - \langle H(\mathsf{x})\rangle^2 && (4.31)
\end{aligned}$$

This relation, which holds for the dispersion of any quantity under any type of average, is worth memorizing. Furthermore it's easy to memorize: The only thing that might trip you up is whether the result is $\langle H^2\rangle - \langle H\rangle^2$ or $\langle H\rangle^2 - \langle H^2\rangle$, but the result must be positive (it is equal to ΔE^2) and it's easy to see that the average of the squares must exceed the square of the averages (consider a list of data containing both positive and negative numbers).

Now it remains to find $\langle H^2(\mathsf{x})\rangle$. Recall that we evaluated $\langle H(\mathsf{x})\rangle$ through a slick trick ("parametric differentiation") involving the derivative

$$\left.\frac{\partial \ln Z}{\partial \beta}\right)_{\text{parameters}}, \tag{4.32}$$

namely

$$\frac{\partial \ln Z}{\partial \beta} = \frac{1}{Z}\frac{\partial Z}{\partial \beta} = \frac{1}{Z}\frac{\partial}{\partial \beta}\left(\sum_{\mathsf{x}} e^{-\beta H(\mathsf{x})}\right) = -\frac{\sum_{\mathsf{x}} H(\mathsf{x}) e^{-\beta H(\mathsf{x})}}{\sum_{\mathsf{x}} e^{-\beta H(\mathsf{x})}} = -E \tag{4.33}$$

The essential part of the trick was that the derivative with respect to β pulls down an $H(\mathsf{x})$ from the exponent in the Boltzmann factor. In order to pull down two factors of $H(\mathsf{x})$, we will need to take two derivatives. Thus the average $\langle H^2(\mathsf{x})\rangle$ must be related to the second-order derivative

$$\left.\frac{\partial^2 \ln Z}{\partial \beta^2}\right)_{\text{parameters}}. \tag{4.34}$$

To see precisely how this works out, we take

$$\begin{aligned}
\frac{\partial^2 \ln Z}{\partial \beta^2} &= -\frac{\left(\sum_{\mathsf{x}} e^{-\beta H(\mathsf{x})}\right)\left(-\sum_{\mathsf{x}} H^2(\mathsf{x}) e^{-\beta H(\mathsf{x})}\right) - \left(\sum_{\mathsf{x}} H(\mathsf{x}) e^{-\beta H(\mathsf{x})}\right)\left(-\sum_{\mathsf{x}} H(\mathsf{x}) e^{-\beta H(\mathsf{x})}\right)}{\left(\sum_{\mathsf{x}} e^{-\beta H(\mathsf{x})}\right)^2} \\
&= \frac{\sum_{\mathsf{x}} H^2(\mathsf{x}) e^{-\beta H(\mathsf{x})}}{\sum_{\mathsf{x}} e^{-\beta H(\mathsf{x})}} - \left(\frac{\sum_{\mathsf{x}} H(\mathsf{x}) e^{-\beta H(\mathsf{x})}}{\sum_{\mathsf{x}} e^{-\beta H(\mathsf{x})}}\right)^2 \\
&= \langle H^2(\mathsf{x})\rangle - \langle H(\mathsf{x})\rangle^2.
\end{aligned}$$

For our purposes, this result is better than we could ever possibly have hoped. It tells us that

$$\Delta E^2 = \frac{\partial^2 \ln Z}{\partial \beta^2} = -\frac{\partial E}{\partial \beta} = -\frac{\partial E}{\partial T}\frac{\partial T}{\partial \beta}. \tag{4.35}$$

To simplify the rightmost expression, note that

$$\frac{\partial E}{\partial T} = C_V \quad \text{and} \quad \frac{\partial \beta}{\partial T} = \frac{\partial (1/k_B T)}{\partial T} = -\frac{1}{k_B T^2},$$

whence

$$\Delta E^2 = k_B T^2 C_V$$

or

$$\Delta E = T\sqrt{k_B C_V}. \tag{4.36}$$

This result is called a "fluctuation-susceptibility theorem".

Analogy: "susceptibility" means "herd instinct". A herd of goats is highly susceptible to external influence, because a small influence (such as a waving handkerchief) will cause the whole herd to stampede. On the other hand a herd of cows is insusceptible to external influence... indeed, cowherds often talk about how hard it is to get the herd into motion. A politician is susceptible if he or she is readily swayed by the winds of public opinion. The opposite of "susceptible" is "steadfast" or "stalwart". If a herd (or politician) is highly susceptible, you expect to see large fluctuations. A herd of goats runs all over its pasture, whereas a herd of cows stays pretty much in the same place.

How does ΔE behave for large systems, that is "in the thermodynamic limit"? Because C_V is intensive, ΔE goes up like $\sqrt{N}$ as the thermodynamic limit is approached. But of course, we expect things to go to infinity in the thermodynamic limit! There will be infinite number, volume, energy, entropy, free energy, etc. The question is what happens to the relative dispersion in energy, $\Delta E/E$. This quantity goes to zero in the thermodynamic limit.

This resolves the question raised at the end of the previous section. A system in a canonical ensemble is *allowed to* have any energy from ground state to infinity. But most of the systems will not make use of that

option: they will *in fact* have energies falling within a very narrow band about the mean energy. For larger and larger systems, this energy band becomes more and more narrow. *This* is why the canonical entropy is the same as the microcanonical entropy.

4.5 Temperature as a Control Variable for Energy (Canonical Ensemble)

Note distinction!

$\mathcal{P}(\mathsf{x})$ is the probability of falling in a particular microstate x. This happens to be a function only of the energy of the microstate, whence this function is often called $\mathcal{P}(H(\mathsf{x}))$. (For a continuous system, $\mathcal{P}(\mathsf{x})$ corresponds to $p(\Gamma)d\Gamma\ldots$ the probability of the system falling in some phase space point within the volume $d\Gamma$ about point Γ.)

$P(H)$ is the probability of having a particular energy. It is equal to $\mathcal{P}(H(\Gamma))\Omega(H)$. ($\Omega(H)$: the number of microstates with energy H.)

Temperature as persuasion.

Temperature as an energy control parameter.

Analogy: "The Jim Smith effect." A man asks a statistician "What is the probability that a man named Milton Abramowitz will win the lottery?" The questioner, whose name is Milton Abramowitz, is disappointed when the statistician's answer is so tiny. So he tries again: "What is the probability that a man named Jim Smith will win the lottery?" The statistician replies with a number that is still small, but not quite so small as the first reply. In response, the man changes his name to Jim Smith.[1]

Because $\mathcal{P}(H(\Gamma))$ decreases rapidly with H, and $\Omega(H)$ increases rapidly with H, we expect $P(H)$ to be pretty sharply peaked near some $\langle H\rangle$ (recall the arguments of section 2.7.3). We also expect this peak to become sharper and sharper as the system becomes larger and larger (approaching "the thermodynamic limit").

Even at high temperature, the most probable microstate is the ground state. However the most probable energy increases with temperature.

In the canonical ensemble (where all microstates are "accessible") the microstate most likely to be occupied is the ground state, and this is true *at any positive temperature, no matter how high.* The ground state energy is not the most probable energy, nor is the ground state typical, yet the ground state is the most probable microstate. In specific, even at a temperature of 1 000 000 K, a sample of helium is more likely to be in a particular crystalline microstate than in any particular plasma microstate. However, there are so many more plasma than crystalline microstates that (in the thermodynamic limit) the sample occupies a plasma macrostate with probability 1.

[1] Only the most ardent students of American history know that Jim Smith signed the Declaration of Independence.

Economic analogy for temperature as an energy control parameter: strict regulation versus incentives. "Market-based approaches work more efficiently than clumsy command-and-control techniques"—Alan S. Blinder, "Needed: Planet Insurance", New York *Times*, 22 October 1997, page A23.

4.1 **Probability of microstate vs. probability of energy**

$\mathcal{P}(H)$ is the probability of being in a particular microstate with energy H, whereas $P(H)\,dH$ is the probability of being in any microstate whose energy falls within the range H to $H + dH$. The number of microstates with energy from H to $H + dH$ is $\Omega(H)\,dH$.

Following the arguments of section 2.7.1 ("Rapidly increasing character of the $\Omega(H)$ function"), assume that $\Omega(H) = cH^\nu$. (For a classical monatomic ideal gas of N atoms, $\nu = (3/2)N$.) Show that under this assumption

$$Z(T, V, N) = c(k_B T)^{\nu+1}\Gamma(\nu + 1). \tag{4.37}$$

Does Z have the correct dimensions? Show that

$$\mathcal{P}(H) = \frac{e^{-H/k_B T}}{c(k_B T)^{\nu+1}\Gamma(\nu + 1)} \quad \text{while} \quad P(H) = \frac{H^\nu e^{-H/k_B T}}{(k_B T)^{\nu+1}\Gamma(\nu + 1)}. \tag{4.38}$$

Sketch $\Omega(H)$, $\mathcal{P}(H)$, and $P(H)$.

4.6 The Equivalence of Canonical and Microcanonical Ensembles

In a microcanonical ensemble the individual systems are all restricted, by definition, to have a given energy. In a canonical ensemble the individual systems are *allowed* to have any energy, from that of the ground state to that of an ionized plasma, but we have seen that (for large systems) they tend not to use this permission and instead they cluster within a narrow band of energies. We might expect, therefore, that a calculation performed in the microcanonical ensemble for systems at a given energy value, say 5 Joules, would give the same results as a calculation performed in the canonical ensemble for systems at whatever temperature corresponds to an average energy of 5 Joules.

If the canonical and microcanonical ensembles give the same results, why bother with the canonical ensemble? We will see soon (section 5.1) that calculations are much easier to perform in the canonical ensemble.

4.7 The Grand Canonical Ensemble

Grand as in the French "big" rather than as in the English "magnificent". This ensemble is "bigger" than the canonical ensemble in that there are more possible microstates.

Summary

The grand canonical ensemble for a pure classical monatomic fluid:

The probability that the system has N particles and is in the microstate Γ_N is proportional to

$$e^{-\beta(H(\Gamma_N)-\mu N)}, \tag{4.39}$$

where

$$\beta = \frac{1}{k_B T}. \tag{4.40}$$

Writing out all the normalizations correctly gives: the probability that the system has N particles and is in some microstate within the phase space volume element $d\Gamma_N$ about Γ_N is

$$\frac{e^{-\beta(H(\Gamma_N)-\mu N)}}{N! h_0^{3N} \Xi(\beta, V, \mu)} d\Gamma_N, \tag{4.41}$$

where the "grand canonical partition function" is

$$\Xi(\beta, V, \mu) = \sum_{N=0}^{\infty} \frac{1}{N! h_0^{3N}} \int e^{-\beta(H(\Gamma_N)-\mu N)}\, d\Gamma_N \tag{4.42}$$

$$= \sum_{N=0}^{\infty} e^{\beta\mu N} Z(\beta, V, N). \tag{4.43}$$

This sum is expected to converge when μ is negative.

The connection to thermodynamics is that

$$\Pi(T, V, \mu) = -p(T, \mu)V = -k_B T \ln \Xi(T, V, \mu). \tag{4.44}$$

4.8 The Grand Canonical Ensemble in the Thermodynamic Limit

This section uncovers results that are of interest in their own right, but it also serves as an example of a mathematically rigorous argument in statistical mechanics.

Definitions. The grand partition function is

$$\Xi(T, V, \alpha) = \sum_{N=0}^{\infty} e^{-\alpha N} Z(T, V, N), \tag{4.45}$$

the Helmholtz free energy is

$$F(T, V, N) = -k_B T \ln Z(T, V, N), \tag{4.46}$$

and the chemical potential is

$$\mu(T, V, N) = \left.\frac{\partial F(T, V, N)}{\partial N}\right)_{T,V}. \tag{4.47}$$

Theorem. *Consider a grand canonical ensemble of hard-core particles at equilibrium with a bath of temperature T and "number control parameter" α. There is a "most probable value of N", called $\hat{N}$ (a function of T, V, and α), satisfying*

$$\alpha = -\frac{\mu(T, V, \hat{N})}{k_B T}, \tag{4.48}$$

such that, in the thermodynamic limit,

$$-k_B T \ln \Xi(T, V, \alpha) \to -\mu(T, V, \hat{N})\hat{N} + F(T, V, \hat{N}). \tag{4.49}$$

Strategy. The theorem assumes hard-core particles so that there will be a maximum number of particles in any given volume, whence the sum (4.45) is finite rather than infinite, and convergence questions do not arise. (Less restrictive conditions—e.g. that the repulsion between two nearby particles increases rapidly enough as they are brought together—may be used instead of the hard-core condition, but then the proof becomes more technical and less insightful.) The proof works by establishing both upper and lower bounds on sum (4.45), and then showing that, in the thermodynamic limit, these two bounds are equal.

Proof. Assume that each hard-core particle has volume v_0. Then the maximum number of particles that can fit into a container of volume V is $N^* \leq V/v_0$. Thus the sum in equation (4.45) does not go to ∞, but stops at N^*.

We will need a trivial mathematical result, called "the method of the maximum term": If f_N is a sequence of positive terms, $N = 0, 1, \ldots N^*$, and if the maximum element of the sequence is $\max\{f_N\}$, then

$$\max\{f_N\} \leq \sum_{N=0}^{N^*} f_N \leq (N^* + 1) \max\{f_N\}. \tag{4.50}$$

Applying this result to the sum (4.45), we obtain

$$\max\{e^{-\alpha N} Z_N\} \leq \Xi \leq (N^* + 1) \max\{e^{-\alpha N} Z_N\}, \tag{4.51}$$

where Z_N is shorthand for $Z(T, V, N)$. Taking the logarithm of each side, and realizing that $\ln \max\{f\} = \max\{\ln f\}$, gives

$$\max\{-\alpha N + \ln Z_N\} \leq \ln \Xi \leq \ln(N^* + 1) + \max\{-\alpha N + \ln Z_N\}. \tag{4.52}$$

To prepare for taking the thermodynamic limit, we divide both sides by V and employ the definition $F_N = -k_B T \ln Z_N$, resulting in

$$\max\left\{-\alpha \frac{N}{V} - \frac{F_N}{k_B T V}\right\} \leq \frac{\ln \Xi}{V} \leq \frac{\ln(N^* + 1)}{V} + \max\left\{-\alpha \frac{N}{V} - \frac{F_N}{k_B T V}\right\}. \tag{4.53}$$

Consider the difference between these upper and lower bounds. It is clear that

$$0 \leq \frac{\ln(N^* + 1)}{V} \leq \frac{\ln(V/v_0 + 1)}{V}. \tag{4.54}$$

In the limit as $V \to \infty$, the right hand side above approaches 0, so

$$\frac{\ln(N^* + 1)}{V} \to 0. \tag{4.55}$$

Thus in the thermodynamic limit the lower bound approaches the upper bound, and

$$\frac{\ln \Xi}{V} \to \max\left\{-\alpha\frac{N}{V} - \frac{F_N}{k_B T V}\right\}. \tag{4.56}$$

Now we need to find the maximum of the quantity in curly brackets above. The maximum is located at $N = \hat{N}$, which we find by taking the partial derivative with respect to N and setting it equal to zero:

$$\frac{\alpha}{V} = -\frac{1}{k_B T V}\left.\frac{\partial F_N}{\partial N}\right]_{N=\hat{N}}. \tag{4.57}$$

Using the definition (4.47), the equation for $\hat{N}$ becomes

$$\alpha = -\frac{\mu(T, V, \hat{N})}{k_B T}. \tag{4.58}$$

Returning to equation (4.56), this location of $\hat{N}$ implies that, for sufficiently large values of V,

$$\frac{\ln \Xi}{V} = -\frac{\mu(T, V, \hat{N})}{k_B T}\frac{\hat{N}}{V} - \frac{F_{\hat{N}}}{k_B T V}, \tag{4.59}$$

from which the final result (4.49) follows immediately.

Resume. The system under study can exchange energy and particles with a heat bath that has "energy control parameter" $\beta = 1/k_B T$ and "number control parameter" α. The probability that the system contains exactly N particles is

$$P_N = \frac{e^{-\alpha N} Z(\beta, V, N)}{\Xi(\beta, V, \alpha)}. \tag{4.60}$$

The proof shows that, in the thermodynamic limit, there is one particular value of N, namely $\hat{N}$, for which P_N approaches 1. All the other P_N's, of course, approach 0. The condition for locating $\hat{N}$ is just that it gives the maximum value of P_N... this condition gives rise to equation (4.48). Once $\hat{N}$ is located, the result $P_{\hat{N}} \to 1$ becomes equation (4.49).

4.9 Summary of Major Ensembles

	boundary	variables	probability of microstate	p.f.	master function
microcanonical	adiabatic (no-skid)	E, V, N	$\frac{d\Gamma}{N! h_0^{3N}}\frac{1}{\Omega}$ or 0	Ω	$S(E, V, N) = k_B \ln \Omega$
canonical	heat bath	T, V, N	$\frac{d\Gamma e^{-\beta H(\Gamma)}}{N! h_0^{3N}}\frac{1}{Z}$	Z	$F(T, V, N) = -k_B T \ln Z$
grand canonical	heat bath, with holes	T, V, μ	$\frac{d\Gamma_N e^{-\beta H(\Gamma_N) - \alpha N}}{N! h_0^{3N}}\frac{1}{\Xi}$	Ξ	$\Pi(T, V, \mu) = -k_B T \ln \Xi$

In all cases, the partition function (p.f. in the above table) is the normalization factor

$$\text{p.f.} = \sum_{\text{microstates}} \text{unnormalized probability.}$$

4.10 Quantal Statistical Mechanics

How shall we define a partition function for quantal systems? A reasonable first guess is that

$$Z(T, \text{parameters}) = \sum_{\text{all quantal states}} e^{-\beta E}. \qquad [\![\text{first guess}]\!]$$

This guess runs into problems immediately. Most quantal states are not energy eigenstates, so it's not clear how to interpret "E" for such states. Another difficulty—harder to see—arises because there are many more quantal states than classical states, so the guess above is in some sense counting too many states. It's hard to see how it could have the correct classical limit. (Indeed, a mathematical investigation of the above guess would reveal that it doesn't even have the correct thermodynamic limit—if you were to calculate the Helmholtz free energy from the above guess, the resulting $F(T, V, N)$ would increase far faster than linearly with system size.)

A second guess might be

$$Z(T, \text{parameters}) = \sum_{\text{all energy eigenstates}} e^{-\beta E}. \qquad [\![\text{second guess}]\!]$$

This avoids the problem of how to interpret E for non-energy eigenstates, but there still seems to be an over-counting of states in the case of an energy degeneracy: If two energy eigenstates are separate in energy by any amount, no matter how small, those two energy eigenstates will contribute two terms to the above sum. But if the energy separation vanishes, then all linear combinations of the two states are also energy eigenstates, and each of this infinite number of combinations will enter into the sum.

Finally we arrive at a workable definition for quantal partition function, namely

$$Z(T, \text{parameters}) = \sum_{\text{energy eigenbasis } n} e^{-\beta E_n}. \tag{4.61}$$

Note that this is a sum over an *energy eigenbasis*, not over *energy eigenvalues*. These two sums differ through degeneracy: If $E_6 = E_5$, then two equal terms enter into the sum. More formally: if $N(m)$ is the degeneracy of eigenvalue m, that is, the number of linearly independent states with energy E_m, then

$$Z(T, \text{parameters}) = \sum_{\text{energy eigenvalues } m} N(m) e^{-\beta E_m}. \tag{4.62}$$

The above argument has a strange, *ex post facto* character: We define the quantal partition function not from first principles, but to insure that it has properties that we consider desirable (namely, that it avoids subtle interpretation issues, it posses the expected classical limit, and it has desirable behavior at

level crossings). Linear combinations of energy eigenstates are quantal states just as legitimate as energy eigenstates, so why do we ignore them in the sum? The argument reminds me of politics at its worst: "I'm going to pass this law because I want it, even though I can't support it with any rationale." The resulting definition of partition function is strange in that it seems to select the energy eigenbasis as some sort of God-given basis, "better" than any other basis. In fact, the result is not as bad as it seems. It turns out (see problem 4.2) that our definition can be rephrased in a basis-independent manner as

$$Z(T, \text{parameters}) = \sum_{\text{energy eigenbasis } n} e^{-\beta E_n} = \sum_{\text{any basis } j} \langle j|e^{-\beta \hat{H}}|j\rangle. \tag{4.63}$$

If you don't know what it means to exponentiate the Hamiltonian operator $\hat{H}$, don't worry. For this book—and for most practical calculations—the most effective definition is the form on the left. The point is simply that the definition of partition function *can* be cast into the basis-independent form on the right. The energy eigenbasis is *not* special.

In this chapter we have often found, first, results about probabilities, and second, connections to thermodynamics. This section has so far dealt with only the second element. What do these results imply for probabilities? I will often say things like "The probability of having energy E_m is

$$\frac{N(m)e^{-\beta E_m}}{Z}."$$

On the face of it, this statement is absurd, because most of the microstates aren't energy eigenstates so they don't have any energy at all. What I mean is "If the energy is measured, the probability of finding energy E_m is

$$\frac{N(m)e^{-\beta E_m}}{Z}."$$

I will often use the first statement rather than the long and formal second statement. I do so, however, fully realizing that the first statement is wrong and that I use it just as convenient shorthand for the second statement. You may use this shorthand too, as long as you don't mean it.

> The ultimate source of such problems is that the English language was invented by people who did not understand quantum mechanics, hence they never produced concise, accurate phrases to describe quantal phenomena. In the same way, the ancient phrase "Search the four corners of the Earth" is still colorful and practical, and is used today even by those who know that the Earth isn't flat.

Similarly, I will often say "The probability of being in the energy eigenstate $\eta_n(\mathbf{x}_1, \ldots, \mathbf{x}_N)$ with energy E_n is

$$\frac{e^{-\beta E_n}}{Z}."$$

But what I really mean is "If the microstate is projected onto a basis which includes the energy eigenstate $\eta_n(\mathbf{x}_1, \ldots, \mathbf{x}_N)$, then the probability of projecting onto that state is

$$\frac{e^{-\beta E_n}}{Z}."$$

4.2 **The trace**

Show that for any operator $\hat{A}$, the "trace" of $\hat{A}$, defined as

$$\text{trace}\{\hat{A}\} \equiv \sum_{\text{any basis } j} \langle j|\hat{A}|j\rangle, \tag{4.64}$$

is independent of basis. What is the trace in terms of the matrix representation of the operator?

4.3 **Entropy in the canonical ensemble**

From the partition function (4.61), one can find the free energy F and hence the entropy S. Show that the entropy obtained in this manner equals

$$S(T) = -k_B \sum_n p_n \ln p_n, \tag{4.65}$$

where

$$p_n = \frac{e^{-\beta E_n}}{Z(T)} \tag{4.66}$$

is the probability that the system is in its nth energy eigenstate (using the shorthand mentioned above). This expression for the entropy is particularly useful in information theory. (Clue: Work forward from $S = -\partial F/\partial T$ and backwards from equation (4.65) to meet at the middle.)

4.4 **Entropy in the microcanonical ensemble**

[You must work the previous problem before working this one.] What are the state probabilities p_n, analogous to (4.66), for an system in the microcanonical ensemble? Devise a suitable quantal replacement for the microcanonical definition of entropy (2.7). Cast your replacement into the form of a function of the microcanonical probabilities p_n. Is formula (4.65) correct in the microcanonical ensemble?

4.11 Ensemble Problems I

4.5 **Classical monatomic ideal gas in the canonical ensemble**

In section 5.1 we will show that the canonical partition function of a classical monatomic ideal gas is

$$Z(T, V, N) = \frac{1}{N!}\left[\frac{V}{\lambda^3(T)}\right]^N, \tag{4.67}$$

where

$$\lambda(T) \equiv \frac{h_0}{\sqrt{2\pi m k_B T}}. \tag{4.68}$$

a. Show that in the thermodynamic limit, the Helmholtz free energy per particle is

$$\frac{F(T, V, N)}{N} = -k_B T\left[\ln\left(\frac{V/N}{\lambda^3(T)}\right) + 1\right]. \tag{4.69}$$

b. Differentiate with respect to T to find $S(T, V, N)$.

c. Change variables to $S(E, V, N)$ using $E = \frac{3}{2}Nk_BT$, and compare the resulting expression to the entropy derived in class through the microcanonical ensemble.

4.6 **Number fluctuations in the grand canonical ensemble**

Calculate ΔN, the root mean square fluctuations in the particle number, for a system in the grand canonical ensemble. (Clue: Take derivatives with respect to μ.) Show that this result is related to the isothermal compressibility κ_T through

$$\frac{\Delta N}{N} = \sqrt{\frac{k_BT\kappa_T}{V}}. \tag{4.70}$$

(Clue: Use the results of problem 3.33.)

4.7 **Classical monatomic ideal gas in the grand canonical ensemble**

a. Use the fact that

$$Z(T, V, N) = \frac{1}{N!}\left[\frac{V}{\lambda^3(T)}\right]^N, \tag{4.71}$$

to find $\Xi(T, V, \mu)$ for the classical monatomic ideal gas.

b. Use the connection to thermodynamics for the grand canonical ensemble, namely

$$p(T, \mu)V = k_BT \ln \Xi(T, V, \mu), \tag{4.72}$$

to show that for any substance

$$N(T, V, \mu) = k_BT \left.\frac{\partial \ln \Xi}{\partial \mu}\right)_{T,V}. \tag{4.73}$$

c. Derive the ideal gas equation of state in the grand canonical ensemble.

4.8 **Isobaric ensemble**

This problem encourages you to think about an ensemble with a specified temperature, pressure, and particle number. A physical model to keep in mind is a collection of systems each with the same number of particles and all connected to the same heat bath but enclosed in balloons rather than in rigid boxes. I want you to guess the answers to the following questions rather than produce mathematical or verbal arguments. If you are uncomfortable making guesses in a physics problem assignment then do only the first two parts.

a. Given that T, p, and N are fixed, what quantities fluctuate?

b. In going from the canonical to the isobaric ensemble, what control parameter (comparable to $\beta = 1/k_BT$ or $\alpha = -\mu/k_BT$) must be introduced, and how is it related to the pressure?

c. What is the (unnormalized) probability of finding the system in microstate x with energy $H(x)$ and volume $V(x)$?

d. What is the relevant state sum (comparable to Z or Ξ), and how is it related to the thermodynamic master function $G(T, p, N)$?

e. The relevant fluctuation-susceptibility relation connects the fluctuations in volume to what susceptibility?

f. Write out the fluctuation-susceptibility relation, using B to represent a product of several unguessable intensive quantities but writing out the extensive quantities explicitly.

g. What are the dimensions of B? Can you use this knowledge to guess at the unguessable?

4.9 **Fluctuation-susceptibility relation for a magnetic system**

In the Ising model for a magnet, the spins are located at fixed sites (labeled by i), and can point either up or down ($s_i = \pm 1$). The Ising Hamiltonian is

$$\mathcal{H} = \mathcal{H}_0(s_1, \ldots, s_N) - mH \sum_{i=1}^{N} s_i, \tag{4.74}$$

where m is the magnetic moment of each spin, H is the applied magnetic field, and $\mathcal{H}_0$ represents some spin-spin interaction energy, the details of which are not important here except for the fact that it is independent of H. The microscopic magnetization, which varies from configuration to configuration, is

$$\mathcal{M}(s_1, \ldots, s_N) = m \sum_i s_i, \tag{4.75}$$

while the macroscopic (or thermodynamic) magnetization is the canonical

average over all such microscopic magnetizations:

$$M(T, H) = \langle \mathcal{M} \rangle. \tag{4.76}$$

The magnetic susceptibility—easily measured experimentally—is

$$\chi_T(T, H) = \left.\frac{\partial M}{\partial H}\right)_T. \tag{4.77}$$

Show that in the canonical ensemble the fluctuations in magnetization are related to the susceptibility by

$$\Delta M = \sqrt{k_B T \chi_T}. \tag{4.78}$$

4.10 **Two definitions of magnetization**

The previous problem gave a statistical mechanical definition of magnetization:

$$M(T, H) = \langle \mathcal{M} \rangle, \tag{4.79}$$

while equation (3.100) gave a thermodynamic definition:

$$M(S, H) = -\left.\frac{\partial E}{\partial H}\right)_S. \tag{4.80}$$

Show that these two definitions give equivalent results.

4.11 **Values for the approach to the thermodynamic limit**

In problem 2.13, "The approach to the thermodynamic limit," you were given a value of $EV^{2/3}/h_0^2$ and told that it was appropriate "for a gas at room temperature and atmospheric pressure." Justify this value as

$$EV^{2/3}/h_0^2 = [\tfrac{3}{2}(k_BT)^{5/3}/(p^{2/3}h^2)]N^{5/3} \tag{4.81}$$

4.12 **Integrals by parametric differentiation**

To find the mean energy or energy dispersion in the canonical ensemble, we introduced the "slick trick" of differentiating with respect to β. This trick, called parametric differentiation, is in fact useful in a number of circumstances. For example, in a graduate electrodynamics class I was once assigned a problem (Jackson, *Classical Electrodynamics*, problem 14.5(b)) for which I needed to evaluate

$$\int_0^{2\pi} \frac{\cos^2\theta}{(1+b\sin\theta)^5}d\theta \tag{4.82}$$

where b is a constant with $|b| < 1$. My classmates and I tried everything: substitutions, partial fractions, conversion to a contour integral in the complex plane, and, most popular of all, "look it up in reference books". I puzzled over it for a dozen hours before giving up. When the solutions were handed out, I expected to find many pages devoted to evaluating the integral. Instead I found a two-line argument that began with the known integral (see Dwight 858.546 or Gradshteyn and Ryzhik 3.644.4)

$$\int_0^{2\pi} \frac{\cos^2\theta}{a+b\sin\theta}d\theta = \frac{2\pi}{a+\sqrt{a^2-b^2}} \tag{4.83}$$

and then took a fourth derivative. What was the argument, and what is the value of the integral?

4.13 **Parametric differentiation in quantum mechanics**

The previous problem showed how parametric differentiation could be useful in electrodynamics. This one shows how it can be useful in quantum mechanics. In quantum mechanics, an infinite square well of width π has energy eigenfunctions

$$\eta_n(x) = \sqrt{\frac{2}{\pi}}\sin(nx) \quad n = 1, 2, 3, \ldots. \tag{4.84}$$

One frequently needs to evaluate matrix elements such as

$$\langle n|x^2|m\rangle = \frac{2}{\pi}\int_0^{\pi} \sin(nx)x^2\sin(mx)\,dx. \tag{4.85}$$

Show how this integral—as well as a host of other useful matrix elements—can be obtained easily from the well known result

$$\int_0^{\pi} \sin(ax)\sin(bx)\,dx = \frac{1}{2}\left[\frac{\sin(a-b)\pi}{a-b} - \frac{\sin(a+b)\pi}{a+b}\right] \quad a \neq \pm b. \tag{4.86}$$

4.14 **Polymers**

A primitive model for a polymer is a random walk on a simple cubic lattice. A random walk consists of n steps (or "links") starting at ("anchored to") the origin. (In this model a polymer unit can step back onto a lattice site already occupied by a different polymer unit. This unrealistic feature is corrected in a more sophisticated model, the so-called "self avoiding walk.")

a. Show that the number of distinct walks consisting of n links is $N_n = 6^n$. Does this formula hold when $n = 0$?

For many purposes it is valuable to consider the ensemble of *all* random walks, regardless of their size. In this ensemble there is a "size control parameter" α such that the probability of finding a walk x consisting of $n(\mathsf{x})$ links is proportional to $e^{-\alpha n(\mathsf{x})}$. (Thus *longer* walks are *less* probable in this ensemble, but there are more of them.) The partition function associated with this model is

$$\Xi(\alpha) = \sum_{\text{walks } \mathsf{x}} e^{-\alpha n(\mathsf{x})}. \tag{4.87}$$

b. Show that the mean walk size in this ensemble is a function of α given through

$$\langle n \rangle = -\frac{\partial \ln \Xi(\alpha)}{\partial \alpha}. \tag{4.88}$$

c. Show that

$$\Xi(\alpha) = \frac{1}{1 - 6e^{-\alpha}} \quad \text{and that} \quad \langle n \rangle = \frac{6}{e^{\alpha} - 6}. \tag{4.89}$$

Clue: The geometric series sums to

$$1 + x + x^2 + x^3 + \cdots = \frac{1}{1 - x} \quad \text{when} \quad |x| < 1.$$

d. What is the smallest possible value of the control parameter α? Does large α correspond to long polymers or short polymers?

e. Show that the dispersion in n is given through

$$(\Delta n)^2 = \frac{\partial^2 \ln \Xi}{\partial \alpha^2}, \tag{4.90}$$

whence

$$\frac{\Delta n}{\langle n \rangle} = \sqrt{\frac{e^{\alpha}}{6}} = \sqrt{\frac{1}{\langle n \rangle} + 1}. \tag{4.91}$$

Thus the relative dispersion decreases for longer polymers.

(See also loops: S. Leibler, R.R.P. Singh, and M.E. Fisher, "Thermodynamic behavior of two-dimensional vesicles," *Phys. Rev. Lett.* **59** (1987) 1989–1992; C. Richard, A.J. Guttmann, and I. Jensen, "Scaling function and universal amplitude combinations for self-avoiding polygons," *J. Phys.* A **34** (2001) L495–501.)

4.15 **A class of new ensembles**

Any mechanical parameter in the Hamiltonian can be "traded in" for a control parameter. This problem details the procedure if the mechanical parameter is intensive and an intrinsically positive quantity. Examples to keep in mind are 1) the particle mass, 2) the spacing between planes of a high temperature superconductor, or 3) the natural frequency of vibration for a diatomic molecule modeled as a simple harmonic oscillator. In honor of this last example I will call the parameter ω. In the "micro-ω" ensemble, all of the systems in the ensemble have a particular value of the parameter. In the "grand-ω" ensemble, the systems in the ensemble have a variety of values for the parameter, but larger values are less probable. In fact, the probability is proportional to $e^{-\gamma N\omega}$, where γ is a "control parameter" which can be adjusted to set the mean value of ω to any desired positive value. We suspect that, in the thermodynamic limit, the results of the two ensembles will be identical when the mean value of ω in the grand-ω ensemble is the same as the fixed value of ω in the micro-ω ensemble.

a. The thermodynamic assembly corresponding to the micro-ω ensemble has as master function the Helmholtz free energy $F(T, V, N, \omega)$. Using the definition

$$B(T, V, N, \omega) = -\left.\frac{\partial F}{\partial \omega}\right)_{T,V,N}, \tag{4.92}$$

write out the differential equation for dF.

b. Execute a Legendre transformation to an assembly with master function

$$K(T, V, N, B) = F + B\omega, \tag{4.93}$$

and show that in this assembly, the function (not variable!) ω is

$$\omega(T, V, N, B) = \left.\frac{\partial K}{\partial B}\right)_{T,V,N}. \tag{4.94}$$

In the grand-ω ensemble, the probability that a system has a parameter falling within $d\omega$ about ω is

$$\frac{e^{-\gamma N\omega} Z(T, V, N, \omega)\, d\omega}{\int_0^\infty e^{-\gamma N\omega'} Z(T, V, N, \omega')\, d\omega'}. \tag{4.95}$$

Define the partition function

$$Y(T, V, N, \gamma) = \frac{1}{\omega_0}\int_0^\infty e^{-\gamma N\omega} Z(T, V, N, \omega)\, d\omega, \tag{4.96}$$

where ω_0 is an arbitrary constant with the dimensions of ω whose only purpose is to make Y dimensionless. Presumably, the relationship between thermodynamics and statistical mechanics in the grand-ω ensemble is

$$K = -k_B T \ln Y, \tag{4.97}$$

by analogy to the similar relation in the canonical ensemble. We will assume this relationship and use it, first to find the relationship between the statistical mechanical γ and the thermodynamic B and second to show that the statistical mechanical equation (4.96) is consistent with the thermodynamic equation (4.93).

c. Show that, in the grand-ω ensemble,

$$\langle\omega\rangle = -\frac{1}{N}\frac{\partial \ln Y}{\partial \gamma} \tag{4.98}$$

and

$$\Delta\omega^2 = \frac{1}{N^2}\frac{\partial^2 Y}{\partial \gamma^2}. \tag{4.99}$$

Note that $\langle\omega\rangle$ and $\Delta\omega$ are functions of T, V, N, and γ.

d. Compare equations (4.94) and (4.98) and conclude that, if (4.97) is to hold, then

$$B = Nk_BT\gamma. \tag{4.100}$$

e. Show that

$$\Delta\omega = \sqrt{k_BT\left.\frac{\partial\omega}{\partial B}\right)_{T,V,N}}. \tag{4.101}$$

f. We suspect that, in the thermodynamic limit $N, V \to \infty$, $\ln Y \sim N$. If true, show that $\langle\omega\rangle$ is intensive (independent of N) whereas $\Delta\omega$ falls like $1/\sqrt{N}$.

g. To evaluate $Y(\gamma)$ in the thermodynamic limit, write

$$Y(T,V,N,\gamma) = \frac{1}{\omega_0}\int_0^\infty e^{-\gamma N\omega} Z(T,V,N,\omega)\, d\omega \approx e^{-\gamma N\langle\omega\rangle} Z(T,V,N,\langle\omega\rangle)\frac{\Delta\omega}{\omega_0}. \tag{4.102}$$

Take logarithms to show that

$$\ln Y \approx -\gamma N\langle\omega\rangle + \ln Z + \ln(\Delta\omega/\omega_0), \tag{4.103}$$

and argue that in the thermodynamic limit the last term becomes negligible. Finally conclude that

$$K = k_BT\gamma N\langle\omega\rangle + F \tag{4.104}$$

or, in light of relationship (4.100),

$$K = F + B\langle\omega\rangle, \tag{4.105}$$

which should be compared to equation (4.93).

We have shown only that this ensemble scheme is "not inconsistent". It is not obviously wrong, but of course this does not prove that it is right. A flawless general proof would require detailed mathematical reasoning, but proofs for particular situations are outlined in the next problem and in problem 4.18.

4.16 **The grand-ω ensemble for the classical monatomic ideal gas**

Apply the concepts of the previous problem to the classical monatomic ideal gas where the mechanical parameter ω is the particle mass. In particular, find expressions

a. for $B(T,V,N,m)$,

b. for $m(T,V,N,B)$,

c. and for $\langle m \rangle$ and Δm as functions of T, V, N, and γ.

4.17 **An ensemble of your own**

Any mechanical parameter in the Hamiltonian can be traded in for a control parameter. Can you think up a good problem *using* this fact where the parameter is, say, an applied magnetic field? Or the spring constant (or relaxed length) between two atoms in a diatomic molecule (or in the Einstein model)? Or the moment of inertia of a diatomic molecule modeled as a dumbbell? Or a parameter in the Lennard-Jones potential? Or the size of hard-sphere atoms? Or the spacing between atoms in a crystal?

4.12 Ensemble Problems II

The remaining problems in this chapter deal with the same "principles" issues that the others do, but they assume some familiarity with physical and mathematical topics that we have not yet treated. I place them here because of their character, but I do not expect you to do them at this moment. Instead I list their prerequisites and I hope you will come back to them later.

4.18 **The grand-ω ensemble for the simple harmonic oscillator**

(Prerequisite: Problems 4.15 and 5.7.) Problem 4.15 introduced a class of ensembles in which mechanical parameters in the Hamiltonian were not fixed but rather allowed to fluctuate under the control of some parameters γ. Apply this concept to the diatomic molecules of problem 5.7, using the natural frequency of vibration as the mechanical parameter. Develop the suggestive arguments of problem 4.15 into a mathematically rigorous proof.

4.19 **Particles in a bowl**

(Prerequisite: Chapter 5.) I can't count how many times I've waved my hands and told you that the precise character of the boundaries should be irrelevant to the bulk properties found by statistical mechanics. Now you have a chance to prove it (for a restricted situation of course).

a. Consider N non-interacting, classical point particles moving subject to the potential energy function $U(\mathbf{r}) = Kr^2/2$. Evaluate the classical partition function $Z_b(T, K, N)$ and compare it to the partition function $Z_h(T, V, N)$ for N particles in a hard-walled container of volume V.

b. For what V (as a function of K) will the two partition functions in part (a.) be the same?

c. The "bowl" container has no stiff walls, so it is possible for a particle to be located at any distance from the origin. But is it not likely that a particle will get too far away. Calculate the rms radius $\sqrt{\langle r^2 \rangle}$. (Use equipartition?)

d. Suppose the hard-walled container is spherical with radius R. Place the origin at the center of the sphere and find the rms radius $\sqrt{\langle r^2 \rangle}$.

e. For the V corresponding to K in the sense of part (b.), compare the two rms radii.

Extra: Do this problem quantum mechanically. (Clue: Notice the mathematical similarity between this problem and the Einstein model of lattice vibrations.)

4.20 **Quantal monatomic ideal gas**

(Prerequisite: Chapter 6.) We have many times considered the problem of N non-interacting classical point particles in a box of edge length $L = V^{1/3}$. Now we will solve the problem for quantum mechanical particles.

The partition function is $Z = \sum e^{-\beta E}$, where the sum is taken over the many-particle energy eigenstates $\Psi(\mathbf{r}_1, \ldots, \mathbf{r}_N)$. Ignore symmetry-under-interchange requirements so that such states are specified by a simple listing of the component one-particle levels (or orbitals) $\eta_{\mathbf{k}}(\mathbf{r})$. Thus Ψ can be specified by simply listing the relevant $\mathbf{k}$ values of its component levels. Show then that

$$Z = \frac{1}{N!}\left(\sum_{\mathbf{k}} e^{-\beta E_k}\right)^N \tag{4.106}$$

where $E_k = \hbar^2 k^2/2m$ and the sum is over all $\mathbf{k}$ permitted by periodic boundary conditions. Evaluate the sum (in the thermodynamic limit) by turning it into an integral. Compare the resulting partition function with that of a classical monatomic ideal gas.

4.21 **Quantal monatomic ideal gas in the microcanonical ensemble**

Consider the situation of the above problem but calculate the microcanonical partition function $\Omega(E, V, N)$. Compare with the classical result.

Chapter 5

Classical Ideal Gases

End of principles, beginning of applications. There is still more that could be done with principles, but "the proof of the pudding is in the eating". And in this case the pudding is excellent eating indeed... calculations made from statistical mechanics theory give results that are actually more accurate than experiment.

This chapter considers ideal (i.e. non-interacting) gases made up of atoms or molecules that may have internal structure (i.e. not point particles). The internal degrees of freedom will be treated either classically or quantally, but the translational degrees of freedom will always be treated classically.

5.1 Classical Monatomic Ideal Gases

We have already found the Helmholtz free energy of the classical monatomic ideal gas (section 4.3). I think you will agree that the canonical calculation is considerably easier than the corresponding microcanonical calculation. Here we will review the calculation, then go back and investigate just what caused the solution to be so much easier, and this investigation will lead to an important theorem.

5.1.1 Solving the problem

The energy

$$H(\Gamma) = \frac{1}{2m}(p_{x,1}^2 + p_{y,1}^2 + p_{z,1}^2 + p_{x,2}^2 + \cdots + p_{z,N}^2) + U(\mathbf{x}_1) + U(\mathbf{x}_2) + \cdots + U(\mathbf{x}_N) \tag{5.1}$$

is a sum of terms, each involving different variables, so the Boltzmann factor

$$e^{-\beta H(\Gamma)} = e^{-\beta(p_{x,1}^2/2m)} e^{-\beta(p_{y,1}^2/2m)} \times \cdots \times e^{-\beta(p_{z,N}^2/2m)} e^{-\beta U(\mathbf{x}_1)} e^{-\beta U(\mathbf{x}_2)} \times \cdots \times e^{-\beta U(\mathbf{x}_N)} \tag{5.2}$$

is a product of factors, each involving different variables. Thus the partition function, which in general is an integral in a $6N$-dimensional space, factors into 3- and 1-dimensional integrals

$$\begin{aligned} Z(T,V,N) &= \frac{1}{N!h^{3N}}\left[\int_{-\infty}^{+\infty} dx_1 \int_{-\infty}^{+\infty} dy_1 \int_{-\infty}^{+\infty} dz_1\, e^{-\beta U(\mathbf{x}_1)}\right] \times \cdots \times \left[\int_{-\infty}^{+\infty} dx_N \int_{-\infty}^{+\infty} dy_N \int_{-\infty}^{+\infty} dz_N\, e^{-\beta U(\mathbf{x}_N)}\right] \\ &\times \left[\int_{-\infty}^{+\infty} dp_{x,1}\, e^{-\beta(p_{x,1}^2/2m)}\right] \times \cdots \times \left[\int_{-\infty}^{+\infty} dp_{z,N}\, e^{-\beta(p_{z,N}^2/2m)}\right]. \end{aligned}$$

This doesn't happen in the microcanonical calculation!

Moreover, we found it easy to evaluate the N three-dimensional position integrals and the $3N$ one-dimensional momentum integrals. The result was

$$Z(T,V,N) = \frac{V^N}{N!h^{3N}}\left[\sqrt{2\pi m k_B T}\right]^{3N}. \tag{5.3}$$

The result above is correct, but it is not the most convenient form for the partition function. There are lots of variables and constants floating around, many of them with hard-to-remember dimensions. The quantity Z is in fact dimensionless, but that's not obvious from the expression. For this reason is customary to define the quantity

$$\lambda(T) = \frac{h}{\sqrt{2\pi m k_B T}}, \tag{5.4}$$

called the "thermal de Broglie wavelength". Like any wavelength, it has the dimensions of length, and the notation $\lambda(T)$ suggests this. It is, of course, an intensive quantity. In terms of the thermal de Broglie wavelength, the partition function for the classical monatomic ideal gas is

$$Z(T,V,N) = \frac{1}{N!}\left(\frac{V}{\lambda^3(T)}\right)^N. \tag{5.5}$$

If a particle of mass m has kinetic energy $k_B T$, then it has momentum $\sqrt{2mk_BT}$ and thus a quantum-mechanical de Broglie wavelength of

$$\lambda = \frac{h}{p} = \frac{h}{\sqrt{2mk_BT}}. \tag{5.6}$$

The definition (5.4) above slips in an extra dimensionless factor of $\sqrt{\pi} = 1.77$ just to make the result (5.5) easier to remember and to work with. Physically, the thermal de Broglie wavelength represents the quantal smearing of each individual point particle. Thus if the mean separation between atoms is much greater than $\lambda(T)$, we expect that the classical approximation will be a good one. If the mean separation is less than or about equal to $\lambda(T)$, we expect that a quantal treatment will be required.

After doing any calculation is it useful to perform a "debriefing" to figure out why—in general rather than technical terms—the calculation went the way it did. Almost everyone agrees that this canonical calculation is a lot easier than the corresponding microcanonical calculation to find $\Omega(E,V,N)$ for the classical monatomic ideal gas. But what was the particular feature that made it so much easier? Surely it was the factorization of the Boltzmann factor and the subsequent factorization of the partition function. This enabled us to perform one- and three-dimensional integrals rather than the $3N$-dimensional integral required (see appendix D) for the microcanonical calculation. Is there anything that can be said more generally about this feature?

5.1.2 Theorem on decoupling Hamiltonians

Indeed there is. Suppose a Hamiltonian is a sum of two pieces

$$H(\Gamma) = H_1(\Gamma_1) + H_2(\Gamma_2), \tag{5.7}$$

where Γ_1 and Γ_2 are *exclusive.* (That is, the phase space variables in the list Γ_1 and the phase space variables in the list Γ_2 together make up the whole list Γ, and no variable appears in both lists.) The Hamiltonian $H(\Gamma)$ is then said to "decouple" into the two parts $H_1(\Gamma_1)$ and $H_2(\Gamma_2)$.[1] Then it is easy to see that the partition function will factor into two parts:

$$Z(T,V,N) = \frac{1}{N!h^{3N}} \int d\Gamma\, e^{-\beta H(\Gamma)} = \frac{1}{N!h^{3N}} \int d\Gamma_1\, e^{-\beta H(\Gamma_1)} \int d\Gamma_2\, e^{-\beta H(\Gamma_2)}. \tag{5.8}$$

If the Hamiltonian decouples into three parts, then this reasoning can be used to show that the partition function breaks into three factors. And so forth.

For example, the Hamiltonian of the classical monatomic ideal gas that we have just investigated decouples into $4N$ pieces,

$$H(\Gamma) = \frac{p_{x,1}^2}{2m} + \frac{p_{y,1}^2}{2m} + \cdots + \frac{p_{z,N}^2}{2m} + U(\mathbf{x}_1) + U(\mathbf{x}_2) + \cdots + U(\mathbf{x}_N). \tag{5.9}$$

Each term of the form $p^2/2m$ in the Hamiltonian leads to a factor of $h/\lambda(T)$ in the partition function... each term of the form $U(\mathbf{x})$ leads to a factor of V. Thus our theorem on decoupling Hamiltonians shows that the partition function is

$$Z(T,V,N) = \frac{1}{N!h^{3N}} \left(\frac{h}{\lambda(T)}\right)^{3N} V^N. \tag{5.10}$$

If the particles *do* interact, then the potential energy part of the Hamiltonian does not break up into exclusive pieces, but the kinetic energy part still does. Thus for an interacting gas with potential energy of interaction

$$U_N(\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_N), \tag{5.11}$$

the partition function is

$$Z(T,V,N) = \frac{1}{N!\lambda^{3N}(T)} \int_{-\infty}^{+\infty} dx_1 \cdots \int_{-\infty}^{+\infty} dz_N\, e^{-\beta U_N}. \tag{5.12}$$

Although the potential energy part of the problem is now excruciatingly difficult, our theorem on decoupling Hamiltonians assures us that the kinetic energy part is just as easy to handle for an interacting as for a non-interacting gas.

5.1 **Gas with pair interactions**

For a pure classical monatomic gas that interacts only through pair interactions (i.e. for which there are no three-body or higher interactions) the Hamiltonian is

$$H(\Gamma) = \sum_{i=1}^{N} \frac{\mathbf{p}_i^2}{2m} + \sum_{i=1}^{N} U_A(\mathbf{x}_i) + \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} U_B(\mathbf{x}_i, \mathbf{x}_j), \tag{5.13}$$

[1] Alternatively, the two parts of the Hamiltonian are called "uncoupled". These terms derive from the ordinary English word "to couple", meaning to join or to connect.

where $U_A(\mathbf{x})$ is the potential energy due to the interaction between a particle at point $\mathbf{x}$ and the container, and $U_B(\mathbf{x}, \mathbf{y})$ is the potential energy due to the interaction between a particle at point $\mathbf{x}$ and a particle at point $\mathbf{y}$. The potential energy thus breaks up into a sum of $N + N(N-1)/2$ pieces. Does the decoupling theorem apply?

5.2 Classical Diatomic Ideal Gases

Rigid linear model. The partition function is

$$Z(T, V, N) = \frac{1}{N!}\left[\frac{V}{\lambda^3(T)}\left(\frac{4\pi^3 I k_B T}{h^2}\right)\right]^N \tag{5.14}$$

General results for separation of Z into translational (center of mass) and internal degrees of freedom. Term "internal specific heat".

5.3 Heat Capacity of an Ideal Gas

5.3.1 The equipartition theorem

Equipartition theorem. (For classical systems.) Suppose the Hamiltonian $H(\Gamma)$ decouples into one piece involving a single phase space variable—call it H_1, plus another piece which involves all the other phase space variables—call it $H_2(\Gamma_2)$. Suppose further that the energy depends quadratically upon this single phase space variable, and that this variable may take on values from $-\infty$ to $+\infty$. Then, in classical statistical mechanics, the mean contribution to the energy due to that single variable is

$$\langle H_1 \rangle = \tfrac{1}{2} k_B T. \tag{5.15}$$

Notice how general this theorem is. The remaining piece of the Hamiltonian, $H_2(\Gamma_2)$, might decouple further or it might not. The phase space variable entering H_1 might be a momentum,

$$H_1(p) = \frac{p^2}{2m}, \tag{5.16}$$

or an angular momentum,

$$H_1(\ell) = \frac{\ell^2}{2I}, \tag{5.17}$$

or even a position coordinate, as in the simple harmonic oscillator energy

$$H_1(x) = \tfrac{1}{2} k x^2. \tag{5.18}$$

Furthermore, in all these circumstances, the mean energy is *independent* of the particular parameters m or I or k...it depends only upon the temperature. This explains the origin of the name "equipartition": the

mean translational energy due to motion in the x direction is equal to the mean rotational energy due to the change of θ, and this holds true even if the gas is a mixture of molecules with different masses and different moments of inertia. The energy is equally partitioned among all these different ways of holding energy.

Proof. We will write

$$H_1(p) = ap^2, \tag{5.19}$$

although the variable might not be a linear momentum. Then the average of H_1 is

$$\langle H_1 \rangle = \frac{\int d\Gamma\, H_1 e^{-\beta H(\Gamma)}}{\int d\Gamma\, e^{-\beta H(\Gamma)}} = \frac{\int_{-\infty}^{+\infty} dp\, H_1 e^{-\beta H_1} \int d\Gamma_2\, e^{-\beta H_2(\Gamma_2)}}{\int_{-\infty}^{+\infty} dp\, e^{-\beta H_1} \int d\Gamma_2\, e^{-\beta H_2(\Gamma_2)}}. \tag{5.20}$$

Clearly, the integrals over Γ_2 cancel in this last expression. (This explains why the form of H_2 is irrelevant to the theorem.) We are left with

$$\langle H_1 \rangle = \frac{\int_{-\infty}^{+\infty} dp\, ap^2 e^{-\beta a p^2}}{\int_{-\infty}^{+\infty} dp\, e^{-\beta a p^2}}. \tag{5.21}$$

These two integrals could be evaluated in terms of Gamma functions (see appendix C), but they don't need to be evaluated yet. Think for a moment about our "slick trick" of parametric differentiation... using it we can write

$$\langle H_1 \rangle = -\frac{d}{d\beta} \ln \left[\int_{-\infty}^{+\infty} dp\, e^{-\beta a p^2} \right]. \tag{5.22}$$

The integral that remains is of Gaussian character and we *could* evaluate it using the results of Appendix B. But before rushing in to integrate, let's employ the substitution $u = \sqrt{\beta a}\, p$ to find

$$\langle H_1 \rangle = -\frac{d}{d\beta} \ln \left[\frac{1}{\sqrt{\beta a}} \int_{-\infty}^{+\infty} du\, e^{-u^2} \right] = -\frac{d}{d\beta} \left\{ \ln \left[\frac{1}{\sqrt{\beta}} \right] + \ln \left[\frac{1}{\sqrt{a}} \int_{-\infty}^{+\infty} du\, e^{-u^2} \right] \right\}. \tag{5.23}$$

This last expression shows that there's no need to evaluate the integral. Whatever its value is, it is some number, not a function of β, so when we take the derivative with respect to β the term involving that number will differentiate to zero. Similarly for the constant a, which explains why the equipartition result is independent of that prefactor. We are left with

$$\langle H_1 \rangle = -\frac{d}{d\beta} \ln \frac{1}{\sqrt{\beta}} = \frac{1}{2} \frac{d}{d\beta} \ln \beta = \frac{1}{2} \frac{1}{\beta} \tag{5.24}$$

or, finally, the desired equipartition result

$$\langle H_1 \rangle = \tfrac{1}{2} k_B T. \tag{5.25}$$

5.3.2 Applications of equipartition; Comparison with experiment

5.3.3 Crossover between classical and quantal behavior; Freeze out

At high temperatures, typical thermal energies are much greater than level spacings. Transitions from one level to another are very easy to do and the granular character of the quantized energies can be ignored. This is the classical limit, and equipartition holds!

At low temperatures, typical thermal energies are less that the level spacing between the ground state and the first excited state. There is so little thermal energy around that the molecule cannot even be excited out of its ground state. Virtually all the molecules are in their ground states, and the excited states might as well just not exist.

In classical mechanics, a diatomic molecule offered a small amount of rotational energy will accept that energy and rotate slowly. But in quantum mechanics, a diatomic molecule offered a small amount of rotational energy will reject that energy and remain in the ground state, because the energy offered is not enough to lift it into the first excited state.[2] The quantal diatomic molecule does not rotate at all at low temperatures, so it behaves exactly like a monatomic molecule with only center-of-mass degrees of freedom.

In short, we explain the high-temperature rotational specific heat ($c_V^{\rm rot} = k_B$) through equipartition. We explain the low-temperature rotational specific heat ($c_V^{\rm rot}$ vanishes) through difficulty of promotion to the first quantal excited state. This fall-off of specific heat as the temperature is reduced is called "freeze out".

The crossover between the high-temperature and low-temperature regimes occurs in the vicinity of a characteristic temperature θ at which the typical thermal energy is equal to energy separation between the ground state and the first excited state. If the energies of these two states are ϵ_0 and ϵ_1 respectively, then we define the characteristic crossover temperature through

$$k_B\theta \equiv \epsilon_1 - \epsilon_0. \tag{5.26}$$

5.2 Generalized equipartition theorem and the ultra-relativistic gas

a. Suppose the Hamiltonian $H(\Gamma)$ decouples into two pieces

$$H(\Gamma) = a|p|^n + H_2(\Gamma_2) \tag{5.27}$$

where p is some phase space variable that may take on values from $-\infty$ to $+\infty$, and where Γ_2 represents all the phase space variables except for p. (Note that the absolute value $|p|$ is needed in order to avoid, for example, taking the square root of a negative number in the case $n = 1/2$.) Show that, in classical statistical mechanics, the mean contribution to the energy due to that single variable is

$$\langle a|p|^n\rangle = \tfrac{1}{n}k_BT. \tag{5.28}$$

b. In special relativity, the energy of a free (i.e. non-interacting) particle is given by

$$\sqrt{(mc^2)^2 + (pc)^2}, \tag{5.29}$$

where c is the speed of light. As you know, when $v \ll c$ this gives the non-relativistic kinetic energy KE $\approx mc^2 + p^2/2m$. In the "ultra-relativistic" limit, where v is close to c, the energy is approximately pc. What is the heat capacity of a gas of non-interacting ultra-relativistic particles?

[2] This paragraph is written in the "shorthand" language discussed on page 113, as if energy eigenstates were the only allowed quantal states.

c. Estimate the crossover temperature between the non-relativistic and ultra-relativistic regimes.

5.3 **Another generalization of equipartition**
Consider the same situation as the equipartition theorem in the text, but now suppose the single phase space variable takes on values from 0 to $+\infty$. What is the corresponding result for $\langle H_1 \rangle$?

5.4 **Equipartition and the virial theorem**
Look up the term "virial theorem" in a classical mechanics textbook. Is there any relation between the virial theorem of classical mechanics and the equipartition theorem of classical statistical mechanics?

Box: The $\mathcal{O}$ Notation

Approximations are an important part of physics, and an important part of approximation is to ensure their reliability and consistency. The $\mathcal{O}$ notation (pronounced "the big-oh notation") is an important and practical tool for making approximations reliable and consistent.

The technique is best illustrated through an example. Suppose you desire an approximation for

$$f(x) = \frac{e^{-x}}{1-x} \tag{5.30}$$

valid for small values of x, that is, $x \ll 1$. You know that

$$e^{-x} = 1 - x + \tfrac{1}{2}x^2 - \tfrac{1}{6}x^3 + \cdots \tag{5.31}$$

and that

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots, \tag{5.32}$$

so it seems that reasonable approximations are

$$e^{-x} \approx 1 - x \tag{5.33}$$

and

$$\frac{1}{1-x} \approx 1 + x, \tag{5.34}$$

whence

$$\frac{e^{-x}}{1-x} \approx (1-x)(1+x) = 1 - x^2. \tag{5.35}$$

Let's try out this approximation at $x_0 = 0.01$. A calculator shows that

$$\frac{e^{-x_0}}{1-x_0} = 1.0000503\ldots \tag{5.36}$$

while the value for the approximation is

$$1 - x_0^2 = 0.9999000. \tag{5.37}$$

This is a very poor approximation indeed... the deviation from $f(0) = 1$ is even of the wrong sign!

Let's do the problem over again, but this time keeping track of exactly how much we've thrown away while making each approximation. We write

$$e^{-x} = 1 - x + \tfrac{1}{2}x^2 - \tfrac{1}{6}x^3 + \cdots \tag{5.38}$$

as

$$e^{-x} = 1 - x + \tfrac{1}{2}x^2 + \mathcal{O}(x^3), \tag{5.39}$$

where the notation $\mathcal{O}(x^3)$ stands for the small terms that we haven't bothered to write out explicitly. The symbol $\mathcal{O}(x^3)$ means "terms that are about the magnitude of x^3, or smaller" and is pronounced "terms of order x^3". The $\mathcal{O}$ notation will allow us to make controlled approximations in which we keep track of exactly how good the approximation is.

Similarly, we write

$$\frac{1}{1-x} = 1 + x + x^2 + \mathcal{O}(x^3), \tag{5.40}$$

and find the product

$$\begin{aligned} f(x) &= \left[1 - x + \tfrac{1}{2}x^2 + \mathcal{O}(x^3)\right] \times \left[1 + x + x^2 + \mathcal{O}(x^3)\right] && (5.41)\\ &= \left[1 - x + \tfrac{1}{2}x^2 + \mathcal{O}(x^3)\right] && (5.42)\\ &\quad + \left[1 - x + \tfrac{1}{2}x^2 + \mathcal{O}(x^3)\right] x && (5.43)\\ &\quad + \left[1 - x + \tfrac{1}{2}x^2 + \mathcal{O}(x^3)\right] x^2 && (5.44)\\ &\quad + \left[1 - x + \tfrac{1}{2}x^2 + \mathcal{O}(x^3)\right] \mathcal{O}(x^3). && (5.45) \end{aligned}$$

Note, however, that $x \times \frac{1}{2}x^2 = \mathcal{O}(x^3)$, and that $x^2 \times \mathcal{O}(x^3) = \mathcal{O}(x^3)$, and so forth, whence

$$\begin{aligned} f(x) &= \left[1 - x + \tfrac{1}{2}x^2 + \mathcal{O}(x^3)\right] && (5.46)\\ &\quad + \left[x - x^2 + \mathcal{O}(x^3)\right] && (5.47)\\ &\quad + \left[x^2 + \mathcal{O}(x^3)\right] && (5.48)\\ &\quad + \mathcal{O}(x^3) && (5.49)\\ &= 1 + \tfrac{1}{2}x^2 + \mathcal{O}(x^3). && (5.50) \end{aligned}$$

Thus we have the approximation

$$f(x) \approx 1 + \tfrac{1}{2}x^2. \tag{5.51}$$

Furthermore, we know that this approximation is accurate to terms of order $\mathcal{O}(x^2)$ (i.e. that the first neglected terms are of order $\mathcal{O}(x^3)$). Evaluating this approximation at $x_0 = 0.01$ gives

$$1 + \tfrac{1}{2}x_0^2 = 1.0000500, \tag{5.52}$$

far superior to our old approximation (5.35).

What went wrong on our first try? The $-x^2$ in approximation (5.35) is the same as the $-x^2$ on line (5.47). However, lines (5.46) and (5.48) demonstrate that there were other terms of about the same size (that is, other "terms of order x^2") that we neglected in our first attempt.

The $\mathcal{O}$ notation is superior to the "dot notation" (such as $\cdots$) in that dots stand for "a bunch of small terms", but the dots don't tell you just how small they are. The symbol $\mathcal{O}(x^3)$ also stands for "a bunch of small terms", but in addition it tells you precisely how small those terms are. The $\mathcal{O}$ notation allows us to approximate in a consistent manner, unlike the uncontrolled approximations where we ignore a "small term" without knowing whether we have already retained terms that are even smaller.

End of Box: The $\mathcal{O}$ Notation

5.4 Specific Heat of a Hetero-nuclear Diatomic Ideal Gas

The classical partition function is

$$Z = \frac{1}{N!}\left[\frac{V}{\lambda^3(T)}\frac{1}{h^2}\int_0^{\pi} d\theta \int_0^{2\pi} d\varphi \int_{-\infty}^{+\infty} d\ell_\theta \int_{-\infty}^{+\infty} d\ell_\varphi \, e^{-\beta(\ell_\theta^2+\ell_\varphi^2)/2I}\right]^N. \tag{5.53}$$

Let's make sure we have the quantum mechanics of one molecule straight before we jump into treating 6.02×10^{23} molecules: First of all, we consider *hetero*-nuclear rather than homo-nuclear diatomic molecules (such as CO or HCl rather than H_2 or N_2) because in the latter there are additional issues due to the Pauli principle. Let's just avoid these issues on our first encounter. The quantal energy eigenvalues are

$$\epsilon_{\rm rot} = \ell(\ell+1)\frac{\hbar^2}{2I} \qquad \ell = 0, 1, 2, \ldots, \tag{5.54}$$

with

$$\text{degeneracy}_\ell = 2\ell + 1. \tag{5.55}$$

Thus the quantal "characteristic temperature" θ (or $\Theta_{\rm rot}$) is

$$k_B\theta \equiv \epsilon_1 - \epsilon_0 = \frac{\hbar^2}{I}. \tag{5.56}$$

And, finally, the quantal partition function is

$$Z = \frac{1}{N!}\left[\frac{V}{\lambda^3(T)}\sum_{\ell=0}^{\infty}(2\ell+1)e^{-\beta\ell(\ell+1)\hbar^2/2I}\right]^N. \tag{5.57}$$

It saves a lot of writing to define the "rotational partition function"

$$\zeta(T) \equiv \zeta^{\rm rot}(T) \equiv \sum_{\ell=0}^{\infty}(2\ell+1)e^{-\beta\ell(\ell+1)\hbar^2/2I} = \sum_{\ell=0}^{\infty}(2\ell+1)e^{-\ell(\ell+1)\theta/2T}. \tag{5.58}$$

The expression on the far right is the one that I like to use... instead of having a lot of variables like T, $\hbar$, and I floating around just waiting to get misplaced, everything is neatly packaged into the ratio θ/T, which is obviously dimensionless.

Now, for *any* partition function of the form

$$Z = \frac{1}{N!}\left[\frac{V}{\lambda^3(T)}\zeta(T)\right]^N = Z_{\text{mono}}\zeta^N(T). \tag{5.59}$$

We have

$$F = -k_BT\ln Z = \underbrace{-k_BT\ln Z_{\text{mono}}}_{F_{\text{mono}}}\underbrace{-k_BTN\ln\zeta}_{Nf^{\text{rot}}} \tag{5.60}$$

and

$$e^{\text{rot}} = \frac{\partial(f^{\text{rot}}/T)}{\partial(1/T)} = -\frac{\partial\ln\zeta}{\partial\beta} \qquad c_V^{\text{rot}} = \frac{\partial e^{\text{rot}}}{\partial T}. \tag{5.61}$$

Because the rotational partition function depends on temperature only through the combination θ/T, it makes sense to use that variable for derivatives:

$$e^{\text{rot}} = \frac{\partial(f^{\text{rot}}/T)}{\partial(1/T)} = \frac{\partial(-k_BT\ln\zeta/T)}{\partial(1/T)} = -k_B\frac{\partial\ln\zeta}{\partial(1/T)} = -k_B\theta\frac{\partial\ln\zeta}{\partial(\theta/T)} \tag{5.62}$$

and

$$c_V^{\text{rot}} = \frac{\partial e^{\text{rot}}}{\partial T} = \frac{\partial e^{\text{rot}}}{\partial(\theta/T)}\frac{\partial(\theta/T)}{\partial T} = -\frac{\theta}{T^2}\frac{\partial e^{\text{rot}}}{\partial(\theta/T)}. \tag{5.63}$$

Now back to the evaluation of $\zeta(T)$. It is quite easy to prove that the infinite sum for $\zeta(T)$ converges. You will then be tempted to find an expression for $\zeta(T)$ in terms of well-known functions like polynomials or exponentials. If you try this, you will not find any simple expression. Instead of manipulating $\zeta(T)$ into some combination of familiar functions, we will have to become familiar with it on our own. It can be evaluated and plotted on a computer just by adding up a finite number of terms in the defining infinite sum. But doing so is not the most productive way to approach an understanding of this function... given a plot of $\zeta(T)$, how would you find the specific heat due to rotation? A better way to approach the problem is to find how the partition function $\zeta(T)$ (and hence the specific heat $c_V^{\text{rot}}(T)$) behaves at low and high temperatures.

At low temperatures, we expand in terms of the small variable $e^{-\theta/T}$ to find

$$\begin{aligned}\zeta(T) &= 1 + 3e^{-\theta/T} + 5e^{-3\theta/T} + 7e^{-6\theta/T} + \cdots \\ &= 1 + 3e^{-\theta/T} + 5e^{-3\theta/T} + \mathcal{O}(e^{-6\theta/T}).\end{aligned} \tag{5.64}$$

The notation $\mathcal{O}(x^N)$ is read "terms of order x^N". (See the box on page 129.) Now for small variables ϵ the logarithm is

$$\ln(1+\epsilon) = \epsilon - \tfrac{1}{2}\epsilon^2 + \tfrac{1}{3}\epsilon^3 - \tfrac{1}{4}\epsilon^4 + \mathcal{O}(\epsilon^5), \tag{5.65}$$

whence

$$\begin{aligned}
\ln\zeta(T) &= 3e^{-\theta/T} + 5e^{-3\theta/T} + \mathcal{O}(e^{-6\theta/T}) \qquad (5.66)\\
&\quad -\tfrac{1}{2}[3e^{-\theta/T} + 5e^{-3\theta/T} + \mathcal{O}(e^{-6\theta/T})]^2\\
&\quad +\tfrac{1}{3}[3e^{-\theta/T} + 5e^{-3\theta/T} + \mathcal{O}(e^{-6\theta/T})]^3\\
&\quad +\mathcal{O}(e^{-4\theta/T})\\
&= 3e^{-\theta/T} + 5e^{-3\theta/T} + \mathcal{O}(e^{-6\theta/T}) \qquad (5.67)\\
&\quad -\tfrac{1}{2}[9e^{-2\theta/T} + 30e^{-4\theta/T} + \mathcal{O}(e^{-6\theta/T})]\\
&\quad +\tfrac{1}{3}[27e^{-3\theta/T} + 135e^{-5\theta/T} + \mathcal{O}(e^{-6\theta/T})]\\
&\quad +\mathcal{O}(e^{-4\theta/T})\\
&= 3e^{-\theta/T} - \tfrac{9}{2}e^{-2\theta/T} + 14e^{-3\theta/T} + \mathcal{O}(e^{-4\theta/T}). \qquad (5.68)
\end{aligned}$$

Note that in going from equation (5.67) to equation (5.68) it would have been impossible to make sure that all of the several expansions were kept to equivalent orders of accuracy had we used $+\cdots$ rather than the $\mathcal{O}$ notation.

Now it is easy to use equations (5.62) and (5.63) to find

$$e^{\rm rot} = -k_B\theta[-3e^{-\theta/T} - 9e^{-2\theta/T} - 42e^{-3\theta/T} + \mathcal{O}(e^{-4\theta/T})] \qquad (5.69)$$

and

$$c_V^{\rm rot} = k_B\left(\frac{\theta}{T}\right)^2[3e^{-\theta/T} - 18e^{-2\theta/T} + 126e^{-3\theta/T} + \mathcal{O}(e^{-4\theta/T})]. \qquad (5.70)$$

(Note that if we had used equations (5.61) instead, it would have been a lot more work.) The specific heat vanishes at zero temperature, and it increases *exponentially* slowly with increasing temperature. This is a very slow growth indeed... not only is the slope zero, but the second derivative is zero, the third derivative is zero, indeed *all* orders of derivative vanish at the origin.

What about high temperatures? We have discussed the fact that at high temperatures the rotational specific heat will approach the classical equipartition value of k_B, but *how* will it approach the limiting value? In other words, what are the deviations from the high temperature limit? We can answer these questions by approximating the sum by an integral. The *Euler-MacLaurin formula* states that if $f(x) \to 0$, $f'(x) \to 0$, $f''(x) \to 0$, etc. as $x \to \infty$, then

$$\sum_{\ell=0}^{\infty} f(\ell) \approx \int_0^\infty f(x)\,dx + \tfrac{1}{2}f(0) - \tfrac{1}{12}f'(0) + \tfrac{1}{720}f'''(0) - \tfrac{1}{30240}f^{(\rm v)}(0) + \cdots. \qquad (5.71)$$

In our case,

$$f(x) = (2x+1)e^{-x(x+1)\theta/2T}, \qquad (5.72)$$

so

$$\int_0^\infty f(x)\,dx = 2\frac{T}{\theta}$$

$$\begin{aligned} f(0) &= 1 \\ f'(0) &= 2 - \frac{\theta}{2T} \\ f'''(0) &= -6\frac{\theta}{T} + \mathcal{O}\left(\frac{\theta}{T}\right)^2 \\ f^{(\mathrm{v})}(0) &= \mathcal{O}\left(\frac{\theta}{T}\right)^2. \end{aligned}$$

Thus we have

$$\zeta(T) \approx 2\frac{T}{\theta} + \frac{1}{3} + \frac{1}{30}\frac{\theta}{T} + \mathcal{O}\left(\frac{\theta}{T}\right)^2 \tag{5.73}$$

$$\ln\zeta(T) \approx -\ln\frac{\theta}{2T} + \frac{1}{6}\frac{\theta}{T} + \frac{1}{360}\left(\frac{\theta}{T}\right)^2 + \mathcal{O}\left(\frac{\theta}{T}\right)^3 \tag{5.74}$$

$$e^{\mathrm{rot}} \approx -k_B\theta\left[-\frac{T}{\theta} + \frac{1}{6} + \frac{1}{180}\frac{\theta}{T} + \mathcal{O}\left(\frac{\theta}{T}\right)^2\right] \tag{5.75}$$

$$c_V^{\mathrm{rot}} \approx k_B\left[1 + \frac{1}{180}\left(\frac{\theta}{T}\right)^2 + \mathcal{O}\left(\frac{\theta}{T}\right)^3\right]. \tag{5.76}$$

For high temperatures the specific heat is nearly equal to its classical value, and the quantity grows slightly larger as the temperature decreases.

Note: The $\approx$ sign in the above formulas represents "asymptotic equality". The infinite series on the right does not necessarily converge, and even if it does then it might not converge to the quantity on the left. However a finite truncation of the sum can be a good approximation for the quantity on the left, and that approximation grows better and better with increasing temperature (i.e. smaller values of θ/T).

5.5 Chemical Reactions Between Gases

5.6 Problems

5.5 **Decoupling quantal Hamiltonians**
Prove the "decoupling Hamilton implies factoring partition function" theorem of section 5.1.2 for quantal systems.

5.6 **Schottky anomaly**
A molecule can be accurately modeled by a quantal two-state system with ground state energy 0 and excited state energy ϵ. Show that the internal specific heat is

$$c_V^{\mathrm{int}}(T) = k_B\left(\frac{\epsilon}{k_BT}\right)^2 \frac{e^{-\epsilon/k_BT}}{(1+e^{-\epsilon/k_BT})^2}. \tag{5.77}$$

Sketch this specific heat as a function of k_BT/ϵ. How does the function behave when $k_BT \ll \epsilon$ and $k_BT \gg \epsilon$?

5.7 **Simple harmonic oscillator**

Suppose a molecule can be accurately modeled as a harmonic oscillator of natural frequency ω.

a. Find the expected internal energy of one such molecule, written as a sum of the ground state energy plus a temperature-dependent part.

b. Show that the internal specific heat is

$$c_V^{\text{int}}(T) = k_B \left(\frac{\hbar\omega}{k_B T}\right)^2 \frac{e^{-\hbar\omega/k_B T}}{(1 - e^{-\hbar\omega/k_B T})^2}. \tag{5.78}$$

c. Show that at low temperatures ($k_B T \ll \hbar\omega$),

$$c_V^{\text{int}}(T) \approx k_B \left(\frac{\hbar\omega}{k_B T}\right)^2 e^{-\hbar\omega/k_B T}, \tag{5.79}$$

whereas at high temperatures ($k_B T \gg \hbar\omega$),

$$c_V^{\text{int}}(T) \approx k_B. \tag{5.80}$$

d. (Optional.) Show that the leading quantal correction to the high-temperature specific heat is

$$c_V^{\text{int}}(T) = k_B[1 - \tfrac{1}{12}x^2 + \mathcal{O}(x^3)], \quad \text{where} \quad x = \frac{\hbar\omega}{k_B T}. \tag{5.81}$$

e. Sketch the internal specific heat as a function of $k_B T/\hbar\omega$.

5.8 **Simple harmonic oscillator — entropy**

I was speaking with someone who claimed that thermodynamics didn't apply to systems of interacting atoms, or to living things, or to anything other than the ideal gas. He challenged me "What's the entropy of a pendulum? The answer is that entropy doesn't apply to a pendulum!" Answer his question for a pendulum in the simple harmonic oscillator approximation, with natural frequency ω and temperature T.

5.9 **Kinetic energy in air**

What is the kinetic energy is a cubic liter of air (mostly nitrogen and oxygen) at one atmosphere pressure and room temperature? At one atmosphere pressure and temperature 600 K? Use classical mechanics, and express your answer both in Joules and in the commercial unit kilowatt-hours. (*Clue:* If you calculate the average kinetic energy of one molecule, you're working way too hard.) Why doesn't anyone exploit all this valuable energy lying around in every cubic meter of air?

5.10 **Conceptual comparison**

a. Explain qualitatively why the results of the two previous problems are parallel at low temperatures.

b. (Harder.) Explain qualitatively both high temperature results. (Clue: At high temperatures, the average energy per particle in the Schottky case approaches $\epsilon/2$. Why?)

5.11 **Compressibility of a diatomic gas**

Find the isothermal compressibility κ_T for an ideal diatomic gas, where each molecule is modeled as a dumbbell with moment of inertia I.

5.12 **Systems with a small number of states**

(This problem requires *no calculation!* All the answers can be found in your head.) A collection of non-interacting particles is in thermal equilibrium. Each particle has only three energy eigenvalues, namely 0, ϵ, and 4ϵ.

a. What is the criterion for "high temperature" in this situation?

b. Suppose there are three non-degenerate energy eigenstates. At high temperatures, what is the average energy of each particle? (Clue: The answer is *not* 4ϵ.)

c. Now suppose that the lower two energy eigenstates are non-degenerate, but that there are two independent states with energy 4ϵ. What is the average energy per particle at high temperatures in this case?

5.13 **Anharmonic oscillator**

The energy eigenvalues of a simple harmonic oscillator are equally spaced, and we have explored the consequences of this for the heat capacity of a collection of harmonic oscillators. Suppose an anharmonic oscillator is approximately harmonic (with natural frequency ω_0) for small energies, but that for large energies (greater than, say, E_x) the eigenvalues become *more* closely spaced as energy increases. At temperatures greater than E_x/k_B, will the heat capacity of a collection of such anharmonic oscillators be greater than or less than that of a collection of harmonic oscillators with the same natural frequency ω_0? Why?

5.14 **Descriptive features of models**

(This problem is stolen from a GRE Physics test.)

Two possible models for a diatomic ideal gas are the rigid dumbbell (model R; two point particles connected by a rigid rod) and the springy dumbbell (model S; two point particles connected by a spring). In classical statistical mechanics, which of the following statements is true?

a. Model R has a specific heat $c_V = \frac{3}{2}k_B$.

b. Model S has a smaller specific heat than model R.

c. Model S is always correct.

d. Model R is always correct.

e. The choice between models R and S depends on the temperature.

5.15 **An n-state system, qualitatively**

A model molecule has n equally-spaced energy levels, all of them non-degenerate, with energy spacing ϵ. Thus as n varies from 2 to ∞ this model interpolates between the Schottky system of problem 5.6 and the simple harmonic oscillator of problem 5.7.

a. Find a low-temperature approximation for the specific heat that is independent of n.

b. At high temperatures, the specific heat approaches zero. What is the criterion for "high temperature"?

c. At high temperatures, what is the expected energy of this model?

d. There is a theorem stating that at any fixed positive temperature, the specific heat must increase with increasing n. Assume this theorem and use it to prove that as n increases, the maximum in the specific heat versus temperature curve becomes higher.

5.16 **An n-state system, quantitatively**

Show that the system of the previous problem has internal specific heat

$$c_V^{\text{int}}(T) = k_B \left[\left(\frac{\epsilon}{k_B T} \right)^2 \frac{e^{-\epsilon/k_B T}}{(1 - e^{-\epsilon/k_B T})^2} - \left(\frac{n\epsilon}{k_B T} \right)^2 \frac{e^{-n\epsilon/k_B T}}{(1 - e^{-n\epsilon/k_B T})^2} \right]. \tag{5.82}$$

Does this expression have the proper limits when $n = 2$ and when $n \to \infty$?

Chapter 6

Quantal Ideal Gases

6.1 Introduction

In the previous chapter, we found that at high temperatures, an ideal gas of diatomic molecules with spring interactions has a heat capacity of $\frac{7}{2}k_B$ per molecule: $\frac{3}{2}k_B$ from the translational degrees of freedom, k_B from the rotational degrees of freedom, and k_B from the spring degrees of freedom. If the temperature is decreased, the spring degrees of freedom become governed by quantum mechanics rather than classical mechanics, the equipartition theorem no longer holds, and eventually these degrees of freedom "freeze out" and contribute nothing to the heat capacity: the total heat capacity per molecule becomes $\frac{5}{2}k_B$. If the temperature is decreased still further, the story is repeated for the rotational degrees of freedom and eventually they freeze out. What happens if the temperature is decreased yet again? Do the translational degrees of freedom then freeze out as well? The answer is "sort of", but the crossover from the classical to the quantal regime is complicated in this case by the quantal requirement of interchange symmetry. This requirement gives rise to a much richer and more interesting crossover behavior than is provided by simple freeze out.

6.2 The Interchange Rule

Before turning to statistical mechanics, let us review the quantal "interchange rule". The wavefunction for a system of three particles is a function of three variables: $\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C)$. [The symbol x represents whatever is needed to specify the state: For a spinless particle in one dimension x represents the coordinate x or, if you are working in momentum space, the coordinate p. For a particle with spin moving in three dimensions, x represents (x, y, z, m_z), or perhaps (p_x, p_y, p_z, m_x).] If the three particles are identical, then the wavefunction must be either symmetric under the interchange of any pair of variables,

$$\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C) = +\psi(\mathsf{x}_B, \mathsf{x}_A, \mathsf{x}_C) = +\psi(\mathsf{x}_B, \mathsf{x}_C, \mathsf{x}_A), \quad \text{etc.,} \tag{6.1}$$

or else antisymmetric under the interchange of any pair of variables,

$$\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C) = -\psi(\mathsf{x}_B, \mathsf{x}_A, \mathsf{x}_C) = +\psi(\mathsf{x}_B, \mathsf{x}_C, \mathsf{x}_A), \qquad \text{etc.} \tag{6.2}$$

The foregoing assertion is an empirical rule that cannot be derived from any of the other principles of quantum mechanics. (Indeed there is currently considerable interest in *anyons*, hypothetical particles that obey all the principles of quantum mechanics except the interchange rule.) The rule holds for all quantal states, not just energy eigenstates. It holds for interacting as well as for non-interacting particles. And the rule has a number of surprising consequences, both within the domain of quantum mechanics and atomic physics, and, as well shall soon see in detail, within the domain of statistical mechanics.

The sign of the interchange symmetry, either + or −, is governed only by the type of particle involved: for pions it is always +, for electrons it is always −. Particles for which the sign is always + are called *bosons*, and those for which it is always − are called *fermions*. It is an experimental fact that particles with integral spin s are bosons and those with half integral spin s are fermions.

6.2.1 Problems

6.1 **The interchange rule in another representation**
Full information about the state of three identical spinless particles is contained not only in the configurational wave function $\psi(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C)$ but also in the momentum space wave function

$$\tilde{\psi}(\mathbf{p}_A, \mathbf{p}_B, \mathbf{p}_C) = \frac{1}{(2\pi\hbar)^{9/2}} \int d^3x_A \int d^3x_B \int d^3x_C \psi(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C) e^{-i(\mathbf{p}_A\cdot\mathbf{x}_A + \mathbf{p}_B\cdot\mathbf{x}_B + \mathbf{p}_C\cdot\mathbf{x}_C)/\hbar}. \tag{6.3}$$

Show that if one representation is symmetric (or antisymmetric), then the other one is as well.

6.2 **Symmetrization**
Show that for any function $f(x, y, z)$, the function

$$f_S(x, y, z) = f(x, y, z) + f(x, z, y) + f(z, x, y) + f(z, y, x) + f(y, z, x) + f(y, x, z) \tag{6.4}$$

is symmetric under the interchange of any pair of variables. How many such pairs are there? Is there a similar algorithm for building up an antisymmetric function from any garden-variety function?

6.3 Quantum Mechanics of Independent Identical Particles

This chapter considers collections of independent (i.e. non-interacting) identical monatomic particles. It does not treat mixtures or diatomic molecules. Notice that "independent" means only that the particles do not interact with each other. In contrast, each particle individually may interact with some background potential, such as a square well or harmonic oscillator potential. Later on (in section 6.5) we will restrict our consideration to particles that interact only with the walls of their container ("free particles"), but for now we treat the more general case.

Doing quantal statistical mechanics requires a basis of states to sum over. In this section we consider the only the quantum mechanics of our situation in order to produce an energy eigenbasis... we defer all questions of statistical mechanics to the next section.

Most of our discussion in this section is phrased in terms of the three-particle problem rather than the N-particle problem. The use of a specific number lends the advantage of concreteness, and three is the smallest number that generalizes readily to the N-particle case.

6.3.1 A basis for non-identical particles

Consider first a single particle moving in the background potential. Elementary quantum mechanics deals with this situation, and it tells us that there will be a certain number M of (possibly degenerate) energy eigenstates. (Usually M will be infinite, but there are advantages to calling it M and maintaining the ability to take the limit $M \to \infty$.) The rth energy eigenstate has energy ϵ_r and is represented by the wavefunction $\eta_r(\mathsf{x})$, where x denotes the arguments of the wavefunction: Thus for a spinless particle, x could stand for x, y, z or p_x, p_y, p_z or even p_x, y, p_z. For a particle with spin s, x could stand for expressions like x, y, z, s_z or p_x, p_y, p_z, s_x.

We will soon use these one-particle energy eigenstates as building blocks to construct energy eigenstates for the N-particle situation, i.e. energy eigenstates for the entire system. Thus we will frequently be talking about both "one-particle energy eigenstates" and "N-particle energy eigenstates" in the same sentence, and both phrases are mouthfulls. To make our sentences shorter and clearer—and to emphasize the dramatic distinction between these two entities—we will adopt the common convention of referring to the one-particle energy eigenstates as "levels" and to the N-particle energy eigenstates as "states". (Some books use the term "orbitals" rather than "levels".) Thus we have for the one-particle energy eigenproblem:

$$\begin{array}{rcccccc} \text{levels:} & \eta_1(\mathsf{x}) & \eta_2(\mathsf{x}) & \cdots & \eta_r(\mathsf{x}) & \cdots & \eta_M(\mathsf{x}) \\ \text{energies:} & \epsilon_1 & \epsilon_2 & \cdots & \epsilon_r & \cdots & \epsilon_M \end{array}$$

Now consider more than one particle—let's say three particles—moving independently and all subject to the same background potential. If the particles are not identical (say one helium, one argon, and one neon atom), then the three-body energy eigenbasis will contain elements ("states") such as

$$\eta_r(\mathsf{x}_A)\eta_s(\mathsf{x}_B)\eta_t(\mathsf{x}_C) \quad \text{with energy} \quad \epsilon_r + \epsilon_s + \epsilon_t. \tag{6.5}$$

(If the particles interacted, then this would still be a state of the system, but it would not be an energy eigenstate.) We will represent this state by the notation $|r, s, t\rangle$:

$$\eta_r(\mathsf{x}_A)\eta_s(\mathsf{x}_B)\eta_t(\mathsf{x}_C) \quad \Longleftrightarrow \quad |r, s, t\rangle. \tag{6.6}$$

Of course, degeneracies now *must* exist: the energy of $|r, s, t\rangle$ is the same as the energy of $|s, r, t\rangle$. The entire basis consists of M^3 such states (more generally, for the case of N independent particles, M^N), namely:

$$
\begin{array}{c}
|\,1\,,\ 1\ ,\ 1\ \rangle \\
|\,2\,,\ 1\ ,\ 1\ \rangle \\
|\,1\,,\ 2\ ,\ 1\ \rangle \\
|\,1\,,\ 1\ ,\ 2\ \rangle \\
|\,3\,,\ 1\ ,\ 1\ \rangle \\
\vdots \\
|\,3\,,\ 2\ ,\ 1\ \rangle \\
|\,3\,,\ 1\ ,\ 2\ \rangle \\
\vdots \\
|\,4\,,\ 2\ ,\ 1\ \rangle \\
\vdots \\
|\mathrm{M},\ \mathrm{M},\ \mathrm{M}\ \rangle
\end{array}
$$

Out of these M^3 states we can build up (by addition and scalar multiplication) any wavefunction, i.e. any normalized function of $(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C)$. This is exactly what we want for non-identical particles. But for identical particles we don't want to be able to build *any* wavefunction. Because of the interchange rule, the only relevant wavefunctions are those symmetric (or antisymmetric) under the interchange of any pair of coordinates, for example

$$\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C) = \pm\psi(\mathsf{x}_C, \mathsf{x}_B, \mathsf{x}_A), \tag{6.7}$$

and hence these are the only kind of wavefunctions we want to be able to build. In other words, we need an energy eigenbasis consisting only of symmetric (or antisymmetric) wavefunctions.

6.3.2 A basis for identical particles

Fortunately, there exists a general procedure for constructing symmetric (or antisymmetric) functions out of garden-variety (or "non-symmetrized") functions, and this procedure will enable us to build the two desired basis sets. For functions of three variables the procedure (which is readily generalized to N variables) works like this: Given a function $f(x_A, x_B, x_C)$, the new function

$$f(x_A, x_B, x_C) + f(x_A, x_C, x_B) + f(x_C, x_A, x_B) + f(x_C, x_B, x_A) + f(x_B, x_C, x_A) + f(x_B, x_A, x_C) \tag{6.8}$$

is surely symmetric under the interchange of any pair of variables, and the new function

$$f(x_A, x_B, x_C) - f(x_A, x_C, x_B) + f(x_C, x_A, x_B) - f(x_C, x_B, x_A) + f(x_B, x_C, x_A) - f(x_B, x_A, x_C) \tag{6.9}$$

is surely antisymmetric. These two procedures are called "symmetrization" and "antisymmetrization" (or sometimes "alternation") respectively. It is of course possible that the resulting function vanishes, but this does not invalidate the procedure, because zero functions are both symmetric *and* antisymmetric!

When applied to a quantal wavefunction $\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C)$, these processes result in the symmetric wavefunction

$$\hat{S}\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C) \equiv A_s[\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C) + \psi(\mathsf{x}_A, \mathsf{x}_C, \mathsf{x}_B) + \psi(\mathsf{x}_C, \mathsf{x}_A, \mathsf{x}_B) + \psi(\mathsf{x}_C, \mathsf{x}_B, \mathsf{x}_A) + \psi(\mathsf{x}_B, \mathsf{x}_C, \mathsf{x}_A) + \psi(\mathsf{x}_B, \mathsf{x}_A, \mathsf{x}_C)] \tag{6.10}$$

and in the antisymmetric wavefunction

$$\hat{A}\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C) \equiv A_a[\psi(\mathsf{x}_A, \mathsf{x}_B, \mathsf{x}_C) - \psi(\mathsf{x}_A, \mathsf{x}_C, \mathsf{x}_B) + \psi(\mathsf{x}_C, \mathsf{x}_A, \mathsf{x}_B) - \psi(\mathsf{x}_C, \mathsf{x}_B, \mathsf{x}_A) + \psi(\mathsf{x}_B, \mathsf{x}_C, \mathsf{x}_A) - \psi(\mathsf{x}_B, \mathsf{x}_A, \mathsf{x}_C)] \tag{6.11}$$

where the constants A_s and A_a are inserted simply to insure normalization. Note that the operators $\hat{S}$ and $\hat{A}$ defined above are *linear*, whence *a basis for symmetric wavefunctions may be obtained by symmetrizing every element of an ordinary, non-symmetrized, basis*, and similarly for antisymmetric wavefunctions.

For functions of three variables that factorize into three functions of one variable, i.e. functions of the form (6.5), these definitions can be expressed even more simply because, for example, swapping η_s and η_t is the same as swapping x_B and x_C. Thus, if $|r, s, t\rangle$ is an energy eigenstate, then

$$\hat{S}|r, s, t\rangle \equiv A_s[|r, s, t\rangle + |r, t, s\rangle + |t, r, s\rangle + |t, s, r\rangle + |s, t, r\rangle + |s, r, t\rangle] \tag{6.12}$$

is a symmetric state with the same energy, while

$$\hat{A}|r, s, t\rangle \equiv A_a[|r, s, t\rangle - |r, t, s\rangle + |t, r, s\rangle - |t, s, r\rangle + |s, t, r\rangle - |s, r, t\rangle] \tag{6.13}$$

in an antisymmetric state with the same energy.

It is easy to see that the symmetrization process is unaffected by the order of the letters, i.e. that

$$\hat{S}|r, s, t\rangle = \hat{S}|r, t, s\rangle \quad \text{etc.}, \tag{6.14}$$

whence, for example, the states $|1, 1, 2\rangle$, $|1, 2, 1\rangle$, and $|2, 1, 1\rangle$ all symmetrize to the same state. Similarly in antisymmetrization the order of the letters affects only the sign of the result,

$$\hat{A}|r, s, t\rangle = -\hat{A}|r, t, s\rangle = \hat{A}|t, r, s\rangle \quad \text{etc.}, \tag{6.15}$$

but this result is considerably more profound: It shows not only that, for example, $\hat{A}|1, 2, 3\rangle$ is the same state as $\hat{A}|3, 2, 1\rangle$, but also that

$$\hat{A}|3, 1, 1\rangle = -\hat{A}|3, 1, 1\rangle \tag{6.16}$$

whence $\hat{A}|3, 1, 1\rangle = 0$, and, in general, that $|r, s, t\rangle$ antisymmetrizes to zero unless r, s, and t are all distinct.

Now we construct a basis for symmetric wavefunctions by applying the operator $\hat{S}$ to every element of our non-symmetrized basis on page 141. We obtain

$$
\begin{array}{c}
\hat{S}|\,1\,,\ 1\ ,\ 1\ \rangle \\
\hat{S}|\,2\,,\ 1\ ,\ 1\ \rangle \\
\cancel{\hat{S}|\,1\,,\ 2\ ,\ 1\ \rangle} \\
\cancel{\hat{S}|\,1\,,\ 1\ ,\ 2\ \rangle} \\
\cancel{\hat{S}|\,3\,,\ 1\ ,\ 1\ \rangle} \\
\vdots \\
\hat{S}|\,3\,,\ 2\ ,\ 1\ \rangle \\
\cancel{\hat{S}|\,3\,,\ 1\ ,\ 2\ \rangle} \\
\vdots \\
\hat{S}|\,4\,,\ 2\ ,\ 1\ \rangle \\
\vdots \\
\hat{S}|\mathrm{M},\ \mathrm{M},\ \mathrm{M}\rangle
\end{array}
$$

where the crossed out elements are those that have already appeared earlier in the list. It is clear that there are fewer elements in this basis than there are in the non-symmetrized basis, although it is a challenge to count them exactly. The number of elements turns out to be

$$\frac{(M)(M+1)(M+2)}{3!}, \tag{6.17}$$

or, for the N-particle rather than the three-particle case,

$$\begin{pmatrix} M+N-1 \\ N \end{pmatrix}. \tag{6.18}$$

where the binomial coefficient symbol is defined through

$$\begin{pmatrix} m \\ n \end{pmatrix} \equiv \frac{m!}{n!(m-n)!}. \tag{6.19}$$

We construct a basis for antisymmetric wavefunctions in the same way. The result is

$$
\begin{array}{c}
\cancel{\hat{A}|\,1\,,\ 1\ ,\ 1\,\rangle} \\
\cancel{\hat{A}|\,2\,,\ 1\ ,\ 1\,\rangle} \\
\cancel{\hat{A}|\,1\,,\ 2\ ,\ 1\,\rangle} \\
\cancel{\hat{A}|\,1\,,\ 1\ ,\ 2\,\rangle} \\
\cancel{\hat{A}|\,3\,,\ 1\ ,\ 1\,\rangle} \\
\vdots \\
\hat{A}|\,3\,,\ 2\ ,\ 1\,\rangle \\
\cancel{\hat{A}|\,3\,,\ 1\ ,\ 2\,\rangle} \\
\vdots \\
\hat{A}|\,4\,,\ 2\ ,\ 1\,\rangle \\
\vdots \\
\cancel{\hat{A}|\mathrm{M},\ \mathrm{M},\ \mathrm{M}\rangle}
\end{array}
$$

where now the crossed out elements have either appeared earlier in the list or else are equal to zero. There are even fewer elements in this basis than there were in the symmetrized basis. In fact there are exactly

$$\frac{(M)(M-1)(M-2)}{3!}, \tag{6.20}$$

such elements or, for the case of N particles,

$$\begin{pmatrix} M \\ N \end{pmatrix} \equiv \frac{M!}{N!(M-N)!} \tag{6.21}$$

elements.

For the case $N = 2$, there are $M(M+1)/2$ elements the symmetric basis and $M(M-1)/2$ elements in the antisymmetric basis, so the sum is M^2, the number of elements in the non-symmetrized basis. In fact, for $N = 2$, the set resulting from the conjunction of the symmetric and antisymmetric bases is a basis for the entire set of any function of two variables. This is related to the fact that any two-variable function can be written as the sum of a symmetric function and an antisymmetric function. I point out these results to emphasize that they apply for the two-variable case only, and are not general properties of symmetric and antisymmetric functions. For $N \geq 3$, the conjunction of the symmetric and antisymmetric bases does *not* span the set of all N-variable functions.

6.3.3 The occupation number representation

We have seen that in order to specify an element of the symmetric or the antisymmetric basis that we have just produced, it is not necessary to specify the order of the one-particle level building blocks. For example $\hat{A}|4, 9, 7\rangle$ is the same state as $\hat{A}|4, 7, 9\rangle$, so there's no need to pay attention to the order in which the 4, 7, and 9 appear. This observation permits the "occupation number" representation of such states, in which

we specify the basis state simply by listing the one-particle levels that are used as building blocks to make up that state. Or, equivalently but more commonly, we specify the basis state by listing the number n_r of one-body levels of each type r that are used as building blocks. (And, of course, we must also specify whether we're considering the symmetric or the antisymmetric basis.) Thus, for example:

level r:	1	2	3	4	5	6	$\cdots$	M
$\hat{S}\|3,3,4\rangle$ has n_r:	0	0	2	1	0	0	$\cdots$	0
$\hat{A}\|1,3,4\rangle$ has n_r:	1	0	1	1	0	0	$\cdots$	0

The second line in this table means that the state $\hat{S}|3,3,4\rangle$ is built by starting with the three levels $\eta_3(\mathsf{x}_A)$, $\eta_3(\mathsf{x}_B)$, and $\eta_4(\mathsf{x}_A)$, multiplying them together, and then symmetrizing. Sometimes you will hear this state described by the phrase "there are two particles in level 3 and one particle in level 4", but that can't be literally true... the three particles are identical, and if they could be assigned to distinct levels they would not be identical! Phrases such as the one above[1] invoke the "balls in buckets" picture of N-particle quantal wavefunctions: The state $\hat{S}|3,3,4\rangle$ is pictured as two balls in bucket number 3 and one ball in bucket number 4. It is all right to use this picture and this phraseology, as long as you don't believe it. Always keep in mind that it is a shorthand for a more elaborate process of building up states from levels by multiplication and symmetrization.

The very term "occupation number" for n_r is a poor one, because it so strongly suggests the balls-in-buckets picture. A somewhat better name for n_r is "occupancy", and I will sometimes use it. If you can think of a better name, please tell me.

To summarize the occupation number representation: an element of the symmetric basis is specified by the list

$$n_r, \quad \text{for} \quad r = 1, 2, \ldots M, \quad \text{where} \quad n_r \text{ is } 0, 1, 2, \ldots, \tag{6.22}$$

and an element of the antisymmetric basis is specified by the list

$$n_r, \quad \text{for} \quad r = 1, 2, \ldots M, \quad \text{where} \quad n_r \text{ is 0 or 1.} \tag{6.23}$$

The total number of particles in such a state is

$$N = \sum_{r=1}^{M} n_r, \tag{6.24}$$

and the energy of the state is

$$E = \sum_{r=1}^{M} n_r \epsilon_r. \tag{6.25}$$

Finally, since we have been devoting so much attention to energy eigenstates, I remind you that there do exist states other than energy states. (Expand on this paragraph and insert a reference to the caveat concerning meaning of $e^{-\beta E}$ for arbitrary states on page 113.)

[1] For example, phrases like "the level is filled" or "the level is empty" or "the level is half-filled".

6.3.4 Problems

6.3 **Bases in quantum mechanics**
We have just produced an energy eigenbasis for independent non-identical particles, one for independent bosons, and one for independent fermions. In each case did we produce the only possible energy eigenbasis or just one of several possible energy eigenbases? If the particles interact, the states in question will no longer constitute an energy eigenbasis. But will they constitute a basis?

6.4 **Symmetrization in the N-variable case**
Equations (6.8) and (6.9) contain algorithms for constructing (anti)symmetric functions of three variables by summing up six terms. How do these equations generalize to functions of N variables and, in particular, how many terms appear in each sum?

6.5 **Symmetrizing the symmetric**
The non-symmetrized state $|r, s, t\rangle$ is already symmetric in the case that $r = s = t$. What happens if you attempt to symmetrize this already symmetric state through the procedure (6.12)? What if you attempt to antisymmetrize it through the procedure (6.13)?

6.6 **Normalization constants**
Find the normalization constants A_s and A_a in equations (6.12) and (6.13). Be sure your formula works for the case of $\hat{S}|3, 3, 5\rangle$ and $\hat{S}|4, 4, 4\rangle$. Generalize your result to the N-particle case, where the answer is a function of $n_1, n_2, \ldots, n_M$ as well as N.

6.7 **Algorithms for permutations**
Invent some problem concerning the Heap algorithm for generating permutations. Better just give a reference to Sedgwick's paper.

6.8 **Algorithms to list basis states**
Come up with some algorithm for producing the three sets of basis states that we have discussed, preferably in some useful order (where "useful" is to be defined by you). Your algorithm should generalize readily to the N-particle case.

6.9 **Number of basis elements**
Perform the counts (6.18) and (6.21). (Clue: Do it first for $N = 3$, then for N arbitrary.)

6.4 Statistical Mechanics of Independent Identical Particles

6.4.1 Partition function

Now that we have an energy eigenbasis, the obvious thing to do is to calculate the canonical partition function

$$Z(\beta) = \sum_{\text{states}} e^{-\beta E}, \tag{6.26}$$

where for fermions and bosons, respectively, the term "state" implies the occupation number lists:

fermions	$(n_1, n_2, \cdots, n_M)$, $n_r = 0, 1$, subject to $\sum_r n_r = N$
bosons	$(n_1, n_2, \cdots, n_M)$, subject to $\sum_r n_r = N$

As we have seen, it is difficult to even count these lists, much less enumerate them and perform the relevant sum! It can be done, but there is a trick that renders it unnecessary. (Don't be ashamed if you don't see the trick... neither did Einstein or Fermi. They both did it the hard, canonical way.)

The trick here, as in so many places in statistical mechanics, is to use the grand canonical ensemble. In this ensemble, the partition function is

$$\Xi(\beta, \mu) = \sum_{\text{states}} e^{-\beta E + \beta\mu N} = \sum_{\text{states}} e^{-\beta \sum_r (n_r \epsilon_r - \mu n_r)} = \sum_{\text{states}} \prod_{r=1}^{M} e^{-\beta n_r (\epsilon_r - \mu)} \tag{6.27}$$

where the term "state" now implies the occupation number lists *without any restriction on total particle number:*

fermions	$(n_1, n_2, \cdots, n_M)$, $n_r = 0, 1$
bosons	$(n_1, n_2, \cdots, n_M)$

Writing out the sum over states explicitly, we have for fermions

$$\Xi(\beta, \mu) = \sum_{n_1=0}^{1} \sum_{n_2=0}^{1} \cdots \sum_{n_M=0}^{1} \prod_{r=1}^{M} e^{-\beta n_r (\epsilon_r - \mu)} \tag{6.28}$$

$$= \left[\sum_{n_1=0}^{1} e^{-\beta n_1 (\epsilon_1 - \mu)}\right] \left[\sum_{n_2=0}^{1} e^{-\beta n_2 (\epsilon_1 - \mu)}\right] \cdots \left[\sum_{n_M=0}^{1} e^{-\beta n_M (\epsilon_1 - \mu)}\right]. \tag{6.29}$$

A typical factor in the product is

$$\left[\sum_{n_r=0}^{1} e^{-\beta n_r (\epsilon_r - \mu)}\right] = 1 + e^{-\beta(\epsilon_r - \mu)}, \tag{6.30}$$

so for fermions

$$\Xi(\beta, \mu) = \prod_{r=1}^{M} 1 + e^{-\beta(\epsilon_r - \mu)}. \tag{6.31}$$

Meanwhile, for bosons, the explicit state sum is

$$\Xi(\beta, \mu) = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \cdots \sum_{n_M=0}^{\infty} \prod_{r=1}^{M} e^{-\beta n_r (\epsilon_r - \mu)} \tag{6.32}$$

$$= \left[\sum_{n_1=0}^{\infty} e^{-\beta n_1 (\epsilon_1 - \mu)}\right] \left[\sum_{n_2=0}^{\infty} e^{-\beta n_2 (\epsilon_1 - \mu)}\right] \cdots \left[\sum_{n_M=0}^{\infty} e^{-\beta n_M (\epsilon_1 - \mu)}\right], \tag{6.33}$$

and a typical factor in the product is

$$\left[\sum_{n_r=0}^{\infty} e^{-\beta n_r (\epsilon_r - \mu)}\right] = 1 + [e^{-\beta(\epsilon_r - \mu)}] + [e^{-\beta(\epsilon_r - \mu)}]^2 + [e^{-\beta(\epsilon_r - \mu)}]^3 + \cdots \tag{6.34}$$

$$= \frac{1}{1 - e^{-\beta(\epsilon_r - \mu)}}, \tag{6.35}$$

where in the last line we have summed the geometric series under the assumption that $\epsilon_r > \mu$. Thus for bosons

$$\Xi(\beta,\mu) = \prod_{r=1}^{M} \frac{1}{1-e^{-\beta(\epsilon_r-\mu)}}. \tag{6.36}$$

The two results are compactly written together as

$$\Xi(\beta,\mu) = \prod_{r=1}^{M} [1 \pm e^{-\beta(\epsilon_r-\mu)}]^{\pm 1}, \tag{6.37}$$

where the + sign refers to fermions and the − sign to bosons.

6.4.2 Mean occupation numbers

In our previous work, we have always found the partition function and worked from there. Surprisingly, however, for the situation of quantal ideal gases it is more useful to find the mean occupation numbers, such as

$$\langle n_5 \rangle = \frac{\sum_{\text{states}} n_5 e^{-\beta(E-\mu N)}}{\sum_{\text{states}} e^{-\beta(E-\mu N)}} = \frac{\sum_{\text{states}} n_5 e^{-\beta \sum_r (n_r\epsilon_r - n_r\mu)}}{\Xi(\beta,\mu)}. \tag{6.38}$$

Note that the averages $\langle n_r \rangle$ are functions of β and μ (as well as of r) but it is notationally clumsy to show that dependence.

How can such averages be evaluated? A slick trick would be helpful here! Consider the derivative

$$\frac{\partial \ln \Xi}{\partial \epsilon_5} = \frac{1}{\Xi}\frac{\partial \Xi}{\partial \epsilon_5} = \frac{1}{\Xi}\sum_{\text{states}} (-\beta n_5) e^{-\beta \sum_r (n_r\epsilon_r - n_r\mu)} = -\beta \langle n_5 \rangle. \tag{6.39}$$

Using the explicit expression (6.37) for Ξ (where the + sign refers to fermions and the − sign to bosons), this gives

$$\begin{aligned}
\langle n_5 \rangle &= -\frac{1}{\beta}\frac{\partial \ln \Xi}{\partial \epsilon_5} && (6.40)\\
&= -\frac{1}{\beta}\left\{\frac{\partial}{\partial \epsilon_5}\sum_{r=1}^{M} \ln[1 \pm e^{-\beta(\epsilon_r-\mu)}]^{\pm 1}\right\} && (6.41)\\
&= -\frac{1}{\beta}\left\{\frac{\partial}{\partial \epsilon_5}\ln[1 \pm e^{-\beta(\epsilon_5-\mu)}]^{\pm 1}\right\} && (6.42)\\
&= -\frac{1}{\beta}\left\{\pm\frac{\pm(-\beta)e^{-\beta(\epsilon_5-\mu)}}{[1 \pm e^{-\beta(\epsilon_5-\mu)}]}\right\} && (6.43)\\
&= \frac{1}{e^{\beta(\epsilon_5-\mu)} \pm 1}, && (6.44)
\end{aligned}$$

leaving us with the final result

$$\langle n_r \rangle = \frac{1}{e^{\beta(\epsilon_r-\mu)} \pm 1}. \tag{6.45}$$

As before, the + sign refers to fermions and the − sign to bosons.

The mean occupation numbers play such an important role that it is easy to forget that they are only averages, that there will be fluctuations, that for a given T and μ not all states will have exactly $\langle n_5 \rangle$ building blocks of level 5 (see problem 6.13). Keep this in mind if you ever find yourself saying "occupation number" rather than "mean occupation number".

In practice, these results from the grand canonical ensemble are used as follows: One uses these results to find quantities of interest as functions of temperature, volume, and chemical potential, such as the pressure $p(T, V, \mu)$. But most experiments are done with a fixed number of particles N, so at the very end of your calculation you will want to find $\mu(T, V, N)$ in order to express your final answer as $p(T, V, N)$. You can find $\mu(T, V, N)$ by demanding that

$$N = \sum_{r=1}^{M} \langle n_r \rangle = \sum_{r=1}^{M} \frac{1}{e^{\beta(\epsilon_r - \mu)} \pm 1}. \tag{6.46}$$

In other words, the quantity μ serves as a parameter to insure normalization, very much as the quantity Z serves to insure normalization in the canonical ensemble through

$$1 = \sum_n \frac{e^{-\beta E_n}}{Z}. \tag{6.47}$$

You might wonder, in fact, about the relation between the canonical probability

$$\frac{e^{-\beta E_n}}{Z}, \tag{6.48}$$

which we have seen many times before, and the recently derived occupancy probability

$$\frac{1}{N} \frac{1}{e^{\beta(\epsilon_r - \mu)} \pm 1}. \tag{6.49}$$

The first result applies to both interacting and non-interacting systems, both classical and quantal. The second applies only to non-interacting quantal systems. Why do we need a new probability? What was wrong with our derivation (in section 4.1) of the canonical probability that requires us to replace it with an occupancy probability? The answer is that nothing was wrong and that the occupancy probability doesn't replace the canonical probability. The canonical probability and the occupancy probability answer different questions. The first finds the probability that the entire system is in the many-body state n. The second finds the probability that the one-body level r is used as a building block in constructing the many-body state. Indeed, although we derived the occupancy probability result through a grand canonical argument, it is also possible to derive the occupancy probabilities from strict canonically arguments, proof that these two probabilities can coexist peacefully.

6.4.3 The Boltzmann limit

This is the limit where particles are far enough apart that overlap of wavefunction is minimal, so we needn't worry about symmetrization or antisymmetrization. Equivalently, it is the limit where $\langle n_r \rangle \ll 1$ for all r.

6.4.4 Problems

6.10 **Evaluation of the grand canonical partition function**

Can you find a simple expression for $\Xi(\beta, \mu)$ for non-interacting particles in a one-dimensional harmonic well? For non-interacting particles in a one-dimensional infinite square well? For any other potential? Can you do anything valuable with such an expression once you've found it?

6.11 **Entropy of quantal ideal gases**

This problem derives an expression for the entropy of a quantal ideal gas in terms of the mean occupation numbers $\langle n_r \rangle$. (Compare problem 4.3.) Throughout the problem, in the symbols $\pm$ and $\mp$, the top sign refers to fermions and the bottom sign refers to bosons.

a. Use the connection between thermodynamics and statistical mechanics to show that, for any system,

$$\frac{S(T, V, \mu)}{k_B} = \ln \Xi - \beta \frac{\partial \ln \Xi}{\partial \beta}. \tag{6.50}$$

b. Show that for the quantal ideal gas,

$$\ln \Xi(T, V, \mu) = \mp \sum_r \ln(1 \mp \langle n_r \rangle). \tag{6.51}$$

c. The mean occupation numbers $\langle n_r \rangle$ are functions of T, V, and μ (although it is notationally clumsy to show this dependence). Show that

$$\begin{aligned} \beta \left. \frac{\partial \langle n_r \rangle}{\partial \beta} \right)_{V,\mu} &= -\beta(\epsilon_r - \mu)\langle n_r \rangle (1 \mp \langle n_r \rangle) \\ &= -[\ln(1 \mp \langle n_r \rangle) - \ln\langle n_r \rangle]\langle n_r \rangle (1 \mp \langle n_r \rangle). \end{aligned} \tag{6.52}$$

d. Finally, show that

$$S(T, V, \mu) = -k_B \sum_r [\langle n_r \rangle \ln\langle n_r \rangle \pm (1 \mp \langle n_r \rangle) \ln(1 \mp \langle n_r \rangle)]. \tag{6.53}$$

e. Find a good approximation for this expression in the Boltzmann limit, $\langle n_r \rangle \ll 1$.

f. (Optional.) Find an expression for C_V in terms of the quantities $\langle n_r \rangle$.

6.12 **Isothermal compressibility of quantal ideal gases**

a. Show that in a quantal ideal gas, the isothermal compressibility is

$$\kappa_T = \frac{1}{\rho k_B T} \left[1 \mp \frac{\sum_r \langle n_r \rangle^2}{\sum_r \langle n_r \rangle} \right], \tag{6.54}$$

where as usual the top sign refers to fermions and the bottom sign to bosons. (Clue: Choose the most appropriate expression for κ_T from those uncovered in problem 3.33.)

b. Compare this expression to that for a classical ("Maxwell-Boltzmann") ideal gas.

c. The negative sign in the expression for fermions opens the possibility that κ_T could be negative. Prove that this potential horror never happens.

d. Do the relative sizes of the three compressibilites (fermion, classical, boson) adhere to your qualitative expectations? (Compare problem 6.29.)

6.13 **Dispersion in occupation number**

Find an expression analogous to (6.45) giving the dispersion in the occupation numbers. (Clue: A slick trick would be helpful here.) Answer:

$$\Delta n_r = \frac{1}{e^{\beta(\epsilon_r-\mu)/2} \pm e^{-\beta(\epsilon_r-\mu)/2}} = \sqrt{\langle n_r\rangle(1 \mp \langle n_r\rangle)} \tag{6.55}$$

6.5 Quantum Mechanics of Free Particles

"Particle in a box." Periodic boundary conditions. k-space. In the thermodynamic limit, the dots in k-space become densely packed, and it seems appropriate to replace sums over levels with integrals over k-space volumes. (In fact, there is at least one situation (see equation 6.83) in which this replacement is *not* correct.)

Density of levels in k-space:

$$\frac{V}{8\pi^3} \qquad \text{(worth memorizing).} \tag{6.56}$$

Energy density of levels:

$$\text{number of one-body levels with } \epsilon_r \text{ from } \mathcal{E} \text{ to } \mathcal{E} + d\mathcal{E} \equiv G(\mathcal{E})\, d\mathcal{E} = V\frac{\sqrt{2m^3}}{2\pi^2\hbar^3}\sqrt{\mathcal{E}}\, d\mathcal{E}. \tag{6.57}$$

How to use energy density of levels:

$$\sum_r f(\epsilon_r) \approx \int_0^\infty G(\mathcal{E}) f(\mathcal{E})\, d\mathcal{E} \tag{6.58}$$

and this approximation (usually) becomes exact in the thermodynamic limit.

6.5.1 Problems

6.14 **Free particles in a box**

We argued that, for a big box, periodic boundary conditions would give the same results as "clamped boundary conditions". Demonstrate this by finding the density of levels for three-dimensional particle in a box problem.

6.15 **Density of levels for bound particles**

What is the density of levels $G(\mathcal{E})$ for a one-dimensional harmonic oscillator with spring constant K? For a three-dimensional isotropic harmonic oscillator?

6.16 **Density of levels in d dimensions**
What is the density of levels $G(\mathcal{E})$ for free particles subject to periodic boundary conditions in a world of d dimensions?

6.6 Fermi-Dirac Statistics

In three dimensions, the chemical potential μ *decreases* with temperature. Why? It is clear that at $T = 0$, $\mu = \mathcal{E}_F > 0$. But as the temperature rises the gas approaches the classical limit, for which $\mu < 0$ (see problem 2.23). This is not proof, but it makes sense that μ should decrease with increasing temperature. A proof is available but surprisingly difficult.

6.6.1 Problems

6.17 **Qualitative origin of $\mathcal{E}_F$**
(This problem is stolen from a GRE Physics test.)
The mean kinetic energy of electrons in metals at room temperature is usually many times the thermal energy k_BT. Which of the following can best be used to explain this fact?

a. The time-energy uncertainty relation.
b. The Pauli exclusion principle.
c. The degeneracy of the energy levels.
d. The Born approximation.
e. The wave-particle duality.

6.18 **Fermion gas in two dimensions**
(This problem is based on one in Ashcroft and Mermin, page 53.)
Consider a gas of free, independent, spin-$\frac{1}{2}$ fermions in two dimensions. The gas is contained within an area (or two dimensional volume) of A.

a. What is the density of one-particle levels in k-space?
b. How does the Fermi energy $\mathcal{E}_F$ depend upon the density N/A?
c. Use $\sum_r \langle n_r \rangle = N$ to show that

$$\mu + k_BT \ln(1 + e^{-\mu/k_BT}) = \mathcal{E}_F. \tag{6.59}$$

Notice that the chemical potential μ decreases with temperature.

6.19 **Dependence of chemical potential on temperature**
Show that for independent (not necessarily free) fermions, the $\mu(T)$ curve has slope (when N and V are constant)

$$\frac{d\mu}{dT} = -\frac{1}{T}\frac{\int_0^\infty G(\mathcal{E})\mathrm{sech}^2(\beta(\mathcal{E}-\mu)/2)(\mathcal{E}-\mu)\,d\mathcal{E}}{\int_0^\infty G(\mathcal{E})\mathrm{sech}^2(\beta(\mathcal{E}-\mu)/2)\,d\mathcal{E}}. \tag{6.60}$$

Can you use this result to show that the chemical potential must decrease with temperature? I can't.

6.20 **Thermodynamics of the fermion gas**

Consider a collection of free and independent spin-$\frac{1}{2}$ fermions. Do *not* assume that the temperature vanishes.

a. Use the fundamental grand canonical result $\Pi = -k_B T \ln \Xi$ to show that

$$p(T,\mu)V = k_B T \int_0^\infty G(\mathcal{E}) \ln(1 + e^{\beta(\mu - \mathcal{E})})\, d\mathcal{E}. \tag{6.61}$$

b. Use the expression for $G(\mathcal{E})$ and the change of variable $x = \beta\mathcal{E}$ to find

$$p(T,\mu) = (k_B T)^{5/2} \left[\frac{\sqrt{2m^3}}{\pi^2 \hbar^3}\right] \int_0^\infty \sqrt{x} \ln(1 + e^{\beta\mu} e^{-x})\, dx. \tag{6.62}$$

c. Integrate by parts to obtain

$$p(T,\mu) = \frac{2}{3}(k_B T)^{5/2} \left[\frac{\sqrt{2m^3}}{\pi^2 \hbar^3}\right] \int_0^\infty \frac{x^{3/2}}{e^x e^{-\beta\mu} + 1} dx. \tag{6.63}$$

(Do not attempt to evaluate the integral that remains.)

d. Meanwhile, show that the total energy

$$E(T,V,\mu) = \int_0^\infty G(\mathcal{E}) f(\mathcal{E}) \mathcal{E}\, d\mathcal{E} \tag{6.64}$$

is given by

$$E(T,V,\mu) = V(k_B T)^{5/2} \left[\frac{\sqrt{2m^3}}{\pi^2 \hbar^3}\right] \int_0^\infty \frac{x^{3/2}}{e^x e^{-\beta\mu} + 1} dx. \tag{6.65}$$

Thus the pressure and energy are related by

$$pV = \tfrac{2}{3} E, \tag{6.66}$$

which is exactly the same relationship they have in the classical monatomic ideal gas! (See problem 6.28.)

6.21 **Pressure at zero temperature**

Use the result of the previous problem to show that at absolute zero the pressure of a gas of free and independent spin-$\frac{1}{2}$ fermions does not vanish (as it does for a classical ideal gas). Instead, the zero-temperature pressure is $\frac{2}{5}\rho\mathcal{E}_F$.

6.22 **Mass-radius relation for white dwarf stars**

(This problem is modified from Kittel and Kroemer, *Thermal Physics*, second edition, page 216.)

A white dwarf star (see Kittel and Kroemer, pp. 196–198) consists of highly compressed hydrogen in which the atoms have ionized into independent protons and electrons. In most cases it is a good model to assume that the electrons are free and independent non-relativistic fermions at zero temperature. Consider a white dwarf of mass M and radius R, containing N electrons.

a. Show that to a very good approximation, $N = M/m_p$, where m_p is the mass of a proton.

b. Show that the gravitational potential energy of a uniform sphere of mass M and radius R is $-cGM^2/R$, where G is the gravitational constant and c is a dimensionless constant. (In fact $c = \frac{3}{5}$ but this is tedious to show.)

c. Show that the kinetic energy of the electrons is

$$\mathrm{KE} = \frac{9}{20}\left(\frac{3\pi^2}{2}\right)^{1/3} \frac{\hbar^2}{m_e m_p^{5/3}} \frac{M^{5/3}}{R^2}, \tag{6.67}$$

where m_e is the mass of an electron.

d. If the potential and kinetic energies satisfy

$$\mathrm{KE} = -\tfrac{1}{2}\mathrm{PE}, \tag{6.68}$$

as required by the virial theorem of mechanics, show that

$$RM^{1/3} = \text{ a constant of approximate value } 10^{17} \text{ m kg}^{1/3}. \tag{6.69}$$

Evaluate the constant assuming that $c = \frac{3}{5}$. Note that the radius *decreases* as the mass *increases*.

e. If the white dwarf has the mass of the sun (2×10^{30} kg), what is its radius (in km)? Compare this to the radius of our sun.

f. Neutron stars (observed as pulsars) are also zero temperature fermion gases, but in this case the fermions are neutrons rather than electrons. Derive the mass-radius relation for a neutron star, and use it to find the radius of a neutron star with the mass of the sun.

6.7 Bose-Einstein Statistics

Perhaps you thought the Fermi-Dirac results were strange: Non-interacting particles forming a collection as hard as steel. . . room temperature being effectively zero. Wait until you see the Bose-Einstein results.

6.7.1 Theory

For independent bosons, whether free or subject to an external potential, the mean occupation number function is

$$b(\mathcal{E}) = \frac{1}{e^{\beta(\mathcal{E}-\mu)} - 1}. \tag{6.70}$$

To begin to understand this function, note that

when:	we have:
$\mathcal{E} < \mu$	$b(\mathcal{E})$ negative
$\mathcal{E} = \mu$	$b(\mathcal{E}) = \infty$
$\mathcal{E} > \mu$	$b(\mathcal{E})$ positive

Thus this function has the general character sketched below:

Although we have written $b(\mathcal{E})$ as a function of the continuous variable $\mathcal{E}$, we will in fact have occasion to evaluate it only at the energy eigenvalues ϵ_1, ϵ_2, ϵ_3,..., ϵ_r,..., eigenvalues which of course differ for different external potentials.

It seems bizarre that $b(\mathcal{E})$ can be negative, and indeed this is only a mathematical artifact: Recall that in our derivation of the Bose function we needed to assume that $\mu < \epsilon_1$ in order to insure convergence (see equation 6.35). Evaluating $b(\mathcal{E})$ at any eigenvalue ϵ_r will always result in a positive mean occupation number.

The character of the Bose function is dominated by the singularity at $\mathcal{E} = \mu$, so in trying to understand the function and its physical implications one must first locate the chemical potential. This section will provide a tour of Bose-Einstein behavior with decreasing temperature, throughout which the chemical potential shifts to the right. (When we investigated Fermi-Dirac behavior, we started at $T = 0$ and toured with increasing temperature, so the chemical potential shifted left.) This rightward shift presents a potential problem, because as the temperature decreases μ might shift right all the way to ϵ_1, and we know that $\mu < \epsilon_1$. We will just have to go ahead and take the tour, being wary because we know that a $\mu = \epsilon_1$ road block might pop up right in front of us as we view the countryside. With any luck μ will not yet have reached ϵ_1 when our tour halts at $T = 0$.

For the case of free and independent bosons (subject to periodic boundary conditions), the ground level energy is $\epsilon_1 = 0$. The natural first step is to find $\mu(T, V, N)$ by demanding that

$$N = \int_0^\infty G(\mathcal{E}) b(\mathcal{E})\, d\mathcal{E}. \tag{6.71}$$

Natural though this may be, caution is in order. Remember that the integral above is an approximation to the sum over discrete energy levels

$$N = \sum_r b(\epsilon_r). \tag{6.72}$$

It is legitimate to replace the sum with the integral when the value of $b(\mathcal{E})$ changes little from one energy level to the next. We saw in section 6.5 that in the thermodynamic limit, the free-particle level spacings approach zero, so for the most part this approximation is excellent... even exact in the thermodynamic limit. But there is one exception: At $\mathcal{E} = \mu$, the Bose function $b(\mathcal{E})$ is infinite, so $b(\mu) = \infty$ is very different from $b(\mu + \delta)$, no matter how small the positive number δ is. Usually we can ignore this caution, because $\mu < \epsilon_1 = 0$. But if μ approaches 0 then we expect the approximation (6.71) to fail.

In summary, the integral (6.71) is a good approximation for the sum (6.72) as long as the integrand varies slowly. Now for any value of $\mathcal{E}$ greater than μ, you can make $b(\mathcal{E} + \delta)$ very close to $b(\mathcal{E})$ simply by choosing $\delta > 0$ small enough. This is what happens in the thermodynamic limit. But for $\mathcal{E} = \mu$, there is *always* a large difference between $b(\mathcal{E}) = \infty$ and $b(\mathcal{E} + \delta)$, which is finite. Thus the integral approximation will be a good one as long as we avoid $\mathcal{E} = \mu$.

In situations where the sum *can* be legitimately replaced with the integral, we have

$$N = \int_0^\infty G(\mathcal{E}) b(\mathcal{E})\, d\mathcal{E} \tag{6.73}$$

$$= V \left[\frac{\sqrt{2m^3}}{2\pi^2 \hbar^3} \right] \int_0^\infty \sqrt{\mathcal{E}} \frac{1}{e^{\beta(\mathcal{E}-\mu)} - 1} d\mathcal{E}. \tag{6.74}$$

Use of the obvious substitution $x = \beta\mathcal{E}$ gives

$$N = V \left[\frac{\sqrt{2m^3}}{2\pi^2 \hbar^3} \right] (k_B T)^{3/2} \int_0^\infty \frac{x^{1/2}}{e^x e^{-\beta\mu} - 1} dx, \tag{6.75}$$

and remembering the definition (5.4) of thermal de Broglie wavelength results in a more convenient expression

$$N = \frac{2}{\sqrt{\pi}} \frac{V}{\lambda^3(T)} \int_0^\infty \frac{x^{1/2}}{e^x e^{-\beta\mu} - 1} dx. \tag{6.76}$$

Note that the definite integral above is *not* "just a number"... it is a function of the product $\beta\mu$. There is no closed-form expression for the integral, although it is readily evaluated numerically and can be found in tabulations of functions. However it is easy to find an analytic upper bound: Because $\mu < 0$, we have $e^{-\beta\mu} > 1$ whence

$$\int_0^\infty \frac{x^{1/2}}{e^x e^{-\beta\mu} - 1} dx < \int_0^\infty \frac{x^{1/2}}{e^x - 1} dx. \tag{6.77}$$

The expression on the right *is* just a number, and a little bit of work (see problem 6.26) shows that it is the number

$$\zeta(\tfrac{3}{2}) \frac{\sqrt{\pi}}{2}, \tag{6.78}$$

where the Riemann zeta function is defined by

$$\zeta(s) = \sum_{n=1}^\infty \frac{1}{n^s} \quad \text{and} \quad \zeta(\tfrac{3}{2}) = 2.612375348\ldots. \tag{6.79}$$

So, how does this upper bound help us? It shows that

$$N = \frac{2}{\sqrt{\pi}} \frac{V}{\lambda^3(T)} \int_0^\infty \frac{x^{1/2}}{e^x e^{-\beta\mu} - 1} dx < \frac{V}{\lambda^3(T)} \zeta(\tfrac{3}{2}), \tag{6.80}$$

but what is the significance of this result? Remember that the upper bound is just the value of the function when $\mu = 0$, which is exactly where we expect a road block due to the breakdown of the integral approximation (6.71). Our hopes that we could avoid the issue have been dashed. The breakdown occurs at the temperature T_0 that satisfies

$$N = \frac{V}{\lambda^3(T_0)}\zeta(\tfrac{3}{2}) \quad \text{or} \quad \lambda^3(T_0) = \zeta(\tfrac{3}{2})/\rho \tag{6.81}$$

or

$$T_0(\rho) = \frac{h^2}{2\pi m k_B \zeta^{2/3}(\frac{3}{2})}\rho^{2/3}. \tag{6.82}$$

For temperatures above T_0, the chemical potential shifts right as the temperature falls, and the integral approximation (6.71) is legitimate (indeed, exact in the thermodynamic limit). But as the temperature decreases below T_0, the chemical potential sticks at $\mu = 0$ and the integral approximation (6.71) must be modified.

What is the proper modification? The function $b(\mathcal{E})$ is slowly varying for all values of $\mathcal{E}$ except $\mathcal{E} = \mu$, so the integral approximation is legitimate for all the energy levels *except* the ground level at $\epsilon_1 = 0 = \mu$. Only for the ground level is some other result needed, so we just add in the ground level occupancy by hand:

$$N = \langle n_1 \rangle + \int_0^\infty G(\mathcal{E}) b(\mathcal{E})\, d\mathcal{E}. \tag{6.83}$$

Now, we have already seen that when $\mu = 0$—the condition for validity of this equation—the integral can be evaluated exactly and we have

$$N = \langle n_1 \rangle + \frac{V}{\lambda^3(T)}\zeta(\tfrac{3}{2}). \tag{6.84}$$

Note that $\langle n_1 \rangle$ here is *not* given by its traditional formula (6.45), because

$$\frac{1}{e^{\beta(\mathcal{E}-\mu)} - 1} = \frac{1}{e^0 - 1} = \infty. \tag{6.85}$$

Instead, equation (6.84) *is* the formula for $\langle n_1 \rangle$ when $T < T_0$. The mean ground level occupancy $\langle n_1 \rangle$ is an intensive quantity when $T > T_0$ but an extensive quantity when $T < T_0$.

In summary, the correct normalization equation breaks into two parts, namely

$$N = \begin{cases} \dfrac{2}{\sqrt{\pi}} \dfrac{V}{\lambda^3(T)} \displaystyle\int_0^\infty \frac{x^{1/2}}{e^x e^{-\beta\mu} - 1} dx & \text{for } T > T_0(\rho) \\ \langle n_1 \rangle + \dfrac{V}{\lambda^3(T)}\zeta(\tfrac{3}{2}) & \text{for } T < T_0(\rho). \end{cases} \tag{6.86}$$

We should expect that each part will behave quite differently, i.e. we expect a sudden change of behavior as the temperature drops through T_0.

What does all this mean physically? A naive reading of equation (6.80) suggests an upper bound on the number of particles that can be placed into the volume V. This would be sensible if the particles were marbles with hard-core repulsions. But these are non-interacting particles! Surely we can add more particles just by throwing them into the container. Indeed we can do so, and the associated excess mean occupation

number is due to the level that particles like best at low temperatures, namely the ground level. The ground level thus has a much higher mean occupancy than the first excited level, and this rapid variation of mean occupancy with energy renders the approximation of sum by integral invalid. The inequality (6.80) does not limit the number of particles in the system: instead it shows the domain within which it is legitimate to approximate the sum (6.72) by the integral (6.71).

The abrupt transition at $T_0(\rho)$ is called Bose-Einstein condensation and the material at temperatures below $T_0(\rho)$ is called the Bose condensate. These terms are unfortunate: they conjure images of a gas condensing into a liquid, in which circumstance the atoms separate into two different classes: those in the liquid and those remaining in the vapor. This suggests that in Bose-Einstein condensation too there are two classes of particles: those in the ground level and those in the excited levels. This picture is totally false. It is incorrect to say "one particle is in the ground level, another is in the fourth excited level". In truth the individual particles are not in individual levels at all: instead the whole system is in a state produced by multiplying together the individual level wavefunctions ("building blocks") and then symmetrizing them. The literature of Bose-Einstein statistics is full of statements like "at temperatures below T_0, any particle added to the system goes into the ground level." Such statements are wrong. They should be replaced with statements like "at temperatures below T_0, any increase in particle number occurs through an increase in $\langle n_1 \rangle$, the mean occupancy of the ground level." Or alternatively, "at temperatures below T_0, it is very likely that many of the building blocks from which the system wavefunction is built are the ground level." Or again, to be absolutely precise, "at temperatures below T_0, if the energy is measured then it is very likely that many of the building blocks from which the resulting energy eigenfunction is built are the ground level." Read again the previous paragraph—the one that begins "What does all this mean physically?". Notice that I never need to say that a particle "is in" or "goes into" a given level.

6.7.2 Experiment

References: M.H. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman, E.A. Cornell, "Observation of Bose-Einstein condensation in a dilute atomic vapor", *Science*, **269** (14 July 1995) 198–201; Randall G. Hullet at Rice; Malcolm W. Browne, "Two groups of physicists produce matter that Einstein postulated", New York *Times*, 14 July 1995, page 1.

6.7.3 Problems

6.23 **Character of the Bose function**

What are the limits of the Bose function $b(\mathcal{E})$ (equation 6.70) as $\mathcal{E} \to \pm\infty$? Is the curvature of the function greater when the temperature is high or when it is low?

6.24 **Thermodynamics of the Bose condensate**

For temperatures less than the Bose condensation temperature T_0, find the energy, heat capacity, and entropy of an ideal gas of spin-zero bosons confined to a volume V. Write your answers in terms of the

dimensionless integral

$$I = \int_0^\infty \frac{x^{3/2}}{e^x - 1}\, dx \tag{6.87}$$

but don't bother to evaluate it. Show that

$$C_V = \frac{5}{2}\frac{E}{T} \quad \text{and} \quad S = \frac{5}{3}\frac{E}{T}. \tag{6.88}$$

6.25 **More thermodynamics of the Bose condensate**

For the system of the previous problem, show that

$$F = -\frac{2}{3}E \quad \text{and} \quad p = \frac{2}{3}\frac{E}{V}. \tag{6.89}$$

From this show that at low temperatures, the pressure of a collection of free and independent bosons goes like $p \sim T^{5/2}$. (This is always less than the classical pressure $p \sim T$.)

6.26 **An integral important for Bose condensation**

Show that

$$\int_0^\infty \frac{x^{1/2}}{e^x - 1} dx = \zeta(\tfrac{3}{2})\frac{\sqrt{\pi}}{2} \tag{6.90}$$

where

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}. \tag{6.91}$$

Clue:

$$\frac{1}{e^x - 1} = \frac{1}{e^x(1 - e^{-x})} = e^{-x} \sum_{n=0}^{\infty} e^{-nx} \quad \text{for} \quad x > 0. \tag{6.92}$$

6.8 Specific Heat of the Ideal Fermion Gas

There is a "set of the pants" heuristic argument showing that at low temperatures,

$$C_V(T) \approx k_B G(\mathcal{E}_F)(k_B T). \tag{6.93}$$

This argument is not convincing in detail as far as the magnitude is concerned and, indeed, we will soon find that it is wrong by a factor of $\pi^2/3$. On the other hand it is quite clear that the specific heat will increase linearly with T at low temperatures. From equipartition, the classical result is

$$C_V^{\text{classical}} = \tfrac{3}{2} N k_B, \tag{6.94}$$

and we expect this result to hold at high temperature. Thus our overall expectation is that the specific heat will behave as sketched below.

How can we be more precise and rigorous? That is the burden of this section.

Reminders:

$$f(\mathcal{E}) = \frac{1}{e^{(\mathcal{E}-\mu)/k_BT}+1} \tag{6.95}$$

$$\mathcal{E}_F = \frac{\hbar^2}{2m}\left(3\pi^2\frac{N}{V}\right)^{2/3} \tag{6.96}$$

$$G(\mathcal{E}) = V\frac{\sqrt{2m^3}}{\pi^2\hbar^3}\sqrt{\mathcal{E}} = N\frac{3}{2}\frac{1}{\mathcal{E}_F^{3/2}}\sqrt{\mathcal{E}} \tag{6.97}$$

The expressions for N and E are

$$\begin{aligned} N(T,V,\mu) &= \int_0^\infty G(\mathcal{E})f(\mathcal{E})\,d\mathcal{E} &&= V\frac{\sqrt{2m^3}}{\pi^2\hbar^3}\int_0^\infty \mathcal{E}^{1/2}f(\mathcal{E})\,d\mathcal{E} \\ E(T,V,\mu) &= \int_0^\infty \mathcal{E}G(\mathcal{E})f(\mathcal{E})\,d\mathcal{E} &&= V\frac{\sqrt{2m^3}}{\pi^2\hbar^3}\int_0^\infty \mathcal{E}^{3/2}f(\mathcal{E})\,d\mathcal{E}. \end{aligned} \tag{6.98}$$

Remember how to use these expressions: The first we will invert to find $\mu(T,V,N)$, then we will plug this result into the second to find $E(T,V,N)$. As a preliminary, recognize that we can clean up some messy expressions in terms of the physically significant constant $\mathcal{E}_F$ by writing these two expressions as

$$1 = \frac{3}{2}\frac{1}{\mathcal{E}_F^{3/2}}\int_0^\infty \mathcal{E}^{1/2}f(\mathcal{E})\,d\mathcal{E} \tag{6.99}$$

$$\frac{E}{N} = \frac{3}{2}\frac{1}{\mathcal{E}_F^{3/2}}\int_0^\infty \mathcal{E}^{3/2}f(\mathcal{E})\,d\mathcal{E}. \tag{6.100}$$

In short, we must evaluate integrals like

$$\int_0^\infty \frac{\mathcal{E}^{\alpha-1}}{e^{(\mathcal{E}-\mu)/k_BT}+1}d\mathcal{E} \tag{6.101}$$

with the dimensions of [energy]$^\alpha$. To make these more mathematical and less physical, convert to the dimensionless quantities

$$x = \mathcal{E}/k_BT \tag{6.102}$$

$$x_0 = \mu/k_BT \tag{6.103}$$

to find

$$\int_0^\infty \frac{\mathcal{E}^{\alpha-1}}{e^{(\mathcal{E}-\mu)/k_BT}+1}d\mathcal{E} = (k_BT)^\alpha \int_0^\infty \frac{x^{\alpha-1}}{e^{x-x_0}+1}dx. \tag{6.104}$$

Define the dimensionless integral

$$A_\alpha(x_0) = \int_0^\infty \frac{x^{\alpha-1}}{e^{x-x_0}+1}dx. \tag{6.105}$$

so that

$$1 = \frac{3}{2}\frac{(k_BT)^{3/2}}{\mathcal{E}_F^{3/2}}A_{3/2}(\mu/k_BT) \tag{6.106}$$

$$\frac{E}{N} = \frac{3}{2}\frac{(k_BT)^{5/2}}{\mathcal{E}_F^{3/2}}A_{5/2}(\mu/k_BT). \tag{6.107}$$

Let us pause before rushing forward. You might think that we should turn to Mathematica (or to a table of integrals) and try to evaluate these integrals in terms of well-known functions like exponentials and Bessel functions. Even if we could do this, it wouldn't help: The result would just be an incomprehensible morass of functions, and to try to understand it we would need to, among other things, look at the low-temperature limit and the high-temperature limit. Since we are particularly interested in the low-temperature behavior, let's just set out to find the low-temperature series in the first place. The meaning of "low temperature" in this context is $k_BT \ll \mu$ or $x_0 = \mu/k_BT \gg 1$, so we suspect a series like

$$E(T) = E(T=0) + \frac{E_1}{x_0} + \frac{E_2}{x_0^2} + \cdots. \tag{6.108}$$

Whenever we have previously searched for such a series (for example, when we investigated internal specific heat of the rigid dumbbell, section 5.4, or the simple harmonic oscillator, problem 5.7) we had good luck using a Taylor series expansion about the origin. In this case, any such technique is doomed to failure, for the following reason: We need a good approximation for the function

$$g(x) = \frac{1}{e^{x-x_0}+1}, \tag{6.109}$$

a function that is very flat near the origin.

In fact, this function is so flat at the origin that, even if we kept dozens of terms in its Taylor series expansion about the origin, the Taylor approximation would be quite poor.

Indeed, all of the *action* associated with this function happens near $x = x_0$. (We saw earlier, in our heuristic discussion, that the low-temperature specific heat was due entirely to promotion of a few electrons near the Fermi energy.) Thus an accurate low-temperature approximation will necessarily involve expansions around $x = x_0$, not around $x = 0$.

Arnold Sommerfeld was aware this situation and had the genius to recognize that a change in focus would solve the problem. Specifically, the focus needs to change from the function $g(x)$ to its derivative $g'(x)$, because the derivative is very small for all x except near $x = x_0$, where the action is. In specific, he noted that

$$\begin{aligned} g'(x) &= -\frac{e^{x-x_0}}{(e^{x-x_0}+1)^2} \\ &= -\frac{1}{(e^{-(x-x_0)/2}+e^{(x-x_0)/2})^2} \\ &= -\frac{1}{4\cosh^2((x-x_0)/2)}. \end{aligned}$$

The last form makes clear the remarkable fact that $g'(x)$ is even under reflections about x_0.

How can we use these facts about $g'(x)$ to solve integrals involving $g(x)$? Through integration by parts.

$$\begin{aligned} A_\alpha(x_0) &= \int_0^\infty x^{\alpha-1} g(x)\, dx \\ &= \left[\frac{x^\alpha}{\alpha} g(x)\right]_0^\infty - \int_0^\infty \frac{x^\alpha}{\alpha} g'(x)\, dx \\ &= \frac{1}{4\alpha}\int_0^\infty \frac{x^\alpha}{\cosh^2((x-x_0)/2)} dx \end{aligned} \tag{6.110}$$

It's now easy to plot the pieces of the integrand, and to see that the integrand nearly vanishes except near $x = x_0$.

Because x^α varies slowly near $x = x_0$, it makes sense to expand it in a Taylor series about $x = x_0$:

$$\begin{array}{rclclclcl} f(x) & = & f(x_0) & + & f'(x_0)(x-x_0) & + & \frac{1}{2}f''(x_0)(x-x_0)^2 & + & \cdots \\ x^\alpha & = & x_0^\alpha & + & \alpha x_0^{\alpha-1}(x-x_0) & + & \frac{1}{2}\alpha(\alpha-1)x_0^{\alpha-2}(x-x_0)^2 & + & \cdots \end{array}$$

So

$$\begin{aligned} 4\alpha A_\alpha(x_0) &= \int_0^\infty \frac{x^\alpha}{\cosh^2((x-x_0)/2)}dx \\ &= x_0^\alpha \int_0^\infty \frac{1}{\cosh^2((x-x_0)/2)}dx + \alpha x_0^{\alpha-1}\int_0^\infty \frac{(x-x_0)}{\cosh^2((x-x_0)/2)}dx \\ &\qquad + \tfrac{1}{2}\alpha(\alpha-1)x_0^{\alpha-2}\int_0^\infty \frac{(x-x_0)^2}{\cosh^2((x-x_0)/2)}dx + \cdots. \end{aligned}$$

Let's think about how to evaluate integrals like

$$\int_0^\infty \frac{(x-x_0)^n}{\cosh^2((x-x_0)/2)}dx. \tag{6.111}$$

The denominator is even in the variable $(x - x_0)$, while the numerator is either even or odd in the variable $(x - x_0)$. If the lower limit were $-\infty$ instead of 0, half of these integrals would vanish... an immense labor savings! Now, of course, the lower limit isn't $-\infty$, but on the other hand if we extended the lower limit from 0 to $-\infty$, we wouldn't pick up much error, because the integrand nearly vanishes when x is negative. In fact, the error introduced is

$$\int_{-\infty}^0 \frac{(x-x_0)^n}{\cosh^2((x-x_0)/2)}dx = \int_{-\infty}^\infty \frac{(x-x_0)^n}{\cosh^2((x-x_0)/2)}dx - \int_0^\infty \frac{(x-x_0)^n}{\cosh^2((x-x_0)/2)}dx, \tag{6.112}$$

and the scale of this error is about the value of the integrand at $x = 0$, namely

$$\frac{(-x_0)^n}{\cosh^2(x_0/2)} \approx (-x_0)^n(4e^{-x_0}).$$

We could find more accurate approximations for this error, but there's no need to. We know the error is of order e^{-x_0} and we're interested in case that $x_0 \gg 1$. This error is utterly negligible. Remember that we expect an answer of the form

$$E(T) = E(T=0) + \frac{E_1}{x_0} + \frac{E_2}{x_0^2} + \cdots, \tag{6.113}$$

and that (in the limit as $x_0 \to \infty$) the error e^{-x_0} is *smaller* than any of these terms...even smaller than E_{492}/x_0^{492}. From now on we will change the lower limit of integration from 0 to $-\infty$, with the understanding that this replacement will introduce a negligible error. Thus, most of the remaining formulas in this section will show not equality, but rather "asymptotic equality", denoted by the symbol $\approx$.

(Perhaps I'm being too glib, because while this error is indeed small, I introduce it an infinite number of times. This is the kind of problem you might want to take to your friendly neighborhood applied mathematician. These folks can usually solve the problem more easily than you can and they often appreciate your bringing neat problems to them.)

Using these ideas, we write

$$\begin{aligned}
4\alpha A_\alpha(x_0) &= \int_0^\infty \frac{x^\alpha}{\cosh^2((x-x_0)/2)} dx \\
&= x_0^\alpha \int_0^\infty \frac{1}{\cosh^2((x-x_0)/2)} dx + \alpha x_0^{\alpha-1} \int_0^\infty \frac{(x-x_0)}{\cosh^2((x-x_0)/2)} dx \\
&\qquad + \tfrac{1}{2}\alpha(\alpha-1) x_0^{\alpha-2} \int_0^\infty \frac{(x-x_0)^2}{\cosh^2((x-x_0)/2)} dx + \cdots \\
&\approx x_0^\alpha \int_{-\infty}^\infty \frac{1}{\cosh^2((x-x_0)/2)} dx + \alpha x_0^{\alpha-1} \int_{-\infty}^\infty \frac{(x-x_0)}{\cosh^2((x-x_0)/2)} dx \\
&\qquad + \tfrac{1}{2}\alpha(\alpha-1) x_0^{\alpha-2} \int_{-\infty}^\infty \frac{(x-x_0)^2}{\cosh^2((x-x_0)/2)} dx + \cdots
\end{aligned}$$

Because the integrals extend from $-\infty$ to $+\infty$, it is easy to change the variable of integration to $x - x_0$, and then symmetry dictates that

$$\int_{-\infty}^\infty \frac{(x-x_0)^n}{\cosh^2((x-x_0)/2)} d(x-x_0) = 0 \qquad \text{for all odd } n.$$

Using the substitution $u = (x-x_0)/2$, we have

$$4\alpha A_\alpha(x_0) \approx x_0^\alpha \left[2\int_{-\infty}^\infty \frac{1}{\cosh^2(u)} du + \frac{4\alpha(\alpha-1)}{x_0^2} \int_{-\infty}^\infty \frac{u^2}{\cosh^2(u)} du + \cdots \right] \tag{6.114}$$

The first integral can be found in Dwight and is

$$\int_{-\infty}^\infty \frac{1}{\cosh^2(u)} du = 2.$$

The second is Gradshteyn 3.527.5, namely

$$\int_{-\infty}^\infty \frac{u^2}{\cosh^2(u)} du = \frac{\pi^2}{6}.$$

Consequently

$$A_\alpha(x_0) \approx \frac{x_0^\alpha}{\alpha}\left[1 + \frac{\pi^2}{6}\frac{\alpha(\alpha-1)}{x_0^2} + \mathcal{O}\left(\frac{1}{x_0^4}\right)\right]. \tag{6.115}$$

This is the desired series in powers of $1/x_0$... and surprisingly, all the odd powers vanish! This result is known as "the Sommerfeld expansion".

Now we apply this formula to equation (6.106) to obtain the function $\mu(T)$:

$$\begin{aligned}
1 &\approx \frac{\mu^{3/2}}{\mathcal{E}_F^{3/2}}\left[1 + \frac{\pi^2}{8}\left(\frac{k_BT}{\mu}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right] \\
\mu^{3/2} &\approx \mathcal{E}_F^{3/2}\left[1 + \frac{\pi^2}{8}\left(\frac{k_BT}{\mu}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right]^{-1} \\
\mu &\approx \mathcal{E}_F\left[1 + \frac{\pi^2}{8}\left(\frac{k_BT}{\mu}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right]^{-2/3} \\
&= \mathcal{E}_F\left[1 - \frac{\pi^2}{12}\left(\frac{k_BT}{\mu}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right]
\end{aligned}$$

This equation tells us a lot: It confirms that at $T = 0$, we have $\mu = \mathcal{E}_F$. It confirms that at T increases from zero, the chemical potential μ decreases. However it doesn't directly give us an expression for $\mu(T)$ because μ appears on both the left and right sides. On the other hand, it does show us that μ deviates from $\mathcal{E}_F$ by an amount proportional to $(k_BT/\mu)^2$, which is small. How small?

$$\begin{aligned}
\left(\frac{k_BT}{\mu}\right)^2 &\approx \frac{(k_BT)^2}{\mathcal{E}_F^2\left[1 - \frac{\pi^2}{12}\left(\frac{k_BT}{\mu}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right]^2} \\
&= \frac{(k_BT)^2}{\mathcal{E}_F^2\left[1 - \frac{\pi^2}{6}\left(\frac{k_BT}{\mu}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right]} \\
&= \left(\frac{k_BT}{\mathcal{E}_F}\right)^2\left[1 + \frac{\pi^2}{6}\left(\frac{k_BT}{\mu}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right]
\end{aligned}$$

In other words, it is small in the sense that

$$\left(\frac{k_BT}{\mu}\right)^2 \approx \left(\frac{k_BT}{\mathcal{E}_F}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4 \tag{6.116}$$

and that

$$\mathcal{O}\left(\frac{k_BT}{\mu}\right)^4 \approx \mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4 \tag{6.117}$$

This gives us the desired formula for μ as a function of T:

$$\mu(T) \approx \mathcal{E}_F\left[1 - \frac{\pi^2}{12}\left(\frac{k_BT}{\mathcal{E}_F}\right)^2 + \mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4\right]. \tag{6.118}$$

This result verifies our previous heuristic arguments: $\mu(T)$ declines as the temperature increases, and it does so slowly (i.e. to second order in $k_BT/\mathcal{E}_F$).

Now apply expansion (6.115) for A_α to equation (6.107) to find the energy $E(T,V,\mu)$:

$$\frac{E}{N} \approx \frac{3}{5}\frac{\mu^{5/2}}{\mathcal{E}_F^{3/2}}\left[1+\frac{5\pi^2}{8}\left(\frac{k_BT}{\mu}\right)^2+\mathcal{O}\left(\frac{k_BT}{\mu}\right)^4\right]. \tag{6.119}$$

Using equations (6.116) and (6.117):

$$\frac{E}{N} \approx \frac{3}{5}\frac{\mu^{5/2}}{\mathcal{E}_F^{3/2}}\left[1+\frac{5\pi^2}{8}\left(\frac{k_BT}{\mathcal{E}_F}\right)^2+\mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4\right]. \tag{6.120}$$

Now it's time to plug $\mu(T,V,N)$ into $E(T,V,\mu)$ to find $E(T,V,N)$: Use equation (6.118) for $\mu(T,V,N)$, and find

$$\begin{aligned}\mu(T)^{5/2} &\approx \mathcal{E}_F^{5/2}\left[1-\frac{\pi^2}{12}\left(\frac{k_BT}{\mathcal{E}_F}\right)^2+\mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4\right]^{5/2}\\ &= \mathcal{E}_F^{5/2}\left[1-\frac{5\pi^2}{24}\left(\frac{k_BT}{\mathcal{E}_F}\right)^2+\mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4\right].\end{aligned} \tag{6.121}$$

Using (6.121) in (6.120) gives

$$\frac{E}{N} \approx \frac{3}{5}\mathcal{E}_F\left[1-\frac{5\pi^2}{24}\left(\frac{k_BT}{\mathcal{E}_F}\right)^2+\mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4\right]\left[1+\frac{5\pi^2}{8}\left(\frac{k_BT}{\mathcal{E}_F}\right)^2+\mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4\right] \tag{6.122}$$

whence

$$E \approx \frac{3}{5}N\mathcal{E}_F\left[1+\frac{5\pi^2}{12}\left(\frac{k_BT}{\mathcal{E}_F}\right)^2+\mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^4\right]. \tag{6.123}$$

Wow! At long last we're able to say

$$C_V = \left.\frac{\partial E}{\partial T}\right)_{N,V} = \left.\frac{\partial E}{\partial(k_BT/\mathcal{E}_F)}\right)_{N,V}\left(\frac{k_B}{\mathcal{E}_F}\right) \tag{6.124}$$

whence

$$C_V \approx \frac{3}{5}Nk_B\left[\frac{5\pi^2}{6}\left(\frac{k_BT}{\mathcal{E}_F}\right)+\mathcal{O}\left(\frac{k_BT}{\mathcal{E}_F}\right)^3\right], \tag{6.125}$$

or, for low temperatures.

$$C_V \approx \frac{\pi^2}{2}Nk_B\left(\frac{k_BT}{\mathcal{E}_F}\right). \tag{6.126}$$

Was all this worth it, just to get a factor of $\pi^2/3$? Perhaps not, but our analysis gives us more than the factor. It gives us a better understanding of the ideal fermion gas and a deeper appreciation for the significance of the thin "action region" near $\mathcal{E} = \mathcal{E}_F$.

6.9 Additional Problems

6.27 **What if there were no interchange rule?**

Suppose that the interchange rule did not apply, so that the basis on page 141 were the correct one for three identical particles. (Alternatively, consider a gas of N non-identical particles.) Find and sketch the heat capacity as a function of temperature.

6.28 **Pressure and energy density**

(This problem was inspired by Reif problem 9.5.)

Any non-relativistic monatomic ideal gas, whether classical or quantal, satisfies

$$p = \frac{2}{3}\frac{E}{V}. \tag{6.127}$$

This remarkable fact can be demonstrated most easily in the canonical ensemble. (Throughout this problem the particle number is fixed so the N dependence of functions is never mentioned explicitly.)

a. Use $E = \frac{3}{2}Nk_BT$ to demonstrate the relation for the classical monatomic ideal gas.

b. From thermodynamics, show that

$$E(T,V) = -\left.\frac{\partial \ln Z}{\partial \beta}\right)_V \quad \text{and} \quad p(T,V) = \frac{1}{\beta}\left.\frac{\partial \ln Z}{\partial V}\right)_T. \tag{6.128}$$

c. Argue that temperature and energy eigenvalues enter into the canonical partition function as quotients:

$$\ln Z(T,V) = \mathcal{F}(\beta\epsilon_1, \beta\epsilon_2, \ldots, \beta\epsilon_M). \tag{6.129}$$

d. Show that in non-relativistic quantum mechanics the free particle energy eigenvalues depend on box volume through

$$\frac{\partial \epsilon_r}{\partial V} = -\frac{2}{3}\frac{\epsilon_r}{V}. \tag{6.130}$$

e. Use the last three items together to prove the pressure-energy density relation.

f. How are the pressure and energy density related for blackbody radiation? At what stage does the above proof break down in this case?

6.29 **Pressure comparison**

(This problem is modified from one in a GRE Physics test.)

Consider three systems of non-interacting identical particles, each with the same T, V, and N. In one system the particles are fermions, in another they are bosons, and in a third they behave classically. Which system has the greatest pressure? Which has the smallest?

6.30 **Challenge**

For many years I suspected that the chemical potential μ of an ideal gas would have to decrease or at least remain constant when the temperature increased (with V and N constant). I tried proving this in a number of ways, and in my attempts I came across several interesting facts (such as the results

of problem 3.36 and part (f.) of problem 6.11) but I was never been able to prove the desired result. That's because the result is false! The chemical potential increases with temperature for ideal fermions in one dimension. Can you show this?

Appendix

Even though we haven't been able to derive everything on this graph, I can't resist showing you the equation of state (that is, pressure as a function of density and temperature), for ideal fermions, ideal classical particles, and ideal bosons.

Chapter 7

Harmonic Lattice Vibrations

7.1 The Problem

7.1 **The harmonic Hamiltonian**

The Hamiltonian for lattice vibrations, in the harmonic approximation, is

$$\mathcal{H} = \tfrac{1}{2}\sum_{i=1}^{3N} m_i \dot{x}_i^2 + \tfrac{1}{2}\sum_{i=1}^{3N}\sum_{j=1}^{3N} x_i A_{ij} x_j. \tag{7.1}$$

Notice that this Hamiltonian allows the possibility that atoms at different lattice sites might have different masses. Accept the fact that any real symmetric matrix S can be diagonalized through an orthogonal transformation, i.e. that for any such S there exists a matrix B whose inverse is its transpose and such that

$$\mathsf{BSB}^{-1} \tag{7.2}$$

is diagonal. Show that the Hamiltonian can be cast into the form

$$\mathcal{H} = \tfrac{1}{2}\sum_{r=1}^{3N}(\dot{q}_r^2 + D_r q_r^2) \tag{7.3}$$

by a linear change of variables. (Clue: As a first step, introduce the change of variable $z_i = \sqrt{m_i}x_i$.)

7.2 Statistical Mechanics of the Problem

7.2 **Thermal energy versus kinetic energy**

According to the Fall 2001 Review Draft of *Ohio's Academic Content Standards for Science* (page 48), eight-grade students should understand that thermal energy "is in the disorderly motion of molecules." Explain how position as well as motion (i.e. potential energy as well as kinetic energy) contribute to thermal energy.

7.3 Normal Modes for a One-dimensional Chain

The matrix A is all zeros except for 2 on the diagonal and -2 on the superdiagonal. But this doesn't really help us solve the problem. The solution comes from physical insight, not mathematical trickery!

Dispersion relation:

$$\omega(k) = 2\sqrt{\frac{K}{m}}|\sin(\tfrac{1}{2}ka)| \tag{7.4}$$

Meaning of term "dispersion relation":

Start with an arbitrary wave packet, break it up into Fourier components.
Each such component moves at a particular speed.
After some time, find how all the components have moved, then sew them back together.
The wave packet will have changed shape (usually broadened... dispersed).

Remember that we haven't done any statistical mechanics in this section, nor even quantum mechanics. This has been classical mechanics!

7.4 Normal Modes in Three Dimensions

If you wanted to study heat flow, or how sound warms up a crystal, or how light warms up a crystal, you'd have to study these time-dependent dispersion effects. This is hard: the classical mechanics is hard (normal modes) and the quantum mechanics is hard. For the purposes of this book we need only examine static effects, and in particular we need only find the heat capacity at low temperatures.

7.5 Low-temperature Heat Capacity

If

$$G(\omega)\,d\omega = \text{ number of normal modes with frequencies from } \omega \text{ to } \omega + d\omega \tag{7.5}$$

then

$$E^{\text{crystal}} = \int_0^\infty G(\omega)e^{\text{SHO}}(\omega)\,d\omega \quad \text{and} \quad C_V^{\text{crystal}} = \int_0^\infty G(\omega)c_V^{\text{SHO}}(\omega)\,d\omega \tag{7.6}$$

and so forth.

Density of modes:

$$\begin{aligned} G(\omega)\,d\omega &= \sum_{\text{branches}} [\text{vol. of shell in } k\text{-space}](\text{density of modes in } k\text{-space}) \\ &= \sum_{\text{branches}} [4\pi(k_b(\omega))^2\,dk_b]\left(\frac{V}{8\pi^3}\right) \end{aligned} \tag{7.7}$$

This formula holds for any isotropic dispersion relation $k_b(\omega)$. For small values of ω the dispersion relation for each branch is linear (with sound speed c_b) so

$$k_b = \frac{\omega}{c_b} \quad \text{and} \quad dk_b = \frac{d\omega}{c_b}, \tag{7.8}$$

whence

$$\begin{aligned} G(\omega)\,d\omega &= \sum_{\text{branches}} \left[4\pi \left(\frac{\omega}{c_b}\right)^2 \frac{d\omega}{c_b} \right] \left(\frac{V}{8\pi^3}\right) \\ &= \frac{V}{2\pi^2} \left(\sum_{b=1}^{3} \frac{1}{c_b^3} \right) \omega^2\,d\omega. \end{aligned} \tag{7.9}$$

If we define the "average sound speed" c_s through the so-called "harmonic cubed average",

$$\frac{1}{c_s^3} \equiv \frac{1}{3} \sum_{b=1}^{3} \frac{1}{c_b^3}, \tag{7.10}$$

then we have the small-ω density of modes

$$G(\omega)\,d\omega = \frac{3V}{2\pi^2} \frac{\omega^2}{c_s^3}\,d\omega. \tag{7.11}$$

At any temperature,

$$C_V^{\text{crystal}} = \int_0^\infty G(\omega) c_V^{\text{SHO}}(\omega)\,d\omega. \tag{7.12}$$

Recall from equation (5.78) that

$$c_V^{\text{SHO}}(\omega) = k_B \left(\frac{\hbar\omega}{k_B T}\right)^2 \frac{e^{-\hbar\omega/k_B T}}{(1-e^{-\hbar\omega/k_B T})^2}, \tag{7.13}$$

and using the small-ω result (7.11), we have the low-temperature result

$$C_V^{\text{crystal}} = \frac{3V}{2\pi^2} \frac{1}{c_s^3} k_B \int_0^\infty \omega^2\,d\omega \left(\frac{\hbar\omega}{k_B T}\right)^2 \frac{e^{-\hbar\omega/k_B T}}{(1-e^{-\hbar\omega/k_B T})^2}. \tag{7.14}$$

For our first step, avoid despair — instead convert to the dimensionless variable

$$x = \frac{\hbar\omega}{k_B T}$$

and find

$$C_V^{\text{crystal}} = \frac{3V}{2\pi^2} \frac{1}{c_s^3} k_B \left(\frac{k_B T}{\hbar}\right)^3 \int_0^\infty \frac{x^4 e^{-x}}{(1-e^{-x})^2}\,dx. \tag{7.15}$$

The integral is rather hard to do, but we don't *need* to do it — the integral is just a number. We have achieved our aim, namely to show that at low temperatures, $C_V \sim T^3$.

However, if you want to chase down the right numbers, after some fiddling you'll find that

$$\int_0^\infty \frac{x^4 e^{-x}}{(1-e^{-x})^2}\,dx = 4\Gamma(4)\zeta(4) = \frac{4}{15}\pi^4.$$

Thus, the low-temperature specific-heat of a solid due to lattice vibration is

$$C_V^{\text{crystal}} = k_B V \frac{2\pi^2}{5}\left(\frac{k_B T}{\hbar c_s}\right)^3. \tag{7.16}$$

7.3 **How far do the atoms vibrate?**
Consider a simplified classical Einstein model in which N atoms, each of mass m, move classically on a simple cubic lattice with nearest neighbor separation of a. Each atom is bound to its lattice site by a spring of spring constant K, and all the values of K are the same. At temperature T, what is the root mean square average distance of each atom from its equilibrium site? (Note: I am asking for an ensemble average, not a time average.)

7.6 More Realistic Models

7.4 **Debye frequency**
In the Debye model, a solid with average sound speed c_s has density of normal-mode frequencies

$$G(\omega) = \begin{cases} \dfrac{3V}{2\pi^2 c_s^3}\,\omega^2 & \text{for } \omega < \omega_D \\ 0 & \text{for } \omega > \omega_D. \end{cases} \tag{7.17}$$

Find a formula for $\omega_D(N, V, c_s)$, and write $G(\omega)$ in terms of ω_D.

7.5 **Debye model energy and heat capacity**
Find $E(T, V, N)$ and $C_V(T, V, N)$ for a harmonic solid in the Debye model, in terms of ω_D and the function

$$D(x) = \frac{3}{x^3}\int_0^x \frac{t^3}{e^t - 1}dt. \tag{7.18}$$

7.7 What is a Phonon?

7.8 Additional Problems

7.6 **Spin waves**
For harmonic lattice vibrations at low frequencies, $\omega = c_s k$. There are analogous excitations of ferromagnets called "spin waves" which, at low frequencies, satisfy $\omega = Ak^2$. Find the temperature dependence of the heat capacity of a ferromagnet at low temperatures. (Do not bother to evaluate constants. . . I am only looking for the functional form of the temperature dependence.)

7.7 **Comparison of models**

(This problem is stolen from a GRE Physics test.)

One feature common to both the Debye and Einstein models for the specific heat of a crystal composed of N identical atoms is that

a. The average energy of each atom is $3k_BT$.

b. The vibrational energy of the crystal is equivalent to the energy of $3N$ independent harmonic oscillators.

c. The crystal is assumed to be continuous for all elastic waves.

d. The speed of longitudinal elastic waves is less than the speed of transverse elastic waves.

e. The upper cutoff frequency of elastic waves is the same.

Chapter 8

Interacting Classical Fluids

8.1 Introduction

The subject of this chapter is also called "real gases", or "dense gases", or "non-ideal gases", or "imperfect gases", or "liquids and dense gases". The many names are a clue that the same problem has been approached by many different scientists from many different points of view, which in turn is a hint that the problem is one of enormous importance. And in this case the hints are correct. We have treated non-interacting classical atoms for long enough in this book. We know that this treatment leads to results in which all substances have the same equation of state (although they can sometimes be distinguished through their heat capacities) and in which there are no phase transitions, no crystals, no life, and very little of interest. It galls me that anyone ever called such a situation "ideal". The problem of interacting atoms is important for industrial and military applications, but it is just as important from the point of view of pure science. We are now going to see what happens when we allow the atoms to interact. Some call the methods and results "messy". I call them fascinating. I would not want to live in the "ideal" world.

To be specific, we will consider the model problem of:

- *Monatomic* atoms with no internal structure, interacting through spherically symmetric pair potentials.
- *Classical mechanics.*

Many undergraduate books state that this is an "important outstanding problem". That is no longer correct. In fact, the problem was solved (except near phase transitions) in the years 1980–85 by Lado, Ashcroft, Foiles, and Rosenfeld (see Talbot, Lebowitz, *et al.* "A comparison of MHNC and MC at high pressures", *J. Chem. Phys.* **85** (1986) 2187–2192, Y. Rosenfeld, "Variational MHNC", *J. Stat. Phys.* **42** (1986) 437–457). The solution is a technique called "modified hyper-netted chain". Unfortunately, I will not have time to tell you what the solution is or even what the words in the name mean. But I can tell you that it is an ingenious

combination of perturbation and variational methods, and I will be telling you about these two techniques in this chapter. (The best references for this chapter are the books by Donald McQuarrie, *Statistical Mechanics*, and by J.P. Hansen and I.R. McDonald, *Theory of Simple Liquids*.)

Problems

8.1 **The Lennard-Jones potential**

The general Lennard-Jones potential is

$$u(r) = -\frac{a}{r^6} + \frac{b}{r^n}. \tag{8.1}$$

a. Given a and b, find $-\epsilon_m$, the minimum energy, and r_0, the distance of separation at that minimum. Write $u(r)$ in terms of the parameters ϵ_m and r_0 rather than the parameters a and b.

b. If σ is the separation at which $u(\sigma) = 0$, find σ in terms of r_0. Why is ϵ_m irrelevant here?

c. Note the simplifications in the above results if $n = 12$. For this case, write $u(r)$ using the parameters ϵ_m and σ.

8.2 **Negative compressibilities**

Look at figure 8.6.1 on page 307 of Reif. Notice that for temperatures below the critical temperature (T_4 in the figure), the van der Waals fluid can have negative values of κ_T. Suppose a fluid with negative compressibility existed, and I had a bottle of it on the table in front of me. What would happen if the volume of the bottle decreased slightly...say if I rapped my fingers on it? (This impossibility is a warning that the van der Waals equation is simply not applicable at temperatures in the vicinity of the critical temperature or below.)

8.3 **Entropy of a real gas**

A real gas obeys the equation of state

$$pV = Nk_BT\left[1 + B\frac{N}{V}\right], \tag{8.2}$$

where B is a positive constant.

a. In one sentence, describe the behavior of this gas in the limit that T and N are constant but $V \to \infty$. (Answers containing mathematical symbols are automatically disqualified.)

b. Using an appropriate Maxwell relation, calculate the difference in entropy between an ideal gas and this real gas when both have the same T, V, and N.

c. At a given T, V, and N, which entropy is larger? Is this consistent with "the entropy inequality"? Can you justify it with a qualitative argument? Is it possible for the constant B to be negative or zero?

8.2 Perturbation Theory

8.2.1 Fluids in the canonical ensemble

The partition function for a fluid is

$$Z(T,V,N) = \frac{1}{h^{3N}N!}\int d\Gamma e^{-\beta H(\Gamma)} \tag{8.3}$$

where

$$H(\Gamma) = \frac{1}{2m}\sum_{i=1}^{N}\mathbf{p}_i^2 + U_N(\mathbf{r}_1,\ldots,\mathbf{r}_N). \tag{8.4}$$

The momentum integrals can be performed precisely as they were for the ideal gas, giving

$$Z(T,V,N) = \frac{1}{\lambda^{3N}(T)N!}\int d\mathbf{r}_1\cdots d\mathbf{r}_N e^{-\beta U_N(\mathbf{r}_1,\ldots,\mathbf{r}_N)}, \tag{8.5}$$

where $\lambda(T)$ is the thermal de Broglie wavelength. It is useful to define the "configurational partition function"

$$Q_N(T,V) = \frac{1}{N!}\int d\mathbf{r}_1\cdots d\mathbf{r}_N e^{-\beta U_N(\mathbf{r}_1,\ldots,\mathbf{r}_N)}. \tag{8.6}$$

For example, in a three-particle system with spherically symmetric pair potential $u(r)$, the configurational partition function is

$$Q_3(T,V) = \frac{1}{6}\int_V d\mathbf{r}_1\int_V d\mathbf{r}_2\int_V d\mathbf{r}_3 e^{-\beta[u(|\mathbf{r}_1-\mathbf{r}_2|)+u(|\mathbf{r}_1-\mathbf{r}_3|)+u(|\mathbf{r}_2-\mathbf{r}_3|)]}. \tag{8.7}$$

Every other time in this book that we have encountered a complicated partition function, like the one above, it has factorized into the product of many similar terms. (This happened even in our treatment of harmonic lattice vibrations, through the use of normal coordinates.) The expression above does not factorize in this neat way. With considerable diligence and patience, it is possible to evaluate $Q_N(T,V)$ for $N = 3$. But you can see that doing it for cases of interest, such as $N = 6.02\times 10^{23}$, is simply out of the question. Instead we attempt to use perturbation theory by expanding about an already solved problem, namely the ideal gas. Such an approximation will be a good one at low densities, where the gas is "nearly ideal". The obvious thing to do is to expand $Q_N(T,V)$ in the small parameter $\rho = N/V$, the number density. But how can we do that? The quantity ρ doesn't appear in these formulas!

8.2.2 The search for a small parameter

We cannot expand $Q_N(T,V)$ in terms of ρ, but we do know that at low densities all fluids behave like an ideal gas, and that for an ideal gas the chemical potential is

$$\mu = k_B T\ln\left(\frac{\rho}{1/\lambda^3(T)}\right), \tag{8.8}$$

so that as $\rho\to 0$, $\mu\to-\infty$ and $e^{\beta\mu}\to 0$. So here is a second small parameter. You might object that $e^{\beta\mu}$ doesn't appear in the partition function any more than ρ does, but it *does* appear in the grand canonical partition function!

8.2.3 Fluids in the grand canonical ensemble

The grand canonical partition function is

$$\begin{aligned}
\Xi(T,V,\mu) &= \sum_{N=0}^{\infty} e^{\beta\mu N} Z(T,V,N) && (8.9)\\
&= \sum_{N=0}^{\infty} \left(\frac{e^{\beta\mu}}{\lambda^3(T)}\right)^N Q_N(T,V) && (8.10)\\
&= \sum_{N=0}^{\infty} z^N Q_N(T,V) && (8.11)\\
&= 1 + Q_1(T,V)z + Q_2(T,V)z^2 + Q_3(T,V)z^3 + \mathcal{O}(z^4), && (8.12)
\end{aligned}$$

where we have defined the *activity*

$$z \equiv \frac{e^{\beta\mu}}{\lambda^3(T)}. \tag{8.13}$$

The activity is the small parameter upon which perturbation theory relies. (For an ideal gas, it is equal to the number density ρ.)

Note that in this approach we still have to calculate Q_N coefficients, but we will probably stop at $N = 4$ or 5 rather than $N = 6.02 \times 10^{23}$. I will write down the first two coefficients:

$$\begin{aligned}
Q_1(T,V) &= \int_V d\mathbf{r}_1 e^{-\beta 0} = V\\
Q_2(T,V) &= \frac{1}{2}\int_V d\mathbf{r}_1 \int_V d\mathbf{r}_2 e^{-\beta u(|\mathbf{r}_1-\mathbf{r}_2|)}. && (8.14)
\end{aligned}$$

Because the interactions are short-ranged, most of the time $u(|\mathbf{r}_1 - \mathbf{r}_2|)$ nearly vanishes, so Q_2 is pretty close to $V^2/2$. It is neither intensive nor extensive.

8.2.4 The Mayer expansion

Now we go from the partition function to an experimentally measurable master function, namely pressure:

$$\begin{aligned}
\frac{p(T,z)}{k_B T} &= \frac{1}{V}\ln\Xi && (8.15)\\
&= \frac{1}{V}\ln[1 + Q_1 z + Q_2 z^2 + Q_3 z^3 + \mathcal{O}(z^4)]. && (8.16)
\end{aligned}$$

Employing

$$\ln(1+\epsilon) = \epsilon - \tfrac{1}{2}\epsilon^2 + \tfrac{1}{3}\epsilon^3 + \mathcal{O}(\epsilon^4) \tag{8.17}$$

we obtain

$$\begin{aligned}
\frac{p(T,z)}{k_B T} &= \frac{1}{V}[Q_1 z + Q_2 z^2 + Q_3 z^3 - \tfrac{1}{2}(Q_1 z + Q_2 z^2)^2 + \tfrac{1}{3}(Q_1 z)^3 + \mathcal{O}(z^4)] && (8.18)\\
&= \frac{1}{V}Q_1 z + \frac{1}{V}(Q_2 - \tfrac{1}{2}Q_1^2)z^2 + \frac{1}{V}(Q_3 - Q_1 Q_2 + \tfrac{1}{3}Q_1^3)z^3 + \mathcal{O}(z^4) && (8.19)\\
&= \sum_{\ell=1}^{\infty} b_\ell z^\ell. && (8.20)
\end{aligned}$$

The last line above defines the "Mayer expansion". The expansion coefficients $b_\ell = b_\ell(T)$ are intensive functions of temperature (in contrast to the coefficients $Q_N(T,V)$).

Using equations (8.14) it is easy to show that $b_1 = 1$ and that

$$\begin{aligned} b_2 &= \frac{1}{2V}\left(\int_V d\mathbf{r}_1 \int_V d\mathbf{r}_2 e^{-\beta u(|\mathbf{r}_1-\mathbf{r}_2|)} - \int_V d\mathbf{r}_1 \int_V d\mathbf{r}_2 1\right) \\ &= \frac{1}{2V}\int_V d\mathbf{r}_1 \int_V d\mathbf{r}_2 (e^{-\beta u(|\mathbf{r}_1-\mathbf{r}_2|)} - 1). \end{aligned} \tag{8.21}$$

In contrast to the situation with equation (8.14), the integrand above is nearly always zero, which makes it much easier to work with. (The same is true for all the the Mayer coefficients.) Indeed, a little more work will show that

$$b_2 = 2\pi \int_0^\infty dr\, r^2 (e^{-u(r)/k_BT} - 1). \tag{8.22}$$

Specialists in the statistical mechanics of fluids spend a lot of time calculating b_ℓ coefficients for a given potential energy of interaction $u(r)$. We will instead emphasize what to do with the b_ℓs once they are calculated.

8.2.5 Expansion for the density

To obtain the number density $\rho = N/V$ from the Mayer expansion, use the thermodynamic connection between ρ and p:

$$\rho = \left.\frac{\partial p}{\partial \mu}\right)_T = \left.\frac{\partial z}{\partial \mu}\right)_T \left.\frac{\partial p}{\partial z}\right)_T = z \left.\frac{\partial(\beta p)}{\partial z}\right)_T. \tag{8.23}$$

Thus

$$\rho(T,z) = \sum_{\ell=1}^{\infty} \ell b_\ell z^\ell. \tag{8.24}$$

8.2.6 Eliminating the activity: the virial expansion

Now we have two expansions, (8.20) and (8.24), in terms of the activity:

$$\beta p(z) = z + b_2 z^2 + b_3 z^3 + \mathcal{O}(z^4) \tag{8.25}$$

$$\rho(z) = z + 2b_2 z^2 + 3b_3 z^3 + \mathcal{O}(z^4). \tag{8.26}$$

(The temperature dependence of p, ρ, and the b_ℓs is not shown explicitly.) What we would really like, however, would be the traditional (and experimentally accessible) equation of state, $p(\rho)$. In other words, we want to eliminate the zs between the two expansions above. To this end we will invert the expansion for $\rho(z)$ to find $z(\rho)$, and plug that expression into the expansion (8.25) for $p(z)$. In the appendix we perform the inversion to find, correct to third order,

$$z(\rho) = \rho - 2b_2\rho^2 + (8b_2^2 - 3b_3)\rho^3 + \mathcal{O}(\rho^4). \tag{8.27}$$

Plugging into (8.25) gives

$$\begin{aligned}
\beta p(\rho) &= z + b_2 z^2 + b_3 z^3 + \mathcal{O}(z^4) && (8.28)\\
&= [\rho - 2b_2\rho^2 + (8b_2^2 - 3b_3)\rho^3] + b_2[\rho - 2b_2\rho^2]^2 + b_3[\rho]^3 + \mathcal{O}(\rho^4) && (8.29)\\
&= [\rho - 2b_2\rho^2 + (8b_2^2 - 3b_3)\rho^3] + b_2[\rho^2 - 4b_2\rho^3] + b_3[\rho]^3 + \mathcal{O}(\rho^4) && (8.30)\\
&= \rho - b_2\rho^2 + (4b_2^2 - 2b_3)\rho^3 + \mathcal{O}(\rho^4). && (8.31)
\end{aligned}$$

This last expression is called the "virial expansion", and it is usually written (with the temperature dependence put back in) as

$$\frac{p(T,\rho)}{k_B T} = \rho + B_2(T)\rho^2 + B_3(T)\rho^3 + \mathcal{O}(\rho^4). \tag{8.32}$$

8.2.7 Appendix: Inverting $\rho(z)$ to find $z(\rho)$

Begin with

$$\rho(z) = z + 2b_2 z^2 + 3b_3 z^3 + \mathcal{O}(z^4), \tag{8.33}$$

then use

$$z = \rho + \mathcal{O}(z^2)$$

$$\begin{aligned}
z^2 &= (\rho + \mathcal{O}(z^2))(\rho + \mathcal{O}(z^2))\\
&= \rho^2 + 2\rho\mathcal{O}(z^2) + \mathcal{O}(z^4)\\
&= \rho^2 + \mathcal{O}(z^3)
\end{aligned}$$

$$\begin{aligned}
z^3 &= (\rho^2 + \mathcal{O}(z^3))(\rho + \mathcal{O}(z^2))\\
&= \rho^3 + \mathcal{O}(z^4)
\end{aligned}$$

$$z = \rho - 2b_2 z^2 + \mathcal{O}(z^3)$$

$$z = \rho - 2b_2\rho^2 + \mathcal{O}(z^3)$$

$$\begin{aligned}
z^2 &= (\rho - 2b_2\rho^2 + \mathcal{O}(z^3))(\rho - 2b_2\rho^2 + \mathcal{O}(z^3))\\
&= \rho^2 - 4b_2\rho^3 + 4b_2^2\rho^4 + \rho\mathcal{O}(z^3)\\
&= \rho^2 - 4b_2\rho^3 + \mathcal{O}(z^4)
\end{aligned}$$

$$z = \rho - 2b_2 z^2 - 3b_3 z^3 + \mathcal{O}(z^4)$$

$$z = \rho - 2b_2[\rho^2 - 4b_2\rho^3 + \mathcal{O}(z^4)] - 3b_3[\rho^3 + \mathcal{O}(z^4)] + \mathcal{O}(z^4),$$

to find

$$z(\rho) = \rho - 2b_2\rho^2 + (8b_2^2 - 3b_3)\rho^3 + \mathcal{O}(\rho^4). \tag{8.34}$$

Problems

8.4 **Virial expansion for the ideal gas**

Find Q_1, Q_2, Q_3, etc. for an ideal gas. Hence show that the virial equation of state for an ideal gas is $pV = Nk_BT$.

8.5 **General properties of expansion coefficients**

a. Show that $Q_1 = V$, whence $b_1 = 1$.

b. Show (informally) that the b_ℓ Mayer coefficients, as well as the $B_n(T)$ virial coefficients, are intensive.

8.3 Variational Methods

8.6 **Bogoliubov's contribution to the Gibbs-Bogoliubov inequality**
Show that the Gibbs-Bogoliubov inequality, which we derived in class for classical systems, also holds for quantal systems.

8.4 Distribution Functions

8.4.1 One-particle distribution functions

What is the mean number of particles in the box of volume d^3r_A about $\mathbf{r}_A$?

The probability that particle 1 is in d^3r_A about $\mathbf{r}_A$ is

$$\frac{d^3r_A \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_A,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{\int d^3r_1 \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}. \tag{8.35}$$

The probability that particle 2 is in d^3r_A about $\mathbf{r}_A$ is

$$\frac{\int d^3r_1 \;\; d^3r_A \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_A,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{\int d^3r_1 \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}. \tag{8.36}$$

And so forth. I could write down N different integrals, but all of them would be equal.

Thus the mean number of particles in d^3r_A about $\mathbf{r}_A$ is

$$\begin{aligned}
&n_1(\mathbf{r}_A)\, d^3r_A \\
&\quad = N \frac{d^3r_A \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_A,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{\int d^3r_1 \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}} && (8.37)\\
&\quad = N \frac{d^3r_A \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_A,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{h^{3N} N! Z(T,V,N)} && (8.38)\\
&\quad = \frac{1}{(N-1)!} \frac{d^3r_A \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \, e^{-\beta U(\mathbf{r}_A,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N)}}{Q(T,V,N)}. && (8.39)
\end{aligned}$$

8.4.2 Two-particle distribution functions

What is the mean number of pairs of particles, such that one member of the pair in a box of volume d^3r_A about $\mathbf{r}_A$ and the other member is in a box of volume d^3r_B about $\mathbf{r}_B$?

The probability that particle 1 is in d^3r_A about $\mathbf{r}_A$ and particle 2 is in d^3r_B about $\mathbf{r}_B$ is

$$\frac{d^3r_A \;\; d^3r_B \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_A,\mathbf{r}_B,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{\int d^3r_1 \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}. \tag{8.40}$$

The probability that particle 2 is in d^3r_A about $\mathbf{r}_A$ and particle 1 is in d^3r_B about $\mathbf{r}_B$ is

$$\frac{d^3r_B \;\; d^3r_A \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_B,\mathbf{r}_A,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{\int d^3r_1 \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}. \tag{8.41}$$

The probability that particle 3 is in d^3r_A about $\mathbf{r}_A$ and particle 1 is in d^3r_B about $\mathbf{r}_B$ is

$$\frac{d^3r_B \int d^3r_2 \;\; d^3r_A \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_B,\mathbf{r}_2,\mathbf{r}_A,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{\int d^3r_1 \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}. \tag{8.42}$$

And so forth. I could write down $N(N-1)$ different integrals, but all of them would be equal.

Thus the mean number of pairs with one particle in d^3r_A about $\mathbf{r}_A$ and the other in d^3r_B about r_B is

$$\begin{aligned} & n_2(\mathbf{r}_A,\mathbf{r}_B)\, d^3r_A d^3r_B \\ &= N(N-1) \frac{d^3r_A \;\; d^3r_B \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_A,\mathbf{r}_B,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}}{\int d^3r_1 \int d^3r_2 \int d^3r_3 \cdots \int d^3r_N \int d^3p_1 \cdots \int d^3p_N \, e^{-\beta H(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3,\ldots,\mathbf{r}_N,\mathbf{p}_1,\ldots,\mathbf{p}_N)}} \qquad (8.43) \\ &= \frac{1}{(N-2)!} \frac{d^3r_A d^3r_B \int d^3r_3 \cdots \int d^3r_N \, e^{-\beta U(\mathbf{r}_A,\mathbf{r}_B,\mathbf{r}_3,\ldots,\mathbf{r}_N)}}{Q(T,V,N)}. \qquad (8.44) \end{aligned}$$

Problems

8.7 **Correlations of nearby particles**

Suppose that (as is usual) at small distances the interatomic potential $u(r)$ is highly repulsive. Argue that at small r,

$$g_2(r) \approx \text{ constant } e^{-u(r)/k_B T}. \tag{8.45}$$

Do not write down a long or elaborate derivation... I'm looking for a simple qualitative argument.

8.8 **Correlations between non-interacting identical quantal particles**

Guess the form of the pair correlation function $g_2(r)$ for ideal (non-interacting) fermions and bosons. Sketch your conjectures, and then compare them to the graphs presented by G. Baym in *Lectures on Quantum Mechanics* (W.A. Benjamin, Inc., Reading, Mass., 1969) pages 428 and 431.

8.9 **Correlation functions and structure factors**

A typical isotropic fluid, at temperatures above the critical temperature, has correlation functions that are complicated at short distances, but that fall off exponentially at long distances. In fact, the long-distance behavior is

$$g_2(r) = 1 + \frac{Ae^{-r/\xi}}{r} \tag{8.46}$$

where ξ, the so-called correlation length, depends on temperature and density. In contrast, at the critical temperature the correlation function falls off much more slowly, as

$$g_2(r) = 1 + \frac{A}{r^{1+\eta}}. \tag{8.47}$$

Find the structure factor

$$S(\mathbf{k}) = \int d^3r\, [g_2(\mathbf{r}) - 1] e^{-i\mathbf{k}\cdot\mathbf{r}} \tag{8.48}$$

associated with each of these correlation functions. Will your results match those of experiments at small values of k or at large values (i.e. at long or short wavelengths)?

8.10 **Long wavelength structure factor**

Show that, for an isotropic fluid, $\dfrac{dS(k)}{dk}$ vanishes at $k = 0$. Here $S(k)$ is the structure factor

$$S(\mathbf{k}) = 1 + \rho \int d^3r\, [g_2(\mathbf{r}) - 1] e^{i\mathbf{k}\cdot\mathbf{r}}. \tag{8.49}$$

8.11 **Correlations in a magnetic system**

In the Ising model for a magnet, described in problem 4.9, the (net) correlation function is defined by

$$G_i \equiv \langle s_0 s_i \rangle - \langle s_0 \rangle^2, \tag{8.50}$$

where the site $j = 0$ is some arbitrary "central spin". Using the results of problem 4.9, show that for a lattice of N sites,

$$\chi_T(T, H) = N \frac{m^2}{k_B T} \sum_i G_i. \tag{8.51}$$

8.5 Correlations and Scattering

8.6 The Hard Sphere Fluid

Consider a fluid in which each atom is modeled by a "hard sphere" of volume v_0. In this system the potential energy vanishes unless two spheres overlap, while if they do overlap it is infinite. The criterion for overlap is that the centers of the two spheres are separated by a distance of $2r_0$ or less, where $v_0 = \frac{4}{3}\pi r_0^3$. This model is certainly oversimplified in that it ignores the attractive part of the interatomic force, but it is an excellent starting point for perturbation theory, and it also has certain features of interest in its own right, as the problems below demonstrate. (The first problem below is required background for all of the others.)

8.12 **Scaling in the hard sphere fluid**

The canonical partition function for a fluid is

$$Z(T,V,N) = \frac{1}{\lambda^{3N}(T)N!}\int d^3r_1 \cdots \int d^3r_N\, e^{-\beta U_N(\mathbf{r}_1,\ldots,\mathbf{r}_N)}. \tag{8.52}$$

For an ideal gas, the integral over configuration space gives just V^N, so it makes sense to write

$$\begin{aligned} Z(T,V,N) &= \left(\frac{V^N/N!}{\lambda^{3N}(T)}\right)\left(\frac{1}{V^N}\int d^3r_1 \cdots \int d^3r_N\, e^{-\beta U_N(\mathbf{r}_1,\ldots,\mathbf{r}_N)}\right) \\ &\equiv Z_{\text{ideal}}(T,V,N)W(T,V,N), \end{aligned} \tag{8.53}$$

where the last line has defined W, the ratio of interacting to non-interacting ("ideal") configurational partition functions.

a. Argue that in the case of hard spheres W is independent of temperature. It does, however, depend on the mechanical parameter v_0, so we will write the ratio as $W(V,N,v_0)$.

b. Argue that $W(V,N,v_0) \leq 1$, where the equality holds only when $v_0 = 0$.

c. Show that the Helmholtz free energy for any fluid satisfies

$$\begin{aligned} F_{\text{HS}}(T,V,N,v_0) &= F_{\text{ideal}}(T,V,N) - k_BT\ln W(V,N,v_0) \\ &\equiv F_{\text{ideal}} + Nf_{\text{ex}}, \end{aligned} \tag{8.54}$$

where we have defined f_{ex}, the excess free energy per atom. Is the free energy of a hard sphere fluid greater than or less than the free energy of an ideal gas at the same T, V, and N?

d. Because f_{ex} is intensive, it cannot depend upon V and N separately, but only through their intensive quotient, the number density $\rho = N/V$. Use a similar argument to show that

$$W(V,N,v_0) = w(\rho,v_0)^N, \tag{8.55}$$

where w is a dimensionless, intensive quantity.

e. Use dimensional analysis to show that the hard sphere w (and hence $f_{\rm ex}$) cannot depend upon ρ and v_0 separately, but only through their product: $w(\rho v_0)$.

This last result is of enormous importance. It implies that there is no need to perform elaborate computer simulations for a variety of densities and a variety of radii. It suffices to either simulate for a variety of densities at one fixed radius or vice versa. It also demonstrates that the quantity of importance in the statistical mechanics of hard spheres is the "scaled density" $\eta \equiv \rho v_0$.

f. Use thermodynamics to show that

$$p_{\rm HS}(T,\rho,v_0) = p_{\rm ideal}(T,\rho) + \rho\left[\eta\frac{\partial f_{\rm ex}(\eta)}{\partial\eta}\right]_{\eta=\rho v_0}. \tag{8.56}$$

8.13 **The hard sphere phase transition**

a. Interpret the scaled density $\eta \equiv \rho v_0$ geometrically. Why is is sometimes called the "packing fraction"?

b. Is it possible to have a fluid with $\eta \geq 1$?

c. Show that if hard spheres are packed into a face-centered cubic lattice (which is probably the densest possible packing spheres, although no one has been able to prove it), $\eta = \pi\sqrt{2}/6 = 0.7405$.

In fact, at scaled densities η well below the limits mentioned above, the system undergoes a phase transition from a fluid to a solid. Computer simulations show that the transition occurs when the scaled density is $\eta_t \approx 0.496$. The conjecture $\eta_t = \frac{1}{2}$ is tempting, but apparently not correct. (On the other hand it is hard to do computer simulations near phase transitions, so the question is not completely closed!)

8.14 **The hard sphere fluid equation of state**

Many long and difficult hours of analytic calculation and computer simulation have gone into studying the hard sphere fluid. In 1964 Ree and Hoover[1] codified much of this work into the following empirical formula which is a good approximation for the hard sphere fluid equation of state:

$$p_{\rm HS}(T,\rho,v_0) = k_BT\rho\left[1+4\eta\frac{1+0.25403\eta+0.27726\eta^2}{1-2.2460\eta+1.3010\eta^2}\right] \quad \text{where} \quad \eta=\rho v_0. \tag{8.57}$$

Only five years later, Carnahan and Starling[2] hit upon a much simpler yet still remarkably accurate equation of state, namely

$$p_{\rm HS}(T,\rho,v_0) = k_BT\rho\frac{1+\eta+\eta^2-\eta^3}{(1-\eta)^3} \quad \text{where} \quad \eta=\rho v_0. \tag{8.58}$$

Both of these formulas apply only to the fluid phase (see the previous problem) and neither one is exact. Answer the following questions for the Carnahan-Starling formula.

[1] F.H. Ree and W.G. Hoover, *J. Chem. Phys.*, **40** (1964) 939–950.
[2] N.F. Carnahan and K.E. Starling, *J. Chem. Phys.*, **51** (1969) 635–636.

a. As the density increases from zero, at what point does the formula become nonsense?

b. On which regions of the (T, ρ) plane does the hard sphere fluid exert a greater pressure than the ideal gas? Can you explain your answer through a simple physical argument?

c. Integrate equation (8.56) to show that for the hard sphere fluid,

$$f_{\text{ex}}(T, \eta) = k_B T \, \eta \frac{4 - 3\eta}{(1 - \eta)^2}. \tag{8.59}$$

Does this formula satisfy the inequality established in part (c.) of problem 8.12?

d. Find the difference between the entropy of the hard sphere fluid and that of an ideal gas. At a given T, V, and N, which entropy is larger? Can you justify this result with a qualitative argument?

8.15 **The hard sphere free energy**

Use the Gibbs-Bogoliubov inequality to show that

$$F_{\text{HS}}(T, V, N) \geq F_{\text{ideal}}(T, V, N). \tag{8.60}$$

Reason that $E_{\text{HS}}(T, V, N) = E_{\text{ideal}}(T, V, N)$, and conclude with a relation between entropies.

8.16 **Virial expansion for hard spheres**

Using the notation of problem 8.12, show that for a hard sphere fluid:

a. $Q_2 = \frac{1}{2} V (V - 8v_0)$.

b. $b_2 = -4v_0$.

c. $B_2(T) = 4v_0$.

d. Show that both the Ree-Hover and the Carnahan-Starling equations of state, introduced in problem 8.14, expand to give the correct first virial coefficient $B_2(T)$.

Chapter 9

Strongly Interacting Systems and Phase Transitions

9.1 Introduction to Magnetic Systems and Models

9.2 Free Energy of the One-Dimensional Ising Model

The N-spin one-dimensional Ising model consists of a horizontal chain of spins, $s_1, s_2, \ldots, s_N$, where $s_i = \pm 1$.

A vertical magnetic field H is applied, and only nearest neighbor spins interact, so the Hamiltonian is

$$\mathcal{H}_N = -J \sum_{i=1}^{N-1} s_i s_{i+1} - mH \sum_{i=1}^{N} s_i. \tag{9.1}$$

For this system the partition function is

$$Z_N = \sum_{\text{states}} e^{-\beta \mathcal{H}_N} = \sum_{s_1=\pm 1} \sum_{s_2=\pm 1} \cdots \sum_{s_N=\pm 1} e^{K \sum_{i=1}^{N-1} s_i s_{i+1} + L \sum_{i=1}^{N} s_i}, \tag{9.2}$$

where

$$K \equiv \frac{J}{k_B T} \quad \text{and} \quad L \equiv \frac{mH}{k_B T}. \tag{9.3}$$

If $J = 0$, (the ideal paramagnet) the partition function factorizes, and the problem is easily solved using the "summing Hamiltonian yields factorizing partition function" theorem. If $H = 0$, the partition function nearly factorizes, and the problem is not too difficult. (See problem 9.3.) But in general, there is no factorization.

We will solve the problem using *induction on the size of the system.* If we add one more spin (spin number $N+1$), then the change in the system's energy depends only upon the state of the new spin and of the previous spin (spin number N). Define $Z_N^{\uparrow}$ as, not the sum over all states, but the sum over all states in which the last (i.e. Nth) spin is up, and define $Z_N^{\downarrow}$ as the sum over all states in which the last spin is down, so that

$$Z_N = Z_N^{\uparrow} + Z_N^{\downarrow}. \tag{9.4}$$

Now, if one more spin is added, the extra term in $e^{-\beta\mathcal{H}}$ results in a factor of

$$e^{Ks_N s_{N+1} + Ls_{N+1}}. \tag{9.5}$$

From this, it is very easy to see that

$$Z_{N+1}^{\uparrow} = Z_N^{\uparrow} e^{K+L} + Z_N^{\downarrow} e^{-K+L} \tag{9.6}$$

$$Z_{N+1}^{\downarrow} = Z_N^{\uparrow} e^{-K-L} + Z_N^{\downarrow} e^{K-L}. \tag{9.7}$$

This is really the end of the physics of this derivation. The rest is mathematics.

So put on your mathematical hats and look at the pair of equations above. What do you see? A matrix equation!

$$\begin{pmatrix} Z_{N+1}^{\uparrow} \\ Z_{N+1}^{\downarrow} \end{pmatrix} = \begin{pmatrix} e^{K+L} & e^{-K+L} \\ e^{-K-L} & e^{K-L} \end{pmatrix} \begin{pmatrix} Z_N^{\uparrow} \\ Z_N^{\downarrow} \end{pmatrix}. \tag{9.8}$$

We introduce the notation

$$\mathbf{w}_{N+1} = \mathsf{T}\mathbf{w}_N \tag{9.9}$$

for the matrix equation. The 2×2 matrix T, which acts to add one more spin to the chain, is called the *transfer matrix.* Of course, the entire chain can be built by applying T repeatedly to an initial chain of one site, i.e. that

$$\mathbf{w}_{N+1} = \mathsf{T}^N \mathbf{w}_1, \tag{9.10}$$

where

$$\mathbf{w}_1 = \begin{pmatrix} e^{L} \\ e^{-L} \end{pmatrix}. \tag{9.11}$$

The fact that we are raising a matrix to a power suggests that we should diagonalize it. The transfer matrix T has eigenvalues λ_A and λ_B (labeled so that $|\lambda_A| > |\lambda_B|$) and corresponding eigenvectors $\mathbf{x}_A$ and $\mathbf{x}_B$. Like any other vector, $\mathbf{w}_1$ can be expanded in terms of the eigenvectors

$$\mathbf{w}_1 = c_A \mathbf{x}_A + c_B \mathbf{x}_B \tag{9.12}$$

and in this form it is very easy to see what happens when $\mathbf{w}_1$ is multiplied by T N times:

$$\begin{aligned}\mathbf{w}_{N+1} = \mathsf{T}^N \mathbf{w}_1 &= c_A \mathsf{T}^N \mathbf{x}_A + c_B \mathsf{T}^N \mathbf{x}_B \\ &= c_A \lambda_A^N \mathbf{x}_A + c_B \lambda_B^N \mathbf{x}_B. \end{aligned} \tag{9.13}$$

So the partition function is

$$Z_{N+1} = Z_{N+1}^{\uparrow} + Z_{N+1}^{\downarrow} = c_A \lambda_A^N (x_A^{\uparrow} + x_A^{\downarrow}) + c_B \lambda_B^N (x_B^{\uparrow} + x_B^{\downarrow}). \tag{9.14}$$

By diagonalizing matrix T (that is, by finding both its eigenvalues and its eigenvectors) we could find every element in the right hand side of the above equation, and hence we could find the partition function Z_N for any N. But of course we are really interested only in the thermodynamic limit $N \to \infty$. Because $|\lambda_A| > |\lambda_B|$, λ_A^N dominates λ_B^N in the thermodynamic limit, and

$$Z_{N+1} \approx c_A \lambda_A^N (x_A^{\uparrow} + x_A^{\downarrow}), \tag{9.15}$$

provided that $c_A(x_A^{\uparrow} + x_A^{\downarrow}) \neq 0$. Now,

$$F_{N+1} = -k_B T \ln Z_{N+1} \approx -k_B T N \ln \lambda_A - k_B T \ln[c_A(x_A^{\uparrow} + x_A^{\downarrow})], \tag{9.16}$$

and this approximation becomes exact in the thermodynamic limit. Thus the free energy per spin is

$$f(K, L) = \lim_{N \to \infty} \frac{F_{N+1}(K, L)}{N+1} = -k_B T \ln \lambda_A. \tag{9.17}$$

So to find the free energy we only need to find the larger eigenvalue of T: we don't need to find the smaller eigenvalue, and we don't need to find the eigenvectors!

It is a simple matter to find the eigenvalues of our transfer matrix T. They are the two roots of

$$\det \begin{pmatrix} e^{K+L} - \lambda & e^{-K+L} \\ e^{-K-L} & e^{K-L} - \lambda \end{pmatrix} = 0 \tag{9.18}$$

$$(\lambda - e^{K+L})(\lambda - e^{K-L}) - e^{-2K} = 0 \tag{9.19}$$

$$\lambda^2 - 2e^K \cosh L \lambda + e^{2K} - e^{-2K} = 0, \tag{9.20}$$

which are

$$\lambda = e^K \left[\cosh L \pm \sqrt{\cosh^2 L - 1 + e^{-4K}} \right]. \tag{9.21}$$

It is clear that both eigenvalues are real, and that the larger one is positive, so

$$\lambda_A = e^K \left[\cosh L + \sqrt{\sinh^2 L + e^{-4K}} \right]. \tag{9.22}$$

Finally, using equation (9.17), we find the free energy per spin

$$f(T, H) = -J - k_B T \ln \left[\cosh \frac{mH}{k_B T} + \sqrt{\sinh^2 \frac{mH}{k_B T} + e^{-4J/k_B T}} \right]. \tag{9.23}$$

Results

Knowing the free energy, we can take derivatives to find any thermodynamic quantity (see problem 9.2). Here I'll sketch and discuss the results obtained through those derivatives.

The heat capacity at constant, vanishing, magnetic field is sketched here:

Do you think an experimentalist needs a magnet to probe magnetic phenomena? This graph shows that a magnet is *not* required: The magnetic effects result in a bump in the heat capacity near $k_BT = J$, even when no magnetic field is applied.

The magnetic susceptibility is sketched here:

We have already seen that for independent spins (paramagnet, $J = 0$) the susceptibility falls like $1/T$ with temperature (the "Curie law"). Interacting spins at high temperature ($k_BT \gg J$) behave approximately the same way. But at low temperatures, the susceptibility for a ferromagnet *exceeds* the susceptibility for a paramagnet, while the susceptibility for a antiferromagnet *undershoots* the susceptibility for a paramagnet. This makes sense: For a paramagnet, the external magnetic field is inducing the spins to align. For a ferromagnet, both the external magnetic field and the tendency of neighboring spins to align are inducing the spins to align. For an antiferromagnet, the external magnetic field is inducing the spins to align, but the tendency of neighboring spins to antialign is opposing that inducement.

Aligning the spins in a paramagnet is like herding cats: the individual spins are independent and don't naturally take to pointing all in the same direction. Aligning the spins in a ferromagnet is like herding cows: the individual spins want to all go in the same direction and don't care which direction it is. Aligning the spins in an antiferromagnet is like herding siblings in a dysfunctional family, where each sibling says "I want to go to the opposite of wherever my brother/sister wants to go."

9.3 The Mean-Field Approximation

Many times, in this book, I have had occasion to make an approximation, but then I argued (sometimes rigorously and sometimes less rigorously) that this approximation would become "exact in the thermodynamic limit." So let me emphasize that the mean-field approximation is *not* exact in the thermodynamic limit. It is usually accurate at high temperatures. It is rarely accurate near phase transitions.

9.4 Correlation Functions in the Ising Model

9.4.1 Motivation and definition

How does a spin at one site influence a spin at another site? This is not a question of thermodynamics, but it's an interesting and useful question in statistical mechanics. The answer is given through correlation functions.

Consider an Ising model with spins $s_i = \pm 1$ on lattice sites i. The figure below shows part of a square lattice, although the discussion holds for any Bravais lattice in any dimension.

Choose a site at the center of the lattice and call it the origin, site 0. Your choice is quite arbitrary, because the lattice is infinite so *any* site is at the center. Now choose another site and call it site i. The product of these two spins is

$$s_0 s_i = \begin{cases} +1 & \text{if the spins point in the same direction} \\ -1 & \text{if the spins point in opposite directions.} \end{cases} \tag{9.24}$$

What is the average of this product over all the configurations in the ensemble? In general, this will be a difficult quantity to find, but in certain special cases we can hold reasonable expectations. For example if the system is ferromagnetic, then at low temperatures we expect that nearest neighbor spins will mostly be pointing in the same direction, whereas the opposite holds if the system is antiferromagnetic. Thus we

expect that, if site i is adjacent to site 0,

$$\langle s_0 s_i \rangle \approx \begin{cases} +1 & \text{for ferromagnetic systems at low temperatures} \\ -1 & \text{for antiferromagnetic systems at low temperatures.} \end{cases} \tag{9.25}$$

Can you think of other expectations? (High temperatures? Next-nearest-neighbors at low temperatures?) Does the average of the product ever exceed 1 in any circumstances? Let me also point out a frequently-held expectation that is *not* correct. One might expect that $\langle s_0 s_i \rangle$ would vanish for sites very far apart. This is true under some circumstances, but not if an external magnetic field is applied, nor at zero field in a low-temperature ferromagnet, because in these situations all the spins in the sample (even those very far apart) tend to point in the same direction.

We have seen that the average of the product $\langle s_0 s_i \rangle$ is not trivial. In contrast the product of the averages $\langle s_0 \rangle \langle s_i \rangle$ is very easy, because all sites are equivalent whence $\langle s_i \rangle = \langle s_0 \rangle$ and

$$\langle s_0 \rangle \langle s_i \rangle = \langle s_0 \rangle^2. \tag{9.26}$$

In fact, $\langle s_0 s_i \rangle$ would be equal to $\langle s_0 \rangle^2$ if there were no correlations between spins... this is essentially the definition of "correlation". This motivates the definition of the *correlation function*

$$G_i(T, H) = \langle s_0 s_i \rangle - \langle s_0 \rangle^2. \tag{9.27}$$

The correlation function is essentially a measure of "peer pressure": How is the spin at site i influenced by the state of the spin at site 0? If there is a large magnetic field, for example, and if sites i and 0 are far apart, then both spins will tend to point up, but this is not because of peer pressure, it is because of "external pressure". (For example, two children in different cities might dress the same way, not because one imitates the other, but because both listen to the same advertisements.) The term $\langle s_0 \rangle^2$ measures external pressure, and we subtract it from $\langle s_0 s_i \rangle$ so that $G_i(T, H)$ will measure only peer pressure and not external pressure.

This definition doesn't say how the influence travels from site 0 to site i: If site 2 is two sites to the right of site 0, for example, then most of the influence will be due to site 0 influencing site 1 and then site 1 influencing site 2. But a little influence will travel up one site, over two sites, then down one site, and still smaller amounts will take even more circuitous routes.

Note that

$$\begin{aligned} \text{if } i = 0 \text{ then} \quad G_i(T, H) &= 1 - \langle s_0 \rangle^2 \\ \text{if } i \text{ is far from 0 then} \quad G_i(T, H) &= 0. \end{aligned} \tag{9.28}$$

I have three comments to make concerning the correlation function. First, realize that the correlation function gives information about the system beyond thermodynamic information. For most of this book we have been using microscopic information to calculate a partition function and then to find macroscopic (thermodynamic) information about the system. We did not use the partition function to ask any question that involved the distance between two spins (or two atoms). The correlation function enables us to probe more deeply into statistical mechanics and ask such important *microscopic* questions.

Second, the correlation function is a measurable quantity. I have produced the definition through conceptual motivations[1] that make it seem impossible to find except by examining individual spins... a conceptual nicety but experimental impossibility. But in fact the correlation function can be found experimentally through neutron scattering. This technique relies upon interference in the de Broglie waves of neutron waves scattered from nearby spins. It is not trivial: indeed two of the inventors of the technique were awarded the Nobel prize in physics for 1994. (It is also worth noting that the work of one of these two was based upon theoretical work done by Bob Weinstock in his Ph.D. thesis.) I will not have enough time to treat this interesting topic, but you should realize that what follows in this section is not mere theoretical fluff... it has been subjected to, and passed, the cold harsh probe of experimental test.

Third, although the definition given here applies only to Ising models on Bravais lattices, analogous definitions can be produced for other magnetic systems, for fluids, and for solids.

9.4.2 Susceptibility from the correlation function

I have emphasized above that the correlation function gives information beyond that given by thermodynamics. But the correlation function contains thermodynamic information as well. In this subsection we will prove that if the correlation function $G_i(T,H)$ is known for all sites in the lattice, then the susceptibility, a purely thermodynamic quantity, can be found through

$$\chi(T,H) = N\frac{m^2}{k_BT}\sum_i G_i(T,H), \tag{9.29}$$

where the sum is over all sites in the lattice (including the origin).

We begin with a few reminders. The microscopic magnetization for a single configuration is

$$\mathcal{M}(s_1,\ldots,s_N) = m\sum_i s_i \tag{9.30}$$

and the macroscopic (thermodynamic) magnetization is its ensemble average

$$M = \langle\mathcal{M}\rangle = mN\langle s_0\rangle. \tag{9.31}$$

The Ising Hamiltonian breaks into two pieces,

$$\begin{aligned}\mathcal{H} &= \mathcal{H}_0(s_1,\ldots,s_N) - mH\sum_i s_i &(9.32)\\ &= \mathcal{H}_0(s_1,\ldots,s_N) - H\mathcal{M}(s_1,\ldots,s_N), &(9.33)\end{aligned}$$

where $\mathcal{H}_0$ is independent of magnetic field. In the nearest-neighbor Ising model the field-independent part of the Hamiltonian is

$$\mathcal{H}_0(s_1,\ldots,s_N) = -J\sum_{<i,j>} s_is_j, \tag{9.34}$$

[1]Namely, by asking "Wouldn't it be nice to know how one spin is influenced by its neighbors?"

but we will not use this formula... the results of this section are true for spin-spin interactions of arbitrary complexity. Finally, the Boltzmann factor is

$$e^{-\beta\mathcal{H}} = e^{-\beta\mathcal{H}_0+\beta H\mathcal{M}}. \tag{9.35}$$

Now we can start the argument by examining the sum of the correlation function over all sites. In what follows, the notation $\sum_{\mathrm{S}}$ means the "sum over states".

$$\begin{aligned}
\sum_i G_i &= \sum_i \langle s_0 s_i\rangle - \langle s_0\rangle^2 && (9.36)\\
&= \sum_i \left[\frac{\sum_{\mathrm{S}} s_0 s_i e^{-\beta\mathcal{H}}}{Z} - \frac{\left(\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}\right)^2}{Z^2}\right] && (9.37)\\
&= \frac{\sum_{\mathrm{S}} s_0(\mathcal{M}/m)e^{-\beta\mathcal{H}}}{Z} - N\frac{\left(\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}\right)^2}{Z^2} && (9.38)\\
&= \frac{Z\sum_{\mathrm{S}} s_0(\mathcal{M}/m)e^{-\beta\mathcal{H}} - N\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}}{Z^2} && (9.39)
\end{aligned}$$

Look at this last equation carefully. What do you see? The quotient rule! Do you remember our slick trick for finding dispersions? The $\mathcal{M}$ appearing in the leftmost summand can be gotten there by taking a derivative of equation (9.35) with respect to H:

$$\frac{\partial e^{-\beta\mathcal{H}}}{\partial H} = \beta\mathcal{M}e^{-\beta\mathcal{H}}. \tag{9.40}$$

Watch carefully:

$$\begin{aligned}
\frac{\partial}{\partial H}\left(\frac{\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}}{Z}\right) &= \frac{Z\sum_{\mathrm{S}} s_0\beta\mathcal{M}e^{-\beta\mathcal{H}} - \sum_{\mathrm{S}}\beta\mathcal{M}e^{-\beta\mathcal{H}}\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}}{Z^2} && (9.41)\\
&= \beta\left[\frac{Z\sum_{\mathrm{S}} s_0\mathcal{M}e^{-\beta\mathcal{H}} - mN\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}}{Z^2}\right] && (9.42)\\
& && (9.43)
\end{aligned}$$

But the quantity in square brackets is nothing more than m times equation (9.39), so

$$\frac{\partial}{\partial H}\left(\frac{\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}}{Z}\right) = \beta m\sum_i G_i. \tag{9.44}$$

But we also know that

$$\frac{\partial}{\partial H}\left(\frac{\sum_{\mathrm{S}} s_0 e^{-\beta\mathcal{H}}}{Z}\right) = \frac{\partial\langle s_0\rangle}{\partial H} = \frac{\partial M/mN}{\partial H} = \frac{1}{mN}\chi \tag{9.45}$$

whence, finally

$$\chi(T,H) = N\frac{m^2}{k_B T}\sum_i G_i(T,H). \tag{9.46}$$

Whew!

Now we can stop running and do our cool-down stretches. Does the above equation, procured at such expense, make sense? The left hand side is extensive. The right hand side has a factor of N and also a

sum over all sites, and usually a sum over all sites is extensive. So at first glance the left hand side scales linearly with N while the right hand side scales with N^2, a sure sign of error! In fact, there is no error. The correlation function G_i vanishes except for sites very close to the origin. ("Distant sites are uncorrelated", see equation (9.28).) Thus the sum over all sites above has a summand that vanishes for most sites, and it is intensive rather than extensive.

9.4.3 All of thermodynamics from the susceptibility

The previous subsection shows us how to find one thermodynamic quantity, $\chi(T, H)$ at some particular values for temperature and field, by knowing the correlation function $G_i(T, H)$ for all sites i, at those same fixed values for temperature and field. In this subsection we will find how to calculate any thermodynamic quantity at all, at any value of T and H at all, by knowing $\chi(T, H)$ for all values of T and H. Thus knowing the correlation $G_i(T, H)$ as a function of all its arguments—i, T, and H—enables us to find any thermodynamic quantity at all.

To find any thermodynamic quantity it suffices to find the free energy $F(T, H)$, so the claim above reduces to saying that we can find $F(T, H)$ given $\chi(T, H)$. Now we can find $\chi(T, H)$ from $F(T, H)$ by simple differentiation, but going backwards requires some very careful integration. We will do so by looking at the difference between quantities for the Ising model and for the ideal paramagnet.

Remember that the thermodynamic state of an Ising magnet is specified by the two variables T and H, as indicated in the figure.

Over much of this diagram, namely the part outside the wavy line, the Ising magnet behaves very much as an ideal paramagnet does, because the typical thermal energy k_BT or the typical magnetic energy $m|H|$ overwhelms the typical spin-spin interaction energy $|J|$. Thus outside of the dashed line (e.g. at the $\times$) the Ising thermodynamic quantities are well-approximated by the paramagnetic thermodynamic quantities

$$\chi_P(T, H) = \left.\frac{\partial M}{\partial H}\right)_T = N\frac{m^2}{k_BT}\mathrm{sech}^2\frac{mH}{k_BT} \tag{9.47}$$

$$M_P(T,H) = -\left.\frac{\partial F}{\partial H}\right)_T = mN\tanh^2\frac{mH}{k_BT} \tag{9.48}$$

$$F_P(T,H) = -k_BTN\ln\left(2\cosh\frac{mH}{k_BT}\right), \tag{9.49}$$

and this approximation becomes better and better as $T\to\infty$ or $H\to\pm\infty$.

The difference between the susceptibility for our Ising model and for the ideal paramagnet is

$$\chi(T,H)-\chi_P(T,H) = \left.\frac{\partial(M-M_P)}{\partial H}\right)_T. \tag{9.50}$$

Integrate this equation along the path indicated in the figure to find

$$\int_{H_0}^{H} dH'\,[\chi(T,H')-\chi_P(T,H')] = \int_{H_0}^{H} dH'\,\left.\frac{\partial(M-M_P)}{\partial H'}\right)_T \tag{9.51}$$

$$= (M-M_P)|_{T,H}-(M-M_P)|_{T,H_0}. \tag{9.52}$$

Then take the limit $H_0\to-\infty$ so that $M(T,H_0)\to M_P(T,H_0)$ and

$$M(T,H) = M_P(T,H)+\int_{-\infty}^{H} dH'\,[\chi(T,H')-\chi_P(T,H')], \tag{9.53}$$

where, remember, $M_P(T,H)$ and $\chi_P(T,H)$ are known functions.

We don't have to stop here. We can consider the difference

$$M(T,H)-M_P(T,H) = -\left.\frac{\partial(F-F_P)}{\partial H}\right)_T \tag{9.54}$$

and then perform exactly the same manipulations that we performed above to find

$$F(T,H) = F_P(T,H)-\int_{-\infty}^{H} dH'\,[M(T,H')-M_P(T,H')]. \tag{9.55}$$

Then we can combine equations (9.55) and (9.53) to obtain

$$F(T,H) = F_P(T,H)-\int_{-\infty}^{H} dH'\int_{-\infty}^{H'} dH''\,[\chi(T,H'')-\chi_P(T,H'')]. \tag{9.56}$$

Using the known results for $F_P(T,H)$ and $\chi_P(T,H)$, as well as the correlation-susceptibility relation (9.46), this becomes

$$F(T,H) = -k_BTN\ln\left(2\cosh\frac{mH}{k_BT}\right)-N\frac{m^2}{k_BT}\int_{-\infty}^{H} dH'\int_{-\infty}^{H'} dH''\left(\sum_i G_i(T,H'')-\operatorname{sech}^2\frac{mH''}{k_BT}\right). \tag{9.57}$$

Let me stress how remarkable this result is. You would think that to find the thermodynamic energy $E(T,H)$ of the Ising model you would need to know the spin-spin coupling constant J. But you don't: You can find $E(T,H)$ without knowing J, indeed without knowing the exact form of the spin-spin interaction Hamiltonian at all, if you do know the purely geometrical information contained within the correlation function.[2] This is one of the most amazing and useful facts in all of physics.

[2]This does not mean that the thermodynamic energy is independent of the spin-spin Hamiltonian, because the Hamiltonian influences the correlation function.

9.4.4 Parallel results for fluids

I cannot resist telling you how the results of this section change when applied to fluid rather than magnetic systems. In that case the correlation function $g_2(\mathbf{r}; T, V, N)$ depends upon the position $\mathbf{r}$ rather than the site index i. Just as the sum of G_i over all sites is related to the susceptibility χ, so the integral of $g_2(\mathbf{r})$ over all space is related to the compressibility κ_T. And just as the susceptibility can be integrated twice (carefully, through comparison to the ideal paramagnet) to give the magnetic free energy $F(T, H)$, so the compressibility can be integrated twice (carefully, through comparison to the ideal gas) to give the fluid free energy $F(T, V, N)$. And just as the magnetic thermodynamic quantities can be found from the correlation function without knowing the details of the spin-spin interaction Hamiltonian, so the fluid thermodynamic quantities can be found from the correlation function without knowing the details of the particle interaction forces. A question for you: How is it possible to get all this information knowing only the correlation function in space rather than the correlation function in phase space? In other words, why don't you need information about the velocities?

I should put in some references to experimental tests and to say who thought this up. Are these available in Joel's review article?

9.5 Computer Simulation

9.5.1 Basic strategy

If we're going to use a computer to solve problems in statistical mechanics, there are three basic strategies that we could take. I illustrate them here by showing how they would be applied to the Ising model.

1. Exhaustive enumeration. Just do it! List all the states (microstates, configurations), their associated probabilities, and average over them! Admittedly, there are a lot of states, but computers are fast, right? Let's see how many states there are. Consider a pretty small problem: a three dimensional Ising model on an $8 \times 8 \times 8$ cube. This system has 512 spins that can be oriented either "up" or "down", so there are $2^{512} \approx 10^{154}$ configurations. Suppose that our computer could generate and examine a million configurations per second... this is about as fast as is currently possible. Then an exhaustive enumeration would require 10^{148} seconds to do the job. By contrast, the universe is about 10^{18} seconds old. So for one computer to do the problem it would take 10^{130} times the age of the universe. Now this is impractical, but maybe we could do it by arranging computers in parallel. Well, there are about 10^{80} protons in the universe, and if every one were to turn into a spiffy Unix workstation, it would still require 10^{50} times the age of the universe to complete the list. And this is for a small problem! Sorry, computers aren't fast, and they never will be fast enough to solve the complete enumeration problem. It is not for nothing that this technique is called *exhaustive* enumeration.

2. Random sampling. Instead of trying to list the entire pool of possible configurations, we will dip into it at random and sample the configurations. This can be implemented by scanning through the lattice and

orienting spins up with probability $\frac{1}{2}$ and down with probability $\frac{1}{2}$. The problem in this strategy is that all configurations are equally likely to be sampled, but because there are many more high energy configurations than low energy configurations, it is very unlikely to sample a low energy configuration (the "poker paradox" again). But in the Boltzmann distribution $e^{-\text{energy}/k_BT}$ the low energy configurations are in fact *more* likely to occur. So you could sample for a long time before encountering even one configuration with non-negligible probability. This strategy is considerably better than the first one, but it would still take about the age of the universe to implement.

3. Importance sampling. This strategy is to sample the pool of possible configurations not completely at random, but in such a way that the most likely configurations are most likely to be sampled. An enormously successful algorithm to perform importance sampling was developed by Nick Metropolis and his coworkers Arianna Rosenbluth, Marshall Rosenbluth, Augusta Teller, and Edward Teller (of hydrogen bomb fame) in 1953. ("Equation of state calculations by fast computing machines", *J. Chem. Phys.* **21** 1087–1092.) It is called "the Metropolis algorithm", or "the M(RT)^2 algorithm", or simply "the Monte Carlo algorithm". The next section treats this algorithm in detail.

An analogy to political polling is worth making here. Exhaustive enumeration corresponds to an election, random sampling corresponds to a random poll, and importance sampling corresponds to a poll which selects respondents according to the likelihood that they will vote.

9.5.2 The Metropolis algorithm

This algorithm builds a chain of configurations, each one modified (usually only slightly) from the one before. For example, if we number the configurations (say, for the $8 \times 8 \times 8$ Ising model, from 1 to 2^{512}) such a chain of configurations might be

$$C_{171} \to C_{49} \to C_{1294} \to C_{1294} \to C_{171} \to C_{190} \to \cdots.$$

Note that it is possible for two successive configurations in the chain to be identical.

To build any such chain, we need some transition probability rule $W(C_a \to C_b)$ giving the probability that configuration C_b will follow configuration C_a. And to be useful for importance sampling, the rule will have to build the chain in such a way that the probability of C_n appearing in the chain is $e^{-\beta E_n}/Z$.

So we need to produce a result of the form: if "rule" then "Boltzmann probability distribution". It is, however, very difficult to come up with such results. Instead we go the other way to produce a pool of plausible transition rules... plausible in that they are not obviously inconsistent with a Boltzmann probability distribution.

Consider a long chain of N_{configs} configurations, in which the probability of a configuration appearing is given by the Boltzmann distribution. The chain must be "at equilibrium" in that

$$\text{number of transitions}(C_a \to C_b) = \text{number of transitions}(C_b \to C_a). \tag{9.58}$$

But the number of C_as in the chain is $N_{\text{configs}} e^{-\beta E_a}/Z$, and similarly for the number of C_bs, so the equation above is

$$N_{\text{configs}} \frac{e^{-\beta E_a}}{Z} W(C_a \to C_b) = N_{\text{configs}} \frac{e^{-\beta E_b}}{Z} W(C_b \to C_a), \tag{9.59}$$

whence

$$\frac{W(C_a \to C_b)}{W(C_b \to C_a)} = e^{-\beta(E_b - E_a)}. \tag{9.60}$$

This condition, called "detailed balance", defines our pool of plausible transition probability rules. It is possible that a rule could satisfy detailed balance and still not sample according to the Boltzmann distribution, but any rule that does *not* satisfy detailed balance certainly *cannot* sample according to the Boltzmann distribution. In practice, all the rules that satisfy detailed balance seem to work.

Two transitions probability rules that do satisfy detailed balance, and the two rules most commonly used in practice, are

$$W(C_a \to C_b) = [\text{normalization constant}] \left\{ \frac{e^{-\beta \Delta E}}{1 + e^{-\beta \Delta E}} \right\} \tag{9.61}$$

$$W(C_a \to C_b) = [\text{normalization constant}] \left\{ \begin{array}{cc} 1 & \text{if } \Delta E \leq 0 \\ e^{-\beta \Delta E} & \text{if } \Delta E > 0 \end{array} \right\} \tag{9.62}$$

where we have taken

$$\Delta E = E_b - E_a \tag{9.63}$$

and where the normalization constant is fixed so that

$$\sum_{C_b} W(C_a \to C_b) = 1. \tag{9.64}$$

If we had to calculate the normalization constant, then we would be back to performing an exhaustive enumeration, so we must find a way to work with unnormalized transition probabilities. To this end we define $w_{a \to b}$ as the factor for the transition probability aside from the normalization constant, i.e. the part in curly brackets in equations (9.61) and (9.62). Note that it is a positive number less than or equal to one:

$$0 < \{ \quad \} \equiv w_{a \to b} \leq 1. \tag{9.65}$$

Notice that for any rule satisfying detailed balance the transition probability must *increase* as the change in energy *decreases*, suggesting a simple physical interpretation: The chain of configurations is like a walker stepping from configuration to configuration on a "configuration landscape". If a step would decrease the walker's energy, he is likely to take it, whereas if it would increase his energy, he is likely to reject it. Thus the walker tends to go downhill on the configuration landscape, but it is not impossible for him to go uphill. This seems like a recipe for a constantly decreasing energy, but it is not, because more uphill steps than downhill steps are available to be taken.[3]

[3]This paragraph constitutes the most densely packed observation in this book. It looks backward to the poker paradox, to the definition of temperature, and to the "cash incentives" interpretation of the canonical ensemble. It looks forward to equilibration and to optimization by Monte Carlo simulated annealing. Why not read it again?

From these ingredients, Metropolis brews his algorithm:

Generate initial `configuration` [at random or otherwise]
Gather data concerning `configuration` [e.g. find $M := n_\uparrow - n_\downarrow$; $M_{\text{sum}} := M$]
`DO Iconfig = 2, Nconfigs`
    Generate `candidate configuration` [e.g. select a spin to flip]
    Compute $w_{a\to b}$ for transition to candidate
    With probability $w_{a\to b}$, make the transition
    Gather data concerning `configuration` [e.g. find $M := n_\uparrow - n_\downarrow$; $M_{\text{sum}} := M_{\text{sum}} + M$]
`END DO`
Summarize and print data [e.g. $M_{\text{ave}} := M_{\text{sum}}/$`Nconfigs`]

I'll make three comments concerning this algorithm. First of all, note that the step "With probability $w_{a\to b}$, make the transition" implies that sometimes the transition is not made, in which case two configurations adjacent in the chain will be identical. It is a common misconception that in this case the repeated configuration should be counted only once, but that's not correct: you must execute the "Gather data concerning `configuration`" step whether the previous candidate was accepted or rejected.

Secondly, I wish to detail how the step

With probability $w_{a\to b}$, make the transition

is implemented. It is done by expanding the step into the two substeps

Produce a random number z $[0 \le z < 1]$
`IF` $z < w_{a\to b}$ `THEN` switch to candidate
    [i.e. `configuration := candidate configuration`]

Finally, I need to point out that this algorithm does not precisely implement either of the transition probability rules (9.61) or (9.62). For example, if the step "Generate `candidate configuration`" is done by selecting a single spin to flip, then the algorithm will never step from one configuration to a configuration three spin-flips away, regardless of the value of ΔE. Indeed, under such circumstances (and if there are N spins in the system), the transition probability rule is

$$W(C_a \to C_b) = \begin{cases} \frac{1}{N} w_{a\to b} & \text{if } C_a \text{ and } C_b \text{ differ by a single spin flip} \\ 0 & \text{if } C_a \text{ and } C_b \text{ differ by more than a single spin flip} \\ 1 - \sum_{C_b \neq C_a} W(C_a \to C_b) & \text{if } C_a = C_b. \end{cases} \tag{9.66}$$

It is easy to see that this transition probability rule satisfies detailed balance.

9.5.3 Implementing the Metropolis algorithm

It is not atypical to run a Monte Carlo program for about 10,000 Monte Carlo steps per site. As such, the program might run for hours or even days. This is considerably less than the age of the universe, but

probably far longer than you are used to running programs. The following tips are useful for speeding up or otherwise improving programs implementing the Metropolis algorithm.

1. Use scaled quantities. The parameters J, m, H, and T do not enter in any possible combination, but only through two independent products. These are usually taken to be

$$\tilde{T} = \frac{k_B T}{J} \quad \text{and} \quad \tilde{H} = \frac{mH}{J}. \tag{9.67}$$

Thus the Boltzmann exponent for a given configuration is

$$-\frac{E}{k_B T} = \frac{1}{\tilde{T}} \left(\sum_{\langle i,j \rangle} s_i s_j + \tilde{H} \sum_i s_i \right), \tag{9.68}$$

where s_i is $+1$ if the spin at site i is up, -1 if it is down, and where $\langle i, j \rangle$ denotes a nearest neighbor pair.

2. Don't find total energies. To calculate $w_{a \to b}$ you must first know ΔE, and the obvious way to find ΔE is to find E_b and E_a (through equation (9.68)) and subtract. This way is obvious but terribly inefficient. Because the change in configuration is small (usually a single spin flip) the change in energy can be found from purely local considerations without finding the total energy of the entire system being simulated. Similarly, if you are finding the average (scaled) magnetization

$$M = n_\uparrow - n_\downarrow = \sum_i s_i \tag{9.69}$$

(as suggested by the square brackets in the algorithm on page 202) you don't need to scan the entire lattice to find it. Instead, just realize that it changes by ± 2 with each spin flip.

3. Precompute Boltzmann factors. It is computationally expensive to find evaluate a exponential, yet we must know the value of $e^{-\beta \Delta E}$. However, usually there are only a few possible values of ΔE. (For example in the square lattice Ising model with nearest neighbor interactions and a field, a single spin flip gives rise to one of only ten possible values of the energy change.) It saves considerable computational time (and often makes the program clearer) to precalculate the corresponding values of $w_{a \to b}$ just once at the beginning of the program, and to store those values in an array for ready reference when they are needed.

4. Average "on the fly". The algorithm on page 202 finds the average (scaled) magnetization by summing the magnetization of each configuration in the chain and then dividing by the number of configurations. Because the chain is so long this raises the very real possibility of overflow in the value of M_{sum}. It is often better to keep a running tally of the average by tracking the "average so far" through

$$M_{\text{ave}} := M_{\text{ave}}(\texttt{Iconfig} - 1)/\texttt{Iconfig} + M/\texttt{Iconfig} \tag{9.70}$$

or (identical mathematically, but preferable for numerical work)

$$M_{\text{ave}} := M_{\text{ave}} + (M - M_{\text{ave}})/\texttt{Iconfig}. \tag{9.71}$$

5. Finite size effects. Use periodic or skew-periodic boundary conditions.

6. Lattice data structures. Suppose we wish to simulate a two-dimensional Ising model on a 4×3 square grid. An obvious data structure to hold the configuration in the computer's memory is an integer-valued *two-dimensional array* declared through the code

```
INTEGER, PARAMETER ::  Nx = 4, Ny = 3
INTEGER ::  Spin (1:Nx, 1:Ny)
```

(I use the notation of Fortran 90. If you are familiar with some other computer language, the intent should nevertheless be clear.) If the spin at site (3,2) is up, then `Spin`(3,2) $= +1$, and if that spin is down, then `Spin`(3,2) $= -1$. The lattice sites are labeled as shown in this figure:

$$\begin{array}{cccc} \overset{\circ}{(1,3)} & \overset{\circ}{(2,3)} & \overset{\circ}{(3,3)} & \overset{\circ}{(4,3)} \\ \\ \overset{\circ}{(1,2)} & \overset{\circ}{(2,2)} & \overset{\circ}{(3,2)} & \overset{\circ}{(4,2)} \\ \\ \overset{\circ}{(1,1)} & \overset{\circ}{(2,1)} & \overset{\circ}{(3,1)} & \overset{\circ}{(4,1)} \end{array}$$

Although this representation is obvious, it suffers from a number of defects. First, it is difficult to generalize to other lattices, such as the triangular lattice in two dimensions or the face-centered cubic lattice in three dimensions. Second, finding the nearest neighbors of boundary sites using periodic or skew-periodic boundary conditions is complicated. And finally because the array is two-dimensional, any reference to an array element involves a multiplication,[4] which slows down the finding of data considerably.

All of these defects are absent in the *folded array* representation of the lattice sites. In this representation the sites are stored as an integer-valued one-dimensional array declared through

```
INTEGER, PARAMETER ::  Nx = 4, Ny = 3, NSites = Nx*Ny
INTEGER ::  Spin (0:NSites-1)
```

Now the lattice sites are labeled as:

[4]In Fortran, the datum `Site(i,j)` is stored at memory location number `i + j*(Nx-1)`, whence the data are stored in the sequence (1,1), (2,1), (3,1), (4,1), (1,2), (2,2),.... In other computer languages the formula for finding the memory location is different, but in all languages it involves an integer multiply.

It is not hard to show that, with skew-periodic boundary conditions, the four neighbors of site number `l` (`l` for location) are

$$(\texttt{l} + 1) \bmod \texttt{NSites},\ (\texttt{l} - 1) \bmod \texttt{NSites},\ (\texttt{l} + \texttt{Nx}) \bmod \texttt{NSites},\ (\texttt{l} - \texttt{Nx}) \bmod \texttt{NSites}.$$

Unfortunately, the Fortran `Mod` function was inanely chosen to differ from the mathematical mod function, so this arithmetic must be implemented through

```
Mod(l + 1, NSites)
Mod(l + (NSites - 1), NSites)
Mod(l + Nx, NSites)
Mod(l + (NSites - Nx), NSites)
```

7. Random number generators. It is hard to find an easy source of high-quality random numbers: this is a problem for physics, also for government and industry.

D.E. Knuth, *Seminumerical Algorithms* (Addison–Wesley, Reading, Massachusetts, 1981) chapter 3.

W.H. Press, S.A. Teukolsky, W.T. Vettering, and B.P. Flannery, *Numerical Recipes* chapter 7. (For a confession and a prize announcement, see also W.H. Press and S.A. Teukolsky, "Portable random number generators", *Computers in Physics* **6** (1992) 522–524.)

T.-W. Chiu and T.-S. Guu, "A shift-register sequence random number generator," *Computer Physics Communications* **47** (1987) 129–137. (See particularly figures 1 and 2.)

A.M. Ferrenberg, D.P. Landau, and Y.J. Wong, "Monte Carlo simulations: Hidden errors from 'good' random number generators," *Phys. Rev. Lett.* **69** (1992) 3382.

S.K. Park and K.W. Miller, "Random number generators: Good ones are hard to find," *Communications of the ACM* **31** (1988) 1192–1201.

8. Initial configurations and equilibration. The Metropolis algorithm does not specify how to come up with the first configuration in the chain. Indeed, selecting this configuration is something of an art. If you are

simulating at high temperatures, it is usually appropriate to begin with a configuration chosen at random, so that about half the spins will be up and half down. But if you are simulating at low temperatures (or at high fields) it might be better to start at the configuration with all spins up. Other possibilities are also possible. But however you select the initial configuration, it is highly unlikely that the one you pick will be "typical" of the configurations for the temperature and magnetic field at which you are simulating.

9. Low temperatures. At low temperatures, most candidate transitions are rejected. BLK algorithm. (A.B. Bortz, J.L. Lebowitz, and M.H. Kalos, "A new algorithm for Monte Carlo simulation of Ising spin systems," *J. Comput. Phys.* **17** (1975) 10–18.)

10. Critical temperatures. "Critical slowing down," response is cluster flipping. (U. Wolff, "Collective Monte Carlo updating for spin systems," *Phys. Rev. Lett.* **62** (1989) 361–364. See also Jian-Sheng Wang and R.H. Swendsen, "Cluster Monte Carlo algorithms," *Physica* A **167** (1990) 565–579.)

11. First-order transitions.

12. Molecular dynamics.

9.5.4 The Wolff algorithm

In 1989 Ulli Wolff proposed an algorithm for Monte Carlo simulation that is particularly effective near critical points. (U. Wolff, *Phys. Rev. Lett.*, **62** (1989) 361–364.) The next page presents the Wolff algorithm as applied to the zero-field ferromagnetic nearest-neighbor Ising model,

$$\mathcal{H} = -J \sum s_i s_j \quad \text{with} \quad s_i = \pm 1, \tag{9.72}$$

on a square lattice or a simple cubic lattice.

```
Generate initial configuration
Gather data concerning this configuration

DO Iconfig = 2, Nconfigs

    Select site j at random

    Flip spin at j
    Put j into cluster
    NFlippedThisGeneration = 1

    DO
    IF (NFlippedThisGeneration = 0) EXIT
        NFlippedPreviousGeneration = NFlippedThisGeneration
        NFlippedThisGeneration = 0

        FOR each previous generation member i of the cluster DO
            FOR each neighbor j of i DO
                IF (spin at j ≠ spin at i) THEN
                    with probability P = 1 − exp(−2J/k_B T)
                        Flip spin at j
                        Put j into cluster
                        NFlippedThisGeneration = NFlippedThisGeneration + 1
                END IF
            END DO
        END DO

    END DO

    Gather data concerning this configuration
    Empty the cluster

END DO

Summarize and print out results
```

Template for example

configuration a								configuration b							
+	−	−	+	+	−	+	−	+	−	−	+	+	−	+	−
−	−	+	+	+	+	−	+	−	−	+	−	−	−	−	+
−	−	−	+	+	−	−	−	−	−	−	−	−	−	−	−
+	−	+	−	+	−	−	−	+	−	+	−	−	−	−	−
−	+	+	+	+	−	−	+	−	+	+	−	−	−	−	+
−	−	−	−	+	−	−	−	−	−	−	−	+	−	−	−
+	+	−	+	+	−	+	+	+	+	−	+	+	−	+	+
+	−	−	+	+	−	−	+	+	−	−	+	+	−	−	+
$w_{a\to b} = \frac{8}{N} P_J^7 (1-P^J)^5$								$w_{b\to a} = \frac{8}{N} P_J^7 (1-P^J)^{11}$							

Demonstration of detailed balance. In these formulas,

$$P_J = 1 - e^{-2J/k_BT}. \tag{9.73}$$

The integer 8 is the number of sites in the cluster, 7 is the number of internal bonds in the skeleton, 5 is the number of external bonds leading to a + spin, and 11 is the number of external bonds leading to a − spin. The energy difference between configurations a and b depends only upon these last two integers. It is

$$\Delta E = +2J(5) - 2J(11) = -2J(11-5). \tag{9.74}$$

Thus

$$\frac{w_{a\to b}}{w_{b\to a}} = (1-P^J)^{5-11} = e^{-2J(5-11)/k_BT} = e^{-\Delta E/k_BT} \tag{9.75}$$

and detailed balance is insured.

9.6 Additional Problems

9.1 **Denaturation of DNA**

(This problem is modified from one posed by M.E. Fisher. It deals with a research question born in the 1960s that is still of interest today: Douglas Poland and Harold A. Scheraga, "Occurrence of a phase transition in nucleic acid models," *J. Chem. Phys.* **45** (1966) 1464–1469; M.E. Fisher, "Effect of excluded volume on phase transitions in biopolymers," *J. Chem. Phys.* **45** (1966) 1469–1473; Yariv Kafri, David Mukamel, and Luca Peliti, "Why is the DNA denaturation transition first order?" *Phys. Rev. Lett.* **85** (2000) 4988–4991.)

The biological polymer DNA consists of two polymeric strands which, in their usual state, are twisted together to form a rather rigid double helix. However as the temperature is raised regions of the molecule may "melt", breaking the weak bonds that hold the two strands together. In this case the two strands separate and coil randomly in the solution. If the entire molecule melts then the DNA is said to be "denatured". A simple model to aid in understanding this phenomenon is the following.

Successive segments of the molecule are regarded as being in either the *helical state* or in the *melted state*. A segment in the helical state has a lower energy than one in the melted state ($-\epsilon$ *vs.* 0) but a segment in the melted state has q different possible orientations (due to coiling) whereas a segment in the helical state has only one. However, the first or last segment of a run of melted segments has a smaller number of configurations, rq, where r can be as small as 10^{-3}. (A melted segment that is both the first and last segment of a run is doubly restricted and has only r^2q configurations.) Sample DNA configurations and their associated "Boltzmann factors" are shown below.

a. Introduce a variable h_i for the ith segment such that

$$h_i = \begin{cases} 1 & \text{if the } i\text{th segment is helical} \\ 0 & \text{if the } i\text{th segment is melted.} \end{cases} \tag{9.76}$$

Show that the canonical partition function for a molecule of N segments is

$$Z_N(T,\epsilon,r) = \sum_{h_1=0}^{1}\sum_{h_2=0}^{1}\cdots\sum_{h_N=0}^{1}\prod_{i=1}^{N} e^{\beta\epsilon h_i} q^{(1-h_i)} r^{(h_i-h_{i-1})^2}. \tag{9.77}$$

b. If $F(T,\epsilon,r)$ is the Helmholtz free energy, show that the mean fraction of helical segments is

$$\theta(T) = -\frac{1}{N}\frac{\partial F}{\partial \epsilon} \tag{9.78}$$

while the mean number of junctions between melted and helical runs is

$$J(T) = -\frac{r}{k_BT}\frac{\partial F}{\partial r}. \tag{9.79}$$

c. By considering partial partition functions, Z_N^h and Z_N^m, for a molecule of N segments in which the last segment is in a helical or melted state, respectively, construct recursion relations from which the total partition function, Z_N, can be found.

d. Find the eigenvalues of the corresponding 2×2 matrix and conclude that the free energy per segment of a very long DNA molecule is

$$f(T, \epsilon, r) = -k_B T \ln \tfrac{1}{2} \left[1 + w + \sqrt{(1-w)^2 + 4wr^2} \right] - \epsilon, \tag{9.80}$$

where $w = qe^{-\beta\epsilon}$. Does this result take on the correct value at zero temperature?

e. In the limit that r vanishes show that the molecule undergoes a sharp melting transition at temperature

$$T_m = \frac{\epsilon}{k_B \ln q}. \tag{9.81}$$

Sketch the energy, entropy, and helical fraction $\theta(T)$ as a function of temperature and discuss the character of the transition.

f. If r is small but nonzero, sketch the behavior you expect for $\theta(T)$.

g. (Optional.) Show that when r is small but nonzero the sharp transition is "smeared out" over a temperature range of approximate width

$$\Delta T \approx \frac{4\epsilon r}{k_B \ln^2 q}. \tag{9.82}$$

h. (Optional.) Show that the number of junctions $J(T)$ defined in (b.) varies as Ar^α both at $T = T_m$ and at low temperatures. Find the values of A and α in each case.

9.2 **Thermodynamics of the one-dimensional Ising model**

The Helmholtz free energy of the one-dimensional, nearest-neighbor Ising magnet is

$$F(T, H) = -N \left\{ J + k_B T \ln \left[\cosh L + \sqrt{\sinh^2 L + e^{-4K}} \right] \right\}, \tag{9.83}$$

where

$$L = \frac{mH}{k_B T} \quad \text{and} \quad K = \frac{J}{k_B T}. \tag{9.84}$$

Use this result to show that

$$E(T, H = 0) = -JN \tanh K, \tag{9.85}$$

$$C_H(T, H = 0) = k_B N \frac{K^2}{\cosh^2 K}, \tag{9.86}$$

$$M(T, H) = mN \frac{\sinh L}{\sqrt{\sinh^2 L + e^{-4K}}}, \tag{9.87}$$

$$\chi_T(T, H = 0) = N \frac{m^2}{k_B T} e^{2K}. \tag{9.88}$$

9.3 **Zero-field one-dimensional Ising model**

In class we found the free energy of a one-dimensional, nearest-neighbor Ising magnet with arbitrary magnetic field using the transfer matrix method. This problem finds the free energy in a much simpler way, which unfortunately works only in zero magnetic field.

a. Show that the partition function can be written as

$$Z_N = \sum_{s_1=\pm1}\sum_{s_2=\pm1} e^{Ks_1s_2} \sum_{s_3=\pm1} e^{Ks_2s_3} \cdots \sum_{s_{N-1}=\pm1} e^{Ks_{N-2}s_{N-1}} \sum_{s_N=\pm1} e^{Ks_{N-1}s_N}. \tag{9.89}$$

b. Show that the last sum is

$$\sum_{s_N=\pm1} e^{Ks_{N-1}s_N} = 2\cosh K, \tag{9.90}$$

regardless of the value of s_{N-1}.

c. Conclude that

$$Z_N = 2^N \cosh^{N-1} K \tag{9.91}$$

and that the zero-field Helmholtz free energy per site is

$$f(T) = -k_BT \ln\left(2\cosh\frac{J}{k_BT}\right). \tag{9.92}$$

d. Find the zero-field heat capacity, and show that it gives the expected behavior at very low temperatures.

9.4 **Magnetization near the critical point: mean field approximation**

Analyze the Ising model in the mean field approximation to show that the zero-field magnetization near the critical temperature is

$$M(T) = \begin{cases} mNa\left(\dfrac{T_c - T}{T_c}\right)^{1/2} & \text{for } T < T_c \\ 0 & \text{for } T > T_c. \end{cases} \tag{9.93}$$

Be sure to specify the value of the numerical constant a. (Hint: $\tanh x = x - \frac{1}{3}x^3 + \frac{2}{15}x^5 + \cdots$.) (Remark: Experiment shows that the magnetization is not precisely of the character predicted by mean field theory: While it does approach zero like $(T_c - T)^\beta$, the exponent β is not 1/2 — for two-dimensional systems $\beta = 1/8$, while for three-dimensional systems $\beta \approx 0.326$.)

9.5 **Correlation functions for paramagnets**

In a paramagnet (see problem 2.9) the spins are uncorrelated, i.e. the correlation function is

$$G_i(T,H) = 0 \quad \text{for} \quad i \neq 0. \tag{9.94}$$

Use this fact to verify the correlation sum rule

$$\chi_T(T,H) = N\frac{m^2}{k_BT}\sum_i G_i, \tag{9.95}$$

derived in problem 8.11.

9.6 **Lattice gas—Ising magnet**

In class we showed that for a lattice gas

$$\Xi(T,\mu) = \sum_{n_1=0}^{1}\sum_{n_2=0}^{1}\cdots\sum_{n_N=0}^{1}\left(\frac{v_0}{\lambda^3(T)}\right)^N e^{\beta\mu\sum_i n_i + \beta\epsilon\sum_{\text{nn}} n_in_j}, \tag{9.96}$$

whereas for an Ising magnet

$$Z(T,H) = \sum_{s_1=\pm 1} \sum_{s_2=\pm 1} \cdots \sum_{s_{\mathcal{N}}=\pm 1} e^{\beta m H \sum_i s_i + \beta J \sum_{\rm nn} s_i s_j}, \tag{9.97}$$

and we concluded that the lattice gas and Ising magnet were more-or-less equivalent, with chemical potential and magnetic field playing sort of analogous roles. These conclusions are indeed correct, but this problem renders them precise.

a. Show that an up spin is associated with an occupied site, and a down spin with a vacant site, through

$$n_i = \tfrac{1}{2}(s_i + 1). \tag{9.98}$$

b. Write out the sum for $\Xi(T,\mu)$ in terms of the variables s_i, and show that the correspondence

$$\epsilon = 4J \tag{9.99}$$

$$\mu = 2mH - k_B T \ln(v_0/\lambda^3(T)) - 4qJ \tag{9.100}$$

(where q is the number of nearest neighbors for each site on the lattice) leads to

$$p(T,\mu)v_0 = mH - \tfrac{3}{2}qJ - f(T,H), \tag{9.101}$$

where $p(T,\mu)$ is the pressure of the gas and $f(T,H)$ is the free energy per spin of the magnet.

c. Interpret equation (9.100) using the phrase "escaping tendency". (Is this a sensible question?)

d. (Hard.) Take derivatives of equation (9.101) to show that

$$\rho(T,\mu)v_0 = \tfrac{1}{2}[1 + M(T,H)/m\mathcal{N}], \tag{9.102}$$

$$4\rho^2\kappa_T(T,\mu)v_0 = \chi(T,H)/m^2\mathcal{N}, \tag{9.103}$$

$$S_{\rm gas} - k_B N[\ln(v_0/\lambda^3(T)) + \tfrac{3}{2}] = S_{\rm mag}, \tag{9.104}$$

$$C_V - \tfrac{3}{2}k_B N = C_M. \tag{9.105}$$

9.7 **Estimates concerning exhaustive enumeration**

Consider an Ising model with 6.02×10^{23} sites. How many distinct configurations can this system take on? Estimate the mass of paper and ink required to write out a single character. What mass would be required to write out this number of configurations in decimal notation? Compare your estimate to the mass of the earth.

9.8 **Fluctuations in Monte Carlo simulation**

When the mean energy of a system is determined through Monte Carlo simulation, the result will necessarily have an uncertainty due to the statistical character of the simulation. (The same is true of any other measured quantity.) So if mean energy is plotted as a function of temperature, the data points will not fall on a sharp line but instead within a broad band. Using a fluctuation-susceptibility relation, show that the width of the band depends on its slope. Will steep bands be wide or narrow?

9.9 **The folded array representation**

Implement the folded array representation of a triangular lattice subject to skew-periodic boundary conditions. Clue: The piece of lattice under study will be a parallelogram.

9.10 **Monte Carlo project**

Perform a Monte Carlo simulation on the Ising model. The precise form and direction of the simulation is up to you: here are some suggestions that you are free to use, combine, extend, or ignore.

a. Draw movies showing the spin configuration of a two-dimensional Ising model as the simulation proceeds. Discuss the qualitative character of the model at high temperatures, low temperatures, and near the critical temperature (which is known, from Onsager's exact solution, to be $k_B T_c/J = -2/\ln(\sqrt{2}-1)$).

b. Simulate the one-dimensional Ising model and compare your results to the analytic solution obtained in class. Consider using the BKL algorithm[5] at low temperatures.

c. Simulate the DNA denaturation model of problem 9.1, and compare your results to the analytic results of that problem. You will need to modify the Metropolis algorithm.

d. Simulate a two- or three-dimensional Ising ferromagnet at zero field, using an initial configuration of all spins up. Monitor the magnetization for a number of temperatures ranging from $k_B T \approx 5J$ down to $k_B T = 0$. Record both the equilibrium magnetization and the number of steps required to reach equilibrium, and plot these as a function of temperature. Can you locate the critical temperature?

e. Monitor both the mean energy (or magnetization) and the fluctuations in energy (or magnetization). Use a fluctuation-susceptibility relation to find the heat capacity (or susceptibility). In particular, show how a simulation at zero magnetic field can still give information about the magnetic susceptibility.

f. Verify the relation discussed in problem 9.8.

g. Implement the Wolff algorithm[6] for improved simulation near the critical point.

h. Use finite size scaling[7] to obtain accurate results near the critical point.

i. Compare the BKL and Wolff algorithms at low temperature.

j. Find the correlation function

$$G_i = \langle s_0 s_i \rangle - \langle s_0 \rangle^2 \tag{9.106}$$

for a few values of T and H. How does it behave as the critical point is approached?

k. (Very ambitious.) Verify the correlation-susceptibility relation (9.46).

[5] A.B. Bortz, J.L. Lebowitz, and M.H. Kalos, *J. Comput. Phys.*, **17** (1975) 10–18.

[6] U. Wolff, *Phys. Rev. Lett.* **62** (1989) 361–364. J.-S. Wang and R.H. Swendsen, *Physica* A **167** (1990) 565–579.

[7] H. Gould and J. Tobochnik, *An Introduction to Computer Simulation Methods: Applications to Physical Systems*, part I (Addison-Wesley, Reading, Mass., 1988) sections 12.4 and 16.5

Appendix A

Series and Integrals

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \cdots \quad \text{for } |x| < 1 \quad \text{(the "geometric series")} \tag{A.1}$$

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \cdots \tag{A.2}$$

$$\ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \cdots \quad \text{for } |x| < 1 \tag{A.3}$$

$$\int_{-\infty}^{+\infty} e^{-ax^2+bx}\,dx = \sqrt{\frac{\pi}{a}}e^{b^2/4a} \quad \text{for } a > 0 \quad \text{(the "Gaussian integral")} \tag{A.4}$$

Appendix B

Evaluating the Gaussian Integral

The integral

$$\int_{-\infty}^{+\infty} e^{-x^2}\,dx, \tag{B.1}$$

called the Gaussian integral, does not fall to any of the methods of attack that you learned in elementary calculus. But it can be evaluated quite simply using the following trick.

Define the value of the integral to be A. Then

$$A^2 = \int_{-\infty}^{+\infty} e^{-x^2}\,dx \int_{-\infty}^{+\infty} e^{-y^2}\,dy = \int_{-\infty}^{+\infty}\int_{-\infty}^{+\infty} dx\,dy\,e^{-(x^2+y^2)}. \tag{B.2}$$

At the last step we have written A^2 as a two-variable integral over the entire plane. This seems perverse, because most of the times we work hard to reduce two-dimensional integrals to one-dimensional integrals, whereas here we are going in reverse. But look at the integrand again. When regarded as an integral on the plane, it is clear that we can regard x^2+y^2 as just r^2, and this suggests we should convert the integral from Cartesian (x, y) to polar (r, θ) coordinates:

$$A^2 = \int_0^{\infty} dr \int_0^{2\pi} r\,d\theta\,e^{-r^2} = 2\pi \int_0^{\infty} re^{-r^2}\,dr. \tag{B.3}$$

The last integral immediately suggests the substitution $u = r^2$, giving

$$A^2 = \pi \int_0^{\infty} e^{-u}\,du = -\pi\,e^{-u}\big|_0^{\infty} = \pi. \tag{B.4}$$

We conclude that

$$\int_{-\infty}^{+\infty} e^{-x^2}\,dx = \sqrt{\pi}. \tag{B.5}$$

B.1 (I) **Problem: Another integral**

Show that

$$\int_0^{\infty} \frac{e^{-x}}{\sqrt{x}}\,dx = \sqrt{\pi}. \tag{B.6}$$

(Clue: Use the substitution $y = \sqrt{x}$.)

Appendix C

Clinic on the Gamma Function

The gamma function $\Gamma(s)$ is defined, for $s > 0$, by

$$\Gamma(s) = \int_0^\infty x^{s-1} e^{-x}\, dx. \tag{C.1}$$

Upon seeing any integral, your first thought is to evaluate it. Stay calm... first make sure that the integral exists. A quick check shows that the integral above converges when $s > 0$.

There is no simple formula for the gamma function for arbitrary s. But for $s = 1$,

$$\Gamma(1) = \int_0^\infty e^{-x}\, dx = -e^{-x}\Big|_0^\infty = 1. \tag{C.2}$$

For $s > 1$ we may integrate by parts,

$$\int_0^\infty x^{s-1} e^{-x}\, dx = -x^{s-1} e^{-x}\Big|_0^\infty + (s-1)\int_0^\infty x^{s-2} e^{-x}\, dx, \tag{C.3}$$

giving

$$\Gamma(s) = (s-1)\Gamma(s-1) \quad \text{for } s > 1. \tag{C.4}$$

Apply equation (C.4) repeatedly for n a positive integer,

$$\Gamma(n) = (n-1)\Gamma(n-1) = (n-1)(n-2)\Gamma(n-2) = (n-1)(n-2)\cdots 2\cdot 1\cdot \Gamma(1) = (n-1)!, \tag{C.5}$$

to find a relation between the gamma function and the factorial function. Thus the gamma function generalizes the factorial function to non-integer values, and can be used to define the factorial function through

$$x! = \Gamma(x+1) \quad \text{for any } x > -1. \tag{C.6}$$

In particular,

$$0! = \Gamma(1) = 1. \tag{C.7}$$

(It is a deep and non-obvious result that the gamma function is in fact the simplest generalization of the factorial function.)

The gamma function can be simplified for half-integral arguments. For example

$$\Gamma(\tfrac{1}{2}) = \int_0^\infty x^{-1/2} e^{-x}\, dx = \int_0^\infty y^{-1} e^{-y^2} (2y\, dy) = \int_{-\infty}^\infty e^{-y^2}\, dy = \sqrt{\pi} \tag{C.8}$$

where we used the substitution $y = \sqrt{x}$. Thus

$$\Gamma(\tfrac{3}{2}) = \tfrac{1}{2}\Gamma(\tfrac{1}{2}) = \frac{\sqrt{\pi}}{2} = (\tfrac{1}{2})!, \tag{C.9}$$

$$\Gamma(\tfrac{5}{2}) = \tfrac{3}{2}\Gamma(\tfrac{3}{2}) = \tfrac{3}{4}\sqrt{\pi}, \tag{C.10}$$

and so forth.

Appendix D

Volume of a Sphere in d Dimensions

I will call the volume of a d-dimensional sphere, as a function of radius, $V_d(r)$. You know, of course, that

$$V_2(r) = \pi r^2 \tag{D.1}$$

(two-dimensional volume is commonly called "area") and that

$$V_3(r) = \tfrac{4}{3}\pi r^3. \tag{D.2}$$

But what is the formula for arbitrary d? There are a number of ways to find it. I will use induction on dimensionality d. That is, I will use the formula for $d = 2$ to find the formula for $d = 3$, the formula for $d = 3$ to find the formula for $d = 4$, and in general use the formula for d to find the formula for $d + 1$. This is not the most rigorous formal method to derive the formula, but it is very appealing and has much to recommend it.

To illustrate the process, I will begin with a well-known and easily visualized stage, namely deriving $V_3(r)$ from $V_2(r)$. Think of a 3-dimensional sphere (of radius r) as a stack of pancakes of various radii, but each with infinitesimal thickness dz. The pancake on the very bottom of the stack ($z = -r$) has zero radius. The one above it is slightly broader. They get broader and broader until we get to the middle of the stack ($z = 0$), where the pancake has radius r. The pancakes stacked still higher become smaller and smaller, until they vanish again at the top of the stack ($z = +r$). Because the equation for the sphere is

$$x^2 + y^2 + z^2 = r^2, \tag{D.3}$$

the radius of the pancake at height z_0 is

$$\sqrt{r^2 - z_0^2}. \tag{D.4}$$

This whole process shows that

$$V_3(r) = \int_{-r}^{+r} dz\, V_2(\sqrt{r^2 - z^2}). \tag{D.5}$$

It is easy to check this integral against the known result for $V_3(r)$:

$$\begin{aligned} V_3(r) &= \int_{-r}^{+r} dz\, \pi(r^2 - z^2) && \text{(D.6)} \\ &= \pi \left[r^2 z - \tfrac{1}{3} z^3 \right]_{-r}^{+r} && \text{(D.7)} \\ &= \pi \left[2r^3 - \tfrac{2}{3} r^3 \right] && \text{(D.8)} \\ &= \tfrac{4}{3} \pi r^3. && \text{(D.9)} \end{aligned}$$

So we haven't gone wrong yet.

Now, how to derive $V_4(r)$ from $V_3(r)$? This requires a more vivid imagination. Last time we started with a two-dimensional disk of radius r_0 in (x, y) space and thickened it a bit into the third dimension (z) to form a pancake of three-dimensional volume $dz\, V_2(r_0)$. Stacking an infinite number of such pancakes in the z direction, from $z = -r$ to $z = +r$, gave us a three-dimensional sphere. Now we begin with a three-dimensional sphere of radius r_0 in (w, x, y) space and thicken it a bit into the fourth dimension (z) to form a thin four-dimensional pancake of four-dimensional volume $dz\, V_3(r_0)$. Stacking an infinite number of such pancakes in the z direction, from $z = -r$ to $z = +r$, gives a four-dimensional sphere. Because the equation for the four-sphere is

$$w^2 + x^2 + y^2 + z^2 = r^2, \tag{D.10}$$

the radius of the three-dimensional sphere at height z_0 is

$$\sqrt{r^2 - z_0^2}, \tag{D.11}$$

and the volume of the four-sphere is

$$V_4(r) = \int_{-r}^{+r} dz\, V_3(\sqrt{r^2 - z^2}). \tag{D.12}$$

In general, the volume of a $(d+1)$-sphere is

$$V_{d+1}(r) = \int_{-r}^{+r} dz\, V_d(\sqrt{r^2 - z^2}). \tag{D.13}$$

If we guess that the formula for $V_d(r)$ takes the form

$$V_d(r) = C_d r^d \tag{D.14}$$

(which is certainly true for two and three dimensions, and which is reasonable from dimensional analysis), then

$$\begin{aligned} V_{d+1}(r) &= \int_{-r}^{+r} dz\, C_d (r^2 - z^2)^{d/2} && \text{(D.15)} \\ &= \int_{-1}^{+1} r\, du\, C_d (r^2 - r^2 u^2)^{d/2} && \text{(D.16)} \\ &= r^{d+1} C_d \int_{-1}^{+1} du\, (1 - u^2)^{d/2}. && \text{(D.17)} \end{aligned}$$

This proves our guess and gives us a recursive formula for C_d:

$$C_{d+1} = C_d \int_{-1}^{+1} du\,(1-u^2)^{d/2}. \tag{D.18}$$

The problem below shows how to build this recursive chain up from $C_2 = \pi$ to

$$C_d = \frac{\pi^{d/2}}{\Gamma(\frac{d}{2}+1)} = \frac{\pi^{d/2}}{(d/2)!}. \tag{D.19}$$

Thus the volume of a d-dimensional sphere of radius r is

$$V_d(r) = \frac{\pi^{d/2}}{(d/2)!} r^d. \tag{D.20}$$

D.1 (I) **Problem: Volume of a d-dimensional sphere**

Before attempting this problem, you should read the material concerning beta functions in an applied mathematics textbook, such as George Arfken's *Mathematical Methods for Physicists* or Mary Boas's *Mathematical Methods in the Physical Sciences.* (Or in the Digital Library of Mathematical Functions.)

a. Show that

$$\int_{-1}^{+1} (1-u^2)^{d/2}\,du = B(\tfrac{1}{2}, \tfrac{d}{2}+1). \tag{D.21}$$

b. Use

$$B(p,q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)} \tag{D.22}$$

and $C_2 = \pi$ to conclude that

$$C_d = \frac{\pi^{d/2}}{\Gamma(\frac{d}{2}+1)}. \tag{D.23}$$

D.2 (I) **Problem: Volume of a d-dimensional ellipse**

Show that the volume of the d-dimensional ellipse described by the equation

$$\left(\frac{x_1}{a_1}\right)^2 + \left(\frac{x_2}{a_2}\right)^2 + \left(\frac{x_3}{a_3}\right)^2 + \cdots + \left(\frac{x_d}{a_d}\right)^2 = 1 \tag{D.24}$$

is

$$V_d(r) = \frac{\pi^{d/2}}{(d/2)!} a_1 a_2 a_3 \cdots a_d. \tag{D.25}$$

Appendix E

Stirling's Approximation

The Stirling formula is an approximation for $n!$ that is good at large values of n.

$$n! = 1 \cdot 2 \cdot 3 \cdots (n-1) \cdot n \tag{E.1}$$

$$\ln(n!) = \underbrace{\ln 1}_{0} + \ln 2 + \ln 3 + \cdots + \ln(n-1) + \ln(n) \tag{E.2}$$

Note that the function $\ln x$ is nearly flat for large values of x. For example, $\ln 10^{23}$ is about equal to 23.

From the figure

$$\ln(6!) = \text{area under the staircase} > \int_1^6 \ln x \, dx, \tag{E.3}$$

and in general

$$\ln(n!) > \int_1^n \ln x \, dx = \Big[x \ln x - x \Big]_1^n = n \ln n - n + 1. \tag{E.4}$$

For large values of n, where the $\ln n$ function is nearly flat, the two expressions above become quite close. Also, the 1 becomes negligible. We conclude that

$$\ln(n!) \approx n \ln n - n \quad \text{for } n \gg 1. \tag{E.5}$$

This is Stirling's formula. For corrections to the formula, see M. Boas, *Mathematical Methods in the Physical Sciences*, sections 9-10 and 9-11.

You know that

$$A^n \tag{E.6}$$

increases rapidly with n for positive A, but

$$n! \approx \left(\frac{n}{e}\right)^n \tag{E.7}$$

increases a bit more rapidly still.

E.1 **Problem: An upper bound for the factorial function**

Stirling's approximation gives a rigorous lower bound for $n!$.

a. Use the general ideas presented in the derivation of that lower bound to show that

$$\int_1^n \ln(x+1)\,dx > \ln n!. \tag{E.8}$$

b. Conclude that

$$(n+1)\ln(n+1) - n + 1 - 2\ln 2 > \ln n! > n \ln n - n + 1. \tag{E.9}$$

Appendix F

The Euler-MacLaurin Formula and Asymptotic Series

You know that a sum can be approximated by an integral. How accurate is that approximation? The *Euler-MacLaurin formula* gives the corrections.

$$\begin{aligned}\sum_{k=0}^{n-1} f(k) \quad &\approx \quad \int_0^n f(x)\,dx \\ &-\frac{1}{2}[f(n)-f(0)]+\frac{1}{12}[f'(n)-f'(0)]-\frac{1}{720}[f'''(n)-f'''(0)]+\frac{1}{30\,240}[f^{(\mathrm{v})}(n)-f^{(\mathrm{v})}(0)]-\frac{1}{1\,209\,600}[f^{(\mathrm{vii})}(n)-\end{aligned}$$

This series is asymptotic. If the series is truncated at any point, it can give a highly accurate approximation. But the series may be either convergent or divergent, so adding additional terms to the truncated series might give rise to a poorer approximation. The Stirling approximation is a truncation of an asymptotic series.

References

C.M. Bender and S.A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers*, (McGraw-Hill, New York, 1978).

M. Abramowitz and I. Stegum, *Handbook of Mathematical Functions*, (National Bureau of Standards, Washington, D.C., 1964).

Appendix G

Ramblings on the Riemann Zeta Function

The function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad \text{for} \quad s > 1,$$

called the Riemann zeta function, has applications to number theory, statistical mechanics, and quantal chaos.

History

From Simmons:

> No great mind of the past has exerted a deeper influence on the mathematics of the twentieth century than Bernhard Riemann (1826–1866), the son of a poor country minister in northern Germany. He studied the works of Euler and Legendre while he was still in secondary school, and it is said that he mastered Legendre's treatise on the theory of numbers in less than a week. But he was shy and modest, with little awareness of his own extraordinary abilities, so at the age of nineteen he went to the University of Göttingen with the aim of pleasing his father by studying theology and becoming a minister himself.

Instead he became a mathematician. He made contributions to the theory of complex variables ("Cauchy-Riemann equations", "Riemann sheet", "Riemann sphere"), to Fourier series, to analysis ("Riemann integral"), to hypergeometric functions, to convergence of infinite sums ("Riemann rearrangement theorem"), to geometry ("Riemannian curved space"), and to classical mechanics, the theory of fields and other areas of physics. In his thirties he became interested in the prime number theorem.

The first few prime numbers are 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43,.... It is clear that the primes are distributed among all the positive integers in a rather irregular way; for as we move out, they seem to occur less and less frequently, and yet there are many adjoining pairs separated by a single even number ["twin primes"]. ... Many attempts have been made to find simple formulas for the nth prime and for the exact number of primes among the first n positive integers. All such efforts have failed, and real progress was achieved only when mathematicians started instead to look for information about the average distribution of the primes among the positive integers. It is customary to denote by $\pi(x)$ the number of primes less than or equal to a positive number x. Thus $\pi(1) = 0$, $\pi(2) = 1$, $\pi(3) = 2$, $\pi(\pi) = 2$, $\pi(4) = 2$, and so on. In his early youth Gauss studied $\pi(x)$ empirically, with the aim of finding a simple function that seems to approximate it with a small relative error for large x. On the basis of his observations he conjectured (perhaps at the age of fourteen or fifteen) that $x/\log x$ is a good approximating function, in the sense that

$$\lim_{x\to\infty} \frac{\pi(x)}{x/\log x} = 1.$$

This statement is the famous *prime number theorem*; and as far as anyone knows, Gauss was never able to support his guess with even a fragment of proof.

Chebyshev, unaware of Gauss's conjecture, was the first mathematician to establish any firm conclusions about this question. In 1848 and 1850 he proved that

$$0.9213\ldots < \frac{\pi(x)}{x/\log x} < 1.1055\ldots$$

for all sufficiently large x, and also that if the limit exists, then its value must be 1. (The numbers in the inequality are $A = \log 2^{1/2}3^{1/3}5^{1/5}30^{-1/30}$ on the left, and $\frac{6}{5}A$ on the right.) ...

In 1859 Riemann published his only work on the theory of numbers, a brief but exceedingly profound paper of less than 10 pages devoted to the prime number theorem. ... His starting point was a remarkable identity discovered by Euler over a century earlier: if s is a real number greater than 1, then

$$\sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_p \frac{1}{1-(1/p^s)},$$

where the expression on the right denotes the product of the numbers $(1-p^{-s})^{-1}$ for all primes p. To understand how this identity arises, we note that $1/(1-x) = 1 + x + x^2 + \cdots$ for $|x| < 1$, so for each p we have

$$\frac{1}{1-(1/p^s)} = 1 + \frac{1}{p^s} + \frac{1}{p^{2s}} + \cdots.$$

On multiplying these series for all primes p and recalling that each integer $n > 1$ is uniquely expressible as a product of powers of different primes, we see that

$$\begin{aligned} \prod_p \frac{1}{1-(1/p^s)} &= \prod_p \left(1 + \frac{1}{p^s} + \frac{1}{p^{2s}} + \cdots\right) \\ &= 1 + \frac{1}{2^s} + \frac{1}{3^s} + \cdots + \frac{1}{n^s} + \cdots \end{aligned}$$

$$= \sum_{n=1}^{\infty} \frac{1}{n^s},$$

which is the desired identity. The sum of this series is evidently a function of the real variable $s > 1$, and the identity establishes a connection between the behavior of this function and properties of the primes. Euler himself exploited this connection in several ways, but Riemann perceived that access to the deeper features of the distribution of primes can only be gained by allowing s to be a complex variable. ... In his paper he proved several important properties of this function, and in a sovereign way simply stated a number of others without proof. During the century since his death, many of the finest mathematicians in the world have exerted their strongest efforts and created rich new branches of analysis in attempts to prove these statements. The first success was achieved in 1893 by J. Hadamard, and with one exception every statement has since been settled in the sense Riemann expected. (Hadamard's work led him to his 1896 proof of the prime number theorem.) The one exception is the famous Riemann hypothesis: that all the zeros of $\zeta(s)$ in the strip $0 \leq \Re e\{s\} \leq 1$ lie on the central line $\Re e\{s\} = \frac{1}{2}$. It stands today as the most important unsolved problem in mathematics, and is probably the most difficult problem that the mind of man has ever conceived. In a fragmentary note found among his posthumous papers, Riemann wrote that these theorems "follow from an expression for the function $\zeta(s)$ which I have not yet simplified enough to publish". ...

At the age of thirty-nine Riemann died of tuberculosis in Italy, on the last of several trips he undertook in order to avoid the cold, wet climate of northern Germany.

General character of the zeta function

It is easy to see that $\zeta(s)$ diverges at $s = 1$ and that it decreases monotonically to 1 as s increases from 1. As the historical note suggests, its behavior for complex values of s is much more elaborate.

4
3
2
1
10
20
30
40

12
10
8
6
4
2
110
120
130
140

These graphs were made using *Mathematica* commands like

```
Plot[Abs[Zeta[I*y]], {y, 100, 140}].
```

It's fun to try them with other values of the range.

Exact values for the zeta function

The zeta function can be evaluated exactly for positive even integer arguments:

$$\zeta(m) = \frac{(2\pi)^m |B_m|}{2m!} \quad \text{for} \quad m = 2, 4, 6, \ldots.$$

Here the B_m are the Bernoulli numbers, defined through

$$\frac{x}{e^x - 1} = \sum_{m=0}^{\infty} \frac{B_m}{m!} x^m.$$

The first few Bernoulli numbers are

$$B_0 = 1, \quad B_1 = -\tfrac{1}{2}, \quad B_2 = \tfrac{1}{6}, \quad B_4 = -\tfrac{1}{30}, \quad B_6 = \tfrac{1}{42}, \quad B_8 = -\tfrac{1}{30}, \quad B_{10} = \tfrac{5}{66}, \quad B_{12} = -\tfrac{691}{2730}$$

whence

$$\zeta(2) = \frac{\pi^2}{6}, \quad \zeta(4) = \frac{\pi^4}{90}, \quad \zeta(6) = \frac{\pi^6}{945}, \quad \zeta(8) = \frac{\pi^8}{9450}.$$

The zeta function near $s = 1$

The zeta function diverges with a simple pole at $s = 1$. In the vicinity of the divergence it is approximately equal to

$$\zeta(s) \approx C + \frac{1}{s-1}$$

where C is Euler's constant, 0.5772.... (The rest of this section is stolen from Bender and Orszag.)

When s is near 1, the defining series

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

is *very* slowly converging. About 10^{20} terms are needed to compute $\zeta(1.1)$ accurate to 1 percent!

Fortunately, there is an expression for the difference between $\zeta(s)$ and the Nth partial sum:

$$\zeta(s) = \sum_{n=0}^{N} \frac{1}{s^n} + \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{u^{s-1} e^{-Nu}}{e^u - 1}\, du.$$

The integral cannot be evaluated analytically but it can be expanded in an asymptotic series, giving

$$\zeta(s) \approx \sum_{n=0}^{N} \frac{1}{s^n} + \frac{1}{(s-1)N^{s-1}} - \frac{1}{2N^s} + \frac{B_2 s}{N^{s+1}} + \frac{B_4 s(s+1)(s+2)}{2N^{s+3}} + \cdots$$

where the B_m are again the Bernoulli numbers.

Truncating this formula can give extremely accurate results. For example, using $N = 9$, only 38 terms of the series are needed to find $\zeta(1.1)$ accurate to 26 decimal places.

References

G.F. Simmons, *Differential Equations with Applications and Historical Notes*, (McGraw-Hill, New York, 1972) pp. 210–211, 214–218.

C.M. Bender and S.A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers*, (McGraw-Hill, New York, 1978) p. 379.

Jahnke, Emde, and Lösch, *Tables of Higher Functions*, 6th ed. (McGraw-Hill, New York, 1960) chap. IV. (Note particularly the beautiful relief plot of $|\zeta(s)|$ for complex values of s.)

J. Spanier and K. Oldham, *An Atlas of Functions*, (Hemisphere Publishing, Washington, D.C., 1987) chap. 3. (Presents an algorithm purporting to find $\zeta(s)$ accurately for any real s.)

M. Abramowitz and I. Stegum, *Handbook of Mathematical Functions*, (National Bureau of Standards, Washington, D.C., 1964) chap. 23.

M.C. Gutzwiller, *Chaos in Classical and Quantum Mechanics*, (Springer-Verlag, New York, 1990) section 17.9.

Appendix H

Tutorial on Matrix Diagonalization

You know from as far back as your introductory mechanics course that some problems are difficult given one choice of coordinate axes and easy or even trivial given another. (For example, the famous "monkey and hunter" problem is difficult using a horizontal axis, but easy using an axis stretching from the hunter to the monkey.) The mathematical field of linear algebra is devoted, in large part, to systematic techniques for finding coordinate systems that make problems easy. This tutorial introduces the most valuable of these techniques. It assumes that you are familiar with matrix multiplication and with the ideas of the inverse, the transpose, and the determinant of a square matrix. It is also useful to have a nodding acquaintance with the inertia tensor.

This presentation is intentionally non-rigorous. A rigorous, formal treatment of matrix diagonalization can be found in any linear algebra textbook,[1] and there is no need to duplicate that function here. What is provided here instead is a heuristic picture of what's going on in matrix diagonalization, how it works, and why anyone would want to do such a thing anyway. Thus this presentation complements, rather than replaces, the logically impeccable ("bulletproof") arguments of the mathematics texts.

Essential problems in this tutorial are marked by asterisks (*).

Warning: This tutorial is still in **draft** form.

H.1 What's in a name?

There is a difference between an entity and its name. For example, a tree is made of wood, whereas its name "tree" made of ink. One way to see this is to note that in German, the name for a tree is "Baum", so the name changes upon translation, but the tree itself does not change. (Throughout this tutorial, the term "translate" is used as in "translate from one language to another" rather than as in "translate by moving in a straight line".)

The same holds for mathematical entities. Suppose a length is represented by the number "2" because it is two feet long. Then the same length is represented by the number "24" because it is twenty-four inches long. The same length is represented by two different numbers, just as the same tree has two different names. The representation of a length as a number depends not only upon the length, but also upon the coordinate system used to measure the length.

H.2 Vectors in two dimensions

One way of describing a two-dimensional vector $\mathbf{V}$ is by giving its x and y components in the form of a 2×1 column matrix

$$\begin{pmatrix} V_x \\ V_y \end{pmatrix}. \tag{H.1}$$

Indeed it is sometimes said that the vector $\mathbf{V}$ is *equal* to the column matrix (H.1). This is not precisely correct—it is better to say that the vector is *described by* the column matrix or *represented by* the column matrix or that its *name* is the column matrix. This is because if you describe the vector using a different set of coordinate axes you will come up with a different column matrix to describe the same vector. For example, in the situation shown below the descriptions in terms of the two different coordinate systems are related through the matrix equation

$$\begin{pmatrix} V_{x'} \\ V_{y'} \end{pmatrix} = \begin{pmatrix} \cos\phi & \sin\phi \\ -\sin\phi & \cos\phi \end{pmatrix} \begin{pmatrix} V_x \\ V_y \end{pmatrix}. \tag{H.2}$$

The 2×2 matrix above is called the "rotation matrix" and is usually denoted by $\mathsf{R}(\phi)$:

$$\mathsf{R}(\phi) \equiv \begin{pmatrix} \cos\phi & \sin\phi \\ -\sin\phi & \cos\phi \end{pmatrix}. \tag{H.3}$$

One interesting property of the rotation matrix is that it is always invertible, and that its inverse is equal to its transpose. Such matrices are called *orthogonal*.[1] You could prove this by working a matrix multiplication, but it is easier to simply realize that the inverse of a rotation by ϕ is simply a rotation by $-\phi$, and noting that

$$\mathsf{R}^{-1}(\phi) = \mathsf{R}(-\phi) = \mathsf{R}^{\dagger}(\phi). \tag{H.4}$$

(The dagger represents matrix transposition.)

There are, of course, an infinite number of column matrix representations for any vector, corresponding to the infinite number of coordinate axis rotations with ϕ from 0 to 2π. But one of these representations is special: It is the one in which the x'-axis lines up with the vector, so the column matrix representation is just

$$\begin{pmatrix} V \\ 0 \end{pmatrix}, \tag{H.5}$$

where $V = |\mathbf{V}| = \sqrt{V_x^2 + V_y^2}$ is the magnitude of the vector. This set of coordinates is the preferred (or "canonical") set for dealing with this vector: one of the two components is zero, the easiest number to deal with, and the other component is a physically important number. You might wonder how I can claim that this representation has full information about the vector: The initial representation (H.1) contains two independent numbers, whereas the preferred representation (H.5) contains only one. The answer is that the preferred representation contains one number (the magnitude of the vector) explicitly while another number (the polar angle of the vector relative to the initial x-axis) is contained implicitly in the rotation needed to produce the preferred coordinate system.

H.1 **Problem: Right angle rotations**

Verify equation (H.2) in the special cases $\phi = 90°$, $\phi = 180°$, $\phi = 270°$, and $\phi = 360°$.

H.2 **Problem: The rotation matrix**

a. Derive equation (H.2) through purely geometrical arguments.

b. Express $\hat{\mathbf{i}}'$ and $\hat{\mathbf{j}}'$, the unit vectors of the (x', y') coordinate system, as linear combinations of $\hat{\mathbf{i}}$ and $\hat{\mathbf{j}}$. Then use

$$V_{x'} = \mathbf{V}\cdot\hat{\mathbf{i}}' \quad \text{and} \quad V_{y'} = \mathbf{V}\cdot\hat{\mathbf{j}}' \tag{H.6}$$

to derive equation (H.2).

c. Which derivation do you find easier?

H.3 **Problem: Rotation to the preferred coordinate system***

In the preferred coordinate system, $V_{y'} = 0$. Use this requirement to show that the preferred system is rotated from the initial system by an angle ϕ with

$$\tan\phi = \frac{V_y}{V_x}. \tag{H.7}$$

[1] Although all rotation matrices are orthogonal, there are orthogonal matrices that are not rotation matrices: see problem H.4.

For any value of V_y/V_x, there are *two* angles that satisfy this equation. What is the representation of $\mathbf{V}$ in each of these two coordinate systems?

H.4 **Problem: A non-rotation orthogonal transformation**
In one coordinate system the y-axis is vertical and the x-axis points to the right. In another the y'-axis is vertical and the x'-axis points to the left. Find the matrix that translates vector coordinates from one system to the other. Show that this matrix is orthogonal but not a rotation matrix.

H.5 **Problem: Other changes of coordinate***
Suppose vertical distances (distances in the y direction) are measured in feet while horizontal distances (distances in the x direction) are measured in miles. (This system is not perverse. It is used in nearly all American road maps.) Find the matrix that changes the representation of a vector in this coordinate system to the representation of a vector in a system where all distances are measured in feet. Find the matrix that translates back. Are these matrices orthogonal?

H.6 **Problem: Other special representations**
At equation (H.5) we mentioned one "special" (or "canonical") representation of a vector. There are three others, namely

$$\begin{pmatrix} 0 \\ -V \end{pmatrix}, \qquad \begin{pmatrix} -V \\ 0 \end{pmatrix}, \qquad \begin{pmatrix} 0 \\ V \end{pmatrix}. \tag{H.8}$$

If coordinate-system rotation angle ϕ brings the vector representation into the form (H.5), then what rotation angle will result in these three representations?

H.3 Tensors in two dimensions

A tensor, like a vector, is a geometrical entity that may be described ("named") through components, but a d-dimensional tensor requires d^2 rather than d components. Tensors are less familiar and more difficult to visualize than vectors, but they are neither less important nor "less physical". We will introduce tensors through the concrete example of the inertia tensor of classical mechanics (see, for example, reference [2]), but the results we present will be perfectly general.

Just as the two components of a two-dimensional vector are most easily kept track of through a 2×1 matrix, so the four components of two-dimensional tensor are most conveniently written in the form of a 2×2 matrix. For example, the inertia tensor $\mathbf{T}$ of a point particle with mass m located[2] at (x, y) has components

$$\mathsf{T} = \begin{pmatrix} my^2 & -mxy \\ -mxy & mx^2 \end{pmatrix}. \tag{H.9}$$

(Note the distinction between the tensor $\mathbf{T}$ and its matrix of components, its "name", T.) As with vector components, the tensor components are different in different coordinate systems, although the tensor itself

[2]Or, to be absolutely precise, the particle located at the point represented by the vector with components (x, y).

does not change. For example, in the primed coordinate system of the figure on page 231, the tensor components are of course

$$\mathsf{T}' = \begin{pmatrix} my'^2 & -mx'y' \\ -mx'y' & mx'^2 \end{pmatrix}. \tag{H.10}$$

A little calculation shows that the components of the inertia tensor in two different coordinate systems are related through

$$\mathsf{T}' = \mathsf{R}(\phi)\mathsf{T}\mathsf{R}^{-1}(\phi). \tag{H.11}$$

This relation holds for *any* tensor, not just the inertia tensor. (In fact, one way to define "tensor" is as an entity with four components that satisfy the above relation under rotation.) If the matrix representing a tensor is symmetric (i.e. the matrix is equal to its transpose) in one coordinate system, then it is symmetric in all coordinate systems (see problem H.7). Therefore the symmetry is a property of the tensor, not of its matrix representation, and we may speak of "a symmetric tensor" rather than just "a tensor represented by a symmetric matrix".

As with vectors, one of the many matrix representations of a given tensor is considered special (or "canonical"): It is the one in which the lower left component is zero. Furthermore if the tensor is symmetric (as the inertia tensor is) then in this preferred coordinate system the upper right component will be zero also, so the matrix will be all zeros except for the diagonal elements. Such a matrix is called a "diagonal matrix" and the process of finding the rotation that renders the matrix representation of a symmetric tensor diagonal is called "diagonalization".[3] We may do an "accounting of information" for this preferred coordinate system just as we did with vectors. In the initial coordinate system, the symmetric tensor had three independent components. In the preferred system, it has two independent components manifestly visible in the diagonal matrix representation, and one number hidden through the specification of the rotation.

H.7 **Problem: Representations of symmetric tensors***

Show that if the matrix S representing a tensor is symmetric, and if B is any orthogonal matrix, then all of the representations

$$\mathsf{B}\mathsf{S}\mathsf{B}^\dagger \tag{H.12}$$

are symmetric. (Clue: If you try to solve this problem for rotations in two dimensions using the explicit rotation matrix (H.3), you will find it solvable but messy. The clue is that this problem asks you do prove the result in any number of dimensions, and for any orthogonal matrix B, not just rotation matrices. This more general problem is considerably easier to solve.)

H.8 **Problem: Diagonal inertia tensor**

The matrix (H.9) represents the inertia tensor of a point particle with mass m located a distance r from the origin. Show that the matrix is diagonal in four different coordinate systems: one in which the x'-axis points directly toward the particle, one in which the y'-axis points directly away from the particle, one in which the x'-axis points directly away from the particle, and one in which the y'-axis

[3] An efficient algorithm for diagonalization is discussed in section H.8. For the moment, we are more interested in knowing that a diagonal matrix representation must exist than in knowing how to most easily find that preferred coordinate system.

points directly toward the particle. Find the matrix representation in each of these four coordinate systems.

H.9 **Problem: Representations of a certain tensor**

Show that a tensor represented in one coordinate system by a diagonal matrix with equal elements, namely

$$\begin{pmatrix} d_0 & 0 \\ 0 & d_0 \end{pmatrix}, \tag{H.13}$$

has the same representation in *all* orthogonal coordinate systems.

H.10 **Problem: Rotation to the preferred coordinate system***

A tensor is represented in the initial coordinate system by

$$\begin{pmatrix} a & b \\ b & c \end{pmatrix}. \tag{H.14}$$

Show that the tensor is diagonal in a preferred coordinate system which is rotated from the initial system by an angle ϕ with

$$\tan(2\phi) = \frac{2b}{a-c}. \tag{H.15}$$

This equation has four solutions. Find the rotation matrix for $\phi = 90°$, then show how the four different diagonal representations are related. You do not need to find any of the diagonal representations in terms of a, b and c...just show what the other three are given that one of them is

$$\begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}. \tag{H.16}$$

H.11 **Problem: Inertia tensor in outer product notation**

The discussion in this section has emphasized the tensor's matrix representation ("name") T rather than the tensor $\mathbf{T}$ itself.

a. Define the "identity tensor" $\mathbf{1}$ as the tensor represented in some coordinate system by

$$\mathsf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \tag{H.17}$$

Show that this tensor has the same representation in any coordinate system.

b. Show that the inner product between two vectors results in a scalar: Namely

$$\text{if vector } bfa \text{ is represented by } \begin{pmatrix} a_x \\ a_y \end{pmatrix} \text{ and vector } bfb \text{ is represented by } \begin{pmatrix} b_x \\ b_y \end{pmatrix}$$

then the inner product $\mathbf{a} \cdot \mathbf{b}$ is given through

$$\begin{pmatrix} a_x & a_y \end{pmatrix} \begin{pmatrix} b_x \\ b_y \end{pmatrix} = a_x b_x + a_y b_y,$$

and this inner product is a scalar. (A 1×2 matrix times a 2×1 matrix is a 1×1 matrix.) That is, the vector $\mathbf{a}$ is represented by different coordinates in different coordinate systems, and the vector $\mathbf{b}$ is represented by different coordinates in different coordinate systems, but the inner product $\mathbf{a} \cdot \mathbf{b}$ is the same in all coordinate systems.

c. In contrast, show that the outer product of two vectors is a tensor: Namely

$$\mathbf{ab} \doteq \begin{pmatrix} a_x \\ a_y \end{pmatrix} \begin{pmatrix} b_x & b_y \end{pmatrix} = \begin{pmatrix} a_x b_x & a_x b_y \\ a_y b_x & a_y b_y \end{pmatrix}.$$

(A 2×1 matrix times a 1×2 matrix is a 2×2 matrix.) That is, show that the representation of $\mathbf{ab}$ transforms from one coordinate system to another as specified through (H.11).

d. Show that the inertia tensor for a single particle of mass m located at position $\mathbf{r}$ can be written in coordinate-independent fashion as

$$\mathbf{T} = m\mathbf{1}r^2 - m\mathbf{rr}. \tag{H.18}$$

H.4 Tensors in three dimensions

A three-dimensional tensor is represented in component form by a 3×3 matrix with nine entries. If the tensor is symmetric, there are six independent elements... three on the diagonal and three off-diagonal. The components of a tensor in three dimensions change with coordinate system according to

$$\mathsf{T}' = \mathsf{R}\mathsf{T}\mathsf{R}^\dagger, \tag{H.19}$$

where R is the 3×3 rotation matrix.

A rotation in two dimension is described completely by giving a single angle. In three dimensions more information is required. Specifically, we need not only the amount of the rotation, but we must also know the plane in which the rotation takes place. We can specify the plane by giving the unit vector perpendicular to that plane. Specifying an arbitrary vector in three dimensions requires three numbers, but specifying a unit vector in three dimensions requires only two numbers because the magnitude is already fixed at unity. Thus three numbers are required to specify a rotation in three dimensions: two to specify the rotation's plane, one to specify the rotation's size. (One particularly convenient way to specify a three-dimensional rotation is through the three *Euler angles*. Reference [3] defines these angles and shows how to write the 3×3 rotation matrix in terms of these variables. For the purposes of this tutorial, however, we will not need an explicit rotation matrix... all we need is to know is the number of angles required to specify a rotation.)

In two dimensions, any symmetric tensor (which has three independent elements), could be represented by a diagonal tensor (with two independent elements) plus a rotation (one angle). We were able to back up this claim with an explicit expression for the angle.

In three dimensions it seems reasonable that any symmetric tensor (six independent elements) can be represented by a diagonal tensor (three independent elements) plus a rotation (three angles). The three

angles just have to be selected carefully enough to make sure that they cause the off-diagonal elements to vanish. This supposition is indeed correct, although we will not pause for long enough to prove it by producing explicit formulas for the three angles.

H.5 Tensors in d dimensions

A d-dimensional tensor is represented by a $d \times d$ matrix with d^2 entries. If the tensor is symmetric, there are d independent on-diagonal elements and $d(d-1)/2$ independent off-diagonal elements. The tensor components will change with coordinate system in the now-familiar form

$$\mathsf{T}' = \mathsf{R}\mathsf{T}\mathsf{R}^\dagger, \tag{H.20}$$

where R is the $d \times d$ rotation matrix.

How many angles does it take to specify a rotation in d dimensions? Remember how we went from two dimensions to three: The three dimensional rotation took place "in a plane", i.e. in a two-dimensional subspace. It required two (i.e. $d-1$) angles to specify the orientation of the plane plus one to specify the rotation within the plane... a total of three angles.

A rotation in four dimensions takes place within a three-dimensional subspace. It requires $3 = d-1$ angles to specify the orientation of the three-dimensional subspace, plus, as we found above, three angles to specify the rotation within the three-dimensional subspace... a total of six angles.

A rotation in five dimensions requires $4 = d-1$ angles to specify the four-dimensional subspace in which the rotation occurs, plus the six angles that we have just found specify a rotation within that subspace... a total of ten angles.

In general, the number of angles needed to specify a rotation in d dimensions is

$$A_d = d - 1 + A_{d-1} = d(d-1)/2. \tag{H.21}$$

This is exactly the number of independent off-diagonal elements in a symmetric tensor. It seems reasonable that we can choose the angles to ensure that, in the resulting coordinate system, all the off-diagonal elements vanish. The proof of this result is difficult and proceeds in a very different manner from the plausibility argument sketched here. (The proof involves concepts like eigenvectors and eigenvalues, and it gives an explicit recipe for constructing the rotation matrix. It has the advantage of rigor and the disadvantage of being so technical that it's easy to lose track of the fact that that all you're doing is choosing a coordinate system.)

H.12 **Problem: Non-symmetric tensors***

Argue that a non-symmetric tensor can be brought into a "triangular" representation in which all the elements below the diagonal are equal to zero and all the elements on and above the diagonal are independent. (This is indeed the case, although in general some of the non-zero elements remaining will be complex-valued, and some of the angles will involve rotations into complex-valued vectors.)

H.6 Linear transformations in two dimensions

Section H.3 considered 2×2 matrices as representations of tensors. This section gains additional insight by considering 2×2 matrices as representations of linear transformations. It demonstrates how diagonalization can be useful and gives a clue to an efficient algorithm for diagonalization.

A linear transformation is a function from vectors to vectors that can be represented in any given coordinate system as

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \tag{H.22}$$

If the equation above represents ("names") the transformation in one coordinate system, what is its representation in some other coordinate system? We assume that the two coordinate systems are related through an orthogonal matrix B such that

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \mathsf{B} \begin{pmatrix} u \\ v \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x' \\ y' \end{pmatrix} = \mathsf{B} \begin{pmatrix} x \\ y \end{pmatrix}. \tag{H.23}$$

(For example, if the new coordinate system is the primed coordinate system of the figure on page 231, then the matrix B that translates from the original to the new coordinates is the rotation matrix $\mathsf{R}(\phi)$.) Given this "translation dictionary", we have

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \mathsf{B} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \tag{H.24}$$

But B is invertible, so

$$\begin{pmatrix} x \\ y \end{pmatrix} = \mathsf{B}^{-1} \begin{pmatrix} x' \\ y' \end{pmatrix} \tag{H.25}$$

whence

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \mathsf{B} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \mathsf{B}^{-1} \begin{pmatrix} x' \\ y' \end{pmatrix}. \tag{H.26}$$

Thus the representation of the transformation in the primed coordinate system is

$$\mathsf{B} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \mathsf{B}^{-1} \tag{H.27}$$

(compare equation H.11). This equation has a very direct physical meaning. Remember that the matrix B translates from the old (x, y) coordinates to the new (x', y') coordinates, while the matrix B^{-1} translates in the opposite direction. Thus the equation above says that the representation of a transformation in the new coordinates is given by translating from new to old coordinates (through the matrix B^{-1}), then applying the old representation (the "a matrix") to those old coordinates, and finally translating back from old to new coordinates (through the matrix B).

The rest of this section considers only transformations represented by symmetric matrices, which we will denote by

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \tag{H.28}$$

Let's try to understand this transformation as something more than a jumble of symbols awaiting a plunge into the calculator. First of all, suppose the vector $\mathbf{V}$ maps to the vector $\mathbf{W}$. Then the vector $5\mathbf{V}$ will be mapped to vector $5\mathbf{W}$. In short, if we know how the transformation acts on vectors with magnitude unity, we will be able to see immediately how it acts on vectors with other magnitudes. Thus we focus our attention on vectors on the unit circle:

$$x^2 + y^2 = 1. \tag{H.29}$$

A brief calculation shows that the length of the output vector is then

$$\sqrt{u^2 + v^2} = \sqrt{a^2x^2 + b^2 + c^2y^2 + 2b(a+c)xy}, \tag{H.30}$$

which isn't very helpful. Another brief calculation shows that if the input vector has polar angle θ, then the output vector has polar angle φ with

$$\tan\varphi = \frac{b + c\tan\theta}{a + b\tan\theta}, \tag{H.31}$$

which is similarly opaque and messy.

Instead of trying to understand the transformation in its initial coordinate system, let's instead convert (rotate) to the special coordinate system in which the transformation is represented by a diagonal matrix. In this system,

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix} \begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} d_1x' \\ d_2y' \end{pmatrix}. \tag{H.32}$$

The unit circle is still

$$x'^2 + y'^2 = 1, \tag{H.33}$$

so the image of the unit circle is

$$\left(\frac{u'}{d_1}\right)^2 + \left(\frac{v'}{d_2}\right)^2 = 1, \tag{H.34}$$

namely an ellipse! This result is transparent in the special coordinate system, but almost impossible to see in the original one.

Note particularly what happens to a vector pointing along the x' coordinate axis. For example, the unit vector in this direction transforms to

$$\begin{pmatrix} d_1 \\ 0 \end{pmatrix} = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \tag{H.35}$$

In other words, the when the vector is transformed it changes in magnitude, but *not* in direction. Vectors with this property are called *eigenvectors*. It is easy to see that any vector on either the x' or y' coordinate axes are eigenvectors.

H.7 What does "eigen" mean?

If a vector $\mathbf{x}$ is acted upon by a linear transformation B, then the output vector

$$\mathbf{x}' = \mathsf{B}\mathbf{x} \tag{H.36}$$

will usually be skew to the original vector $\mathbf{x}$. However, for some very special vectors it might just happen that $\mathbf{x}'$ is parallel to $\mathbf{x}$. Such vectors are called "eigenvectors". (This is a terrible name because (1) it gives no idea of what eigenvectors are or why they're so important and (2) it sounds gross. However, that's what they're called.) We have already seen, in the previous section, that eigenvectors are related to coordinate systems in which the transformation is particularly easy to understand.

If $\mathbf{x}$ is an eigenvector, then

$$\mathsf{B}\mathbf{x} = \lambda\mathbf{x}, \tag{H.37}$$

where λ is a scalar called "the eigenvalue associated with eigenvector $\mathbf{x}$". If $\mathbf{x}$ is an eigenvector, then any vector parallel to $\mathbf{x}$ is also an eigenvector with the same eigenvalue. (That is, any vector of the form $c\mathbf{x}$, where c is any scalar, is also an eigenvector with the same eigenvalue.) Sometimes we speak of a "line of eigenvectors".

The vector $\mathbf{x} = \mathbf{0}$ is never considered an eigenvector, because

$$\mathsf{B}\mathbf{0} = \lambda\mathbf{0}, \tag{H.38}$$

for any value of λ for any linear transformation. On the other hand, if

$$\mathsf{B}\mathbf{x} = 0\mathbf{x} = \mathbf{0} \tag{H.39}$$

for some non-zero vector $\mathbf{x}$, then $\mathbf{x}$ is an eigenvector with eigenvalue $\lambda = 0$.

H.13 **Problem: Plane of eigenvectors**

Suppose $\mathbf{x}$ and $\mathbf{y}$ are two non-parallel vectors with the same eigenvalue. (In this case the eigenvalue is said to be "degenerate", which sounds like an aspersion cast upon the morals of the eigenvalue but which is really just poor choice of terminology again.) Show that any vector of the form $c_1\mathbf{x} + c_2\mathbf{y}$ is an eigenvector with the same eigenvalue.

H.8 How to diagonalize a symmetric matrix

We saw in section H.3 that for any 2×2 symmetric matrix, represented in its initial basis by, say,

$$\begin{pmatrix} a & b \\ b & c \end{pmatrix}, \tag{H.40}$$

a simple rotation of axes would produce a new coordinate system in which the matrix representation is diagonal:

$$\begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}. \tag{H.41}$$

These two matrices are related through

$$\begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix} = \mathsf{R}(\phi) \begin{pmatrix} a & b \\ b & c \end{pmatrix} \mathsf{R}^{-1}(\phi), \tag{H.42}$$

where $\mathsf{R}(\phi)$ is the rotation matrix (H.3). Problem H.10 gave a direct way to find the desired rotation. However this direct technique is cumbersome and doesn't generalize readily to higher dimensions. This section presents a different technique, which relies on eigenvalues and eigenvectors, that is more efficient and that generalizes readily to complex-valued matrices and to matrices in any dimension, but that is somewhat sneaky and conceptually roundabout.

We begin by noting that any vector lying along the x'-axis (of the preferred coordinate system) is an eigenvector. For example, the vector $5\hat{\mathbf{i}}'$ is represented (in the preferred coordinate system) by

$$\begin{pmatrix} 5 \\ 0 \end{pmatrix}. \tag{H.43}$$

Multiplying this vector by the matrix in question gives

$$\begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix} \begin{pmatrix} 5 \\ 0 \end{pmatrix} = d_1 \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \tag{H.44}$$

so $5\hat{\mathbf{i}}'$ is an eigenvector with eigenvalue d_1. The same holds for any scalar multiple of $\hat{\mathbf{i}}'$, whether positive or negative. Similarly, any scalar multiple of $\hat{\mathbf{j}}'$ is an eigenvector with eigenvalue d_2. In short, the two elements on the diagonal in the preferred (diagonal) representation are the two eigenvalues, and the two unit vectors $\hat{\mathbf{i}}'$ and $\hat{\mathbf{j}}'$ of the preferred coordinate system are two of the eigenvectors.

Thus finding the eigenvectors and eigenvalues of a matrix gives you the information needed to diagonalize that matrix. The unit vectors $\hat{\mathbf{i}}'$ and $\hat{\mathbf{j}}'$ constitute an "orthonormal basis of eigenvectors". The eigenvectors even give the rotation matrix directly, as described in the next paragraph.

Let's call the rotation matrix

$$\mathsf{B} = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, \tag{H.45}$$

so that the inverse (transpose) matrix is

$$\mathsf{B}^{-1} = \mathsf{B}^{\dagger} = \begin{pmatrix} b_{11} & b_{21} \\ b_{12} & b_{22} \end{pmatrix}. \tag{H.46}$$

The representation of $\hat{\mathbf{i}}'$ in the preferred basis is

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \tag{H.47}$$

so its representation in the initial basis is (see equation H.2)

$$\mathsf{B}^{\dagger} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} b_{11} & b_{21} \\ b_{12} & b_{22} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} b_{11} \\ b_{12} \end{pmatrix}. \tag{H.48}$$

Similarly, the representation of $\hat{\mathbf{j}}'$ in the initial basis is

$$\mathsf{B}^{\dagger} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} b_{11} & b_{21} \\ b_{12} & b_{22} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} b_{21} \\ b_{22} \end{pmatrix}. \tag{H.49}$$

Thus the rotation matrix is

$$\mathsf{B} = \begin{pmatrix} \text{initial rep. of } \hat{\mathbf{i}}', \text{ on its side} \\ \text{initial rep. of } \hat{\mathbf{j}}', \text{ on its side} \end{pmatrix}. \tag{H.50}$$

Example

Suppose we need to find a diagonal representation for the matrix

$$\mathsf{T} = \begin{pmatrix} 7 & 3 \\ 3 & 7 \end{pmatrix}. \tag{H.51}$$

First we search for the special vectors—the eigenvectors—such that

$$\begin{pmatrix} 7 & 3 \\ 3 & 7 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \lambda \begin{pmatrix} x \\ y \end{pmatrix}. \tag{H.52}$$

At the moment, we don't know either the eigenvalue λ or the associated eigenvector (x, y). Thus it seems that (bad news) we are trying to solve two equations for three unknowns:

$$\begin{aligned} 7x + 3y &= \lambda x \\ 3x + 7y &= \lambda y \end{aligned} \tag{H.53}$$

Remember, however, that there is not one *single* eigenvector: any multiple of an eigenvector is also an eigenvector. (Alternatively, any vector on the line that extends the eigenvector is another eigenvector.) We only need one of these eigenvectors, so let's take the one that has $x = 1$ (i.e. the vector on the extension line where it intersects the vertical line $x = 1$). (This technique will fail if we have the bad luck that our actual eigenvector is vertical and hence never passes through the line $x = 1$.) So we really have two equations in two unknowns:

$$\begin{aligned} 7 + 3y &= \lambda \\ 3 + 7y &= \lambda y \end{aligned}$$

but note that they are not *linear* equations... the damnable product λy in the lower right corner means that all our techniques for solving linear equations go right out the window. We *can* solve these two equations for λ and y, but there's an easier, if somewhat roundabout, approach.

Finding eigenvalues

Let's go back to equation (H.52) and write it as

$$\begin{pmatrix} 7 & 3 \\ 3 & 7 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} - \lambda \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{H.54}$$

Then

$$\begin{pmatrix} 7 & 3 \\ 3 & 7 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \tag{H.55}$$

or

$$\begin{pmatrix} 7-\lambda & 3 \\ 3 & 7-\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{H.56}$$

Let's think about this. It says that for some matrix $\mathsf{M} = \mathsf{T} - \lambda \mathsf{1}$, we have

$$\mathsf{M} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{H.57}$$

You know right away one vector (x, y) that satisfies this equation, namely $(x, y) = (0, 0)$. And most of the time, this is the *only* vector that satisfies the equation, because

$$\begin{pmatrix} x \\ y \end{pmatrix} = \mathsf{M}^{-1} \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{H.58}$$

We appear to have reached a dead end. The solution is $(x, y) = (0, 0)$, but the zero vector is not, by definition, considered an eigenvector of any transformation. (Because it always gives eigenvalue zero for any transformation.)

However, if the matrix M is *not invertible*, then there will be other solutions to

$$\mathsf{M} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{H.59}$$

in addition to the trivial solution $(x, y) = (0, 0)$. Thus we must look for those special values of λ such that the so-called *characteristic matrix* M is not invertible. These values come if and only if the determinant of M vanishes. For this example, we have to find values of λ such that

$$\det \begin{pmatrix} 7-\lambda & 3 \\ 3 & 7-\lambda \end{pmatrix} = 0. \tag{H.60}$$

This is a quadratic equation in λ

$$(7-\lambda)^2 - 3^2 = 0 \tag{H.61}$$

called the *characteristic equation*. Its two solutions are

$$7 - \lambda = \pm 3 \tag{H.62}$$

or

$$\lambda = 7 \pm 3 = 10 \text{ or } 4. \tag{H.63}$$

We have found the two eigenvalues of our matrix!

Finding eigenvectors

Let's look now for the eigenvector associated with $\lambda = 4$. Equation (H.53)

$$\begin{aligned} 7x + 3y &= \lambda x \\ 3x + 7y &= \lambda y \end{aligned}$$

still holds, but no longer does it look like two equations in three unknowns, because we are now interested in the case $\lambda = 4$:

$$\begin{aligned} 7x + 3y &= 4x \\ 3x + 7y &= 4y \end{aligned}$$

Following our nose gives

$$\begin{aligned} 3x + 3y &= 0 \\ 3x + 3y &= 0 \end{aligned}$$

and when we see this our heart skips a beat or two. . . a degenerate system of equations! Relax and rest your heart. This system has an infinite number of solutions and it's supposed to have an infinite number of solutions, because any multiple of an eigenvector is also an eigenvector. The eigenvectors associated with $\lambda = 4$ are any multiple of

$$\begin{pmatrix} 1 \\ -1 \end{pmatrix}. \tag{H.64}$$

An entirely analogous search for the eigenvectors associated with $\lambda = 10$ finds any multiple of

$$\begin{pmatrix} 1 \\ 1 \end{pmatrix}. \tag{H.65}$$

Tidying up

We have the two sets of eigenvectors, but which shall we call $\hat{\mathbf{i}}'$ and which $\hat{\mathbf{j}}'$? This is a matter of individual choice, but my choice is usually to make the transformation be a rotation (without reflection) through a small positive angle. Our new, preferred coordinate system is related to the original coordinates by a simple rotation of 45° if we choose

$$\hat{\mathbf{i}}' = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \text{and} \quad \hat{\mathbf{j}}' = \tfrac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix}. \tag{H.66}$$

(Note that we have also "normalized the basis", i.e. selected the basis vectors to have magnitude unity.) Given this choice, the orthogonal rotation matrix that changes coordinates from the original to the preferred system is (see equation H.50)

$$\mathsf{B} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \tag{H.67}$$

and the diagonalized matrix (or, more properly, the representation of the matrix in the preferred coordinate system) is

$$\begin{pmatrix} 10 & 0 \\ 0 & 4 \end{pmatrix}. \tag{H.68}$$

You don't believe me? Then multiply out

$$\mathsf{B}\begin{pmatrix} 7 & 3 \\ 3 & 7 \end{pmatrix}\mathsf{B}^\dagger \tag{H.69}$$

and see for yourself.

Problems

H.14 **Problem: Diagonalize a 2×2 matrix***

Diagonalize the matrix

$$\begin{pmatrix} 26 & 12 \\ 12 & 19 \end{pmatrix}. \tag{H.70}$$

a. Find its eigenvalues.

b. Find its eigenvectors, and verify that they are orthogonal.

c. Sketch the eigenvectors, and determine the signs and sequence most convenient for assigning axes. (That is, should the first eigenvector you found be called $\hat{\mathbf{i}}'$, $-\hat{\mathbf{i}}'$, or $\hat{\mathbf{j}}'$?)

d. Find the matrix that translates from the initial basis to the basis of eigenvectors produced in part (c.).

e. Verify that the matrix produced in part (d.) is orthogonal.

f. Verify that the representation of the matrix above in the basis of eigenvectors is diagonal.

g. (Optional.) What is the rotation angle?

H.15 **Problem: Eigenvalues of a 2×2 matrix**

Show that the eigenvalues of

$$\begin{pmatrix} a & b \\ b & c \end{pmatrix} \tag{H.71}$$

are

$$\lambda = \tfrac{1}{2}\left[(a+c) \pm \sqrt{(a-c)^2 + 4b^2}\right]. \tag{H.72}$$

Under what circumstances is an eigenvalue complex valued? Under what circumstances are the two eigenvalues the same?

H.16 **Problem: Diagonalize a 3×3 matrix**

Diagonalize the matrix

$$\frac{1}{625}\begin{pmatrix} 1182 & -924 & 540 \\ -924 & 643 & 720 \\ 540 & 720 & -575 \end{pmatrix}. \tag{H.73}$$

a. Find its eigenvalues by showing that the characteristic equation is

$$\lambda^3 - 2\lambda^2 - 5\lambda + 6 = (\lambda - 3)(\lambda + 2)(\lambda - 1) = 0. \tag{H.74}$$

b. Find its eigenvectors, and verify that they are orthogonal.

c. Show that the translation matrix can be chosen to be

$$\mathsf{B} = \frac{1}{25}\begin{pmatrix} 20 & -15 & 0 \\ 9 & 12 & -20 \\ 12 & 16 & 15 \end{pmatrix}. \tag{H.75}$$

Why did I use the phrase "the translation matrix can be chosen to be" rather then "the translation matrix is"?

H.17 **Problem: A 3×3 matrix eigenproblem**

Find the eigenvalues and associated eigenvectors for the matrix

$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{pmatrix}. \tag{H.76}$$

H.9 A glance at computer algorithms

Anyone who has worked even one of the problems in section H.8 knows that diagonalizing a matrix is no picnic: there's a lot of mundane arithmetic involved and it's very easy to make mistakes. This is a problem ripe for computer solution. One's first thought is to program a computer to solve the problem using the same technique that we used to solve it on paper: first find the eigenvalues through the characteristic equation, then find the eigenvectors through a degenerate set of linear equations.

This turns out to be a very poor algorithm for automatic computation. The effective algorithm is to choose a matrix B such that the off-diagonal elements of

$$\mathsf{B}\mathsf{A}\mathsf{B}^{-1} \tag{H.77}$$

are smaller than the off-diagonal elements of A. Then choose another, and another. Go through this process again and again until the off-diagonal elements have been ground down to machine zero. There are many strategies for choosing the series of B matrices. These are well-described in any edition of *Numerical Recipes*.[4]

When you need to diagonalize matrices numerically, I urge you to look at *Numerical Recipes* to see what's going on, but I urge you *not* to code these algorithms yourself. These algorithms rely in an essential way on the fact that computer arithmetic is approximate rather than exact, and hence they are quite tricky to implement. Instead of coding the algorithms yourself, I recommend that you use the implementations in either LAPACK[5] (the Linear Algebra PACKage) or EISPACK.[6] These packages are probably the finest computer software ever written, and they are free. They can be obtained through the "Guide to Available Mathematical Software" (GAMS) at http://gams.nist.gov.

H.10 A glance at non-symmetric matrices and the Jordan form

Many of the matrices that arise in applications are symmetric and hence the results of the previous sections are the only ones needed. But every once in a while you do encounter a non-symmetric matrix and this section gives you a guide to treating them. It is just an introduction and treats only 2×2 matrices.

Given a non-symmetric matrix, the first thing to do is rotate the axes to make the matrix representation triangular, as discussed in problem H.12:

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}. \tag{H.78}$$

Note that $b \neq 0$ because otherwise the matrix would be symmetric and we would already be done. In this case vectors on the x-axis are eigenvectors because

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = a \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \tag{H.79}$$

Are there any other eigenvectors? The equation

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \lambda \begin{pmatrix} x \\ y \end{pmatrix} \tag{H.80}$$

tells us that

$$\begin{aligned} ax + by &= \lambda x \\ cy &= \lambda y \end{aligned}$$

whence $\lambda = c$ and the eigenvector has polar angle θ where

$$\tan\theta = \frac{c-a}{b}. \tag{H.81}$$

Note that if $c = a$ (the "degenerate" case: both eigenvalues are the same) then $\theta = 0$ or $\theta = \pi$. In this case *all* of the eigenvectors are on the x-axis.

Diagonal form

We already know that that a rotation of orthogonal (Cartesian) coordinates will not diagonalize this matrix. We must instead transform to a skew coordinate system in which the axes are not perpendicular.

Note that in with oblique axes, the coordinates are given by

$$\mathbf{V} = V_{x'}\hat{\mathbf{i}}' + V_{y'}\hat{\mathbf{j}}' \tag{H.82}$$

but, because $\hat{\mathbf{i}}'$ and $\hat{\mathbf{j}}'$ are not perpendicular, it is *not true* that

$$V_{x'} = \mathbf{V} \cdot \hat{\mathbf{i}}'. \qquad \textbf{NO!} \tag{H.83}$$

A little bit of geometry will convince you that the name of the vector $\mathbf{V}$ changes according to

$$\begin{pmatrix} V_{x'} \\ V_{y'} \end{pmatrix} = \mathsf{B} \begin{pmatrix} V_x \\ V_y \end{pmatrix}, \tag{H.84}$$

where

$$\mathsf{B} = \frac{1}{\sin\varphi} \begin{pmatrix} \sin\varphi & -\cos\varphi \\ 0 & 1 \end{pmatrix}. \tag{H.85}$$

This matrix is *not* orthogonal. In fact its inverse is

$$\mathsf{B}^{-1} = \begin{pmatrix} 1 & \cos\varphi \\ 0 & \sin\varphi \end{pmatrix}. \tag{H.86}$$

Finally, note that we cannot have $\varphi = 0$ or $\varphi = \pi$, because then both $V_{x'}$ and $V_{y'}$ would give information about the horizontal component of the vector, and there would be no information about the vertical component of the vector.

What does this say about the representations of tensors (or, equivalently, of linear transformations)? The "name translation" argument of equation (H.27) still applies, so

$$\mathsf{T}' = \mathsf{B}\mathsf{T}\mathsf{B}^{-1}. \tag{H.87}$$

Using the explicit matrices already given, this says

$$\mathsf{T}' = \frac{1}{\sin\varphi}\begin{pmatrix} \sin\varphi & -\cos\varphi \\ 0 & 1 \end{pmatrix}\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}\begin{pmatrix} 1 & \cos\varphi \\ 0 & \sin\varphi \end{pmatrix} = \begin{pmatrix} a & (a-c)\cos\varphi + b\sin\varphi \\ 0 & c \end{pmatrix}. \tag{H.88}$$

To make this diagonal, we need only choose a skew coordinate system where the angle φ gives

$$(a-c)\cos\varphi + b\sin\varphi = 0, \tag{H.89}$$

that is, one with

$$\tan\varphi = \frac{c-a}{b}. \tag{H.90}$$

Comparison with equation (H.81) shows that this simply means that the skew coordinate system should have its axes pointing along two eigenvectors. We have once again found an intimate connection between diagonal representations and eigenvectors, a connection which is exploited fully in abstract mathematical treatments of matrix diagonalization.

Once again we can do an accounting of information. In the initial coordinate system, the four elements of the matrix contain four independent pieces of information. In the diagonalizing coordinate system, two of those pieces are explicit in the matrix, and two are implicit in the two axis rotation angles needed to implement the diagonalization.

This procedure works almost all the time. But, if $a = c$, then it would involve $\varphi = 0$ or $\varphi = \pi$, and we have already seen that this is not an acceptable change of coordinates.

Degenerate case

Suppose our matrix has equal eigenvalues, $a = c$, so that it reads

$$\begin{pmatrix} a & b \\ 0 & a \end{pmatrix}. \tag{H.91}$$

If $b = 0$, then the matrix is already diagonal. (Indeed, in this case *all* vectors are eigenvectors with eigenvalue a, and the linear transformation is simply multiplication of each vector by a).

But if $b \neq 0$, then, as we have seen, the only eigenvectors are on the x-axis, and it is impossible to make a basis of eigenvectors. Only one thing can be done to make the matrix representation simpler than it stands in equation (H.91), and that is a shift in the scale used to measure the y-axis.

For example, suppose that in the (x, y) coordinate system, the y-axis is calibrated in inches. We wish to switch to the (x', y') system in which the y'-axis is calibrated in feet. There is no change in axis orientation or in the x-axis. It is easy to see that the two sets of coordinates are related through

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1/12 \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 12 \end{pmatrix}\begin{pmatrix} x' \\ y' \end{pmatrix} \tag{H.92}$$

This process is sometimes called a “stretching” or a “scaling” of the y-axis.

The transformation represented by matrix (H.91) in the initial coordinate system is represented in the new coordinate system by

$$\begin{pmatrix} 1 & 0 \\ 0 & 1/12 \end{pmatrix} \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 12 \end{pmatrix} = \begin{pmatrix} a & 12b \\ 0 & a \end{pmatrix}. \tag{H.93}$$

The choice of what to do now is clear. Instead of scaling the y-axis by a factor of 12, we can scale it by a factor of $1/b$, and produce a new matrix representation of the form

$$\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}. \tag{H.94}$$

Where is the information in this case? In the initial coordinate system, the four elements of the matrix contain four independent pieces of information. In the new coordinate system, two of those pieces are explicit in the matrix, one is implicit in the rotation angle needed to implement the initial triangularization, and one is implicit in the y-axis scale transformation.

The Jordan form

Remarkably, the situation discussed above for 2×2 matrices covers all the possible cases for $n \times n$ matrices. That is, in n dimensional space, the proper combination of rotations, skews, and stretches of coordinate axes will bring the matrix representation (the "name") of any tensor or linear transformation into a form where every element is zero except on the diagonal and on the superdiagonal. The elements on the diagonal are eigenvalues, and each element on the superdiagonal is either zero or one: zero if the two adjacent eigenvalues differ, either zero or one if they are the same. The warning of problem H.12 applies here as well: The eigenvalues on the diagonal may well be complex valued, and the same applies for the elements of the new basis vectors.

References

[1]For example, Kenneth Hoffman and Ray Kunze, *Linear Algebra*, second edition (Prentice-Hall, Englewood Cliffs, New Jersey, 1971).

[2]For example, Jerry Marion and Stephen Thorton, *Classical Dynamics of Particles and Systems*, fourth edition (Saunders College Publishing, Fort Worth, Texas, 1995) section 11.2.

[3]For example, Jerry Marion and Stephen Thorton, *Classical Dynamics of Particles and Systems*, fourth edition (Saunders College Publishing, Fort Worth, Texas, 1995) section 11.7.

[4]W.H. Press, S.A. Teukolsky, W.T. Vetterling, B.P. Flannery, *Numerical Recipes* (Cambridge University Press, Cambridge, U.K., 1992).

[5]E. Anderson, *et al.*, *LAPACK Users' Guide* (SIAM, Philadelphia, 1992).

[6]B.T. Smith, *et al.*, *Matrix Eigensystem Routines—EISPACK Guide* (Springer-Verlag, Berlin, 1976).

Appendix I

Catalog of Misconceptions

Effective teaching does not simply teach students what is correct—it also insures that students do not believe what is incorrect. There are a number of prevalent misconceptions in statistical mechanics. For example, an excellent history of statistical mechanics is titled *The Kind of Motion We Call Heat*. This title is wonderfully memorable and historically justifiable, but it embodies no fewer than three different physics misconceptions: heat is confused with thermal energy; thermal energy is confused with kinetic energy; and kinetic energy is confused with motion. Below is a list of misconceptions that are explicitly mentioned in this book, together with the page number where that misconception is pointed out and corrected.

a microstate (a configuration) has an entropy, 32
all gases are ideal, 44
at high temperature, the most probable microstate has high energy, 107
balls-in-buckets picture of quantal states, 145, 158
Boltzmann distribution is for molecules within systems, 99, 104
Bose-Einstein "condensation", 158
canonical probability for a quantal state that doesn't have a definite energy, 113, 128
chemical potential increases with temperature, 46, 152, 153, 155, 168
computers are fast, 199
concerning (anti)symmetric functions, 144
concerning Monte Carlo simulation, 202
concerning multivariate calculus, 57
concerning quantum mechanics, 48
configurational entropy differs from thermal entropy, 52
C_p is $\partial E/\partial T)_p$, 81
definition of partition function, 100
density is uniform at equilibrium, 7, 16, 45, 64
energy must be a function of (S, V, N), 68, 78

ensemble is a collection of atoms, 16
entropy as (moral?) decay, 30
entropy as disorder, 34
entropy doesn't apply to a pendulum, 135
equation of state contains all thermodynamic information, 69, 88
heat is thermal energy, 50, 51
highest temperature implies maximum energy, 135, 136
ideal gas is hard-sphere gas, 21
ideal gas potential energy, 21, 102
in Reif, 48
occupancy probability vs. canonical probability, 149
one molecule is an ensemble element, 99, 104
phase diagram plots pressure as a function of temperature, 7
probability of atom having energy E is proportional to e^{-E/k_BT}, 104
sum over typical microstates, not over all microstates, 23, 100
temperature is energy per particle, 42
the most probable energy is the energy of the most probable microstate, 107
thermal energy is kinetic energy, 50, 51, 107

Appendix J

Thermodynamic Master Equations

$E(S, V, N)$

$$dE = T\,dS - p\,dV + \mu\,dN$$

$F(T, V, N) = E - TS$

$$dF = -S\,dT - p\,dV + \mu\,dN$$

$H(S, p, N) = E + pV$

$$dH = T\,dS + V\,dp + \mu\,dN$$

$G(T, p, N) = F + pV$

$$dG = -S\,dT + V\,dp + \mu\,dN$$

$\Pi(T, V, \mu) = F - \mu N = -pV$

$$d\Pi = -S\,dT - p\,dV - N\,d\mu$$

$p(T, \mu)$ [intensive quantities only]

$$dp = \mathcal{S}\,dT + \rho\,d\mu \qquad [\mathcal{S} = S/V, \quad \rho = N/V]$$

Appendix K

Useful Formulas

Isothermal compressibility:

$$\kappa_T = -\frac{1}{V}\,\frac{\partial V}{\partial p}\Big)_{T,N} = \frac{1}{\rho}\,\frac{\partial \rho}{\partial p}\Big)_T = \frac{1}{\rho^2}\,\frac{\partial^2 p}{\partial \mu^2}\Big)_T$$

Master thermodynamic equation:

$$dF = -S\,dT - p\,dV - M\,dH + \sum_i \mu_i\,dN_i$$

Gibbs-Helmholtz equation:

$$E(T,V,N) = \frac{\partial(F/T)}{\partial(1/T)}\Big)_{V,N} = -\,\frac{\partial \ln Z}{\partial \beta}\Big)_{V,N}$$

Free energy from partition function:

$$F(T,V,N) = -k_B T \ln Z(T,V,N)$$

Classical pure point-particle partition function:

$$Z(T,V,N) = \frac{1}{h^{3N}N!}\int d^{3N}p \int d^{3N}r\, e^{-H(r,p)/k_BT}$$

$$Z(T,V,N) = \frac{1}{\lambda^{3N}(T)}\int d^{3N}r\, e^{-U(r)/k_BT} \quad \text{where} \quad \lambda(T) = \frac{h}{\sqrt{2\pi m k_B T}}$$

Quantal ideal gases (+ for fermions, − for bosons):

$$\Xi(\beta,\mu) = \prod_{r=1}^{M}[1 \pm e^{-\beta(\epsilon_r-\mu)}]^{\pm 1}$$

$$\langle n_r \rangle = \frac{1}{e^{\beta(\epsilon_r-\mu)} \pm 1}$$

Index

adiabatic compressibility, 82, 96
analogy
    booty argument, 27, 49
    correlations and peer pressure, 194
    entropy as disorder, 34
    entropy as freedom, 35
    fluctuation-susceptibility as cows *vs.* goats, 106
    Jim Smith effect, 107
    poker paradox, 34, 201
    sampling algorithm to election poll, 200
    susceptibility and herding, 192
    susceptibility as herd instinct, 106
    temperature and cash incentives, 108, 201
    temperature as persuasion, 108
    typical vs. average, 33
approximation
    controlled, 131
    uncontrolled, 131
asteroids, 16

baum, 230
blackbody radiation, 91
Boltzmann constant, 19
Boltzmann factor, 100
boson, 139

Carnot cycle, 56
cavity radiation, 91
characteristic equation, 243
characteristic matrix, 243
chemical potential, 44–46, 73, 82, 83, 109, 152, 167
Clausius-Clapeyron equation, 83
    magnetic, 85
    second-order, 86
compressibility
    adiabatic, 82, 96
    isothermal, 8, 65, 82–83, 115, 136, 150, 176
conceptual definition, 49
configuration work, 50
controlled approximation, 131
couple, 125
cows
    compared to goats, 106
critical point, 26
crossover temperature, 128

de Broglie wavelength, 124
decouple, 125
definition
    conceptual, 49
    operational, 49
degrees of freedom, 126
densities, 64
density of levels, 151
density of levels in k-space, 151
derivative
    as an experiment, 69
dispersion, 105
dispersion relation, 171
dissipative work, 50

eigenvalues, 239, 241
eigenvectors, 239, 241
energy density of levels, 151
ensemble, 16
ensembles, summary of, 111

enthalpy, 70
entropy, 19–46, 66, 114, 150
    of monatomic ideal gas, 25
equilibrium constant, 90
equipartition theorem, 126
    generalized, 128
escaping tendency, 45
Euler-MacLaurin formula, 133, 223
evolution, 9, 97, 98
expansion and compression, 8
extensive quantities, 64, 76

fermion, 139
flat Earth, 113
fluctuation-susceptibility theorem, 106
freedom, degrees of, 126
freeze out, 128

gamma, 78, 83
Gibbs free energy, 70
Gibbs potential, 70
Gibbs-Duhem equation, 72
Gibbs-Helmholtz equation, 68
goats
    compared to cows, 106
gross national product, 24

heat, 50
heat capacity, 8, 65, 66, 81–82, 126–134, 158
    internal, 126
heat capacity ratio, 78, 83

ideal, 21
ideal gas, 8, 21–27, 46, 66, 74, 78–81, 96, 114, 115, 123–169, 180, 184
ideal paramagnet, 20, 25, 46, 47
identical particles, 18
information theory, 114
intensive quantities, 24, 64, 76
interchange rule, 138, 167
internal heat capacity, 126
internal specific heat, 126
Ising model, 20, 25, 46, 47, 116, 188–213
isothermal compressibility, 8, 82–83, 115, 136, 150, 176

Jim Smith effect, 107

knowledge, 26

law of mass action, 90
Legendre transformation, 60, 68
level (*vs.* state), 140
light, thermodynamics of, 90–95

macroscopic, 11
magnetic systems, 20, 25, 46, 47, 75, 82, 97, 116, 173, 188–213
master equation, 60
master function, 60
mathematical functions, 20, 215–229
mathematical rigor, 38, 109
matrix mathematics, 230–250
microscopic, 11
misconceptions, catalog of, 251
mixtures, 73
monatomic, 21
monkeys, 9
most probable value, 41, 110
mountain climbing, 95

never, 9
number densities, 64

$\mathcal{O}$ notation, 129–132
operational definition, 49
orbital, 140
orthogonal matrix, 232
outer product, 236

parametric differentiation, 105, 117, 148, 196
partial derivative
    as an experiment, 69

partition function, 100
    quantal, 112
perturbation theory, 26
poker, 18
politics, 113
polymer, 118
pressure, 6, 43, 167
problem solving tips
    check dimensions, 19, 23, 64, 132
    check extensivity, 64
    check limiting behavior, 133–135, 149
    check special cases, 176
    debrief, 24, 124
    distinguish variables and functions, 59, 64
    list expectations, 22
    poor strategy, 75
    preview, 22
    use scaled quantities, 185
    write variable list, 59

quantities
    extensive, 64
    intensive, 64

radiation, thermodynamics of, 90–95
random walk, 118
relativistic gas, 128
rotation matrix, 231

Sackur-Tetrode formula, 25, 101
say it, don't think it, 113, 145
second law of thermodynamics, 50
shorthand, 113, 145
slick trick, 105, 117, 118, 127, 148, 151, 196
Smith effect, 107
sound, 96
specific heat, 65
    internal, 126
sphere, volume of, 218
spin, 139
spin systems, 20, 25, 46, 47, 75, 82, 97, 116, 173, 188–213
stars, 153
state (*vs.* level), 140
Stefan-Boltzmann law, 92
sum over all states, 100
susceptibility, 8, 116

temperature, 40, 48
thermal de Broglie wavelength, 45, 124
thermodynamic limit, 24, 103, 106, 107
thoughtless invertibility, principle of, 68
trace, 114
trading in variables, 68
tree, 230
Truesdell, Clifford, 68

uncontrolled approximation, 131
uncoupled, 125
universe
    age of, 9, 199
    size of, 199

virial theorem, 129
volume densities, 64

Wien's law, 93
work
    configuration, 50
    dissipative, 50

Zustandsumme, 100