## Aprendizagem 2023 Homework IV – Group 28

Gonçalo Bárias (ist1103124) & Raquel Braunschweig (ist1102624)

## Part I: Pen and Paper

Given the following observations, 
$$\left\{ \begin{pmatrix} 1\\0.6\\0.1 \end{pmatrix}, \begin{pmatrix} 0\\-0.4\\0.8 \end{pmatrix}, \begin{pmatrix} 0\\0.2\\0.5 \end{pmatrix}, \begin{pmatrix} 1\\0.4\\-0.1 \end{pmatrix} \right\}$$
.

Consider a Bayesian clustering that assumes  $\{y_1\} \perp \{y_2, y_3\}$ , two clusters following a Bernoulli distribution on  $y_1$  ( $p_1$  and  $p_2$ ), a multivariate Gaussian on  $\{y_2, y_3\}$  ( $N_1$  and  $N_2$ ), and the following initial mixture:

$$\pi_1 = 0.5 \quad , \quad \pi_2 = 0.5$$

$$p_1 = P(y_1 = 1) = 0.3 \quad , \quad p_2 = P(y_1 = 1) = 0.7$$

$$N_1 \left( \mu_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \Sigma_1 = \begin{pmatrix} 2 & 0.5 \\ 0.5 & 2 \end{pmatrix} \right) \quad , \quad N_2 \left( \mu_2 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \Sigma_2 = \begin{pmatrix} 1.5 & 1 \\ 1 & 1.5 \end{pmatrix} \right)$$

1. Perform one epoch of the EM clustering algorithm and determine the new parameters. *Hint:* we suggest you to use numpy and scipy, however disclose the intermediary results step by step.

Gonçalo

- 2. Given the new observation,  $x_{new} = \begin{bmatrix} 1 & 0.3 & 0.7 \end{bmatrix}^T$ , determine the cluster memberships (posteriors).
- 3. Performing a hard assignment of observations to clusters under a ML assumption, identify the silhouette of both clusters under a Manhattan distance.

Raquel

Raquel

4. Knowing the purity of the clustering solution is 0.75, identify the number of possible classes (ground truth).

Raquel

## Part II: Programming and critical analysis

Recall the column\_diagnosis.arff dataset from previous homeworks. For the following exercises, normalize the data using sklearn's MinMaxScaler.

1. Using sklearn, apply k-means clustering fully unsupervisedly on the normalized data with  $k \in \{2, 3, 4, 5\}$  (random = 0 and remaining parameters as default). Assess the silhouette and purity of the produced solutions.

Gonçalo

- 2. Consider the application of PCA after the data normalization:
  - (a) Identify the variability explained by the top two principal components.

Raquel

(b) For each one of these two components, sort the input variables by relevance by inspecting the absolute weights of the linear projection.

Raquel

3. Visualize side-by-side the data using: i) the ground diagnoses, and ii) the *previously* learned k=3 clustering solution. To this end, projected the normalized data onto a 2-dimensional data space using PCA and then color observations using the reference and cluster annotations.

Raquel

4. Considering the results from questions (1) and (3), identify two ways on how clustering can be used to characterize the population of ill and healthy individuals.

Raquel