A history criterion for the non-axisymmetric part of a viscous flow

September 27, 2026

Abstract

We give a sufficient regularity criterion when the angular mean of a three-dimensional Navier–Stokes velocity satisfies prescribed anisotropic bounds. The additional hypothesis is finiteness of the critical $L^2_t L^\infty_x$ norm of the angular fluctuation. Its Reynolds stress vanishes in the anisotropic ancient limit without differentiating the fluctuation. A circulation barrier for that limit eliminates the swirl; a cylindrical stretching identity then closes the full-velocity enstrophy estimate using small mean derivatives. We also give a separate quantitative gradient-history estimate and exact finite-energy examples of both signs of kinetic transfer between the mean and the fluctuation. The proof adapts the axisymmetric argument of Constantin, Ignatova, and Vicol. This is an analytic conditional criterion, not a Lean theorem or an arbitrary-data regularity proof.

1 Statement

Let $Q = B(1) \times (-1, 0)$ and let (u, p) be a suitable weak solution of

$$u_t + (u \cdot \nabla)u - \Delta u + \nabla p = 0, \qquad \operatorname{div} u = 0,$$

smooth on compact subsets of Q. Assume the usual local energy class $u \in L^\infty_t L^2_x \cap L^2_t H^1_x$ and $p \in L^{3/2}(Q)$. For rotations Q_ϕ about the fixed third axis put

$$\mathcal{P}u(x) = \frac{1}{2\pi} \int_0^{2\pi} Q_\phi u(Q_\phi^{-1} x)\, \mathrm{d}\phi, \qquad v = \mathcal{P}u, \qquad w = u - v.$$

The same notation denotes the induced action on tensors. Write $r = |x'|$, $z = x_3$, and $\tau = -t$. Unless a smaller ball is specified, suprema are on $B(1)$.

Theorem 1. *Suppose $0 < h < 1/2$ and, for every $a + b \le 2$,*

$$|\partial_r^a \partial_z^b v_r| \le C \tau^{-1/2 - a/2 - (1/2-h)b}, \tag{1}$$

$$|\partial_r^a \partial_z^b v_z| + |\partial_r^a \partial_z^b v_\theta| \le C \tau^{-1/2 - h - a/2 - (1/2-h)b}. \tag{2}$$

Set $K(t) = \|w(t)\|_\infty$. If

$$\int_{-1}^0 K(t)^2 \, \mathrm{d}t < \infty, \tag{3}$$

then $(0, 0)$ is a regular point of u.

The mean bounds are those used in [1]. Their validity and (3) are hypotheses, not consequences of the energy inequality. The axis and the ball are fixed; the theorem makes no assertion about propagating approximate symmetry from a shrinking core. No derivative or torque integral of w is assumed. The proof uses the full second-order swirl bounds in (2); it does not extend the meridional-only variant of [1, Remark 5.2] to this setting.

Corollary 2. *In Theorem 1, either of the following bounds near zero suffices in place of (3):*

$$K(t) \le C\tau^{-\beta}, \quad 0 \le \beta < \tfrac{1}{2}; \qquad K(t) \le C\tau^{-1/2}\log(e/\tau)^{-p}, \quad p > \tfrac{1}{2}.$$

These implications follow by integrating K^2 and restricting to a final cylinder. In particular no gradient estimate is inferred from an amplitude bound. A stronger weighted amplitude condition does pay a gradient history in the whole-space setting; that independent estimate is in Section 6.

2 The mean equation and its rescaled stress

Rotation covariance and $\mathcal{P}w = 0$ give the exact equation

$$v_t + (v\cdot\nabla)v - \Delta v + \nabla\mathcal{P}p = F, \qquad F = -\mathcal{P}[(w\cdot\nabla)w] = -\operatorname{div}\mathcal{P}(w\otimes w). \tag{4}$$

The proof retains this divergence form and does not require an L^∞ bound on F.

The original solution has a regular spherical annulus inside any prescribed interval of radii. Indeed, partial regularity makes the top-slice singular set have zero $\mathcal{H}^1$ measure. Its image under $x \mapsto |x|$ has measure zero, so some intervening sphere is regular. Compactness and interior estimates give an annulus on which u, ∇u, and $\nabla^2 u$ are bounded through the endpoint, as in the first part of [1, Lemma 3.2]. Its rotational invariance gives the same bounds for v and w. This argument uses suitability of u; it does not require a partial-regularity theorem for (4).

Write the meridional entries of $\mathcal{P}(w\otimes w)$ in the cylindrical frame as $a = R_{rr}$, $b = R_{\theta\theta}$, $c = R_{zz}$, and $d = R_{rz}$. Their absolute values are at most K^2. Direct differentiation yields

$$F_r = -a_r - (a-b)/r - d_z, \qquad F_z = -d_r - d/r - c_z, \tag{5}$$

$$\frac{(\operatorname{curl} F)_\theta}{r} = \left(\partial_{rr} + \frac{3}{r}\partial_r - \partial_{zz}\right)\frac{d}{r} + \frac{1}{r}\partial_{rz}(c-a) - \frac{1}{r^2}\partial_z(a-b). \tag{6}$$

The quantities here are smooth at the axis in the appropriate lifted variables. For a distributional estimate it suffices that $d = O(r)$.

Consider the rescaling used in [1, Section 4]: for $t_n \uparrow 0$,

$$\tau_n = -t_n, \quad \lambda_n = \tau_n^{1/2}, \quad \delta_n = \tau_n^h, \quad \mu_n = \lambda_n/\delta_n, \quad r = \lambda_n R, \quad z = z_n + \mu_n Z, \quad t = \tau_n s.$$

The mean components are normalized as $V_n = \lambda_n v_r$, $W_n = \lambda_n\delta_n v_z$, $S_n = \lambda_n\delta_n v_\theta$, and the potential vorticity as $\Omega_n = \lambda_n^3\delta_n(\operatorname{curl} v)_\theta/r$. For a stress component a, let a_n denote its composition with this change of variables, without an amplitude normalization.

Lift $R = |X|$ to four radial dimensions. Since

$$\frac{1}{R}\partial_R H = \operatorname{div}_X\left(\frac{XH}{R^2}\right) - \frac{2H}{R^2},$$

the rescaled forcing in the potential-vorticity equation is

$$\begin{aligned} G_n = \lambda_n^2\delta_n\Big[&(\Delta_X - \delta_n^2\partial_{ZZ})(d_n/R) \\ &+ \delta_n\partial_Z\left\{\operatorname{div}_X\frac{X(c_n-a_n)}{R^2} - \frac{2(c_n-a_n)+(a_n-b_n)}{R^2}\right\}\Big]. \end{aligned} \tag{7}$$

Both R^{-1} and R^{-2} are locally integrable in $\mathbb{R}^4$. The divergence identity has no axis-supported mass: the flux through $R = \epsilon$ is $O(\epsilon^2)$ for bounded H. Move all derivatives in (7) onto a compactly

supported smooth test function with time support $J \Subset (-\infty, 0)$. The resulting absolute value is bounded by

$$C_J \lambda_n^2 \delta_n \int_J K(\tau_n s)^2 \, \mathrm{d}s = C_J \delta_n \int_{\tau_n J} K(t)^2 \, \mathrm{d}t \longrightarrow 0. \tag{8}$$

Thus no derivative of the fluctuation is needed in this part of the limiting argument. In a recentered zoom with $r = \lambda_n(A_n + R)$ and $A_n \to \infty$, the azimuthal-vorticity force has the same prefactor $\lambda_n^2 \delta_n$ multiplying at most two derivatives of the stress; its additional coefficients $(A_n + R)^{-1}$ and $(A_n + R)^{-2}$ are uniformly bounded on compact sets. Estimate (8) therefore applies there too.

3 Eliminating the ancient swirl

Lemma 3. *Let S on $[0, \infty) \times \mathbb{R} \times (-\infty, -1]$ have uniform spatial $W^{2,\infty}$ bounds on compact time intervals, be continuous through the upper time, and satisfy $S(0, Z, s) = 0$. The drifts V, W may be only measurable in time, but are uniformly bounded and spatially Lipschitz on each such interval. Suppose, for $R > 0$,*

$$S_s + V S_R + W S_Z + (V/R) S = S_{RR} + R^{-1} S_R - R^{-2} S \tag{9}$$

in distributions. If $h > 0$ and

$$|S(R, Z, s)| \le C(-s)^{-1/2-h}, \qquad |V(R, Z, s)| \le C_r(-s)^{-1/2},$$

then $S = 0$.

Proof. Put $\Gamma = RS$. Its equation on $R > 0$ is

$$\mathcal{L}\Gamma := \Gamma_s + V\Gamma_R + W\Gamma_Z - \Gamma_{RR} + R^{-1}\Gamma_R = 0.$$

For $L > 1$ let $B_L = CL^{-1/2-h}$ and

$$\Phi_L(R, s) = B_L R + C_r B_L \int_{-L}^{s} (-q)^{-1/2} \, \mathrm{d}q.$$

It dominates $|\Gamma|$ at $s = -L$, and

$$\mathcal{L}\Phi_L = B_L\big[C_r(-s)^{-1/2} + V + R^{-1}\big] \ge 0.$$

Comparison is valid on the unbounded spatial domain. On a fixed slab $[-L, s_1]$, perturb Φ_L by

$$\eta e^{k(s+L)}(1 + R^2 + Z^2).$$

The radial diffusion of R^2 is zero for this operator, so k can be chosen in terms of the bounded drifts to make the added function a supersolution. Its quadratic growth dominates the linear growth of Γ at infinity. Near the axis, $\Gamma = O(R^2)$ uniformly on this slab, while $\Phi_L \ge B_L R$. Thus one can excise a sufficiently small axis cylinder, on whose boundary comparison already holds. On the remaining bounded region the positive-part energy argument applies: spatial Lipschitz bounds control the divergence of the drift, and no diffusion in Z is needed. The distributional equation and spatial second-derivative bounds give local bounded time derivatives, so the time traces used in this argument are legitimate. This also justifies the argument for time-measurable drifts, as in [1, Lemma 3.3]. Let $\eta \downarrow 0$ and apply the comparison to both signs of Γ. For a fixed (R, Z, s) it yields

$$|\Gamma(R, Z, s)| \le CRL^{-1/2-h} + 2CC_r L^{-h} \longrightarrow 0 \qquad (L \longrightarrow \infty).$$

The upper time is included by continuity. □

To apply the lemma, first take a limit of the actual mean swirl equation. Write $e = R_{r\theta}$ and $f = R_{z\theta}$ for the corresponding entries of $\mathcal{P}(w \otimes w)$. They obey $|e| + |f| \le 2K^2$, and

$$F_\theta = -\partial_r e - 2e/r - \partial_z f.$$

In the preceding rescaling the swirl equation is

$$\begin{aligned}(S_n)_s + V_n(S_n)_R + W_n(S_n)_Z + (V_n/R)S_n = (S_n)_{RR} + R^{-1}(S_n)_R - R^{-2}S_n + \delta_n^2(S_n)_{ZZ} + H_n,\\ H_n = -\lambda_n^2\delta_n[\partial_R e_n + 2e_n/R + \delta_n\partial_Z f_n].\end{aligned} \tag{10}$$

For a rectangle $D \Subset \{R > 0\}$, a closed time slab $J = [-T, -1]$, and finite p, this gives

$$\|H_n\|_{L^1(J;W^{-1,p}(D))} \le C_{D,J}\delta_n \int_{\tau_n J} K(t)^2 \, \mathrm{d}t \longrightarrow 0. \tag{11}$$

The mean hypotheses uniformly bound V_n, W_n, S_n in spatial C^2 on this slab. All other terms of (10) are bounded in a negative Sobolev norm. Thus the time increments of S_n in that norm are bounded by $C|s - s'| + \epsilon_n$, where $\epsilon_n \to 0$. Interpolation with the spatial bounds gives asymptotic equicontinuity in C^0; spatial interpolation then gives locally uniform convergence of the first spatial derivatives too. Finitely many initial members are smooth, so this yields the usual compactness after a subsequence. The selected time $s = -1$ is included. The bounds $S_n(0, Z, s) = 0$ and $\|\partial_R S_n\|_\infty \le C_J$ extend the convergence to the axis. Also $\partial_Z S_n(0, Z, s) = 0$ and the mixed second derivative is uniformly bounded.

The drifts have weak-* limits, with the same global spatial Lipschitz bounds on each compact time slab. The strong convergence of S_n and its first derivatives permits passage to their products with these drifts. Equations (10)–(11) give (9) in the limit. The backward bounds are

$$|S(R, Z, s)| \le C|s|^{-1/2-h}, \qquad |V(R, Z, s)| \le C|s|^{-1/2}.$$

Their constants are independent of the compact spatial set. Lemma 3 therefore makes this limit zero.

For a recentered zoom with $r = \lambda_n(A_n + R)$ and $A_n \to \infty$, replace every curvature factor $1/R$ in (10) by $1/(A_n + R)$. The force estimate still holds on each fixed rectangle, and the limiting equation on $\mathbb{R}^2$ is simply

$$S_s + VS_R + WS_Z = S_{RR}.$$

Its bounded solution tends uniformly to zero in the remote past, so [1, Lemma 3.3] makes it zero as well.

In particular, on every smaller original ball,

$$(-t)\,\|\partial_z v_\theta(t)\|_\infty \longrightarrow 0. \tag{12}$$

Otherwise select points and times where this fails and use the same finite-axis or receding-axis alternatives. At the selected point $\partial_Z S_n = \lambda_n^2 \partial_z v_\theta$ tends to zero. If its scaled radial coordinate tends to zero, the conclusion instead follows directly from $\partial_Z S_n(0, Z, s) = 0$ and the mixed second-derivative bound. This proves (12).

3.1 A necessary stress input on late time intervals

The same compactness argument also gives a requirement for sustaining a nonzero rescaled mean swirl, without assuming (3). It concerns the actual radial and axial fluxes of angular momentum.

Proposition 4. *Fix $0 < b < 1$, $\kappa > 0$, $0 < h < 1/2$, and constants $C, F_0 \geq 0$. Consider smooth divergence-free solutions on $B(1) \times (-1, 0)$ whose mean satisfies (1)–(2) with constant C, allowing an original force of supremum at most F_0. Let $e = \mathcal{P}(w_r w_\theta)$, $f = \mathcal{P}(w_z w_\theta)$ and define*

$$\mathcal{B}_\tau(T) = \tau^h \int_{-T\tau}^{-\tau} \left(\|e(t)\|_\infty + \tau^h \|f(t)\|_\infty \right) \mathrm{d}t.$$

There are $T > 1$, $\varepsilon > 0$, and $0 < \tau_0 < 1/T$, depending only on the listed constants, such that, for every such solution and $0 < \tau < \tau_0$,

$$\tau^{1/2+h} \|v_\theta(-\tau)\|_{L^\infty(B(b))} \geq \kappa \quad \Longrightarrow \quad \mathcal{B}_\tau(T) \geq \varepsilon. \tag{13}$$

The stress norms are over $B(1)$. Infinite integrals satisfy the conclusion automatically. No local energy or suitability hypothesis is needed for this swirl statement.

Proof. If no such constants existed, choose solutions and scales with $T_n = n$, $\tau_n < n^{-2}$, and $\mathcal{B}_{\tau_n}(n) < 1/n$, while the left side of (13) is at least κ. Choose spatial points realizing at least half this supremum. On every fixed rescaled time slab the Reynolds source in (10) then tends to zero in $L^1_t W^{-1,p}_x$ away from the scaled axis. The original force contributes at most $\lambda_n^3 \delta_n F_0$, which also vanishes. The uniform mean bounds give precisely the spatial compactness, time increment bounds, and weak-* drift limits used above.

If the selected radii divided by λ_n stay bounded, the limit is the ancient swirl of Lemma 3, hence zero. A selected scaled radius tending to zero is already excluded by $S_n(0, Z, s) = 0$ and the uniform radial derivative bound. If those radii tend to infinity, recenter there; the limiting bounded scalar equation is the receding-axis equation above, whose ancient solution is also zero. In either case convergence at $s = -1$ contradicts the selected value of size at least $\kappa/2$. The fixed distance of $B(b)$ from the boundary permits every compact rescaled spatial set in these limits. □

Since $|e|, |f| \leq K^2$, persistent nonzero rescaled mean swirl therefore requires, on every sufficiently late interval of this fixed ratio,

$$\int_{-T\tau}^{-\tau} K(t)^2 \, \mathrm{d}t \geq \frac{\varepsilon}{2} \tau^{-h}. \tag{14}$$

Its root-mean-square defect amplitude on that interval is at least a constant times $\tau^{-1/2-h/2}$. This is an interval estimate; it permits intermittency and gives no pointwise lower bound at every time.

The signed quantity preceding this norm estimate has a useful exact form. Write $\Gamma_n = RS_n$ and let ψ be a smooth test function with compact spatial support away from $R = 0$. Set

$$\mathcal{A}_n^* \psi = \psi_{RR} + \delta_n^2 \psi_{ZZ} + \partial_R((V_n + R^{-1})\psi) + \partial_Z(W_n \psi).$$

For $s_0 < s_1$, with spatial integrals in $\mathrm{d}R\,\mathrm{d}Z$, the circulation equation gives

$$\left[\int \Gamma_n \psi \right]_{s_0}^{s_1} - \int_{s_0}^{s_1} \int \Gamma_n (\psi_s + \mathcal{A}_n^* \psi) = \lambda_n^2 \delta_n \int_{s_0}^{s_1} \int \left[e_n (R\psi_R - \psi) + \delta_n f_n R \psi_Z \right]. \tag{15}$$

Here the original force is zero; otherwise its circulation work is added on the right. Both sides retain the signs of the actual stress. The identity follows by integrating $-R\partial_R e_n - 2e_n - \delta_n R \partial_Z f_n$ against ψ. For a backward adjoint solution the bulk mean term would cancel; (15) itself needs no such kernel construction. Neither this identity nor (14) estimates the recurrent work from initial data.

4 Meridional smallness

Define, on a smaller ball,

$$G_v(t) = \| |\partial_r v_r| + |v_r/r| + |\partial_z v_z| + |\partial_z v_r| \|_\infty .$$

We verify the two changes needed in the proof of [1, Proposition 1.5] to conclude

$$(-t) G_v(t) \longrightarrow 0. \tag{16}$$

First, the preceding swirl argument gives $S_n \to 0$ on compact rescaled time slabs. The mean bounds also give $|S_n/R| \le C$ on those slabs, including the axis. Splitting at $R = \epsilon$ shows that S_n^2/R^2 tends to zero locally in L^1 in the lifted space. In the receding-axis case the source is $\partial_Z(S_n^2/(A_n + R))$, which likewise tends to zero in distributions.

Second, the force is small in a negative Sobolev norm integrated in time, rather than uniformly small as a function. Away from the axis, (7) and (8) give

$$\|G_n\|_{L^1(J;W^{-2,p}(D))} \longrightarrow 0$$

for a fixed rectangle D and any finite p. The analogous assertion holds in the receding-axis case. The other terms in the rescaled vorticity equation are uniformly bounded in this negative norm. Together with the uniform spatial Lipschitz bound supplied by (1)–(2), this gives an increment bound $C|s - s'| + \epsilon_n$ with $\epsilon_n \to 0$. Interpolation gives asymptotic uniform time equicontinuity, also at the selected upper time. Finitely many initial members are smooth, so Arzelà–Ascoli still applies. Boundedness and the small volume of $R < \epsilon$ give convergence across the axis.

The remaining ancient-limit argument is unchanged: the force and swirl source vanish, and the mean bounds supply the backward decay and bounded Lipschitz drift required by [1, Lemma 3.3]. The limiting vorticity vanishes. At the selected time, incompressibility and boundedness of the limiting meridional velocity then contradict a nonzero selected derivative, in either axis case. This proves (16). No bound on $\partial_r v_z$ is asserted.

5 Full-velocity enstrophy

Choose a radial cutoff χ whose derivative is supported in a regular annulus, and write

$$Y = \|\chi \operatorname{curl} u\|_2^2 , \quad M = \|\chi \nabla \operatorname{curl} u\|_2^2 , \quad A_v = \|\partial_z v_\theta\|_\infty .$$

Here A_v and G_v are taken on a ball containing the support of χ, with its boundary in the regular annulus; K is still taken on $B(1)$. All cutoff errors are uniformly bounded. Weighted L^2 contraction of angular averaging, including its tensor action on gradients, gives $Y_v \le Y$ and $M_v \le M$. Local div–curl estimates consequently bound the first derivatives of u, v by $C(Y+1)^{1/2}$ and their second derivatives by $C(M+1)^{1/2}$ in these weighted norms.

For $\Pi_\chi(q) = \int \chi^2 (\operatorname{curl} q \cdot \nabla) q \cdot \operatorname{curl} q$, put $\omega_q = \operatorname{curl} q$. There is an exact telescoping identity

$$\begin{aligned} \Pi_\chi(u) - \Pi_\chi(v) = \int \chi^2 \big[&(\omega_w \cdot \nabla) u \cdot \omega_u + (\omega_v \cdot \nabla) w \cdot \omega_u \\ &+ (\omega_v \cdot \nabla) v \cdot \omega_w \big]. \end{aligned} \tag{17}$$

Integrate the derivative on w off that factor in each term. The remaining factors have derivative orders one and two, apart from bounded cutoff errors. Hence, for every $\epsilon > 0$,

$$\begin{aligned} |\Pi_\chi(u) - \Pi_\chi(v)| &\le CK(Y+1)^{1/2}(M+1)^{1/2} + C \\ &\le \epsilon M + C_\epsilon (K^2+1)(Y+1). \end{aligned}$$

For the mean's stretching, write $a = (\omega_v)_r = -\partial_z v_\theta$, $b = (\omega_v)_\theta = \partial_z v_r - \partial_r v_z$, and $c = (\omega_v)_z = \partial_r v_\theta + v_\theta/r$. The cylindrical gradient identity underlying [1, Equation (5.7)] can be rewritten exactly as

$$(\omega_v \cdot \nabla) v \cdot \omega_v = a^2 \partial_r v_r + b^2 v_r/r + c^2 \partial_z v_z + 2ac\partial_z v_r - abc - 2ab\, v_\theta/r. \tag{18}$$

In particular the large derivative $\partial_r v_z$ has been replaced using the vorticity relation, and the last two terms have the coefficient a, whose supremum is A_v.

A local radial integration by parts gives

$$\|\chi v_\theta/r\|_2^2 \le \|\chi(\omega_v)_z\|_2^2 + C. \tag{19}$$

Indeed, after expanding $(\partial_r v_\theta + v_\theta/r)^2 r$, its cross term is $\partial_r(v_\theta^2)$. The axis boundary term vanishes because $v_\theta = O(r)$; the cutoff error is $2\int \chi \partial_r \chi\, v_\theta^2 \,\mathrm{d}r\,\mathrm{d}z\,\mathrm{d}\theta$. It is bounded on the regular annulus, including its intersection with the axis, where $\partial_r \chi/r$ is bounded. Applying Cauchy–Schwarz to the last two terms of (18), and using (19), yields

$$\Pi_\chi(v) \le C(G_v + A_v)(Y_v + 1).$$

Testing the original unforced vorticity equation and absorbing the small dissipation term from the full-velocity difference now gives

$$Y' + cM \le C_*[G_v + A_v + K^2 + 1](Y + 1). \tag{20}$$

Fix $\eta > 0$. Equations (16) and (12) allow a final interval on which $G_v + A_v \le \eta/\tau$. Since $K \in L_t^2$, Gronwall in (20) gives

$$Y(t) + 1 \le C_\eta(-t)^{-C_*\eta}.$$

Here C_* was fixed before η. Choose η with $2C_*\eta < 1$. Local Sobolev now places u in $L_t^4 L_x^6$ on a smaller cylinder. The local criterion of [2, Theorem 1.1(i)], applied on backward cylinders ending at zero, makes $(0,0)$ regular. This finishes the proof of Theorem 1.

6 A separate quantitative gradient-history estimate

Regularity in Theorem 1 requires only (3). For applications which additionally require an absolute gradient or torque history, the actual defect equation gives the following stronger estimate. The whole-space assumptions here control the pressure produced by localization; they are not assumptions of Theorem 1.

Theorem 5. *Let u be a smooth unforced solution on $\mathbb{R}^3 \times (-1,0)$ with $u \in L^\infty((-1,0); L^2(\mathbb{R}^3))$ and canonical pressure, whose restriction to Q satisfies the local suitability assumptions above. For some $0 < h < 1/2$, assume only the following mean bounds on $B(1)$:*

$$\|v(-\tau)\|_\infty \le C_M \tau^{-1/2-h}, \qquad \|\nabla v(-\tau)\|_\infty \le C_N \tau^{-1-h}, \tag{21}$$

with Cartesian derivatives. Suppose, for some $0 < a < 1$,

$$\int_0^a \tau^{-h/2}\sqrt{\log(e/\tau)}\, \|w(-\tau)\|_{L^\infty(B(1))}^2 \,\mathrm{d}\tau < \infty. \tag{22}$$

Then, on a smaller ball and final time interval,

$$\int \sqrt{-t}\, \|w(t)\|_\infty \|\nabla w(t)\|_\infty \,\mathrm{d}t < \infty. \tag{23}$$

It also bounds the same weighted integral of $\|(\mathcal{P}[(w\cdot\nabla)w])_\theta\|_\infty$. *For this additional conclusion, the amplitude power* $\|w(-\tau)\|_\infty \le C\tau^{-\beta}$ *suffices when* $\beta < 1/2 - h/4$. *At equality, a factor* $\log(e/\tau)^{-p}$ *suffices for* $p > 3/4$. *If the full mean geometry of Theorem 1 also holds, regularity already follows from the weaker condition (3).*

The theorem does not assert (22) from initial data. Its proof has a whole-space smoothing estimate and a localization step. On a sufficiently short final interval it bounds (23) by a constant times the integral in (22), plus a bounded annular-source contribution. Write $\mathbb{L}$ for the Leray projection, to distinguish it from angular averaging $\mathcal{P}$.

Lemma 6. *Let a smooth divergence-free* U *on* $\mathbb{R}^3 \times [0,d]$ *satisfy*

$$U_t - \Delta U = -\mathbb{L}\,\mathrm{div}(U\otimes U) + g, \qquad V = \mathcal{P}U, \quad W = U - V, \quad G = (1-\mathcal{P})g.$$

Assume the fields and the spatial derivatives used below are bounded on compact time prefixes, and that $\|V\|_\infty \le M$, $\|\nabla V\|_\infty \le N$, $\|G\|_\infty \le G_0$ *there, with* $M, N > 0$. *Put* $K(q) = \|W(q)\|_\infty$, $L(q) = \|\nabla W(q)\|_\infty$, *and*

$$Z_d = \sup_{0<q\le d}\{K(q) + \sqrt{q}L(q)\}, \qquad \log_+ x = \max(\log x, 0).$$

There are universal constants such that

$$Z_d \le C_0(K(0) + dG_0) + C_1 N d\left[1 + \log_+ \frac{M}{N\sqrt{d}}\right] Z_d + C_2\sqrt{d}Z_d^2. \tag{24}$$

Consequently $Z_d \le 2C_0(K(0)+dG_0)$ *if the linear coefficient is at most* $1/4$ *and* $\sqrt{d}(K(0)+dG_0)$ *is sufficiently small.*

Proof. Rotation covariance gives the exact homogeneous-in-W equation

$$W_t - \Delta W = -\mathbb{L}\,\mathrm{div}\left[V\otimes W + W\otimes V + (1-\mathcal{P})(W\otimes W)\right] + G. \tag{25}$$

In particular the complete defect self-interaction is retained. For the heat kernel H_s, the kernel of $e^{s\Delta}\mathbb{L}\,\mathrm{div}$ is

$$(B_s)_{ijk} = \delta_{ij}\partial_k H_s + \int_s^\infty \partial_{ijk} H_b \,\mathrm{d}b.$$

Differentiating the Gaussian and integrating in b gives

$$|B_s(y)| \le C(\sqrt{s}+|y|)^{-4}, \qquad |\nabla B_s(y)| \le C(\sqrt{s}+|y|)^{-5},$$
$$\|B_s\|_1 \le Cs^{-1/2}, \qquad \int |y|\,|\nabla B_s(y)|\,\mathrm{d}y \le Cs^{-1/2}.$$

The first absolute moment of B_s is not integrable at infinity. Splitting the radial integral at $\sqrt{s}$ and M/N instead gives

$$\int |B_s(y)|\min(N|y|, 2M)\,\mathrm{d}y \le CN\left[1 + \log_+\frac{M}{N\sqrt{s}}\right], \tag{26}$$

$$\int |\nabla B_s(y)|\min(N|y|, 2M)\,\mathrm{d}y \le CNs^{-1/2}. \tag{27}$$

For each endpoint (x,q) follow the backward characteristic $z'(s) = V(z(s),s)$, $z(q) = x$, and set $a(s) = V(z(s),s)$. Move $a(s)\cdot\nabla W$ to the left side of (25); its two linear tensors now contain $V - a$

instead of V, since $\operatorname{div} W = 0$. This identity holds everywhere with this one spatially constant $a(s)$. The translated Duhamel kernels at the chosen endpoint are centered at $z(s)$, where

$$|V(y,s) - a(s)| \le \min(N|y - z(s)|, 2M).$$

For the gradient estimate, differentiate that representation with $a(s)$ held fixed. No derivative of the selected characteristic is introduced: the same representation is valid at every nearby endpoint with this fixed a. Estimates (26)–(27) bound the linear amplitude and gradient terms.

For the quadratic gradient term move the derivative onto the tensor: $\|\nabla(1-\mathcal{P})(W\otimes W)\|_\infty \le CKL$. Angular averaging and its tensor action are contractions in the Euclidean sup norms. The time integral $\int_0^q (q-s)^{-1/2} s^{-1/2}\,\mathrm{d}s = \pi$ then bounds this term by CZ_d^2, before multiplication by $\sqrt{q}$. The quadratic amplitude term is bounded by $C\sqrt{q}Z_d^2$. The initial term and the divergence-free source use the ordinary translated heat kernel. Finally

$$\int_0^q \left[1 + \log_+ \frac{M}{N\sqrt{s}}\right] \mathrm{d}s \le Cq \left[1 + \log_+ \frac{M}{N\sqrt{q}}\right].$$

This proves (24). The estimate holds on every shorter prefix, whose linear coefficient is no larger. A continuity bootstrap gives the final claim; smoothness supplies the initial continuity. If $K(0) + dG_0 = 0$, use uniqueness or a positive bootstrap bound tending to zero. □

Proof of Theorem 5. First suppose a globally bounded defect equation as in Lemma 6 has been obtained, with G uniformly bounded and

$$\|V(-\tau)\|_\infty \le C_M \tau^{-1/2-h}, \qquad \|\nabla V(-\tau)\|_\infty \le C_N \tau^{-1-h}. \tag{28}$$

Let $\ell(\tau) = \log(e/\tau)$ and choose

$$d(\tau) = \frac{c\tau^{1+h}}{\ell(\tau)}, \qquad 0 < c < 1. \tag{29}$$

On $[-\tau - d, -\tau]$, the linear coefficient in (24) is at most $Cc(1 + |\log c| + h)$: here $Nd \le C_N c/\ell$ and $M/(N\sqrt{d})$ is bounded by a constant times $c^{-1/2}\tau^{-h/2}\sqrt{\ell}$. Take c small enough in terms of the mean bounds.

Write $K = \|W\|_\infty$, $L = \|\nabla W\|_\infty$, and

$$E(\tau) = \int_\tau^{\tau + d(\tau)} K(-s)^2\,\mathrm{d}s.$$

Condition (22), with the localized global norm, implies $K \in L_t^2$ and $E(\tau) \to 0$. There is a time $s_0 \in [-\tau - d, -\tau - d/2]$ at which $K(s_0)^2 \le 2E(\tau)/d$. Applying Lemma 6 on $[s_0, -\tau]$ is therefore legitimate for all sufficiently small τ; its small parameter is at most $C\sqrt{E(\tau)} + d^{3/2}G_0 \to 0$. We obtain

$$K(-\tau) + \sqrt{d}L(-\tau) \le C[d^{-1/2}\sqrt{E(\tau)} + dG_0]. \tag{30}$$

Consequently

$$\sqrt{\tau}K(-\tau)L(-\tau) \le C\sqrt{\tau}d^{-3/2}E(\tau) + C\sqrt{\tau}d^{3/2}G_0^2. \tag{31}$$

The second term is integrable. For the first, Tonelli gives

$$\int_0^{a/2} \sqrt{\tau} d(\tau)^{-3/2} E(\tau)\,\mathrm{d}\tau \le C \int_0^a s^{-h/2}\sqrt{\ell(s)}K(-s)^2\,\mathrm{d}s. \tag{32}$$

Indeed, $\tau \le s \le \tau + d(\tau)$ implies $s/(1+c) \le \tau \le s$. Since d increases, these τ lie in an interval of length at most $d(s)$, while the integrand's coefficient is at most $Cs^{-1-3h/2}\ell(s)^{3/2}$. This proves (32), hence the gradient history on a final interval. The angularly averaged torque is at most KL.

It remains to construct the localization used above. Choose a radial cutoff χ, equal to one on an interior ball and with its transition inside a regular annulus of the original solution. Put

$$\Psi = (-\Delta)^{-1} \operatorname{curl} u, \qquad U = \operatorname{curl}(\chi\Psi) = \chi u + b, \qquad b = \nabla\chi \times \Psi.$$

The L^2 hypothesis and incompressibility give $\operatorname{curl}\Psi = u$; the potential is well defined by its order-minus-one kernel. On the annulus, its separated kernel and every required derivative are bounded against $\|u\|_2$, while its local part is controlled by interior spatial regularity. For canonical pressure, the same decomposition for its order-zero transform of $u \otimes u$ bounds p and its spatial derivatives on the annulus. Likewise

$$\Psi_t = -\operatorname{curl} u - (-\Delta)^{-1} \operatorname{curl}\operatorname{div}(u \otimes u)$$

has bounded spatial derivatives there: its bilinear operator is of order zero, with separated kernels controlled by $\|u\|_2^2$. These statements use only annular regularity and the uniform energy bound, not regularity of the interior through time zero.

With pressure χp, the exact force of U is

$$\begin{aligned} f_U = b_t - 2\nabla\chi\cdot\nabla u - (\Delta\chi)u - \Delta b + p\nabla\chi \\ + (U\cdot\nabla)U - \chi(u\cdot\nabla)u. \end{aligned} \tag{33}$$

It is supported in the regular annulus and uniformly bounded in spatial C^1. Its Leray projection is uniformly bounded, as follows by subtracting $f_U(x)$ in the local principal value and using its L^1 norm for the far part. Thus the equation for U has $g = \mathbb{L}f_U$ and bounded $G = (1-\mathcal{P})g$, including pressure.

Rotations preserve the interior ball, so $U = u$, $\mathcal{P}U = v$, and $U - \mathcal{P}U = w$ there. The transition region contributes bounded fields and derivatives; outside the cutoff $U = 0$. The mean assumptions therefore imply (28) globally. The global weighted amplitude integral is bounded by (22) plus the finite contribution of a bounded function. Equations (30)–(32) now supply (23) on a smaller cylinder of the original flow. The power and logarithmic consequences follow by integrating $\tau^{-h/2-2\beta}\ell(\tau)^{1/2-2p}$. □

7 The sign of kinetic mean–defect transfer

For a smooth finite-energy whole-space unforced flow, orthogonality of angular averaging and integration by parts give

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\|w\|_2^2 + \nu\|\nabla w\|_2^2 = -T, \qquad \frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\|v\|_2^2 + \nu\|\nabla v\|_2^2 = T, \quad T = \int w\cdot(w\cdot\nabla)v. \tag{34}$$

We display viscosity explicitly in this section. Defect self-advection and pressure do no kinetic work on w; the averaging correction is orthogonal to w. In particular the signed net kinetic transfer obeys

$$-\int_0^t T(s)\,\mathrm{d}s = \tfrac{1}{2}\|v(0)\|_2^2 - \tfrac{1}{2}\|v(t)\|_2^2 - \nu\int_0^t \|\nabla v\|_2^2\,\mathrm{d}s \le \tfrac{1}{2}\|v(0)\|_2^2 .$$

This does not bound its positive variation or the critical amplitude integral (3). The following example also excludes an automatic viscous damping argument based just on small relative angular defect.

Proposition 7. *For every $\nu > 0$, $C > 0$, and $0 < \epsilon \le 1$, there are real Schwartz solenoidal data u_+ and u_- with $\|u_\pm\|_2 = 1$ and the same angular mean v. Their defects $w_\pm = u_\pm - v$ have equal squared L^2 and gradient norms, and $\|w_\pm\|_2 = (3\epsilon/2)\,\|v\|_2$. Their nonlinear kinetic transfers have opposite signs, and the unforced solution from u_+ satisfies at its initial time*

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\,\|w_+(t)\|_2^2 > C\nu\,\|\nabla w_+\|_2^2\,.$$

The mean satisfies the spatial bounds (1)–(2) at a time $t = -\tau_\lambda$, with $h = 1/6$ and a constant independent of the concentration parameter λ and of $\epsilon \in (0, 1]$. This last assertion concerns that time slice, not an entire evolution.

Proof. Set $\rho = x^2 + y^2 + z^2$, $r^2 = x^2 + y^2$, and, for $\sigma = \pm 1$,

$$V = \operatorname{curl}(e^{-\rho} e_z), \qquad W_\sigma = \operatorname{curl}\left((x^2 - y^2 + 2\sigma x y r^2) e^{-\rho} e_z\right).$$

Both fields are horizontal, divergence free and Schwartz; $\mathcal{P}V = V$ and $\mathcal{P}W_\sigma = 0$. Write $g_a = (\pi/a)^{3/2}$. Gaussian moments give

$$\begin{aligned} \|V\|_2^2 &= 2g_2, & \|W_\sigma\|_2^2 &= \tfrac{9}{2} g_2, \\ \|\nabla_h W_\sigma\|_2^2 &= 30 g_2, & \|\partial_z W_\sigma\|_2^2 &= \tfrac{9}{2} g_2, \\ \int W_\sigma \cdot (W_\sigma \cdot \nabla) V &= -\sigma \tfrac{16}{9} g_3. \end{aligned}$$

For example, the radial–azimuthal entry of $\mathcal{P}(W_\sigma \otimes W_\sigma)$ is $2\sigma r^4 e^{-2\rho}$, while $\partial_r V_\theta - V_\theta / r = -4 r^2 e^{-\rho}$; their product directly gives the last integral.

Let $\mu = \lambda^{2/3}$, $\lambda \ge 1$, and put

$$C_\epsilon = g_2(2 + \tfrac{9}{2}\epsilon^2), \qquad A = \frac{\lambda\sqrt{\mu}}{\sqrt{C_\epsilon}}, \qquad u_\sigma(x, y, z) = A(V + \epsilon W_\sigma)(\lambda x, \lambda y, \mu z).$$

The absence of an axial component preserves incompressibility under this anisotropic scaling. Orthogonality proves the stated unit norm and relative defect norm. The positive nonlinear injection $I_+ = -T_+$ and its viscous cost νD_+ obey

$$\frac{I_+}{\nu D_+} = \frac{A\lambda(16 g_3/9)}{\nu g_2(30\lambda^2 + \frac{9}{2}\mu^2)} \longrightarrow \infty.$$

Indeed this ratio grows as a positive constant times $\lambda^{1/3}/\nu$, with ϵ fixed. Choose it larger than $C + 1$ and apply (34). For $\sigma = -1$ the injection changes sign while the dissipation is unchanged. These are jets of actual local unforced solutions from Schwartz data; no imposed strain or truncated source is used. Continuity also gives a nonzero initial interval with the strict inequality, but no uniform length of that interval is asserted.

Finally take $\tau_\lambda = \lambda^{-2}$. The mean has only an azimuthal component, whose (a, b) derivative is bounded by

$$C_{ab}\lambda^{4/3 + a + (2/3)b} = C_{ab}\tau_\lambda^{-1/2 - 1/6 - a/2 - (1/2 - 1/6)b}.$$

The constants are uniform for $0 < \epsilon \le 1$ because $C_\epsilon \ge 2g_2$. This proves the asserted spatial bounds. □

The example refutes zero-allowance viscous absorption of kinetic mean–defect transfer, even with small relative defect and the stated spatial geometry. It does not refute a data-dependent history allowance, control of repeated interactions, or Theorems 1 and 5. In particular these transfers are kinetic energy transfers, not vorticity stretching integrals.

There is also an exact circulation test. The initial mean circulation $\Gamma = rv_\theta$ has its global maximum at $r = 1/\lambda$, $z = 0$, with value $2A/(e\lambda)$. For the base Gaussian defect,

$$rF_\theta = 4\sigma r^4(2r^2 - 3)e^{-2\rho}, \qquad F = -\mathcal{P}[(W_\sigma \cdot \nabla)W_\sigma].$$

At that maximum the radial and axial diffusion of the base circulation are $-8/e$ and $-4/e$. Consequently the actual initial mean equation gives

$$\Gamma_t(1/\lambda, 0) = -\frac{4\nu A}{e\lambda}(2\lambda^2 + \mu^2) - \frac{4\sigma\epsilon^2 A^2}{e^2}. \tag{35}$$

For $\sigma = -1$, choose λ still larger so that $\epsilon^2 A/(\nu\lambda) > e[2 + (\mu/\lambda)^2]$. The value at the old maximum then increases initially, and hence exceeds the initial global maximum. Its energy comes from the defect in (34). Thus the maximum principle for exactly axisymmetric circulation cannot be extended by dropping the Reynolds torque, even for arbitrarily small relative angular defect. This is an initial growth example, not repeated spin-up or singularity formation.

Proposition 8. *There are constants $c_0, c_1, c_2, C_0 > 0$ with the following property. Fix $\nu > 0$, $0 < \epsilon \le 1$, and $0 < h \le 1/6$. Set*

$$\mu = \lambda^{1-2h}, \qquad A = \lambda^{1+2h}/\sqrt{C_\epsilon}, \qquad u_\lambda(x_h, z) = A(V + \epsilon W_-)(\lambda x_h, \mu z).$$

For every sufficiently large λ, the actual unforced solution from u_λ exists at least until

$$t_\lambda = \frac{c_0}{A\lambda}.$$

These data have $\|u_\lambda\|_2^2 = \lambda^{-1+6h} \le 1$ and relative L^2 defect $3\epsilon/2$. For $h = 1/6$ they are the unit-energy data u_- of Proposition 7. Its mean circulation has a smooth branch of local maxima in the meridional variables, at $(r, z) = (r_\lambda(t), 0)$, starting at $(\lambda^{-1}, 0)$. Along this branch the radius strictly decreases and the circulation strictly increases. At the last time,

$$r_\lambda(t_\lambda) \le \lambda^{-1}(1 - c_1\epsilon^2), \tag{36}$$

$$\Gamma(r_\lambda(t_\lambda), 0, t_\lambda) \ge \frac{2A}{e\lambda}(1 + c_1\epsilon^2). \tag{37}$$

On this whole interval the mean satisfies (1)–(2) with this h, constant C_0, and $\tau = \lambda^{-2} - t > 0$. Furthermore

$$\int_0^{t_\lambda} \|u(t) - \mathcal{P}u(t)\|^2_{L^\infty(B(1))}\, \mathrm{d}t \ge c_2\epsilon^2\lambda^{2h}. \tag{38}$$

All constants are independent of $\nu, \epsilon, h, \lambda$; the required lower threshold on λ may depend on ν, ϵ, h. The number λ^{-2} specifies a comparison scale, not an asserted singular time.

Proof. Put $y_h = \lambda x_h$, $Z = \mu z$, $s = A\lambda t$, $U = u/A$, and

$$\delta = \mu/\lambda = \lambda^{-2h}, \qquad \eta = \nu\lambda/A, \qquad D_\delta = (\partial_{y_1}, \partial_{y_2}, \delta\partial_Z), \qquad \Delta_\delta = \partial_{y_1y_1} + \partial_{y_2y_2} + \delta^2\partial_{ZZ}.$$

The full projected equation, including pressure, is

$$U_s = -\mathbb{P}_\delta[(U \cdot D_\delta)U] + \eta\Delta_\delta U, \qquad D_\delta \cdot U = 0, \tag{39}$$

with initial field $V + \epsilon W_-$. Here $\mathbb{P}_\delta$ has Fourier multiplier $I - k_\delta \otimes k_\delta/|k_\delta|^2$, $k_\delta = (k_1, k_2, \delta k_3)$. It is an orthogonal projection of norm at most one in every isotropic Sobolev space and commutes with derivatives. No estimate of an inverse power of δ is used.

For $0 < \delta \le 1$ and $0 \le \eta \le 1$, differentiated energy estimates for (39) give

$$\frac{\mathrm{d}}{\mathrm{d}s} \|U\|_{H^{10}}^2 \le C \|U\|_{H^{10}}^3 .$$

Indeed the leading transport cancels by $D_\delta \cdot U = 0$, the commutators have the usual Sobolev product bounds with $|k_\delta| \le |k|$, and the diffusion is nonpositive in the energy. Friedrichs regularization, this estimate, and the corresponding difference estimate construct a common interval $[0, s_0]$ with $\|U\|_{H^{10}} \le M$, uniformly in these parameters and $0 < \epsilon \le 1$. The initial norm is bounded by $\|V\|_{H^{10}} + \|W_-\|_{H^{10}}$. Higher derivative estimates propagate smoothness on that interval. The equation also bounds U_s in H^8 and, after one time derivative, U_{ss} in H^6, uniformly. Sobolev embedding therefore gives uniform spatial bounds for the needed time derivatives.

The fluctuation $w^* = U - \mathcal{P}U$ satisfies the exact equation

$$w_s^* = -\mathbb{P}_\delta\big[(v^* \cdot D_\delta)w^* + (w^* \cdot D_\delta)v^* + (1 - \mathcal{P})((w^* \cdot D_\delta)w^*)\big] + \eta \Delta_\delta w^*, \qquad v^* = \mathcal{P}U. \tag{40}$$

Angular averaging is an orthogonal projection in isotropic H^8, commutes with the isotropic Sobolev multiplier and with $\mathbb{P}_\delta$, and annihilates w^*. Thus the mean correction in the quadratic term does no H^8 work on w^*. Transport cancellation and product bounds, using the uniform H^{10} bound on U, give

$$\frac{\mathrm{d}}{\mathrm{d}s} \|w^*\|_{H^8}^2 \le C_M \|w^*\|_{H^8}^2 , \qquad \|w^*\|_{H^8} + \|w_s^*\|_{H^6} \le C\epsilon \quad (0 \le s \le s_0).$$

The time-derivative estimate follows directly from (40). In particular the stress $\mathcal{P}(w^* \otimes w^*)$ and its time derivative have spatial C^3 norm at most $C\epsilon^2$ on any fixed annulus.

Let $\widetilde{\Gamma} = R(\mathcal{P}U)_\theta$, where $R = |y_h|$. Initially $\widetilde{\Gamma} = 2R^2 e^{-R^2 - Z^2}$. At $(1, 0)$ its meridional Hessian is diagonal with entries $-8/e$ and $-4/e$. The actual mean equation and the Gaussian torque above give exactly

$$\begin{aligned} \widetilde{\Gamma}_s(1, 0, 0) &= 4\epsilon^2/e^2 - 4\eta(2 + \delta^2)/e, \\ \widetilde{\Gamma}_{sR}(1, 0, 0) &= -16\epsilon^2/e^2 + 16\eta/e. \end{aligned}$$

Reflection in Z, with the axial velocity component reflected too, is a symmetry of (39) and its initial data. Uniqueness keeps $\widetilde{\Gamma}$ even in Z. The implicit function theorem and the uniform derivative bounds therefore give a uniform short interval with a nearby meridional local maximum at $(R_*(s), 0)$, $R_*(0) = 1$. Its Hessian remains negative definite, with $|\widetilde{\Gamma}_{RR}| \le 12/e$ there.

Axisymmetry and the mean's actual divergence constraint give

$$v_r^*(R, Z) = -\frac{\delta}{R} \int_0^R q \partial_Z v_z^*(q, Z) \, \mathrm{d}q.$$

Cauchy–Schwarz and the H_Z^1 trace bound into $L_{y_h}^2$ bound this by $C\delta$ uniformly in space. With $f = \partial_Z v_z^*$ its first two radial derivatives are

$$\partial_R v_r^* = -\delta f(R) + \delta \int_0^1 q f(qR) \, \mathrm{d}q, \qquad \partial_{RR} v_r^* = -\delta f_R(R) + \delta \int_0^1 q^2 f_R(qR) \, \mathrm{d}q.$$

The same estimates after axial derivatives and the uniform Sobolev bounds give $|\partial_R^a \partial_Z^b v_r^*| \le C\delta$ for $a + b \le 2$. The other two components have uniformly bounded cylindrical derivatives of these

orders. Applying the same argument to v_s^*, bounded in H^8, also gives an $O(\delta)$ bound on the first two derivatives of $v_{r,s}^*$.

On the fixed rectangle about $(1,0)$, the mean circulation equation is

$$\widetilde{\Gamma}_s + v_r^*\widetilde{\Gamma}_R + \delta v_z^*\widetilde{\Gamma}_Z = \eta(\widetilde{\Gamma}_{RR} - R^{-1}\widetilde{\Gamma}_R + \delta^2\widetilde{\Gamma}_{ZZ}) - Re_R - 2e - \delta R f_Z,$$

where $e = \mathcal{P}(w_r^* w_\theta^*)$ and $f = \mathcal{P}(w_z^* w_\theta^*)$. The preceding bounds, and then one time derivative of this equation, give

$$\left\|\widetilde{\Gamma}_s\right\|_{C^2} + \left\|\widetilde{\Gamma}_{ss}\right\|_{C^1} \le C(\delta + \eta + \epsilon^2).$$

For the diffusive term after that derivative, the uniform H^8 bound on U_s supplies the required derivatives of $\widetilde{\Gamma}_s$. Hence, if λ is large enough that $\delta + \eta \le \epsilon^2/(12e)$, the two displayed initial rates, evaluated along the maximum branch, change by at most $C\epsilon^2 s$. After reducing a universal $c_0 > 0$, for every $0 \le s \le c_0$,

$$\frac{\mathrm{d}}{\mathrm{d}s}\widetilde{\Gamma}(R_*(s), 0, s) \ge 2\epsilon^2/e^2, \qquad R_*'(s) = -\frac{\widetilde{\Gamma}_{sR}}{\widetilde{\Gamma}_{RR}} \le -\frac{2\epsilon^2}{3e}.$$

Integrating and using $r_\lambda = R_*/\lambda$ and $\Gamma = (A/\lambda)\widetilde{\Gamma}$ proves (36)–(37) for some universal $c_1 > 0$. The branch starts at the global circulation maximum and its value exceeds that initial maximum. It need not be identified with the global maximizer at later times.

For the stated physical mean geometry, the radial estimates and $A\delta = \lambda/\sqrt{C_\epsilon}$ give exactly the claimed mean bounds at scale $\tau_\lambda = \lambda^{-2}$. For large λ, $t_\lambda < \tau_\lambda/2$; using $\tau_\lambda - t$ only enlarges their right-hand sides.

Finally $W_-(1,0,0) = (-2/e, 0, 0)$ and the bound on w_s^* gives

$$\|U(s) - \mathcal{P}U(s) - \epsilon W_-\|_\infty \le C\epsilon s.$$

Reduce c_0 once more so that the defect at this fixed point has magnitude at least ϵ/e throughout the interval. Its physical point $(\lambda^{-1}, 0, 0)$ belongs to $B(1)$. Consequently the integral in (38) is at least

$$\frac{A^2\epsilon^2}{e^2}\,\frac{c_0}{A\lambda} = \frac{c_0\epsilon^2}{e^2\sqrt{C_\epsilon}}\lambda^{2h}.$$

The uniform upper bound on C_ϵ proves the assertion. □

This is a finite concentrating event in the actual unforced equation, with all nonlinear feedback retained. It excludes a rule that such feedback always prevents inward motion and growth of a mean circulation peak. It also excludes a bound on the critical defect history depending only on kinetic energy, viscosity and an upper bound for the length of the smooth interval: the interval here tends to zero, whereas (38) tends to infinity for fixed ϵ. The initial higher norms grow with λ. No infinite cascade from one fixed datum, data-dependent history obstruction, or singularity follows from this family.

7.1 Fast axial seeds and the mean geometry

The preceding example has slow axial variation in both its mean and its defect. Fast axial oscillations need not remove its finite inward growth. Their phase relation matters, however, for the regularity of the angular mean. In cylindrical coordinates write

$$\Psi_\alpha(R, \phi, Z) = e^{-R^2 - Z^2}\left[R^2\cos(2\phi + \alpha) - R^4\sin(2\phi + \alpha)\right], \qquad W_\alpha = \mathrm{curl}_h(\Psi_\alpha e_z).$$

Here curl_h takes only the two horizontal derivatives and has zero axial component. These expressions extend smoothly across $R = 0$; $W_0 = W_-$. Two defects with a phase variable α are

$$W^{\mathrm{st}}(y_h, Z, \alpha) = W_0(y_h, Z)\cos\alpha, \qquad W^{\mathrm{hel}}(y_h, Z, \alpha) = W_\alpha(y_h, Z).$$

The first is a standing axial wave, while the second rotates its horizontal pattern with its axial phase.

Proposition 9. *Fix $\nu > 0$, $0 < \epsilon \le 1$, and $0 < h \le 1/6$. With $\delta = \lambda^{-2h}$, $\mu = \lambda^{1-2h}$ and $A = \lambda^{1+2h}/\sqrt{C_\epsilon}$ as in Proposition 8, set*

$$u^j_\lambda(x_h, z) = A\big[V(\lambda x_h, \mu z) + \epsilon W^j(\lambda x_h, \mu z, \lambda z)\big], \qquad j \in \{\mathrm{st}, \mathrm{hel}\}.$$

Both are real Schwartz solenoidal data, with the same angular mean, squared energy at most one and relative L^2 defect at most $3\epsilon/2$. Their initial means obey (1)–(2) with $\tau = \lambda^{-2}$ and a constant independent of λ and ϵ.

For all sufficiently large λ, the helical data have all the conclusions of Proposition 8, with universal constants, except that the maximum branch may also move axially: it is $(r_\lambda(t), z_\lambda(t))$, initially $(\lambda^{-1}, 0)$. In particular the mean geometry persists on $0 \le t \le c_0/(A\lambda)$, the branch moves inward and gains circulation by relative amounts at least $c_1\epsilon^2$, and its actual defect history is at least $c_2\epsilon^2\lambda^{2h}$. Its squared energy is exactly one when $h = 1/6$.

For the standing data, there instead exist universal $c_3, c_4 > 0$ such that, at $t_* = c_3\epsilon^2/(A\lambda)$ and $\tau_* = \lambda^{-2} - t_* > 0$, the actual unforced solution satisfies

$$\tau_*^{3/2-h}\left|\partial_{zz}(\mathcal{P}u^{\mathrm{st}}_\lambda)_\theta(\lambda^{-1}, 0, t_*)\right| \ge c_4\epsilon^4\lambda^{4h}. \tag{41}$$

Thus initial mean geometry, a bounded kinetic energy and a small relative defect alone do not propagate these mean bounds with a uniform constant. All thresholds on λ may depend on ν, ϵ, h.

Proof. Introduce an independent periodic variable $\alpha \in \mathbb{R}/(2\pi\mathbb{Z})$ and solve on $\mathbb{R}^2 \times \mathbb{R} \times (\mathbb{R}/(2\pi\mathbb{Z}))$ with

$$D_{\delta,*} = (\partial_1, \partial_2, \delta\partial_Z + \partial_\alpha), \qquad \eta = \nu\lambda/A,$$
$$U_s = -\mathbb{P}_{\delta,*}[(U \cdot D_{\delta,*})U] + \eta\Delta_{\delta,*}U, \qquad D_{\delta,*} \cdot U = 0, \quad U(0) = V + \epsilon W^j.$$

The projection has symbol $I - \kappa \otimes \kappa/|\kappa|^2$, where $\kappa = (k_1, k_2, \delta k_Z + n)$ and $n \in \mathbb{Z}$ is the phase frequency. It has norm at most one in every Sobolev space. The four-dimensional transport field $(U_1, U_2, \delta U_3, U_3)$ is divergence free. Thus the same transport cancellation, commutator estimates and nonpositive diffusion as before give a common smooth interval with a uniform H^{12} bound, for $0 < \delta \le 1$, $0 \le \eta \le 1$, and $0 < \epsilon \le 1$. The first three time derivatives are bounded in H^{10}, H^8, H^6, respectively. These estimates use the full phase derivative, not a truncation of the generated modes.

Restrict this solution to $\alpha = Z/\delta$. The restricted axial derivative $\delta\partial_Z$ is exactly $\delta\partial_Z + \partial_\alpha$ before restriction. Hence the restriction solves the actual three-dimensional projected equation, with its pressure, from the stated initial datum. The elementary H^1 bound in the periodic variable controls this restriction in $L^2_{y_h,Z}$; applying it to derivatives gives smoothness and finite Sobolev norms for each fixed $\delta > 0$. The pressure has the same property, so this is the finite-energy unforced evolution, by uniqueness. Rescaling $y_h = \lambda x_h$, $Z = \mu z$, and $s = A\lambda t$ gives the physical flow. In particular its phase is λz.

For the helical field the extended initial datum obeys the joint symmetry

$$U(R_\beta y_h, Z, \alpha - 2\beta) = R_\beta U(y_h, Z, \alpha),$$

where the rotation fixes the third velocity component. The equation commutes with this action, so uniqueness preserves it. Angular averaging therefore makes $v^* = \mathcal{P}U$ independent of α throughout the interval. This is a nonlinear symmetry assertion, not only a cancellation of the initial quadratic source. Consequently the physical mean's axial derivatives involve only its slow Z derivatives, which are uniformly bounded. Its divergence constraint is

$$\partial_R v_r^* + R^{-1} v_r^* + \delta \partial_Z v_z^* = 0.$$

The radial primitive and slice estimates in the preceding proof give the required $O(\delta)$ radial gain. All the mean bounds thus transfer to the physical scale exactly as in Proposition 8.

Angular averaging is orthogonal in the extended isotropic Sobolev spaces too. The exact homogeneous fluctuation equation therefore gives $\|U - \mathcal{P}U\|_{H^8} + \|\partial_s(U - \mathcal{P}U)\|_{H^6} \le C\epsilon$. Its stress and the time derivative of that stress are $O(\epsilon^2)$ in spatial C^3 on a fixed annulus. Direct angular integration gives

$$\mathcal{P}(W_r^{\mathrm{st}} W_\theta^{\mathrm{st}}) = -2R^4 e^{-2R^2-2Z^2} \cos^2\alpha, \tag{42}$$

$$\mathcal{P}(W_r^{\mathrm{hel}} W_\theta^{\mathrm{hel}}) = -2R^4 e^{-2R^2-2Z^2}. \tag{43}$$

The latter gives the same initial circulation and radial rates as in the preceding proof, and the initial axial rate of the critical point is zero. The mean equation again bounds its circulation time derivatives by $C(\delta + \eta + \epsilon^2)$. Use the two-variable implicit function theorem for the negative definite meridional Hessian. The initial inward radial rate and positive circulation rate persist on a universal interval when $\delta + \eta$ is sufficiently small compared with ϵ^2. This proves the assertions about the moving critical point without assuming that its axial coordinate stays zero. At the fixed point $(y_h, Z, \alpha) = ((1,0),0,0)$ the initial defect is $\epsilon(-2/e, 0, 0)$. Its uniform $O(\epsilon)$ time-derivative bound proves the same critical-history lower bound.

For the standing field put $G = R(\mathcal{P}U)_\theta$ and $J = \delta\partial_Z + \partial_\alpha$. Initially the mean has no meridional component, so its exact equation is

$$G_s = \eta(G_{RR} - R^{-1}G_R + J^2 G) + 4\epsilon^2 R^4(3 - 2R^2)e^{-2R^2-2Z^2}\cos^2\alpha \quad \text{at } s = 0.$$

At $(R, Z, \alpha) = (1, 0, 0)$ define $F(s) = J^2 G(1, 0, 0, s)$. The two exact initial values are

$$F(0) = -4\delta^2/e, \qquad F'(0) = \eta\delta^2(16 + 24\delta^2)/e - \epsilon^2(16\delta^2 + 8)/e^2.$$

The uniform time and space bounds above give $|F''(s)| \le C$. For sufficiently small δ, $F'(0) \le -4\epsilon^2/e^2$. Choosing a universal small $c_3 > 0$, Taylor's theorem at $s_* = c_3\epsilon^2$ gives $F(s_*) \le -c\epsilon^4$. On restriction and physical rescaling,

$$\partial_{zz}(\mathcal{P}u_\lambda^{\mathrm{st}})_\theta(\lambda^{-1}, 0, t_*) = A\lambda^2 F(s_*).$$

Since $\tau_* \ge \lambda^{-2}/2$ for large λ, this is (41).

Finally angular integration makes the helical defect's squared norm independent of its phase; it is $\frac{9}{2}g_2$ before the factor ϵ^2. The standing defect's squared norm is instead

$$\tfrac{9}{4}g_2(1 + e^{-1/(2\delta^2)}).$$

Both are orthogonal to V. This proves the energy and relative-defect claims, while the common initial mean is AV. □

The standing example disproves a uniform propagation claim based only on the displayed initial controls. It does not disprove a bound depending on the full initial datum. The helical example shows why an argument cannot assume that fast axial oscillations must destroy the mean geometry. Its exact phase symmetry concerns one family; superposing families with different phase relations need not preserve it. The uniform Sobolev control proved here still ends after a fixed normalized time. Neither example establishes a recurrent cascade or an arbitrary-data history bound.

7.2 A mean mode generated by incompatible parents

Two individually helical parents need not share a helical symmetry. Their interaction can create a fast mean mode even when their initial mean stress has no fast axial phase. The next example keeps that mode as a physical Fourier projection of the actual solution.

Fix a real even Schwartz function g whose Fourier transform is smooth, nonnegative, supported in $[-1,1]$, and has nonzero mass away from zero. Normalize $g(0)=1$; then $g''(0)<0$. In the horizontal variables put $q=y_1+iy_2$, $a=|y_h|^2$, and

$$V_h=\operatorname{curl}_h(e^{-a}e_z),\qquad W_2=\operatorname{curl}_h(q^2e^{-a}e_z),\qquad W_4=\operatorname{curl}_h\left(q^4(1-2a/5)e^{-a}e_z\right).$$

The last two velocities are complex; their real parts will be used. For $0<h\le 1/6$ define

$$\delta=\lambda^{-2h},\quad \epsilon=\lambda^{-h}=\sqrt{\delta},\quad \mu=\lambda^{1-2h},\quad C_\epsilon=\pi\,\|g\|_2^2\,(1+9\epsilon^2/5),\quad A=\lambda^{1+2h}/\sqrt{C_\epsilon}.$$

In this subsection C_ϵ denotes this normalization. Take the real initial datum

$$u_\lambda(x_h,z)=Ag(\mu z)\left[V_h(\lambda x_h)+\epsilon\,\mathrm{Re}\left((W_2+W_4)(\lambda x_h)e^{2i\lambda z}\right)\right]. \tag{44}$$

Let Q_λ be angular averaging followed by the axial Fourier projection onto $\lambda\le|k_z|\le 3\lambda$, with the Fourier convention e^{ik_zz}. This is an orthogonal projection in L^2, preserves solenoidality and commutes with the Laplacian.

Proposition 10. *For each fixed $\nu>0$ and $0<h\le 1/6$, and all sufficiently large λ, the actual unforced evolution from (44) exists until $t_\lambda=s_*/(A\lambda)$, for a fixed $s_*>0$. Its initial squared energy is $\lambda^{-1+6h}\le 1$, its relative L^2 angular defect is $3\epsilon/\sqrt{5}\to 0$, and its initial mean satisfies (1)–(2) with a uniform constant. On this interval its critical fluctuation history is uniformly bounded:*

$$\int_0^{t_\lambda}\|u(t)-\mathcal{P}u(t)\|_\infty^2\,\mathrm{d}t\le C.$$

Set $Y=Q_\lambda u$ and

$$E_Y=\tfrac12\|\nabla Y\|_2^2,\qquad D_Y=\nu\|\Delta Y\|_2^2,\qquad P_Y=\langle-\Delta Y,Q_\lambda\mathcal{N}(u)\rangle,\quad \mathcal{N}(u)=-\mathbb{P}[(u\cdot\nabla)u].$$

Here $\mathbb{P}$ is the physical Leray projection. Exactly $Y(0)=Y_t(0)=0$, while at the last time

$$c\lambda^{-1}\le\|Y(t_\lambda)\|_2^2\le C\lambda^{-1}, \tag{45}$$

$$\int_0^{t_\lambda}P_Y(t)\,\mathrm{d}t\ge c\lambda,\qquad\qquad \int_0^{t_\lambda}D_Y(t)\,\mathrm{d}t\le C\nu\lambda^{1-2h}. \tag{46}$$

Moreover, with $\tau_\lambda=\lambda^{-2}-t_\lambda>0$,

$$\tau_\lambda^{3/2-h}\left|\partial_{zz}(\mathcal{P}u)_\theta(\lambda^{-1},\pi/(2\lambda),t_\lambda)\right|\ge c\lambda^h. \tag{47}$$

The constants s_*,c,C depend on the fixed envelope and profiles, not on λ,ν,h; the lower threshold on λ may depend on ν,h.

Proof. Use the extended construction above, now with $D = (\partial_1, \partial_2, \delta\partial_Z + 2\partial_\alpha)$, initial field

$$U(0) = g(Z)\left[V_h + \epsilon\,\mathrm{Re}((W_2 + W_4)e^{i\alpha})\right],$$

and $\eta = \nu\lambda/A$. Restrict to $\alpha = 2Z/\delta$ and then rescale as before. Work initially with a uniform H^{30} bound on the extended local interval. The projection remains a contraction, with symbol corresponding to $(k_h, \delta k_Z + 2n)$, so the same existence, trace and uniqueness arguments apply.

We first compute the relevant cubic jet at $\delta = 0$. Label a mode by its horizontal angular index and its phase index (m, n). The parents have labels $(2,1), (4,1)$ and their negatives. The only three-parent sum with label $(0,1)$ is $(2,1) + (2,1) - (4,1)$. No single parent or pair, even including the $(0,0)$ base, reaches it. Linear derivatives preserve these labels. Thus neither the base nor any term containing viscosity contributes to this mode in $U_{ss}(0)$.

The two intermediate pressure equations have explicit decaying solutions. Suppressing the common factor $g(Z)^2$, they are

$$p_{22} = -q^4 e^{-2a}, \qquad p_{2\bar{4}} = \frac{\bar{q}^2}{5}(4a^3 + 18a^2 - 62a + 9)e^{-2a}.$$

The first solves the pressure equation with axial wave number 4, so its elliptic operator is $-\Delta_h + 16$; the second has zero axial wave number and operator $-\Delta_h$. Their sources are the divergences of $(W_2 \cdot D)W_2$ and $(W_2 \cdot D)\overline{W}_4 + (\overline{W}_4 \cdot D)W_2$, respectively. Direct differentiation verifies both equations and the solenoidality of their full quadratic velocity sources, including the axial pressure component. Decay removes a harmonic ambiguity.

Insert these two sources into both orders of the outer convection. The final pressure has zero angular index and hence no azimuthal component. The resulting circulation coefficient is

$$C_\Gamma(R) = \frac{8}{5}R^4(8R^8 - 110R^6 + 346R^4 - 275R^2 + 27)e^{-3R^2}, \qquad C_\Gamma(1) = -\frac{32}{5e^3}. \tag{48}$$

Each real signed parent has amplitude $\epsilon/2$. Including the conjugate output, the fast mean part of $G_{ss}(0)$, where $G = R(\mathcal{P}U)_\theta$, is therefore

$$\frac{\epsilon^3}{4} g(Z)^3 C_\Gamma(R)\cos\alpha. \tag{49}$$

The zero-phase mean can have additional terms; it is not discarded from the evolution. The lab checker verifies the two pressure equations, every contributing convection branch, and the real-field normalization in (49).

A bound uniform in the small amplitude is needed to use this jet. For this purpose hold η fixed while differentiating in ϵ; all estimates will be uniform for $0 \le \eta \le 1$. If $v_j = \partial_\epsilon^j U$, its equation is the linearized projected equation about U, with additional source

$$-\mathbb{P}_{\delta,*}\sum_{k=1}^{j-1}\binom{j}{k}(v_k \cdot D)v_{j-k}.$$

The transport energy estimate gives uniform bounds for v_j in H^{30-2j} for $j \le 3$. Difference quotients justify these finite parameter derivatives in the lower Sobolev spaces. Three time derivatives cost at most six additional spatial derivatives, leaving a uniform H^{18} bound for the mixed derivatives needed here. No analyticity of the solution map in a fixed Sobolev norm is assumed.

At $\epsilon = 0$, the evolving base has label $(0,0)$, so its linearized equation preserves (m, n). The first parameter derivative has only the four parent labels, and the second only their pairwise sums.

Neither has angular index zero with nonzero phase index. Taylor's theorem in ϵ consequently bounds the entire fast angular mean, including its first three time derivatives in the required spatial norms, by $C\epsilon^3$ throughout a common interval.

The physical projection Q_λ also sees high frequencies of the evolving zero-phase mean. Its slow axial Fourier support is not claimed to remain compact. Six bounded Z derivatives instead make this contribution at most $C\delta^6$, also after the needed spatial and time derivatives. Since $\epsilon = \sqrt{\delta}$, this is bounded by $C\epsilon^3$. The phase trace bound transfers the other components to the actual physical restriction.

At the initial time the support assertions are exact. The initial zero phase has axial support at most μ, its quadratic sources at most 2μ, and its cubic sources at most 3μ. When $\delta < 1/3$ these are below the receiving band. The new phases $n = \pm 1$ in the cubic jet lie in $\pm 2\lambda + [-3\mu, 3\mu]$, wholly inside that band. The pressure multipliers do not enlarge axial Fourier support. This proves $Y(0) = Y_t(0) = 0$ and identifies its second derivative.

In normalized physical coordinates, let $\widetilde{Y}$ denote this recipient and $D_\delta = (\partial_1, \partial_2, \delta\partial_Z)$. The preceding bounds give, in the norm controlling up to two D_δ derivatives,

$$\widetilde{Y}(s) = \tfrac{1}{2}\epsilon^3 s^2 C_\delta + O(\epsilon^3 s^3). \tag{50}$$

The coefficient has uniformly bounded norms. Its phase coefficients converge in Sobolev spaces to the nonzero coefficient just computed: the Leray multipliers converge almost everywhere and are uniformly bounded, so dominated convergence applies to the fixed smooth inputs. The two physical axial bands are disjoint, and on them $|\delta k_Z| \geq 1$. It follows that both $\|C_\delta\|_2$ and $\|D_\delta C_\delta\|_2$ have positive uniform lower bounds for small δ.

Choose a fixed small $s_* > 0$ in (50). Then the recipient's squared velocity and gradient norms at s_* are bounded above and below by constant multiples of ϵ^6, and

$$\eta \int_0^{s_*} \left\| \Delta_\delta \widetilde{Y}(s) \right\|_2^2 \, \mathrm{d}s \leq C\eta\epsilon^6.$$

Physical squared velocity norms acquire the factor $A^2/(\lambda^2\mu)$, and squared gradient norms acquire A^2/μ. Since $\epsilon^6 = \lambda^{-6h}$, these factors yield (45), a recipient enstrophy of order λ, and the dissipation bound in (46). The exact projected energy identity is

$$E_Y'(t) + D_Y(t) = P_Y(t).$$

Since $E_Y(0) = 0$, it proves the signed production lower bound without replacing the nonlinear work by its absolute value.

For the mean geometry put $J = \delta\partial_Z + 2\partial_\alpha$ and evaluate $F(s) = J^2 G(1, \pi\delta/2, \pi, s)$. Initially $F(0)$ is negative of order δ^2, because $g''(0) < 0$, and $F'(0) = O(\eta\delta^2)$. Each initial parent has zero mean radial–azimuthal stress, and their cross stress has nonzero angular index; hence there is no initial fast mean torque. Equations (48)–(49) give the fast part

$$F''(0) = -\frac{32}{5e^3}\epsilon^3 + o(\epsilon^3)$$

as $\delta \to 0$; the zero phase contributes only $O(\delta^2)$. Moreover $|F'''| \leq C(\epsilon^3 + \delta^2)$ by the phase-selection estimate. Here $\delta^2 = o(\epsilon^3)$. After reducing the fixed s_*, Taylor's theorem gives $F(s_*) \leq -c\epsilon^3$. The initial negative term absorbs the $O(\eta\delta^2 s_*)$ term uniformly for $\eta \leq 1$. The physical axial derivative equals $A\lambda^2 F(s_*)$ at the point in (47). Since $\lambda^{-2} - t_\lambda \geq \lambda^{-2}/2$ for large λ, the normalized lower bound is $c\epsilon^3\lambda^{4h} = c\lambda^h$.

The first parameter-derivative estimate also gives $\|U - \mathcal{P}U\|_\infty \le C\epsilon$ on the common interval, since the zero-amplitude solution is angularly symmetric. Therefore the physical fluctuation history is bounded by $CA^2\epsilon^2/(A\lambda) = C/\sqrt{C_\epsilon}$, uniformly in the concentration.

Finally the horizontal squared norms of V_h, $\operatorname{Re} W_2$ and $\operatorname{Re} W_4$ are π, $3\pi/4$ and $21\pi/20$. Distinct angular indices make all cross energies zero. This proves the normalization and relative-defect assertions. □

This receiving sector starts empty and is generated by the actual unforced parents. Its signed production exceeds its own viscous cost by an unbounded factor across the family, while its kinetic energy remains small. This does not refute a signed budget for the full high-frequency tail: the parents are in the same axial band, outside the angular mean, and their signed work must also be included. The initial higher norms grow with λ, so a data-dependent history allowance and a recurrent estimate from one fixed datum remain possible. Nor does the example prove that a possible singularity must preserve the mean geometry assumed in Theorem 1.

7.3 Positive total work after including the parents

The preceding receiving-sector calculation does not determine the sign of the total work. A single helical parent gives a separate test of that sign, with its pressure and the response of the base included. Use the envelope g, base V_h, and parent W_2 defined above. Fix $\nu > 0$ and $0 < h \le 1/6$. With a constant $b > 0$ to be chosen, put

$$\delta = \lambda^{-2h}, \quad \epsilon = b\lambda^{-h}, \quad \mu = \lambda\delta, \quad C_\epsilon = \pi\,\|g\|_2^2\,(1 + 3\epsilon^2/4), \quad A = \lambda^{1+2h}/\sqrt{C_\epsilon},$$

and take

$$u_\lambda(x_h, z) = Ag(\mu z)[V_h(\lambda x_h) + \epsilon\,\mathrm{Re}(W_2(\lambda x_h)e^{2i\lambda z})]. \tag{51}$$

Here $\epsilon \le 1$ for all sufficiently large λ.

Proposition 11. *One can choose b depending on ν so that the actual unforced evolutions from (51) exist until $t_\lambda = s_*/(A\lambda)$ and satisfy*

$$\frac{1}{2}\,\|\nabla u(t_\lambda)\|_2^2 - \frac{1}{2}\,\|\nabla u(0)\|_2^2 \ge c\lambda^{1+4h}. \tag{52}$$

Their initial squared energies are $\lambda^{-1+6h} \le 1$, their relative angular defects tend to zero, and their means obey (1)–(2) throughout this interval, with $\tau = \lambda^{-2} - t$ and a constant independent of λ. Also $\int_0^{t_\lambda} \|u - \mathcal{P}u\|_\infty^2\,\mathrm{d}t \le Cb^2$.

Consequently, for every fixed Fourier cutoff $K < \infty$ and $\theta \le 1$, the high-frequency projection $u_{\ge K}$ satisfies

$$\int_0^{t_\lambda} \left[\langle -\Delta u_{\ge K}, \mathcal{N}(u)\rangle - \theta\nu\,\|\Delta u_{\ge K}\|_2^2\right]\mathrm{d}t \longrightarrow +\infty. \tag{53}$$

The projection uses the physical convention $e^{ik\cdot x}$ and $|k| \ge K$. The constants in this proposition are independent of λ; no constant uniform over full initial Sobolev norms is claimed.

Proof. First compute the full signed response at $\delta = \eta = 0$, with $\eta = \nu\lambda/A$, per unit fast axial phase and with the slow envelope temporarily suppressed. More generally let the fast carrier be $\kappa > 0$. Write

$$v(R) = 2Re^{-R^2}, \qquad f(R) = R^2e^{-R^2},$$
$$\Omega = v/R, \quad \zeta = v' + v/R, \quad d = v' - v/R, \quad L_2 = \partial_{RR} + R^{-1}\partial_R - 4R^{-2}.$$

The complex parent has cylindrical components $(2if/R, -f', 0)$ and phase $e^{2i\phi+i\alpha}$. Its linear pressure amplitude p solves

$$(-L_2 + \kappa^2)p = F, \qquad F(R) = 32R^2(R^2-1)e^{-2R^2}. \tag{54}$$

This source is twice the trace of the base and parent velocity-gradient matrices; the cylindrical basis derivatives are included.

Let $\Pi = \int \omega \cdot (\omega \cdot \nabla)u$ denote the total signed enstrophy production, and let $\mathcal{F} = (\omega \cdot \nabla)u$. In an unforced inviscid evolution the material identities give

$$(\partial_t + u \cdot \nabla)\mathcal{F} = -(\nabla^2 p)\omega, \qquad \Pi' = \int (|\mathcal{F}|^2 - \omega \cdot \nabla^2 p\, \omega).$$

Indeed the two products of velocity gradients cancel when differentiating $\mathcal{F}$; transport integrates to zero. The coefficient of ϵ^2 in this identity is

$$c_\kappa = \pi\kappa^2 \int_0^\infty Rd[4\zeta f^2/R^2 - 2\Omega(f')^2]\,\mathrm{d}R + 2\pi\kappa^2 \int_0^\infty Rp\zeta' f'\,\mathrm{d}R.$$

The first integral is exactly zero. In the second, the vertical parent vorticity $-L_2 f$ cancels the term $\zeta L_2 f$ from the horizontal pressure pairing. Omitting that component gives an incorrect coefficient. The second-order pressure makes no further contribution: its pairing with the two base vorticities is an axial second derivative whose phase average is zero.

The remaining sign can be proved explicitly. The angular heat kernel is

$$e^{tL_2}(R^2 e^{-\beta R^2}) = \frac{R^2}{(1+4\beta t)^3} e^{-\beta R^2/(1+4\beta t)}.$$

Differentiate in β to heat F. Gaussian integration gives

$$J(t) := \int_0^\infty R(e^{tL_2}F)\zeta' f'\,\mathrm{d}R = \frac{5 - 24t - 256t^3}{2(1+4t)^6}.$$

Although J changes sign, its primitive is strictly positive:

$$B(t) = \int_0^t J(s)\,\mathrm{d}s = \frac{t(128t^4 + 160t^3 + 112t^2 + 28t + 5)}{2(1+4t)^5} > 0 \quad (t > 0), \qquad B(\infty) = \frac{1}{16}.$$

The decaying resolvent in (54) and integration by parts therefore yield

$$c_\kappa = 2\pi\kappa^2 \int_0^\infty e^{-\kappa^2 t} J(t)\,\mathrm{d}t = 2\pi\kappa^4 \int_0^\infty e^{-\kappa^2 t} B(t)\,\mathrm{d}t > 0. \tag{55}$$

All radial boundary terms vanish by smoothness at the axis and decay. The checker verifies the source, both pressure components, the heat equation and the displayed primitive. These are exact identities; positivity follows from the integral formula, not a numerical sign test.

Return to the actual extended flow with $D_3 = \delta\partial_Z + 2\partial_\alpha$ and initial field $g(Z)[V_h + \epsilon\,\mathrm{Re}(W_2 e^{i\alpha})]$. The preserved joint rotation–phase symmetry from Proposition 9 makes every horizontally integrated rotation-invariant density independent of α. Its physical trace and its phase average consequently have the same integral. In particular the computation above applies to the actual physical enstrophy, not to an ensemble of different solutions.

Let $\mathcal{P}_{\mathrm{tot}}(s)$ be this total normalized production. Initially $\mathcal{P}_{\mathrm{tot}}(0) = 0$ exactly. For a horizontal solenoidal initial field, the integrated production is $-\int(\partial_z u_h)\cdot(\nabla_h u_h)(\partial_z u_h)$. Here the terms with

two envelope derivatives vanish by the planar kinetic identity, the terms with two phase derivatives vanish after angular averaging, and the remaining terms integrate a multiple of $g^2 g'$ over Z, which is zero.

We justify a time interval uniform in ϵ, δ, η. The extended transport estimate and finite parameter derivatives used in Proposition 10 give uniform Sobolev bounds for three time derivatives and two amplitude derivatives, with ample derivative margin. Hold η independent of ϵ in these estimates. Rotation by $\pi/2$ changes ϵ to $-\epsilon$, so the physically integrated production and its time derivatives are even functions of ϵ.

For the zero-amplitude solution, the vertical velocity and its needed time derivatives are $O(\delta)$ in Sobolev norms: their equations have pressure source $-\delta \partial_Z p$ and zero initial vertical velocity. The double-Riesz representation bounds p uniformly. Thus the horizontal vorticity and the mixed axial strains are $O(\delta)$, while $\partial_z U_z = O(\delta^2)$. The same assertions hold after the required time derivatives. Every contribution to production at zero amplitude is therefore $O(\delta^2)$. Evenness and the amplitude estimates imply

$$|\mathcal{P}''_{\mathrm{tot}}(s)| \le C(\delta^2 + \epsilon^2) \tag{56}$$

on a common normalized interval.

At time zero, the quartic inviscid production derivative has expansion

$$\mathcal{P}'_{\mathrm{tot}}(0) = O(\delta^2) + \epsilon^2(c_2 \|g\|_4^4 + o(1)) + O(\epsilon^4) + O(\eta\epsilon^2).$$

The $o(1)$ tends to zero with δ: the Leray multipliers converge almost everywhere, are uniformly bounded, and act on fixed smooth inputs. The purely viscous zero-amplitude term is zero because the heat evolution remains purely azimuthal. All constants are uniform for the parameters at most one. With $\epsilon^2 = b^2\delta$ and $\eta = \nu\sqrt{C_\epsilon}\delta$, it follows that $\mathcal{P}'_{\mathrm{tot}}(0) \ge c\epsilon^2$ for sufficiently large λ. Equation (56) and $\mathcal{P}_{\mathrm{tot}}(0) = 0$ then give, for a fixed small $s_* > 0$,

$$\int_0^{s_*} \mathcal{P}_{\mathrm{tot}}(s)\,\mathrm{d}s \ge c\epsilon^2, \qquad \eta \int_0^{s_*} \|\Delta_D U(s)\|_2^2\,\mathrm{d}s \le C\eta.$$

Both enstrophy integrals acquire the physical factor A^2/μ. Since C_ϵ is bounded above and below, choosing b sufficiently large compared with $\sqrt{\nu}$ makes the difference at least $c\lambda^{1+4h}$. The exact enstrophy identity proves (52). The symmetry, incompressibility, and uniform Sobolev estimates prove the persistent mean geometry as in Proposition 9. The amplitude derivative gives $\|U - \mathcal{P}U\|_\infty \le C\epsilon$, so the history is at most $CA\epsilon^2/\lambda \le Cb^2$. Orthogonality of angular modes gives the asserted initial energy.

Finally write $E_{\ge K} = \|\nabla u_{\ge K}\|_2^2/2$. The kinetic energy identity bounds the complementary gradient energy by $K^2/2$. Hence its subtraction changes the enstrophy increment by at most $K^2/2$. The exact projected identity gives

$$\int_0^{t_\lambda} (P_{\ge K} - \theta D_{\ge K})\,\mathrm{d}t = E_{\ge K}(t_\lambda) - E_{\ge K}(0) + (1-\theta)\int_0^{t_\lambda} D_{\ge K}\,\mathrm{d}t \ge c\lambda^{1+4h} - K^2/2,$$

where $D_{\ge K} = \nu\|\Delta u_{\ge K}\|_2^2$. This proves (53). □

This example includes the full parent response and preserves the displayed mean geometry on its short interval. It excludes a uniform signed-tail allowance based only on the bounded quantities just listed, with a fixed cutoff. It does not exclude a cutoff or allowance depending on the full datum: the initial enstrophy itself is of order λ^{1+6h}, larger than the increase just proved. No sequence of these events within one fixed solution, and no failure of the conditional regularity theorem, is asserted.

7.4 The energy inequality alone permits the stress cost

The time-integrated energy inequality cannot by itself exclude the cost in (14). An explicit path shows this while keeping all the mean geometry bounds. This path is not a Navier–Stokes solution. Use W_+ from Proposition 7, choose $0 < h < 1/4$, and set

$$\alpha = \tfrac{1}{2} - 2h, \qquad c^2 = \frac{160\nu}{3\alpha}, \qquad d^2 = \frac{8\nu}{\alpha}, \qquad A_0 = \left(\tfrac{9}{2} g_2 c^2 d\right)^{-1/2}.$$

For $0 < \tau \le 1$ define the whole-space smooth solenoidal path

$$u(x, -\tau) = A_0 \tau^{-1/2-h/2} W_+ \left(\frac{x_1}{c\sqrt{\tau}}, \frac{x_2}{c\sqrt{\tau}}, \frac{z}{d\tau^{1/2-h}} \right).$$

It has zero angular mean. Its squared velocity norm E and squared gradient norm D are exactly

$$E(-\tau) = \tau^{\alpha}, \qquad D(-\tau) = E(-\tau) \left(\frac{20}{3c^2\tau} + \frac{1}{d^2 \tau^{1-2h}} \right).$$

Since $t = -\tau$, these imply

$$\tfrac{1}{2} E'(t) + \nu D(t) = \frac{\alpha E(t)}{8\tau} (\tau^{2h} - 3) < 0, \qquad \nu \int_{-1}^{0} D(t)\, \mathrm{d}t = \frac{1 - 2h}{4} < \infty.$$

Thus the path starts with unit squared velocity norm and satisfies the energy inequality on every time interval, with finite total dissipation. On sufficiently late slices,

$$K(-\tau) = A_0 \left\| W_+ \right\|_{L^\infty(\mathbb{R}^3)} \tau^{-1/2-h/2},$$

so the critical history is infinite. Its radial–azimuthal stress has supremum $2A_0^2 e^{-2} \tau^{-1-h}$ and its axial–azimuthal stress is zero, so

$$\mathcal{B}_\tau(T) = \frac{2A_0^2 e^{-2}}{h} (1 - T^{-h}) > 0$$

for fixed $T > 1$ and sufficiently small τ. The stress maximum then lies in $B(1)$ throughout the interval.

The strict energy loss already shows why this is not an unforced smooth solution, which would satisfy energy equality. Its zero mean also has nonzero Reynolds torque, so an original force would be needed to keep that mean zero. The example only separates the energy inequality and mean geometry from the actual evolution required to control the history.

8 Scope of the calculation

The theorem allows an unbounded angular fluctuation, but requires its history to satisfy (3) and separately assumes the mean geometry. Neither hypothesis is established here for general data. Theorem 5 separately supplies a quantitative gradient history under a stronger weighted amplitude condition. Its localization constants may depend on the solution on a regular annulus; they are not uniform data-only bounds. In Theorem 1, an original force uniformly bounded in spatial C^2 on Q can also be included: its rescaled swirl and vorticity sources vanish, and its enstrophy contribution is bounded by $C(Y + 1)$. Thus a distinction between forced and unforced flows must enter a proof of the hypotheses, not be inferred from this criterion.

The accompanying symbolic calculation checks the mean equation, cylindrical stress divergence, lifted force identity, and the stretching integration by parts on explicit polynomial fields with all six stress components nonzero. It also checks the general cylindrical stretching and local radial integration identities, and the Gaussian moments and complete convection work in Proposition 7. The polynomial fields are local algebra tests, whereas the Gaussian fields are finite-energy initial data. It also checks the signed dual-work integrations and the scaling of the final kinematic path. That path is explicitly distinct from the actual unforced initial jets. The program does not integrate solutions or certify the analytic proof. No statement in this note has been added as an axiom or a theorem to the Lean library.

References

[1] P. Constantin, M. Ignatova, and V. Vicol, *Regularity of asymptotically axisymmetric solutions to the 3D Navier–Stokes equations with analytic forcing*, arXiv:2609.20803v1, 2026.

[2] S. Gustafson, K. Kang, and T.-P. Tsai, *Interior regularity criteria for suitable weak solutions of the Navier–Stokes equations*, Comm. Math. Phys. **273** (2007), 161–176, arXiv:math/0607114.