Signed Navier–Stokes tail budgets: forcing, low frequencies and unforced activation

September 27, 2026

Abstract

For a smooth finite-energy forced Navier–Stokes evolution, we bound the absolute time integrals of direct force work and low-frequency feeding in the high-frequency enstrophy balance. The bounds use kinetic energy, total energy dissipation, a fixed cutoff, and derivatives of the force; they spend none of the high-frequency viscous dissipation. Consequently, for the forced singularities constructed in [1], the accumulated tail self-stretching minus even the full tail dissipation is unbounded at every fixed cutoff. The full-field obstruction is checked in Lean. The Fourier-tail extension below is an analytic proof, not a formalized projector theorem. We also bound the forcing derivative of a quartic caloric stretching functional using only energy-level velocity controls and smooth force norms. Its signed nonlinear renewal must therefore fail a uniform absorption budget on the same forced construction. A Fourier multiplier estimate separates every low-frequency parent and every low renewal output. Low stretching outputs have a separate energy-paid history bound, leaving a target with every parent and both internal frequencies high. Actual unforced whole-space solutions with unit initial energy and an initially empty tail refute an allowance uniform over the data and the prescribed cutoff. Their initial fields can be planar polarized in an arbitrarily thin Fourier belt. This does not exclude an allowance depending on the full datum or a separately chosen cutoff. A further positive estimate absorbs stretching pairs whose parents are sufficiently far above their output, uniformly over all occupied shells. The corresponding modified caloric balance retains the complete Navier–Stokes source. Its remaining signed history bound is unproved; no arbitrary-data unforced regularity estimate is obtained.

1 An exact split and finite allowances

Let u, p, f solve

$$\partial_t u + (u\cdot\nabla)u = -\nabla p + \nu\Delta u + f, \qquad \nabla\cdot u = 0, \qquad \nu > 0,$$

on $\mathbb{R}^3 \times [a,T)$, where $0 < L = T - a < \infty$. Assume smoothness and, on each compact slab $[a,b] \subset [a,T)$, $u \in C^1([a,b]; H^m(\mathbb{R}^3))$ and $f, \nabla p \in C([a,b]; H^m(\mathbb{R}^3))$ for every integer $m \geq 0$. These hypotheses justify the following strong energy identities and integrations by parts. Write

$$E(t) = \|u(t)\|_2^2, \quad Y(t) = \|\nabla u(t)\|_2^2 = \|\omega(t)\|_2^2, \quad \omega = \operatorname{curl} u.$$

Suppose

$$E(t) \leq M, \quad \|f(t)\|_2 \leq F_0, \quad \|\Delta f(t)\|_2 \leq F_\Delta \qquad (a \leq t < T).$$

All constants in this note are independent of the shorter endpoint t. The kinetic-energy identity gives

$$\int_a^t Y(s)\,\mathrm{d}s \leq B, \qquad B = \frac{E(a)}{2\nu} + \frac{L\sqrt{M}F_0}{\nu}. \tag{1}$$

Indeed, $E'/2 = -\nu Y + \langle u, f\rangle$ and $E(t) \geq 0$.

Fix an angular Fourier cutoff $K \geq 0$, and set

$$u_L = \mathbf{1}_{\{|\xi|<K\}}(D)u, \quad u_H = u - u_L, \quad \omega_H = \operatorname{curl} u_H, \quad S_H = \tfrac{1}{2}(\nabla u_H + \nabla u_H^{\mathsf{T}}).$$

These are orthogonal L^2 projections commuting with derivatives. Define the physical tail quantities

$$Z_H = \tfrac{1}{2}\|\omega_H\|_2^2, \qquad D_H = \|\nabla\omega_H\|_2^2 = \|\Delta u_H\|_2^2, \qquad \Pi_H = \int_{\mathbb{R}^3} \omega_H \cdot S_H\omega_H \,\mathrm{d}x.$$

In particular, Π_H uses the high field in all three factors. With $\Pi = \int \omega \cdot S\omega$, introduce

$$P_L = \langle -\Delta u_L, u \times \omega\rangle, \quad P_H = \langle -\Delta u_H, u \times \omega\rangle, \quad X_H = P_H - \Pi_H, \quad F_H = \langle -\Delta u_H, f\rangle.$$

Then $P_L + P_H = \Pi$, and the exact tail balance is

$$Z_H' = \Pi_H + X_H - \nu D_H + F_H. \tag{2}$$

The filtered velocity is not assumed to solve an autonomous Navier–Stokes equation: X_H retains every lower-frequency input.

Proposition 1. *Let $c_K = (K^3/(6\pi^2))^{1/2}$. Under the preceding assumptions,*

$$\int_a^t |F_H(s)| \,\mathrm{d}s \leq L\sqrt{M}F_\Delta, \tag{3}$$

$$\int_a^t |X_H(s)| \,\mathrm{d}s \leq 3Kc_K\sqrt{M}B + K^2c_KM\sqrt{LB} =: C_L. \tag{4}$$

Neither bound uses a gradient supremum or tail dissipation of the velocity.

Proof. Integration by parts gives $F_H = -\langle u_H, \Delta f\rangle$, so $|F_H| \leq \sqrt{M}F_\Delta$. This proves (3).

For the unitary Fourier convention, Cauchy–Schwarz on the ball gives

$$\|\nabla u_L\|_\infty \leq Kc_K\sqrt{E}, \qquad \|\Delta u_L\|_\infty \leq K^2c_K\sqrt{E}.$$

Here the gradient uses the Frobenius norm, and $c_K^2 = (2\pi)^{-3}|B_K|$. The vector and matrix bounds follow by Cauchy–Schwarz before taking a component sum.

For matrices define the trilinear form

$$b(A,B,C) = \sum_{i,j,k} A_{ji}B_{kj}C_{ki}, \qquad |b(A,B,C)| \leq |A|_F|B|_F|C|_F.$$

The integrated stretching identity is $\Pi = -\int b(\nabla u, \nabla u, \nabla u)$, and the same identity holds for the smooth solenoidal field u_H. Put $A = \nabla u$, $H = \nabla u_H$ and $G = \nabla u_L = A - H$. The exact telescoping identity

$$b(A,A,A) - b(H,H,H) = b(G,A,A) + b(H,G,A) + b(H,H,G)$$

and $\|H\|_2 \leq \|A\|_2$ imply

$$|\Pi - \Pi_H| \leq 3\|\nabla u_L\|_\infty Y \leq 3Kc_K\sqrt{M}\,Y. \tag{5}$$

Also

$$|P_L| \leq \|\Delta u_L\|_\infty \|u\|_2 \|\omega\|_2 \leq K^2c_KM\sqrt{Y}.$$

Now $X_H = \Pi - \Pi_H - P_L$. Integrate these two bounds and use (1) and $\int_a^t \sqrt{Y} \leq \sqrt{LB}$. □

Corollary 2. *If a finite constant A satisfies*

$$\int_a^t (\Pi_H - \nu D_H)\,\mathrm{d}s \le A \qquad (a \le t < T), \tag{6}$$

then the complete gradient energy obeys

$$Y(t) \le 2Z_H(a) + 2A + 2C_L + 2L\sqrt{M}F_\Delta + K^2 M.$$

In particular, a half-dissipation budget implies this conclusion, but the strictly smaller fraction is unnecessary for this fixed-cutoff gradient-energy implication.

Proof. Integrate (2), apply Proposition 1, and use $Y = 2Z_H + \|\nabla u_L\|_2^2 \le 2Z_H + K^2 M$. □

For an unforced solution, one can take $M = E(a)$ and $B = E(a)/(2\nu)$. Thus the low-frequency allowance is controlled by the datum, viscosity, cutoff and horizon. This does not prove (6): its uniform signed high-frequency history remains the open estimate. Allowing the full dissipation simply exposes its relation to the gradient-energy continuation criterion; it is not a new coercive estimate.

2 Application to the actual forced construction

The whole-space construction in [1] has smooth compactly supported forcing, bounded kinetic energy, zero initial velocity, and unbounded speed as $t \uparrow 1$. On slabs exhausting $[1/2, 1)$, all hypotheses above hold with fixed M, F_0, F_Δ.

The module `MettapediaNS.ForcedSignedBudget` in the Lean library applies directly to the construction's `CandidateProperties`. It proves that its gradient energy cannot have a uniform bound: such a bound would control the order-two energy via the forced viscous estimate, and the H^2 embedding would bound speed. It also proves the full-field signed-production obstruction directly, with force work bounded by $6\sqrt{E_0(u)}\sqrt{E_2(f)}$ in the doubled energy convention, where $E_j(v) = \|\nabla^j v\|_2^2$ sums all ordered derivative components. The constructor theorem is imported at the pinned revision `f9e8bc5b38b6e212696e8a30e3e91517af887bbd`.

Corollary 3. *For every such candidate, every fixed finite $K \ge 0$, and every $\beta \le 1$,*

$$\sup_{1/2 \le t < 1} \int_{1/2}^t (\Pi_H(s) - \beta\nu D_H(s))\,\mathrm{d}s = +\infty.$$

Meanwhile $\int_{1/2}^1 |X_H|$ and $\int_{1/2}^1 |F_H|$ are finite.

Proof. For $\beta = 1$, a finite upper bound would give a uniform gradient-energy bound by Corollary 2, contradicting the preceding forced-candidate theorem. For $\beta \le 1$, the accumulated quantity only increases because $D_H \ge 0$. The other assertions follow from Proposition 1. □

Thus the signed tail self-interaction is the term that defeats the budget; an infinite direct forcing-work allowance is not the cause. This conclusion applies to the complete constructed solution, without replacing its oscillatory corrections by a finite numerical model.

Forcing is still essential to that constructed history. With its zero datum and zero force, the kinetic-energy identity forces the smooth finite-energy solution to remain zero. In the construction the force is the smoothly extended momentum residual of the designed field (Section 10 of [1]). Removing it while keeping that field would change the equation. Restarting an unforced evolution from a later nonzero slice is a different problem, whose subsequent history is not settled by this argument. A proposed unforced budget must therefore use more than bounded forcing norms or bounded direct force work.

3 Forcing in a caloric stretching functional

The preceding obstruction can also test a functional that records stretching beyond the instantaneous enstrophy. Let

$$A = -\Delta, \qquad H_s = e^{-\nu s A}, \qquad R = \operatorname{curl} A^{-1} P, \qquad \mathcal{M}(v) = R\big((\operatorname{curl} v \cdot \nabla) v\big),$$

where P is the whole-space Leray projection. The multiplier R is used on a divergence: $\mathcal{M}(v) = R \operatorname{div}(v \otimes \operatorname{curl} v)$. Thus its low-frequency singularity causes no L^2 difficulty. Parseval's identity gives $\|R \operatorname{div} Q\|_2 \le \|Q\|_2$, where the tensor norm is Frobenius. For solenoidal v,

$$\|\mathcal{M}(v)\|_2 \le \|v\|_\infty \|\nabla v\|_2 , \qquad \langle Av, \mathcal{M}(v)\rangle = \Pi(v). \tag{7}$$

Define the nonnegative functional

$$\mathcal{K}(u) = \frac{1}{\nu} \int_0^\infty \|\mathcal{M}(H_s u)\|_2^2 \, \mathrm{d}s. \tag{8}$$

It is finite for the smooth L^2 fields considered here: the integrand is bounded near zero and is $O(s^{-5/2})$ at infinity by heat smoothing.

The effective force in the projected velocity equation is $g = Pf$, not necessarily f itself. Suppose in addition that

$$\|g(t)\|_\infty \le Q_0, \qquad \|\operatorname{curl} g(t)\|_\infty \le Q_1 \qquad (a \le t < T).$$

These constants are finite for the smooth compactly supported force in the construction: P is bounded on every H^m, Sobolev embedding controls Pf, and $\operatorname{curl} Pf = \operatorname{curl} f$. No L^∞ boundedness of P is assumed.

Proposition 4. *Put $c_h = (8\pi)^{-3/4}$. For a fixed smooth solenoidal slice u and direction g as above, with $E = \|u\|_2^2$ and $Y = \|\nabla u\|_2^2$,*

$$|D\mathcal{K}(u)[g]| \le \frac{c_h}{\nu^2} \left(\frac{32}{3} Q_0 E^{3/4} Y^{3/4} + 16 Q_1 E^{5/4} Y^{1/4} \right). \tag{9}$$

Consequently the standing evolution satisfies, uniformly for $t < T$,

$$\int_a^t |D\mathcal{K}(u)[g]| \, \mathrm{d}s \le C_{\mathcal{K}} := \frac{c_h}{\nu^2} \left(\frac{32}{3} Q_0 M^{3/4} L^{1/4} B^{3/4} + 16 Q_1 M^{5/4} L^{3/4} B^{1/4} \right). \tag{10}$$

This payment spends no order-two velocity dissipation.

Proof. Use the auxiliary heat parameter $r = \nu s$, and write $w_r = e^{-rA} u$, $h_r = e^{-rA} g$. Differentiating both parent slots gives

$$D\mathcal{M}(w)[h] = R \operatorname{div}(h \otimes \operatorname{curl} w + w \otimes \operatorname{curl} h).$$

The multiplier bound, heat contraction and (7) imply

$$|D\mathcal{K}(u)[g]| \le \frac{2}{\nu^2} \int_0^\infty \|w_r\|_\infty \|\nabla w_r\|_2 \left(Q_0 \|\nabla w_r\|_2 + Q_1 \sqrt{E} \right) \mathrm{d}r.$$

The Gaussian kernel has L^2 norm $c_h r^{-3/4}$, and its Fourier multiplier gives

$$\|w_r\|_\infty \le c_h r^{-3/4} \sqrt{E}, \qquad \|\nabla w_r\|_2 \le \min\{\sqrt{Y}, r^{-1/2}\sqrt{E}\}.$$

When $E, Y > 0$, split each integral at $r_* = E/Y$. Direct integration yields

$$\int_0^\infty \|w_r\|_\infty \|\nabla w_r\|_2^2 \, \mathrm{d}r \le \frac{16}{3} c_h E^{3/4} Y^{3/4},$$
$$\sqrt{E} \int_0^\infty \|w_r\|_\infty \|\nabla w_r\|_2 \, \mathrm{d}r \le 8 c_h E^{5/4} Y^{1/4}.$$

These bounds also justify differentiation under the heat integral for the smooth slices. If E or Y vanishes, $u = 0$ as an L^2 whole-space field and the result is immediate. This proves (9). Finally apply Hölder in physical time, using $E \le M$, $\int Y \le B$, and the interval length at most L. □

Let $N(u) = -P((u \cdot \nabla)u)$ be the actual unforced velocity source, and set

$$\mathcal{G} = \frac{Y + \mathcal{K}}{2}, \qquad \mathcal{T} = D\mathcal{K}(u)[N(u)], \qquad \mathcal{W} = \nu \|Au\|_2^2 + \frac{1}{\nu} \|\mathcal{M}(u) - \nu Au\|_2^2 .$$

In particular $\mathcal{G} \ge Y/2$ and $\mathcal{W} \ge 0$. Heat translation in (8) gives $D\mathcal{K}(u)[-\nu Au] = -\|\mathcal{M}(u)\|_2^2/\nu$. Combine this with the actual forced enstrophy identity and expand the square in $\mathcal{W}$ to obtain the exact balance

$$2\mathcal{G}' + \mathcal{W} = \mathcal{T} + D\mathcal{K}(u)[g] + 2\langle Au, f\rangle. \tag{11}$$

Both extra force terms have absolutely bounded physical-time integrals: their sum is bounded by $C_{\mathcal{K}} + 2L\sqrt{M} F_\Delta$.

Corollary 5. *On every forced candidate of the preceding section, for every $\beta \le 1$,*

$$\sup_{1/2 \le t < 1} \int_{1/2}^t (\mathcal{T} - \beta \mathcal{W}) \, \mathrm{d}s = +\infty.$$

Proof. Integrating (11) expresses $\int(\mathcal{T} - \mathcal{W})$ as $2\mathcal{G}(t) - 2\mathcal{G}(1/2)$ minus a uniformly bounded force integral. The candidate's gradient energy has no uniform upper bound, and $\mathcal{G} \ge Y/2$. This proves the case $\beta = 1$; decreasing β only increases the integral since $\mathcal{W} \ge 0$. □

Thus the smooth force does not hide an infinite payment in the derivative of this quartic functional. Its signed nonlinear renewal is itself unbounded relative to its positive cost on the forced history. A proof of the corresponding unforced budget must exploit the zero-force evolution; estimating these explicit force terms is insufficient. The estimate does not bound $\mathcal{T}$: substituting $N(u)$ for g would require velocity-dependent supremum norms that are not controlled by the energy inequality.

4 Low parents and high-frequency caloric renewal

The caloric return admits a further frequency separation. Fix $K \ge 0$, write $v = P_{\ge K} u$, $\ell = u - v$, and retain $E = \|u\|_2^2$, $Y = \|\nabla u\|_2^2$, $D = \|\Delta u\|_2^2$. Choose a universal homogeneous Sobolev constant S such that $\|w\|_6 \le S \|\nabla w\|_2$ for real scalar, vector and matrix fields on $\mathbb{R}^3$. The constant c_K is as in Proposition 1.

Proposition 6. *If $E \le M$, then for every $\varepsilon > 0$,*

$$|\mathcal{T}(u) - \mathcal{T}(v)| \le \varepsilon \nu D + C_2 Y^2 + C_{3/2} Y^{3/2}, \tag{12}$$

where all constants depend only on K, ν, M, ε and S:

$$a_K = \frac{2c_K S^3}{\nu^2}, \qquad b_K = \frac{6c_K S^3}{\nu^2}, \qquad C_2 = a_K\sqrt{M} + \frac{3b_K^{4/3}M}{4(4\varepsilon\nu)^{1/3}}, \qquad C_{3/2} = a_K K M.$$

In particular, the two last terms in (12) are at most $2(C_2 Y + C_{3/2}\sqrt{Y})\mathcal{G}$. Their coefficient is integrable in physical time under the energy bounds.

Proof. We first bound the complete quartic Fourier kernel of $\mathcal{K}$. For ordered inputs a, b let $\mathcal{M}_b(a,b) = R((\operatorname{curl} a \cdot \nabla)b)$. At frequencies p, q with output $k = p + q$, its coefficient satisfies

$$|\widehat{\mathcal{M}_b}(p,q)| \le |p|\,|\widehat{a}(p)|\,|\widehat{b}(q)|. \tag{13}$$

Indeed, $p \times \widehat{a}(p)$ is perpendicular to p, so its dot product with q equals its dot product with k. This cancels the $|k|^{-1}$ multiplier of R. The zero output is zero as well. Thus no lower bound on the output frequency is needed.

In an ordered quartic pairing of two such sources, heat integration gives the denominator $\nu^2(|p|^2 + |q|^2 + |r|^2 + |s|^2)$. Its kernel is bounded by

$$\frac{|p|\,|r|}{\nu^2(|p|^2 + |q|^2 + |r|^2 + |s|^2)} \le \frac{1}{2\nu^2}. \tag{14}$$

All parent pairs and all interference between their heat rates are included.

For a real field a, let $a^{\#}$ be the real scalar field with Fourier transform $|\widehat{a}|$. It has the same L^2 norm and homogeneous Sobolev norms as a. Its physical values need not be nonnegative. The absolute ordered quartic pairing is bounded by $1/(2\nu^2)$ times the nonnegative Fourier convolution corresponding to $\int a^{\#}b^{\#}c^{\#}d^{\#}$. Applying Hölder and Sobolev after this Fourier majorization gives

$$|D\mathcal{K}(u)[h]| \le \frac{2}{\nu^2}\left\|u^{\#}\right\|_6^3 \|h\|_2 \le \frac{2S^3}{\nu^2} Y^{3/2}\|h\|_2. \tag{15}$$

There are four differentiated slots. Subtracting the derivatives at u and v telescopes each of the three other slots, giving twelve terms. Since v is a Fourier restriction of u, coefficient majorization yields

$$\begin{aligned} |(D\mathcal{K}(u) - D\mathcal{K}(v))[h]| &\le \frac{6}{\nu^2}\left\|\ell^{\#}\right\|_\infty \left\|u^{\#}\right\|_4^2 \|h\|_2 \\ &\le \frac{6c_K S^{3/2}}{\nu^2} E^{3/4}Y^{3/4}\|h\|_2. \end{aligned} \tag{16}$$

The coefficient comparison takes place inside the Fourier convolution; it does not assert a pointwise inequality between $|v^{\#}|$ and $|u^{\#}|$. Here we used $\left\|\ell^{\#}\right\|_\infty \le c_K\sqrt{E}$ and interpolation $\left\|u^{\#}\right\|_4^2 \le S^{3/2}E^{1/4}Y^{3/4}$.

Write $\mathcal{B}(a,b) = -P((a\cdot\nabla)b)$. Then

$$N(u) - N(v) = \mathcal{B}(\ell, u) + \mathcal{B}(v, \ell), \qquad \|N(u) - N(v)\|_2 \le c_K\sqrt{EY} + Kc_K E.$$

Also Sobolev and interpolation give

$$\|N(v)\|_2 \le \|v\|_6\,\|\nabla v\|_3 \le S^{3/2}Y^{3/4}D^{1/4}.$$

Only the L^2 contraction and derivative commutation of the sharp projector are used; no general L^p bound for the ball multiplier is assumed. Split the return exactly as

$$\mathcal{T}(u) - \mathcal{T}(v) = D\mathcal{K}(u)[N(u) - N(v)] + (D\mathcal{K}(u) - D\mathcal{K}(v))[N(v)].$$

Equations (15)–(16) bound these terms by

$$a_K(E^{1/2}Y^2 + KEY^{3/2}) + b_K E^{3/4}Y^{3/2}D^{1/4}.$$

Young's inequality with exponents 4 and 4/3 bounds the last term by $\varepsilon\nu D + 3b_K^{4/3}EY^2/(4(4\varepsilon\nu)^{1/3})$. Use $E \le M$ and $Y \le 2\mathcal{G}$ to finish. □

Low renewal outputs are paid separately, even when both source parents are arbitrarily high. Set

$$g_K = P_{<K}N(v), \qquad \mathcal{T}_K(u) = D\mathcal{K}(v)[P_{\ge K}N(v)].$$

The full source $N(v)$ is formed before splitting its output. Cauchy–Schwarz in the unitary Fourier convolution gives $|\widehat{N(v)}(\xi)| \le (2\pi)^{-3/2}|\xi|E$. Integrating this over the ball, with an extra $|\xi|$ for curl, proves

$$\|g_K\|_\infty \le \frac{K^4 E}{8\pi^2}, \qquad \|\operatorname{curl} g_K\|_\infty \le \frac{K^5 E}{10\pi^2}. \tag{17}$$

The derivative is the output frequency because the nonlinearity is in divergence form; this estimate assumes no bound on parent gradients. Proposition 4, applied at the slice v, therefore gives

$$\int_a^t |D\mathcal{K}(v)[g_K]|\,\mathrm{d}s \le C_{\mathrm{out}}, \tag{18}$$

where C_{out} is the right side of (10) with $Q_0 = K^4M/(8\pi^2)$ and $Q_1 = K^5M/(10\pi^2)$. Although v is not an autonomous solution, its energy and integrated gradient energy are bounded by those of u, which is all that this application uses.

Corollary 7. *Suppose $A_0 \ge 0$ and $\theta < 1$ satisfy, on every shorter prefix,*

$$\int_a^t \mathcal{T}_K(u)\,\mathrm{d}s \le A_0 + \theta\int_a^t \mathcal{W}(u)\,\mathrm{d}s. \tag{19}$$

Choose $0 < \varepsilon < 1-\theta$, and let $C_f = C_{\mathcal{K}} + 2L\sqrt{M}F_\Delta$ for the forced equation, or $C_f = 0$ when $f = 0$. Then

$$\mathcal{G}(t) \le \left(\mathcal{G}(a) + \frac{A_0 + C_{\mathrm{out}} + C_f}{2}\right)\exp\Big(C_2B + C_{3/2}\sqrt{LB}\Big).$$

Proof. The exact split is $\mathcal{T}(u) = \mathcal{T}_K(u) + \mathcal{T}(u) - \mathcal{T}(v) + D\mathcal{K}(v)[g_K]$. Insert this in (11), integrate, and use Propositions 4 and 6. Since $\nu D \le \mathcal{W}$, the remaining cost has coefficient $1 - \theta - \varepsilon > 0$ and can be dropped. Grönwall applies with coefficient $C_2Y + C_{3/2}\sqrt{Y}$, whose integral is at most $C_2B + C_{3/2}\sqrt{LB}$. □

For an unforced regularity argument, K and A_0 in (19) must be chosen from the full initial datum, viscosity and a prescribed finite horizon, uniformly over every shorter lifespan and prefix. That high-frequency signed-history estimate is not proved here. The result pays low parents and low outputs; it does not pay repeated high-frequency regeneration.

The same estimates also sharpen the forced test: on every actual forced candidate, for each fixed K and every $\theta < 1$,

$$\sup_{1/2\le t<1} \int_{1/2}^{t} (\mathcal{T}_K(u) - \theta \mathcal{W}(u))\,\mathrm{d}s = +\infty.$$

A finite upper bound would give a uniform gradient-energy bound by Corollary 7, contradicting the constructed candidate. Unlike Corollary 5, this argument requires $\theta < 1$ because the low-parent estimate spends an arbitrarily small, positive part of the dissipation.

5 Low stretching outputs along the actual history

The output of $\mathcal{M}$ is distinct from the output of the nonlinear direction N. It too can be separated without restricting its parents. Fix $J \ge 0$, put $Q_- = P_{<J}$, $Q_+ = P_{\ge J}$, and define

$$\mathcal{K}_\pm(u) = \frac{1}{\nu}\int_0^\infty \|Q_\pm \mathcal{M}(H_s u)\|_2^2\,\mathrm{d}s, \qquad \mathcal{Z}_\pm(u) = \frac{1}{\nu}\|Q_\pm \mathcal{M}(u)\|_2^2, \qquad \mathcal{T}_\pm(u) = D\mathcal{K}_\pm(u)[N(u)].$$

Orthogonality gives $\mathcal{K} = \mathcal{K}_- + \mathcal{K}_+$ and $\mathcal{T} = \mathcal{T}_- + \mathcal{T}_+$. All source outputs are formed before the projection is applied.

Proposition 8. *With $c_J^2 = J^3/(6\pi^2)$,*

$$0 \le \mathcal{K}_-(u) \le \frac{c_J^2 E^2}{4\nu^2}, \qquad 0 \le \mathcal{Z}_-(u) \le \frac{c_J^2 EY}{\nu}. \tag{20}$$

Along the actual forced evolution,

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{K}_-(u) = -\mathcal{Z}_-(u) + \mathcal{T}_-(u) + D\mathcal{K}_-(u)[g]. \tag{21}$$

Consequently, uniformly for all shorter prefixes,

$$\left|\int_a^t \mathcal{T}_-(u)\,\mathrm{d}s\right| \le \frac{c_J^2 M^2}{4\nu^2} + \frac{c_J^2 MB}{\nu} + C_{\mathcal{K}}.$$

The last term is zero for an unforced evolution. This bounds the signed primitive, not the integral of $|\mathcal{T}_-|$.

Proof. The ordered bound (13) and Cauchy–Schwarz in the full convolution give, at every nonzero output k,

$$|\widehat{\mathcal{M}(u)}(k)| \le (2\pi)^{-3/2}\int |p|\,|\widehat{u}(p)|\,|\widehat{u}(k-p)|\,\mathrm{d}p \le (2\pi)^{-3/2}\sqrt{EY}.$$

Integrating its square over $|k| < J$ proves the $\mathcal{Z}_-$ bound. No upper bound on the parent frequencies was used. If $E_s = \|H_s u\|_2^2$ and $Y_s = \|\nabla H_s u\|_2^2$, then $E_s' = -2\nu Y_s$, so

$$\mathcal{K}_-(u) \le \frac{c_J^2}{\nu}\int_0^\infty E_s Y_s\,\mathrm{d}s \le \frac{c_J^2 E^2}{4\nu^2}.$$

Heat translation gives $D\mathcal{K}_-(u)[-\nu Au] = -\mathcal{Z}_-(u)$. The actual equation therefore proves (21). Proposition 4 applies to $\mathcal{K}_-$ with unchanged constants: inserting an orthogonal output projection in both factors of its derivative can only improve the two L^2 norm bounds. Integrate the identity, use $|\mathcal{K}_-(u(t)) - \mathcal{K}_-(u(a))| \le c_J^2 M^2/(4\nu^2)$, and use $\int \mathcal{Z}_- \le c_J^2 MB/\nu$. □

For the unforced equation there is a sharper payment, independent even of the horizon. The actual kinetic-energy equality is $E' = -2\nu Y$, so (20) gives

$$\mathcal{K}_-(u(t)) + \int_a^t \mathcal{Z}_-(u)\,\mathrm{d}s \le \frac{c_J^2 E(a)^2}{4\nu^2}. \tag{22}$$

Indeed, the bound for the integral is $c_J^2(E(a)^2 - E(t)^2)/(4\nu^2)$, which complements the bound for the remaining capacity. Therefore every unforced prefix satisfies

$$-\mathcal{K}_-(u(a)) \le \int_a^t \mathcal{T}_-(u)\,\mathrm{d}s \le \frac{c_J^2 E(a)^2}{4\nu^2} - \mathcal{K}_-(u(a)).$$

This is a bound on repeated low-output renewal from the initial kinetic energy; its constant grows with the fixed output cutoff.

There is a more direct way to use this payment in the regularity target. Keep the original, full positive cost $\mathcal{W}$, and set $\mathcal{G}_+ = (Y + \mathcal{K}_+)/2 \ge Y/2$. Subtracting (21) from (11) gives the exact identity

$$2\mathcal{G}_+' + \mathcal{W} = \mathcal{T}_+ + \mathcal{Z}_- + D\mathcal{K}_+(u)[g] + 2\langle Au, f\rangle. \tag{23}$$

Thus only the spent low-output cost, bounded by $C_{\mathrm{src}} = c_J^2 MB/\nu$, is needed on the right. The endpoint $\mathcal{K}_-$ allowance is unnecessary in this identity. In the unforced case one may take $C_{\mathrm{src}} = c_J^2 E(a)^2/(4\nu^2)$ by (22).

With the input cutoff K from the preceding section, define

$$v = P_{\ge K} u, \qquad \mathcal{T}_{K,J}(u) = D\mathcal{K}_+(v)[P_{\ge K} N(v)].$$

The estimates of Proposition 6 hold with $\mathcal{K}_+$ in place of $\mathcal{K}$: its ordered quartic kernel is restricted at the common output, and has the same bound. The force estimate and the low nonlinear-output payment hold as well. Consequently, if fixed $A_0 \ge 0$ and $\theta < 1$ obey

$$\int_a^t \mathcal{T}_{K,J}(u)\,\mathrm{d}s \le A_0 + \theta \int_a^t \mathcal{W}(u)\,\mathrm{d}s \tag{24}$$

on every shorter prefix, the same Grönwall argument gives

$$\mathcal{G}_+(t) \le \left(\mathcal{G}_+(a) + \frac{A_0 + C_{\mathrm{src}} + C_{\mathrm{out}} + C_f}{2}\right) \exp\left(C_2 B + C_{3/2}\sqrt{LB}\right).$$

Here $0 < \varepsilon < 1 - \theta$ defines $C_2, C_{3/2}$ as before. The exact decomposition used in (23) is

$$\begin{aligned}\mathcal{T}_+(u) = \mathcal{T}_{K,J}(u) &+ D\mathcal{K}_+(u)[N(u) - N(v)] \\ &+ (D\mathcal{K}_+(u) - D\mathcal{K}_+(v))[N(v)] + D\mathcal{K}_+(v)[P_{<K} N(v)].\end{aligned}$$

In particular, no autonomous evolution is assigned to the filtered field v.

Taking $J = K$ leaves a quintic term whose five original velocity inputs, nonlinear-direction output, and stretching output are all above the same fixed cutoff. Its physical-time bound (24) remains open, with the same initial-data and shorter-lifespan quantifiers as (19). On the actual forced construction it fails for every fixed K, J and every $\theta < 1$: a finite upper allowance would bound $\mathcal{G}_+ \ge Y/2$, contrary to the candidate's unbounded gradient energy. This further separation pays low stretching outputs; it does not bound regeneration among the remaining high frequencies.

6 Unforced activation with unit initial energy

The allowance in (24) is permitted to depend on the full initial datum. That permission is substantive even when the initial tail is empty. In this section a subscript Q on a projector means $P_Q = \mathbf{1}_{\{|\xi|\ge Q\}}(D)$; the superscript on $\mathcal{T}^{\nu}_{Q,Q}$ or $\mathcal{W}^{\nu}$ specifies the viscosity in the functional. In particular, $\mathcal{W}^{\nu}$ is the original full-field cost, not a cost computed from $P_Q u$ alone.

Proposition 9. *For every $\nu, H, \varepsilon > 0$, $\theta \ge 0$ and finite $B \ge 0$, there are a real solenoidal Schwartz datum u_0 on $\mathbb{R}^3$, a cutoff $Q_* > 0$, and an actual smooth unforced solution from u_0 on a prefix $[0, t_*]$, $0 < t_* < H$, such that*

$$\|u_0\|_2^2 = 1, \qquad (u_0)_3 = 0, \qquad \operatorname{supp} \widehat{u}_0 \subset \{\xi : 0 < |\xi| < Q_*,\ |\xi_3| \le \varepsilon|\xi|\},$$

but

$$\int_0^{t_*} \mathcal{T}^{\nu}_{Q_*,Q_*}(u(t))\,\mathrm{d}t > B + \theta \int_0^{t_*} \mathcal{W}^{\nu}(u(t))\,\mathrm{d}t. \tag{25}$$

Thus an energy-only allowance uniform over the datum and the prescribed cutoff is false. The cutoff varies in this construction; the proposition does not rule out a cutoff-dependent allowance, an allowance using the full datum, or the existence of a different cutoff for that datum.

Proof. We first obtain a positive first-generation coefficient on the torus, then localize it, and finally use a common local smooth time interval.

Discrete coefficient. On the 2π torus with normalized volume, write $Jk = (-k_2, k_1, 0)$. The real trigonometric field with coefficients

$$\widehat{w}(k) = i a_k Jk, \qquad (k, a_k) \in \{((3,1,1), \tfrac{1}{2}), ((1,3,1), 1), ((1,-1,3), \tfrac{1}{2})\},$$

and their reality partners has no tail above $Q^2 = 14$. For its actual first tail derivative $v_1 = P_Q N(w)$, finite Fourier convolution and integration of all parent heat rates give

$$\mathcal{T}^{1}_{Q,Q}(v_1) = \frac{15667851179107579904}{95414925477178125} > 0. \tag{26}$$

Every parent has squared radius 11. The internal output centers have even coordinates and hence squared radii divisible by 4; none lies on the cutoff 14. Using the cutoff 16 produces the same value.

For arbitrary thinness, use instead

$$p_m = (m,3,1), \quad q_m = (3,m,1), \quad r_m = (m,1,3), \qquad Q_m^2 = 3m^2/2,$$

with planar coefficients $i a_k Jk/m$, weights $(2,1,1)$ and reality partners. The parent radii are $\sqrt{m^2+10}$ and their axial ratios are at most $3/m$. Keeping all frequency labels distinct, the same finite rational calculation gives

$$\mathcal{T}^{1}_{Q_m,Q_m}(P_{Q_m} N(w_m)) = \frac{52224}{25m^3} - \frac{7422976}{375m^4} + O(m^{-5}). \tag{27}$$

One way to check this identity is to put $\delta = 1/m$ and represent each scaled frequency as $a + b\delta$. The scaled return has coefficients zero through order eight, followed by $52224/25$ and $-7422976/375$ at orders nine and ten. Restoring the frequencies and the first generated velocity multiplies it by m^6. Every denominator has nonzero constant term. The finitely many projector decisions are stable for small δ: their leading squared radii are integers, whereas the scaled cutoff squared is $3/2$. These are rational identities, not an assumed invariant finite-mode evolution. Fix a sufficiently large m to make the coefficient positive and the axial ratio strictly smaller than ε.

Whole-space localization. Denote the chosen periodic parent by $w_{\mathrm{per}} = J\nabla\psi$, where $J\nabla = (-\partial_2, \partial_1, 0)$, and its cutoff by Q. Choose a nonzero nonnegative Schwartz function ϕ with compact Fourier support. Such a function is the square of the real inverse Fourier transform of a nonzero real even function in C_c^∞. Set

$$w_L(x) = J\nabla(\phi(x/L)\psi(x)), \qquad v_L = P_Q N(w_L).$$

This datum is exactly solenoidal and has zero axial velocity. Its Fourier support lies in $O(L^{-1})$ neighborhoods of the parent centers; for sufficiently large L it lies strictly below Q, away from zero, and within the required Fourier belt.

The degree-ten functional $\mathcal{T}^1_{Q,Q}(v_L)$ has a finite packet expansion. Each sharp projector is evaluated near one of the finitely many internal centers, all separated from its cutoff. Each retained Leray or inverse-curl multiplier is separated from zero. Nonresonant terms vanish for large L, since the sum of their Fourier supports cannot contain zero. In a resonant term, rescale the Fourier offsets by L. The multiplier converges uniformly to its value at the centers. The parent heat frequencies are bounded away from zero, so the heat integrals are dominated by a polynomial in heat time times e^{-cs} for some $c > 0$. The compact envelope spectra then justify dominated convergence term by term. It follows that

$$\begin{aligned} L^{-3}\mathcal{T}^1_{Q,Q}(v_L) &\longrightarrow \mathcal{T}^1_{Q,Q,\mathrm{per}}(P_Q N(w_{\mathrm{per}})) \int_{\mathbb{R}^3} \phi^{10} > 0, \\ L^{-3}\,\|w_L\|_2^2 &\longrightarrow \|w_{\mathrm{per}}\|_{2,\mathrm{per}}^2 \int_{\mathbb{R}^3} \phi^2 > 0. \end{aligned} \tag{28}$$

The extra velocity term from differentiating $\phi(x/L)$ is of order L^{-1} and has zero limit in these normalizations. The tenth power in (28) retains the actual quadratic generation of the child followed by the quintic renewal functional.

Fix a sufficiently large L and normalize w_L by its exact L^2 norm, obtaining a Schwartz datum w_0 with squared norm one and $P_Q w_0 = 0$. Its child $v_1 = P_Q N(w_0)$ satisfies $c = \mathcal{T}^1_{Q,Q}(v_1) > 0$; the normalization multiplies the coefficient by $\|w_L\|_2^{-10}$.

An actual common time interval. For $0 \le \delta \le 1$, let w^δ solve the unforced equation with viscosity δ and initial datum w_0. Local high-Sobolev energy estimates discard the nonpositive viscous term, giving a common smooth interval and uniform high-Sobolev bounds. Applying the equation twice also bounds two time derivatives in H^2, uniformly in δ. Since the initial tail vanishes,

$$P_Q w^\delta(\tau) = \tau v_1 + O(\tau^2) \quad \text{in } H^2,$$

uniformly for $\delta \in [0,1]$. The quartic kernel bound above gives $|D\mathcal{K}_+(v)[h]| \le C\,\|\nabla v\|_2^3\,\|h\|_2$ at viscosity one, and $N : H^2 \to L^2$ is continuous. Hence the quintic $\mathcal{T}^1_{Q,Q}$ is continuous on H^2, with bounded multilinear coefficients there, and

$$\mathcal{T}^1_{Q,Q}(w^\delta(\tau)) = c\tau^5 + O(\tau^6)$$

uniformly in δ. Choose a sufficiently small fixed $\tau_* > 0$. There are $\kappa, C_0 > 0$, independent of δ, such that

$$\int_0^{\tau_*} \mathcal{T}^1_{Q,Q}(w^\delta)\,\mathrm{d}\tau \ge \kappa, \tag{29}$$

$$\int_0^{\tau_*} \left[\delta^2 \left\|Aw^\delta\right\|_2^2 + \left\|M(w^\delta) - \delta A w^\delta\right\|_2^2\right] \mathrm{d}\tau \le C_0. \tag{30}$$

Here $A = -\Delta$. The second assertion follows directly from the common H^2 bound and $\|M(w)\|_2 \le \|w\|_\infty \|\operatorname{curl} w\|_2$. The time τ_* is fixed only after m, L, w_0 have been fixed; no numerical value or uniformity in the thinness parameter is asserted.

Energy-preserving concentration. For sufficiently large λ, set

$$\delta = \nu/\sqrt{\lambda}, \quad u_\lambda(t,x) = \lambda^{3/2} w^\delta(\lambda^{5/2}t, \lambda x), \quad Q_\lambda = \lambda Q, \quad t_\lambda = \tau_*/\lambda^{5/2}.$$

This is the actual unforced equation with physical viscosity ν; its initial squared L^2 norm is one. Its initial tail is empty above Q_λ, and the initial polarization and belt condition persist under this spatial rescaling. Keeping the spatial Jacobian and the physical-time factor gives the exact identities

$$\int_0^{t_\lambda} \mathcal{T}^\nu_{Q_\lambda, Q_\lambda}(u_\lambda)\,\mathrm{d}t = \frac{\lambda^3}{\nu^2} \int_0^{\tau_*} \mathcal{T}^1_{Q,Q}(w^\delta)\,\mathrm{d}\tau, \tag{31}$$

$$\int_0^{t_\lambda} \mathcal{W}^\nu(u_\lambda)\,\mathrm{d}t = \frac{\lambda^{5/2}}{\nu} \int_0^{\tau_*} \left[\delta^2 \left\|Aw^\delta\right\|_2^2 + \left\|M(w^\delta) - \delta A w^\delta\right\|_2^2\right] \mathrm{d}\tau. \tag{32}$$

In particular, the square in the full cost is retained. By (29)–(30), the excess in (25) is at least

$$\frac{\kappa\lambda^3}{\nu^2} - \frac{\theta C_0 \lambda^{5/2}}{\nu} \longrightarrow +\infty.$$

Also $t_\lambda \to 0$, so the prefix lies within the prescribed horizon. This proves the proposition. □

This obstruction concerns a family of different data and different prescribed cutoffs. It neither produces a singularity for one datum nor contradicts (24) with its full-data allowance and existential cutoff. In this family the full initial caloric resource already scales as λ^3/ν^2; retaining it can pay the finite activation that an energy-only or empty-tail-only allowance misses.

7 Absorbing separated parents across all output scales

There is a sector of the stretching source that can be paid without an angular or sign assumption. Fix integers $J \ge 0$, $r \ge 3$, and put $K = 2^J$. For an output $k \in \Omega_j := \{2^j \le |k| < 2^{j+1}\}$, $j \ge J$, write $c_j = \lceil 3j/2 \rceil$. An ordered pair $p + q = k$ belongs to the separated sector if

$$n := \lfloor \log_2 \max\{|p|, |q|\} \rfloor \ge c_j + r.$$

Let $\mathcal{M}_{\mathrm{far}}$ retain these contributions in the ordered Fourier formula for $\mathcal{M}$, before summing the pairs, and define

$$\mathcal{M}_{\mathrm{lo}} = P_{<K}\mathcal{M}, \qquad \mathcal{M}_{\mathrm{keep}} = \mathcal{M} - \mathcal{M}_{\mathrm{lo}} - \mathcal{M}_{\mathrm{far}}.$$

These are fixed bilinear multipliers, with a mask symmetric in the two parent frequencies. Neither the velocity nor its evolution is truncated. In particular, $\mathcal{M}_{\mathrm{keep}}$ is not an orthogonal projection of the already summed field $\mathcal{M}$.

Proposition 10. *For every smooth solenoidal whole-space field, with the unitary Fourier convention and $E = \|u\|_2^2$, $Y = \|\nabla u\|_2^2$, $D = \|\Delta u\|_2^2$,*

$$\|\mathcal{M}_{\mathrm{far}}(u)\|_2 \le 2\sqrt{17}\, 2^{-r} \sqrt{ED} \le 10\, 2^{-r} \sqrt{ED}. \tag{33}$$

The bound is independent of J and the number of occupied shells. Its auxiliary heat capacity satisfies

$$\mathcal{K}_{\mathrm{far}}(u) := \frac{1}{\nu} \int_0^\infty \|\mathcal{M}_{\mathrm{far}}(H_s u)\|_2^2\,\mathrm{d}s \le \frac{50\, 2^{-2r}}{\nu^2} EY. \tag{34}$$

Proof. Since $n \ge j+3$, both parents belong to $B_n = \{2^{n-1} \le |\xi| < 2^{n+1}\}$. Write $E_n = \int_{B_n} |\widehat{u}|^2$ and $d_n = 2^{2n} E_n$. The ordered bound (13), followed by Cauchy–Schwarz in the parent frequency, gives

$$|\widehat{\mathcal{M}_{j,n}}(k)| \le (2\pi)^{-3/2} 2^{n+1} E_n \qquad (k \in \Omega_j).$$

As $|\Omega_j| = (28\pi/3)2^{3j}$,

$$\left\| P_{\Omega_j} \mathcal{M}_{\mathrm{far}} \right\|_2 \le 2\sqrt{\frac{7}{6\pi^2}} \sum_{n \ge c_j + r} 2^{3j/2 - n} d_n \le \sum_n A_{jn} d_n, \qquad A_{jn} = 2^{c_j - n} \mathbf{1}_{\{n \ge c_j + r\}}.$$

Every row sum of A is 2^{1-r}. Every column sum is at most 2^{1-r}, since the integers c_j strictly increase. Applying weighted Cauchy–Schwarz to each row and then the column bound proves

$$\sum_j \left(\sum_n A_{jn} d_n \right)^2 \le 2^{2-2r} \sum_n d_n^2.$$

The argument first applies to finite submatrices, and then passes to the increasing limit of their nonnegative sums. This summation avoids paying the same high-frequency tail separately at every output shell.

Each frequency of dyadic index ℓ belongs to just B_ℓ and $B_{\ell+1}$. Their fourth-order weights sum to $2^{4\ell} + 2^{4\ell+4} \le 17|\xi|^4$. Consequently

$$\sum_n d_n^2 = \sum_n 2^{4n} E_n^2 \le E \sum_n 2^{4n} E_n \le 17 E D.$$

The output shells are orthogonal, proving (33). For (34), apply that bound at $H_s u$ and use $E(H_s u) \le E$ and $\int_0^\infty D(H_s u)\, \mathrm{d}s = Y/(2\nu)$. □

Fix an energy bound $E \le M$, choose $0 < \eta < 1/2$, and choose r so that $10\, 2^{-r} \sqrt{M} \le \eta\nu$. Then

$$2|\langle Au, \mathcal{M}_{\mathrm{far}} \rangle| \le 2\eta\nu D, \qquad \mathcal{K}_{\mathrm{far}} \le \eta^2 Y/2. \tag{35}$$

The separation is fixed from M, ν, η; it is not adjusted from a future solution norm.

To use this source payment, change the auxiliary capacity and its positive cost explicitly:

$$\mathcal{K}_{\mathrm{keep}}(u) = \frac{1}{\nu} \int_0^\infty \|\mathcal{M}_{\mathrm{keep}}(H_s u)\|_2^2 \, \mathrm{d}s, \qquad \mathcal{G}_{\mathrm{keep}} = (Y + \mathcal{K}_{\mathrm{keep}})/2,$$
$$\mathcal{W}_{\mathrm{keep}} = \nu D + \frac{1}{\nu} \|\mathcal{M}_{\mathrm{keep}} - \nu A u\|_2^2, \qquad \mathcal{T}_{\mathrm{keep}} = D\mathcal{K}_{\mathrm{keep}}(u)[N(u)].$$

In particular $\mathcal{G}_{\mathrm{keep}} \ge Y/2$ and $\mathcal{W}_{\mathrm{keep}} \ge \nu D$. This is a modified cost, not an identification with the original $\mathcal{W}$. Heat translation for the fixed mask, the full enstrophy identity, and expansion of the square give, along the actual equation $u_t = N(u) - \nu A u + g$,

$$\begin{aligned} 2\mathcal{G}'_{\mathrm{keep}} + \mathcal{W}_{\mathrm{keep}} = \mathcal{T}_{\mathrm{keep}} + D\mathcal{K}_{\mathrm{keep}}(u)[g] + 2\langle Au, g \rangle \\ + 2\langle Au, \mathcal{M}_{\mathrm{lo}} \rangle + 2\langle Au, \mathcal{M}_{\mathrm{far}} \rangle. \end{aligned} \tag{36}$$

There is no claim that $\mathcal{K}$ is the sum of the three capacities: the retained and separated pairs can interfere at the same output. The identity uses the complete $N(u)$ in the derivative direction.

The low-output estimate $\|\mathcal{M}_{\mathrm{lo}}\|_2 \le c_K \sqrt{EY}$ implies

$$|\langle Au, \mathcal{M}_{\mathrm{lo}} \rangle| \le K^2 c_K E \sqrt{Y}, \qquad 2\int_a^t |\langle Au, \mathcal{M}_{\mathrm{lo}} \rangle| \, \mathrm{d}s \le 2K^2 c_K M \sqrt{LB} =: C_{\mathrm{lo}}.$$

Thus the two last terms of (36) cost a finite allowance and a fraction 2η of $\mathcal{W}_{\mathrm{keep}}$.

For completeness, the other payments must respect the parent mask. Every retained ordered symbol still obeys (13); the two masks in the quartic pairing are bounded by one. Hence Proposition 6 applies to $\mathcal{K}_{\mathrm{keep}}$ with exactly the same constants. For the forcing estimate use the Fourier moments

$$Q_\ell^{\#} = (2\pi)^{-3/2} \int_{\mathbb{R}^3} |\xi|^\ell |\widehat{g}(\xi)| \, \mathrm{d}\xi, \qquad \ell = 0, 1,$$

rather than asserting an L^∞ bound for the masked bilinear operator. Fourier convolution and Young's inequality give, with $w = H_s u$,

$$\|\mathcal{M}_{\mathrm{keep}}(w)\|_2 \le \left\|w^{\#}\right\|_\infty \|\nabla w\|_2 , \qquad \|D\mathcal{M}_{\mathrm{keep}}(w)[H_s g]\|_2 \le Q_0^{\#} \, \|\nabla w\|_2 + Q_1^{\#} \sqrt{E}.$$

Here $\left\|w^{\#}\right\|_\infty = (2\pi)^{-3/2} \int |\widehat{w}|$; the physical values of $w^{\#}$ need not be nonnegative. The heat-splitting proof of Proposition 4 now applies with $Q_0^{\#}, Q_1^{\#}$ in place of Q_0, Q_1. These moments are bounded by the H^2 and H^3 norms of g, respectively. They are finite uniformly in time for the force in the construction.

For $v = P_{\ge K} u$ and $g_K = P_{<K} N(v)$, the Fourier proof of (17) bounds these same moments:

$$Q_0^{\#}(g_K) \le K^4 E/(8\pi^2), \qquad Q_1^{\#}(g_K) \le K^5 E/(10\pi^2).$$

It therefore also pays the low nonlinear output of the retained capacity. Let C_{out} be the resulting time-integral allowance, and let C_f include the two explicit force terms of (36). They are finite under the preceding energy and force assumptions.

The remaining signed term is exactly

$$\mathcal{T}_{\mathrm{core}}(u) = D\mathcal{K}_{\mathrm{keep}}(v)[P_{\ge K} N(v)], \qquad v = P_{\ge K} u.$$

Its five original velocity inputs, nonlinear-direction output and stretching output are all above K. The outer stretching pairs additionally satisfy

$$\max\{|p|, |q|\} < 2^{c_j + r} \le \sqrt{2}\, 2^r |p + q|^{3/2} \qquad (p + q \in \Omega_j).$$

This restriction is imposed on the two slots of each stretching source. In a differentiated slot, $N(v)$ has two further parents; the displayed restriction imposes no corresponding comparability on those two parents. Their transfer and the signed time history remain to be controlled.

If constants $A_0 \ge 0$, $\theta < 1 - 2\eta$ obey

$$\int_a^t \mathcal{T}_{\mathrm{core}}(u) \, \mathrm{d}s \le A_0 + \theta \int_a^t \mathcal{W}_{\mathrm{keep}}(u) \, \mathrm{d}s \tag{37}$$

on every shorter prefix, choose $0 < \varepsilon < 1 - 2\eta - \theta$ in Proposition 6. The same exact low-parent decomposition and integral Grönwall argument then yield

$$\mathcal{G}_{\mathrm{keep}}(t) \le \left(\mathcal{G}_{\mathrm{keep}}(a) + \frac{A_0 + C_{\mathrm{lo}} + C_{\mathrm{out}} + C_f}{2} \right) \exp\Bigl(C_2 B + C_{3/2} \sqrt{LB} \Bigr) .$$

The target (37) remains open. For regularity its cutoffs, separation and allowances must be fixed from the full initial datum, viscosity and horizon, uniformly over shorter solution intervals. The actual forced singularities violate this target: otherwise the displayed bound would control their unbounded gradient energy.

The positive generated-child example (26) is retained at $J = 2$, $r = 3$ with exactly the same coefficient. Thus the absorption does not remove all positive renewal or make a sign claim about the remaining term. It pays one frequency sector uniformly, leaving a smaller physical-time problem.

8 Verification boundary

The Lean theorem `openAI_solution_no_uniform_signed_production_budget` applies to the actual construction at every positive viscosity and concerns the full-field spatial cubic integral. The sharp Fourier-tail extension and all caloric results here are analytic proofs, not Lean theorems or axioms. The companion `labs/caloric_force.py` checks the complete caloric balance by exact rational Fourier arithmetic on the periodic cube, including forces with a gradient part. It also checks the three-part frequency split, including a nonzero low-output contribution, and the ordered mode bound on nearly cancelling parents. Separate source-output checks retain the original positive cost and include parent frequency parameters up to one million that feed low stretching outputs. These checks do not certify the whole-space estimates or a physical-time bound. The checks `labs/generated_caloric_renewal.py` and `labs/thin_caloric_renewal.py` reproduce the finite coefficients used in Proposition 9. `labs/caloric_localization.py` checks the strict cutoff gap, the complete scaling factors and a finite periodic streamfunction envelope. Its rational calculations retain the envelope derivative and all generated modes. They do not certify the Schwartz localization limit or the common local PDE time interval, which are analytic arguments above. The rational checker `labs/separated_caloric.py` tests the Schur matrix bounds, enlarged-band overlap, exact source partition, heat translation, and (36) with the full actual source. An independent quartic polarization checks the masked derivative. It also verifies the unchanged positive generated-child coefficient. Its finite periodic source tests use the conservative lattice constant 200, not the whole-space constant 10: counting at most $125\,2^{3j}$ lattice points in Ω_j replaces $2\sqrt{17}$ by $20\sqrt{85} < 200$ in the proof. The infinite-shell estimate and its whole-space use remain analytic. Arbitrary-data unforced regularity remains open.

References

[1] OpenAI, *Finite time blowup for Navier–Stokes*, manuscript, with formalization in NavierStokesAndEuler, 2026.