



# Inequality in Health

## Lecture IV: Measuring and Decomposing Health Inequality II

Dr Martin Karlsson  
University of Duisburg-Essen  
Winter semester 2022-23

- 1 Recap of Last Lecture
- 2 Introduction
- 3 Correcting the Concentration Index
  - Level of Measurement
  - Issues with Bounded Variables
  - Desirable Properties of Indices
  - Comparison of Indices
- 4 Decomposition of the Concentration Index
- 5 Level-Dependent Indices
- 6 Two-Dimensional Decomposition
- 7 Summary and Conclusions

## Recap of Last Lecture

# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.

# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.
- The decomposition partitions differences into:

# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.
- The decomposition partitions differences into:
  - Gap in **endowments** (explained)

# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.
- The decomposition partitions differences into:
  - Gap in **endowments** (explained)
  - Gap in **coefficients** (unexplained)

# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.
- The decomposition partitions differences into:
  - Gap in **endowments** (explained)
  - Gap in **coefficients** (unexplained)
  - Interaction between gaps.



# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.
- The decomposition partitions differences into:
  - Gap in **endowments** (explained)
  - Gap in **coefficients** (unexplained)
  - Interaction between gaps.
- Machado & Mata propose a decomposition method based on quantile regression.

# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.
- The decomposition partitions differences into:
  - Gap in **endowments** (explained)
  - Gap in **coefficients** (unexplained)
  - Interaction between gaps.
- Machado & Mata propose a decomposition method based on quantile regression.
  - It allows to “go beyond the mean” by performing a **detailed** decomposition by quantiles.

# Recap of Last Lecture

- The **Oaxaca-Blinder decomposition** allows to attribute differences in the outcome variable (ex. health) between two groups to several explanatory factors.
- The decomposition partitions differences into:
  - Gap in **endowments** (explained)
  - Gap in **coefficients** (unexplained)
  - Interaction between gaps.
- Machado & Mata propose a decomposition method based on quantile regression.
  - It allows to “go beyond the mean” by performing a **detailed** decomposition by quantiles.
  - It **simulates counterfactual** distributions of the covariates.

# Introduction

# Introduction

- Decomposition methods like OB and MM allow for a **limited** understanding of socioeconomic-related inequality in health: explain differences between **two groups** only (e.g. rich-poor).

# Introduction

- Decomposition methods like OB and MM allow for a **limited** understanding of socioeconomic-related inequality in health: explain differences between **two groups** only (e.g. rich-poor).
- Alternative decomposition methods explain socioeconomic-related inequality across the **entire distribution** of the SES variable (ex. income distribution).

# Introduction

- Decomposition methods like OB and MM allow for a **limited** understanding of socioeconomic-related inequality in health: explain differences between **two groups** only (e.g. rich-poor).
- Alternative decomposition methods explain socioeconomic-related inequality across the **entire distribution** of the SES variable (ex. income distribution).
- Popular measures of inequality: the Gini coefficient, the concentration curve and the concentration index.

# Introduction

- Decomposition methods like OB and MM allow for a **limited** understanding of socioeconomic-related inequality in health: explain differences between **two groups** only (e.g. rich-poor).
- Alternative decomposition methods explain socioeconomic-related inequality across the **entire distribution** of the SES variable (ex. income distribution).
- Popular measures of inequality: the Gini coefficient, the concentration curve and the concentration index.
- Focus on the **concentration index**: it measures the socioeconomic inequality of health taking into account:



# Introduction

- Decomposition methods like OB and MM allow for a **limited** understanding of socioeconomic-related inequality in health: explain differences between **two groups** only (e.g. rich-poor).
- Alternative decomposition methods explain socioeconomic-related inequality across the **entire distribution** of the SES variable (ex. income distribution).
- Popular measures of inequality: the Gini coefficient, the concentration curve and the concentration index.
- Focus on the **concentration index**: it measures the socioeconomic inequality of health taking into account:
  - the **level of health** of each individual

# Introduction

- Decomposition methods like OB and MM allow for a **limited** understanding of socioeconomic-related inequality in health: explain differences between **two groups** only (e.g. rich-poor).
- Alternative decomposition methods explain socioeconomic-related inequality across the **entire distribution** of the SES variable (ex. income distribution).
- Popular measures of inequality: the Gini coefficient, the concentration curve and the concentration index.
- Focus on the **concentration index**: it measures the socioeconomic inequality of health taking into account:
  - the **level of health** of each individual
  - their socioeconomic **rank**.

## Correcting the Concentration Index

# Introduction

- The **choice** of the inequality indicator to use may influence the results generated by the analysis.

# Introduction

- The **choice** of the inequality indicator to use may influence the results generated by the analysis.
- Over the years researchers proposed alternative ways to compute the **concentration index**, providing **improved** alternatives to tackle some shortcomings of the original CI.

# Introduction

- The **choice** of the inequality indicator to use may influence the results generated by the analysis.
- Over the years researchers proposed alternative ways to compute the **concentration index**, providing **improved** alternatives to tackle some shortcomings of the original CI.
- We consider:

# Introduction

- The **choice** of the inequality indicator to use may influence the results generated by the analysis.
- Over the years researchers proposed alternative ways to compute the **concentration index**, providing **improved** alternatives to tackle some shortcomings of the original CI.
- We consider:
  - ① A modified concentration index

# Introduction

- The **choice** of the inequality indicator to use may influence the results generated by the analysis.
- Over the years researchers proposed alternative ways to compute the **concentration index**, providing **improved** alternatives to tackle some shortcomings of the original CI.
- We consider:
  - 1 A modified concentration index
  - 2 The Wagstaff normalization



# Introduction

- The **choice** of the inequality indicator to use may influence the results generated by the analysis.
- Over the years researchers proposed alternative ways to compute the **concentration index**, providing **improved** alternatives to tackle some shortcomings of the original CI.
- We consider:
  - 1 A modified concentration index
  - 2 The Wagstaff normalization
  - 3 The Erreygers index.

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).
  - ② **Cardinal**: possible to **compare differences** between health states.

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).
  - ② **Cardinal**: possible to **compare differences** between health states.
    - Example: body temperature.

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).
  - ② **Cardinal**: possible to **compare differences** between health states.
    - Example: body temperature.
  - ③ **Ratio-scale**: “zero health” fixed in an unambiguous way (reference).

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).
  - ② **Cardinal**: possible to **compare differences** between health states.
    - Example: body temperature.
  - ③ **Ratio-scale**: “zero health” fixed in an unambiguous way (reference).
    - Example: life expectancy; pulse.



# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).
  - ② **Cardinal**: possible to **compare differences** between health states.
    - Example: body temperature.
  - ③ **Ratio-scale**: “zero health” fixed in an unambiguous way (reference).
    - Example: life expectancy; pulse.
- **Range** of the health variable:

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).
  - ② **Cardinal**: possible to **compare differences** between health states.
    - Example: body temperature.
  - ③ **Ratio-scale**: “zero health” fixed in an unambiguous way (reference).
    - Example: life expectancy; pulse.
- **Range** of the health variable:
  - **Bounded**: have both a finite upper and lower bound.

# Health Variable: Level of Measurement

- According to the type and the amount of **information** available we can express health status through different type of variables:
  - ① **Ordinal**: we are just able to **rank** individuals.
    - Example: Self-Assessed Health – SAH (1 – excellent; 2 – fair; 3 – poor health).
  - ② **Cardinal**: possible to **compare differences** between health states.
    - Example: body temperature.
  - ③ **Ratio-scale**: “zero health” fixed in an unambiguous way (reference).
    - Example: life expectancy; pulse.
- **Range** of the health variable:
  - **Bounded**: have both a finite upper and lower bound.
  - **Unbounded**: we assume here that unbounded variables have a finite lower bound.

# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

# Limitations of $C(h)$ by Health Variable

- 1 **Ordinal** variable:
  - Any **positive monotone** transformation captures the same information.

# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.

# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.
- With **qualitative** health variables  $C(h)$  is to a large extent **arbitrary**, because differences between individuals cannot be compared.

# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.
- With **qualitative** health variables  $C(h)$  is to a large extent **arbitrary**, because differences between individuals cannot be compared.

## 2 Cardinal variable:



# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.
- With **qualitative** health variables  $C(h)$  is to a large extent **arbitrary**, because differences between individuals cannot be compared.

## 2 Cardinal variable:

- Any **positive linear** transformation captures the same information.

# Limitations of $C(h)$ by Health Variable

## ① Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.
- With **qualitative** health variables  $C(h)$  is to a large extent **arbitrary**, because differences between individuals cannot be compared.

## ② Cardinal variable:

- Any **positive linear** transformation captures the same information.
- $C(h)$  is not invariant to positive linear transformations.

# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.
- With **qualitative** health variables  $C(h)$  is to a large extent **arbitrary**, because differences between individuals cannot be compared.

## 2 Cardinal variable:

- Any **positive linear** transformation captures the same information.
- $C(h)$  is not invariant to positive linear transformations.

## 3 Ratio-scale variable:

# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.
- With **qualitative** health variables  $C(h)$  is to a large extent **arbitrary**, because differences between individuals cannot be compared.

## 2 Cardinal variable:

- Any **positive linear** transformation captures the same information.
- $C(h)$  is not invariant to positive linear transformations.

## 3 Ratio-scale variable:

- Any **positive proportional** transformation captures the same information.

# Limitations of $C(h)$ by Health Variable

## 1 Ordinal variable:

- Any **positive monotone** transformation captures the same information.
- $C(h)$  is not invariant to positive monotone transformations.
- With **qualitative** health variables  $C(h)$  is to a large extent **arbitrary**, because differences between individuals cannot be compared.

## 2 Cardinal variable:

- Any **positive linear** transformation captures the same information.
- $C(h)$  is not invariant to positive linear transformations.

## 3 Ratio-scale variable:

- Any **positive proportional** transformation captures the same information.
- $C(h)$  is only invariant to positive proportional transformations, implying that health should be measured on a **ratio scale**.

# Limitations of $C(h)$ : Bounds

- Bounds of  $C(h)$ :  $C(h) \in \left(-\frac{(n-1)}{n}; \frac{(n-1)}{n}\right)$ .

# Limitations of $C(h)$ : Bounds

- Bounds of  $C(h)$ :  $C(h) \in \left(-\frac{(n-1)}{n}; \frac{(n-1)}{n}\right)$ .
- For any **bounded** health variable (with a finite upper value or a positive lower value),  $C(h)$  has **varying bounds**:

$$-\frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)} \leq C(h) \leq \frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)}$$

where

# Limitations of $C(h)$ : Bounds

- Bounds of  $C(h)$ :  $C(h) \in \left(-\frac{(n-1)}{n}; \frac{(n-1)}{n}\right)$ .
- For any **bounded** health variable (with a finite upper value or a positive lower value),  $C(h)$  has **varying bounds**:

$$-\frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)} \leq C(h) \leq \frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)}$$

where

$a_h$  lower bound of the health variable  $h$



# Limitations of $C(h)$ : Bounds

- Bounds of  $C(h)$ :  $C(h) \in \left(-\frac{(n-1)}{n}; \frac{(n-1)}{n}\right)$ .
- For any **bounded** health variable (with a finite upper value or a positive lower value),  $C(h)$  has **varying bounds**:

$$-\frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)} \leq C(h) \leq \frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)}$$

where

$a_h$  lower bound of the health variable  $h$

$b_h$  upper bound of the health variable  $h$

# Limitations of $C(h)$ : Bounds

- Bounds of  $C(h)$ :  $C(h) \in \left(-\frac{(n-1)}{n}; \frac{(n-1)}{n}\right)$ .
- For any **bounded** health variable (with a finite upper value or a positive lower value),  $C(h)$  has **varying bounds**:

$$-\frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)} \leq C(h) \leq \frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)}$$

where

$a_h$  lower bound of the health variable  $h$

$b_h$  upper bound of the health variable  $h$

and with

$$0 \leq a_h < b_h < +\infty.$$

# Limitations of $C(h)$ : Bounds

- Bounds of  $C(h)$ :  $C(h) \in \left(-\frac{(n-1)}{n}; \frac{(n-1)}{n}\right)$ .
- For any **bounded** health variable (with a finite upper value or a positive lower value),  $C(h)$  has **varying bounds**:

$$-\frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)} \leq C(h) \leq \frac{(b_h - \mu_h)(\mu_h - a_h)}{\mu_h(b_h - a_h)}$$

where

$a_h$  lower bound of the health variable  $h$

$b_h$  upper bound of the health variable  $h$

and with

$$0 \leq a_h < b_h < +\infty.$$

- **Comparing** populations with **different mean** health levels is problematic, even if health is measured on a ratio-scale level.

# Limitations of $C(h)$ : Health and Ill-Health

- Example: health levels measure **health**, while malnutrition measures **ill health**.

# Limitations of $C(h)$ : Health and Ill-Health

- Example: health levels measure **health**, while malnutrition measures **ill health**.
- If the health variable  $h$  is **bounded**, define a corresponding ill health variable  $s$ :

$$s_i \equiv b_h - h_i$$

$$\mu_s = b_h - \mu_h$$

# Limitations of $C(h)$ : Health and Ill-Health

- Example: health levels measure **health**, while malnutrition measures **ill health**.
- If the health variable  $h$  is **bounded**, define a corresponding ill health variable  $s$ :

$$s_i \equiv b_h - h_i$$

$$\mu_s = b_h - \mu_h$$

where

# Limitations of $C(h)$ : Health and Ill-Health

- Example: health levels measure **health**, while malnutrition measures **ill health**.
- If the health variable  $h$  is **bounded**, define a corresponding ill health variable  $s$ :

$$s_i \equiv b_h - h_i$$

$$\mu_s = b_h - \mu_h$$

where

$s_i$  ill-health status of individual  $i$ ;

# Limitations of $C(h)$ : Health and Ill-Health

- Example: health levels measure **health**, while malnutrition measures **ill health**.
- If the health variable  $h$  is **bounded**, define a corresponding ill health variable  $s$ :

$$s_i \equiv b_h - h_i$$

$$\mu_s = b_h - \mu_h$$

where

$s_i$  ill-health status of individual  $i$ ;

$\mu_s$  average ill health of the population.



# Limitations of $C(h)$ : Health and Ill-Health

- Example: health levels measure **health**, while malnutrition measures **ill health**.
- If the health variable  $h$  is **bounded**, define a corresponding ill health variable  $s$ :

$$s_i \equiv b_h - h_i$$

$$\mu_s = b_h - \mu_h$$

where

$s_i$  ill-health status of individual  $i$ ;

$\mu_s$  average ill health of the population.

- The ill-health Concentration Index  $C(s)$  is defined by analogy with the health Concentration Index  $C(h)$ :

$$C(s) \equiv 1 - \frac{\sum_{i=1}^n (2\lambda_i - 1)s_i}{n^2 \mu_s}$$

# Health and Ill-Health

- Health and ill-health are *mirrors* of one another  $\Rightarrow C(h)$  and  $C(s)$  should give “mirror images” of inequality.

# Health and Ill-Health

- Health and ill-health are *mirrors* of one another  $\Rightarrow C(h)$  and  $C(s)$  should give “mirror images” of inequality.
- Comparing the two:

$$C(s) = 1 - \frac{(\mu_h + \mu_s)}{\mu_s} + \frac{\mu_h}{\mu_s} \frac{\sum_{i=1}^n (2\lambda_i - 1)h_i}{n^2\mu_h} = -\frac{\mu_h}{\mu_s} C(h)$$

# Health and Ill-Health

- Health and ill-health are *mirrors* of one another  $\Rightarrow C(h)$  and  $C(s)$  should give “mirror images” of inequality.
- Comparing the two:

$$C(s) = 1 - \frac{(\mu_h + \mu_s)}{\mu_s} + \frac{\mu_h}{\mu_s} \frac{\sum_{i=1}^n (2\lambda_i - 1)h_i}{n^2\mu_h} = -\frac{\mu_h}{\mu_s} C(h)$$

- Consider two health distributions:  $h_1$  and  $h_2$ .

# Health and Ill-Health

- Health and ill-health are *mirrors* of one another  $\Rightarrow C(h)$  and  $C(s)$  should give “mirror images” of inequality.
- Comparing the two:

$$C(s) = 1 - \frac{(\mu_h + \mu_s)}{\mu_s} + \frac{\mu_h}{\mu_s} \frac{\sum_{i=1}^n (2\lambda_i - 1)h_i}{n^2\mu_h} = -\frac{\mu_h}{\mu_s} C(h)$$

- Consider two health distributions:  $h_1$  and  $h_2$ .
- If  $\mu_{h_1} = \mu_{h_2}$ , then  $C(h_1) > C(h_2) \Leftrightarrow C(s_1) < C(s_2)$ .

# Health and Ill-Health

- Health and ill-health are *mirrors* of one another  $\Rightarrow C(h)$  and  $C(s)$  should give “mirror images” of inequality.
- Comparing the two:

$$C(s) = 1 - \frac{(\mu_h + \mu_s)}{\mu_s} + \frac{\mu_h}{\mu_s} \frac{\sum_{i=1}^n (2\lambda_i - 1)h_i}{n^2 \mu_h} = -\frac{\mu_h}{\mu_s} C(h)$$

- Consider two health distributions:  $h_1$  and  $h_2$ .
- If  $\mu_{h_1} = \mu_{h_2}$ , then  $C(h_1) > C(h_2) \Leftrightarrow C(s_1) < C(s_2)$ .
- Relative differences** correctly measured:  $\frac{C(h_1)}{C(h_2)} = \frac{C(s_1)}{C(s_2)}$ .

# Health and Ill-Health

- Health and ill-health are *mirrors* of one another  $\Rightarrow C(h)$  and  $C(s)$  should give “mirror images” of inequality.
- Comparing the two:

$$C(s) = 1 - \frac{(\mu_h + \mu_s)}{\mu_s} + \frac{\mu_h}{\mu_s} \frac{\sum_{i=1}^n (2\lambda_i - 1)h_i}{n^2 \mu_h} = -\frac{\mu_h}{\mu_s} C(h)$$

- Consider two health distributions:  $h_1$  and  $h_2$ .
- If  $\mu_{h_1} = \mu_{h_2}$ , then  $C(h_1) > C(h_2) \Leftrightarrow C(s_1) < C(s_2)$ .
- Relative differences** correctly measured:  $\frac{C(h_1)}{C(h_2)} = \frac{C(s_1)}{C(s_2)}$ .
- If  $\mu_{h_1} \neq \mu_{h_2}$ , these properties no longer hold: inequalities in **ill health** may give different rankings than inequalities in **health**.

# Introduction

- Define a generic **family of indices**  $I$  for some distribution  $h$ :

$$I(h) = f(a_h, b_h, \mu_h, n) \sum_{i=1}^n r_i h_i$$

for a continuous function  $f(\cdot)$ .



# Introduction

- Define a generic **family of indices**  $I$  for some distribution  $h$ :

$$I(h) = f(a_h, b_h, \mu_h, n) \sum_{i=1}^n r_i h_i$$

for a continuous function  $f(\cdot)$ .

- For unbounded variables,  $b_h = +\infty$  and  $f(\cdot) = f(a_h, \mu_h, n)$ .

# Introduction

- Define a generic **family of indices**  $I$  for some distribution  $h$ :

$$I(h) = f(a_h, b_h, \mu_h, n) \sum_{i=1}^n r_i h_i$$

for a continuous function  $f(\cdot)$ .

- For unbounded variables,  $b_h = +\infty$  and  $f(\cdot) = f(a_h, \mu_h, n)$ .
- We may impose restrictions on the form of  $f(\cdot)$  depending on the properties that we want  $I(h)$  to satisfy.

# Sign Condition

- By convention positive (negative; 0) values of  $I(h)$  should signal a pro-rich (pro-poor; no) **bias** in the distribution.

# Sign Condition

- By convention positive (negative; 0) values of  $I(h)$  should signal a pro-rich (pro-poor; no) **bias** in the distribution.
- **Sign condition:** the sign of  $I(h)$  coincides with the sign of  $\sum_{i=1}^n r_i h_i$ .

# Sign Condition

- By convention positive (negative; 0) values of  $I(h)$  should signal a pro-rich (pro-poor; no) **bias** in the distribution.
- **Sign condition:** the sign of  $I(h)$  coincides with the sign of  $\sum_{i=1}^n r_i h_i$ .

## Proposition 1

*The sign condition is satisfied if and only if:*

- For  $h$  **unbounded**:  $f(a_h, \mu_h, n) > 0$  for  $n > 0$  and  $a_h < \mu_h < +\infty$ ;
- For  $h$  **bounded**:  $f(a_h, b_h, \mu_h, n) > 0$  for  $n > 0$  and  $a_h < \mu_h < b_h$ .

# Scale Invariance

- $I(h)$  should be independent of the unit of measurement of  $h$ .

# Scale Invariance

- $I(h)$  should be independent of the unit of measurement of  $h$ .
- **Scale invariance:** for a change in the unit of measurement of  $h$  that transform the distribution  $h$  into  $\tilde{h}$ ,  $I(\tilde{h}) = I(h)$ .

# Scale Invariance

- $I(h)$  should be independent of the unit of measurement of  $h$ .
- **Scale invariance:** for a change in the unit of measurement of  $h$  that transform the distribution  $h$  into  $\tilde{h}$ ,  $I(\tilde{h}) = I(h)$ .
  - If  $h$  **cardinal**: a positive linear transformation,

$$\tilde{h}_i = \alpha + \beta h_i; \quad \tilde{a}_h = \alpha + \beta a_h; \quad \tilde{b}_h = \alpha + \beta b_h.$$



# Scale Invariance

- $I(h)$  should be independent of the unit of measurement of  $h$ .
- **Scale invariance:** for a change in the unit of measurement of  $h$  that transform the distribution  $h$  into  $\tilde{h}$ ,  $I(\tilde{h}) = I(h)$ .
  - If  $h$  **cardinal**: a positive linear transformation,

$$\tilde{h}_i = \alpha + \beta h_i; \quad \tilde{a}_h = \alpha + \beta a_h; \quad \tilde{b}_h = \alpha + \beta b_h.$$

- If  $h$  **ratio-scale**: a positive proportional transformation,

$$\tilde{h}_i = \beta h_i; \quad \tilde{a}_h = \beta a_h; \quad \tilde{b}_h = \beta b_h.$$

## Proposition 2

$I(h)$  has the scale invariance property if and only if:

- For  $h$  **unbounded**:  $f(a_h, \mu_h, n) = \frac{1}{\mu_h - a_h} k(n)$ ;
- For  $h$  **bounded**:  $f(a_h, b_h, \mu_h, n) = \frac{1}{b_h - a_h} g\left(\frac{\mu_h - a_h}{b_h - a_h}, n\right)$ .

# Modified Concentration Index

- For  $h$  **unbounded**, if we fix  $I(h)$ 's bounds to  $-1$  and  $1$ , we can write a **modified** version  $\hat{C}(h)$  of the standard **concentration index**  $C(h)$ :

$$\hat{C}(h) = \frac{2}{n^2 (\mu_h - a_h)} \sum_{i=1}^n r_i h_i$$

that satisfies the sign condition and the scale invariance property.

# Modified Concentration Index

- For  $h$  **unbounded**, if we fix  $I(h)$ 's bounds to  $-1$  and  $1$ , we can write a **modified** version  $\hat{C}(h)$  of the standard **concentration index**  $C(h)$ :

$$\hat{C}(h) = \frac{2}{n^2 (\mu_h - a_h)} \sum_{i=1}^n r_i h_i$$

that satisfies the sign condition and the scale invariance property.

- For  $h$  **bounded**, it is more convenient to write a standardized definition  $h_i^*$  of  $h_i$ :

$$h_i^* \equiv \frac{h_i - a_h}{b_h - a_h}$$

where  $h_i^* \in [a_{h^*} = 0; b_{h^*} = 1]$  and  $\mu_{h^*} = \frac{\mu_h - a_h}{b_h - a_h}$ .

# Modified Concentration Index

- For  $h$  **unbounded**, if we fix  $I(h)$ 's bounds to  $-1$  and  $1$ , we can write a **modified** version  $\hat{C}(h)$  of the standard **concentration index**  $C(h)$ :

$$\hat{C}(h) = \frac{2}{n^2 (\mu_h - a_h)} \sum_{i=1}^n r_i h_i$$

that satisfies the sign condition and the scale invariance property.

- For  $h$  **bounded**, it is more convenient to write a standardized definition  $h_i^*$  of  $h_i$ :

$$h_i^* \equiv \frac{h_i - a_h}{b_h - a_h}$$

where  $h_i^* \in [a_{h^*} = 0; b_{h^*} = 1]$  and  $\mu_{h^*} = \frac{\mu_h - a_h}{b_h - a_h}$ .

- We can write a generic scale-invariant, rank dependent index  $I(h^*)$  for bounded **cardinal** and **ratio-scale** variables:

$$I(h^*) = g(\mu_{h^*}, n) \sum_{i=1}^n r_i h_i^*$$

# Mirror Property (Bounded Vars)

- The **ranking** of the distribution of the health index should be the opposite of the distribution of the corresponding ill-health index.

# Mirror Property (Bounded Vars)

- The **ranking** of the distribution of the health index should be the opposite of the distribution of the corresponding ill-health index.
- **Mirror property:** for a health distribution  $h$  and the associated ill health distribution  $s$ ,  $I(h) = -I(s)$ .

# Wagstaff Index (Bounded Vars)

- Wagstaff (2005) suggested a **normalization** formula aimed at remedying the **bounds** issue.

# Wagstaff Index (Bounded Vars)

- Wagstaff (2005) suggested a **normalization** formula aimed at remedying the **bounds** issue.
- We can define the **Wagstaff-normalized CI**  $W(h)$  as:

$$\begin{aligned}
 W(h^*) &\equiv \frac{\mu_{h^*}(b_{h^*} - a_{h^*})}{(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} C(h^*) \\
 &= \frac{2(b_{h^*} - a_{h^*})}{n^2(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^*
 \end{aligned}$$



# Wagstaff Index (Bounded Vars)

- Wagstaff (2005) suggested a **normalization** formula aimed at remedying the **bounds** issue.
- We can define the **Wagstaff-normalized CI**  $W(h)$  as:

$$\begin{aligned} W(h^*) &\equiv \frac{\mu_{h^*}(b_{h^*} - a_{h^*})}{(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} C(h^*) \\ &= \frac{2(b_{h^*} - a_{h^*})}{n^2(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^* \end{aligned}$$

- $W(h^*) \in [-1, 1]$ .

# Wagstaff Index (Bounded Vars)

- Wagstaff (2005) suggested a **normalization** formula aimed at remedying the **bounds** issue.
- We can define the **Wagstaff-normalized CI**  $W(h)$  as:

$$\begin{aligned} W(h^*) &\equiv \frac{\mu_{h^*}(b_{h^*} - a_{h^*})}{(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} C(h^*) \\ &= \frac{2(b_{h^*} - a_{h^*})}{n^2(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^* \end{aligned}$$

- $W(h^*) \in [-1, 1]$ .
- $W(h^*)$  satisfies the **mirror condition**.

# Wagstaff Index (Bounded Vars)

- Wagstaff (2005) suggested a **normalization** formula aimed at remedying the **bounds** issue.
- We can define the **Wagstaff-normalized CI**  $W(h)$  as:

$$\begin{aligned} W(h^*) &\equiv \frac{\mu_{h^*}(b_{h^*} - a_{h^*})}{(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} C(h^*) \\ &= \frac{2(b_{h^*} - a_{h^*})}{n^2(b_{h^*} - \mu_{h^*})(\mu_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^* \end{aligned}$$

- $W(h^*) \in [-1, 1]$ .
- $W(h^*)$  satisfies the **mirror condition**.
- Advantage over  $C(h^*)$ :  $W(h^*)$  is invariant to a positive linear transformation of  $h^*$ , so  $h^*$  can be also measured on a **cardinal scale**.

# Erreygers Index (Bounded Vars)

- Erreygers (2009) proposes the following index:

$$E(h^*) = \frac{8}{n^2(b_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^*$$

# Erreygers Index (Bounded Vars)

- Erreygers (2009) proposes the following index:

$$E(h^*) = \frac{8}{n^2(b_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^*$$

- Formally  $E(h^*)$  and  $W(h^*)$  differ only with respect to the **normalization** applied to the weighted sum of  $h$ .

# Erreygers Index (Bounded Vars)

- Erreygers (2009) proposes the following index:

$$E(h^*) = \frac{8}{n^2(b_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^*$$

- Formally  $E(h^*)$  and  $W(h^*)$  differ only with respect to the **normalization** applied to the weighted sum of  $h$ .
- $E(h^*)$  is the only index that is insensitive to any feasible **equal addition** to the standardized health variable  $h^*$ :

# Erreygers Index (Bounded Vars)

- Erreygers (2009) proposes the following index:

$$E(h^*) = \frac{8}{n^2(b_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^*$$

- Formally  $E(h^*)$  and  $W(h^*)$  differ only with respect to the **normalization** applied to the weighted sum of  $h$ .
- $E(h^*)$  is the only index that is insensitive to any feasible **equal addition** to the standardized health variable  $h^*$ :
  - an equal increment of  $h^*$  for all individuals – **keeping its bounds constant** – does not affect the value of the index, all other things equal.

# Erreygers Index (Bounded Vars)

- Erreygers (2009) proposes the following index:

$$E(h^*) = \frac{8}{n^2(b_{h^*} - a_{h^*})} \sum_{i=1}^n r_i h_i^*$$

- Formally  $E(h^*)$  and  $W(h^*)$  differ only with respect to the **normalization** applied to the weighted sum of  $h$ .
- $E(h^*)$  is the only index that is insensitive to any feasible **equal addition** to the standardized health variable  $h^*$ :
  - an equal increment of  $h^*$  for all individuals – **keeping its bounds constant** – does not affect the value of the index, all other things equal.
- However, there is a debate in the literature on whether one index between  $E(h^*)$  and  $W(h^*)$  is superior to the other.



# Indices and Their Properties: A Recap

Table 1. Concentration indices and their properties by level of measurement of the health variable.

Variable Level	Variable Range			
	<i>Unbounded</i>		<i>Bounded</i>	
	Index	Property	Index	Property
<i>Ordinal</i>	Concentration index and its variants in principle meaningless			
<i>Cardinal</i>	Modified CI	Sign condition Scale invariance	Erreygers and Wagstaff indices	Sign condition Scale invariance
<i>Ratio-scale</i>	Concentration index		Mirror property	

Source: Erreygers and Van Ourti (2011).

# Comparing Indices: An Example

- **Stunting** is a (ill-) health measure reflecting chronic **malnutrition**.

# Comparing Indices: An Example

- **Stunting** is a (ill-) health measure reflecting chronic **malnutrition**.
- Binary **ill-health** variable  $s$ :  $s_i = 1$  if child  $i$  is stunted, 0 otherwise.

# Comparing Indices: An Example

- **Stunting** is a (ill-) health measure reflecting chronic **malnutrition**.
- Binary **ill-health** variable  $s$ :  $s_i = 1$  if child  $i$  is stunted, 0 otherwise.
- **Health** variable  $h$ :  $h_i = 1$  if child  $i$  is not stunted, 0 otherwise.

# Comparing Indices: An Example

- **Stunting** is a (ill-) health measure reflecting chronic **malnutrition**.
- Binary **ill-health** variable  $s$ :  $s_i = 1$  if child  $i$  is stunted, 0 otherwise.
- **Health** variable  $h$ :  $h_i = 1$  if child  $i$  is not stunted, 0 otherwise.

Table 2. Stunting in some developing countries. Source: Erreygers (2009).

Country	$\mu_s$	$\mu_h$	$-C(s)$	$C(h)$	$W(h)$	$E(h)$
Nigeria (2003)	0.3845	0.6155	0.1612	<b>0.1007</b>	<b>0.2619</b>	0.2479
Cameroon (2004)	0.3165	0.6835	0.1698	<b>0.0786</b>	<b>0.2484</b>	0.2150
Kenya (2003)	0.3056	0.6944	0.1265	0.0557	0.1822	0.1546
Ghana (2003)	0.2943	0.7057	0.1743	<b>0.0727</b>	<b>0.2470</b>	0.2052
Cambodia (2000)	0.2943	0.7057	0.0887	0.0370	0.1257	0.1044
Bolivia (2003)	0.0680	0.9320	0.2739	<b>0.0200</b>	<b>0.2939</b>	<b>0.0745</b>
Peru (2000)	0.0501	0.9499	0.3676	<b>0.0194</b>	<b>0.3870</b>	<b>0.0737</b>
Nicaragua (2001)	0.0422	0.9578	0.3304	<b>0.0146</b>	<b>0.3450</b>	<b>0.0558</b>
Colombia (2005)	0.0215	0.9785	0.2699	<b>0.0059</b>	<b>0.2758</b>	<b>0.0232</b>

# Decomposition of the Concentration Index

# Introduction

- Decomposition of the CI: provide a specification for the health outcome:

$$h_{it} = \alpha_t + \sum_{j=1}^J \beta_j X_{jit} + \varepsilon_{it}. \quad (1)$$

# Introduction

- Decomposition of the CI: provide a specification for the health outcome:

$$h_{it} = \alpha_t + \sum_{j=1}^J \beta_j X_{jit} + \varepsilon_{it}. \quad (1)$$

- Consider the Erreygers index and adjust it to include multiple observations per individual (Kjellsson, 2018):

$$E(h) = \frac{8}{(nt)^2} \sum_{t=1}^T \sum_{i=1}^n r_i h_i \quad (2)$$



# Decomposition of the Concentration Index

- Substitute (1) in (2) to obtain

$$E(h) = 4 \sum_{j=1}^J \beta_j V(X_j) + 4V^\varepsilon \quad (3)$$

where

$$V(X_j) = \frac{2}{(nt)^2} \sum_{t=1}^T \sum_{i=1}^n r_i X_{jit} \quad V^\varepsilon = \frac{2}{(nt)^2} \sum_{t=1}^T \sum_{i=1}^n r_i \varepsilon_{it} \quad (4)$$

denote the generalized concentration indices for  $X_j$  and the error term, respectively.

# Decomposition of the Concentration Index

- Substitute (1) in (2) to obtain

$$E(h) = 4 \sum_{j=1}^J \beta_j V(X_j) + 4V^\varepsilon \quad (3)$$

where

$$V(X_j) = \frac{2}{(nt)^2} \sum_{t=1}^T \sum_{i=1}^n r_i X_{jit} \quad V^\varepsilon = \frac{2}{(nt)^2} \sum_{t=1}^T \sum_{i=1}^n r_i \varepsilon_{it} \quad (4)$$

denote the generalized concentration indices for  $X_j$  and the error term, respectively.

- So  $E(h)$  is the weighted sum of the **CI for the  $J$  regressors** plus a **residual component**.

# Decomposition of the Concentration Index

- Substitute (1) in (2) to obtain

$$E(h) = 4 \sum_{j=1}^J \beta_j V(X_j) + 4V^\varepsilon \quad (3)$$

where

$$V(X_j) = \frac{2}{(nt)^2} \sum_{t=1}^T \sum_{i=1}^n r_i X_{jit} \quad V^\varepsilon = \frac{2}{(nt)^2} \sum_{t=1}^T \sum_{i=1}^n r_i \varepsilon_{it} \quad (4)$$

denote the generalized concentration indices for  $X_j$  and the error term, respectively.

- So  $E(h)$  is the weighted sum of the **CI for the  $J$  regressors** plus a **residual component**.
- A similar decomposition applies to the other indices.

# Empirical Application: Kjellsson (2018)

- In a recent study, Kjellsson (2018) investigates **income-related smoking inequality** among Swedish women.

# Empirical Application: Kjellsson (2018)

- In a recent study, Kjellsson (2018) investigates **income-related smoking inequality** among Swedish women.
- The author uses a random effects probit model to determine the  $\beta_j$  coefficients from equation (1).

# Empirical Application: Kjellsson (2018)

- In a recent study, Kjellsson (2018) investigates **income-related smoking inequality** among Swedish women.
- The author uses a random effects probit model to determine the  $\beta_j$  coefficients from equation (1).
- To account for **persistence** in smoking behavior, **lagged outcomes**  $h_{it-1}$  are included in the regressions.

# Empirical Application: Kjellsson (2018)

- In a recent study, Kjellsson (2018) investigates **income-related smoking inequality** among Swedish women.
- The author uses a random effects probit model to determine the  $\beta_j$  coefficients from equation (1).
- To account for **persistence** in smoking behavior, **lagged outcomes**  $h_{it-1}$  are included in the regressions.
- Within-individual means of time-variant variables allow to control for unobserved time-invariant heterogeneity (**Mundlak**-type specification).

# Empirical Application: Kjellsson (2018)

- In a recent study, Kjellsson (2018) investigates **income-related smoking inequality** among Swedish women.
- The author uses a random effects probit model to determine the  $\beta_j$  coefficients from equation (1).
- To account for **persistence** in smoking behavior, **lagged outcomes**  $h_{it-1}$  are included in the regressions.
- Within-individual means of time-variant variables allow to control for unobserved time-invariant heterogeneity (**Mundlak-type** specification).
- The results show an income related smoking inequality in favor of the rich ( $E = -0.084$ ) which is persistent over time.



# Empirical Application: Kjellsson (2018)

- In a recent study, Kjellsson (2018) investigates **income-related smoking inequality** among Swedish women.
- The author uses a random effects probit model to determine the  $\beta_j$  coefficients from equation (1).
- To account for **persistence** in smoking behavior, **lagged outcomes**  $h_{it-1}$  are included in the regressions.
- Within-individual means of time-variant variables allow to control for unobserved time-invariant heterogeneity (**Mundlak-type** specification).
- The results show an income related smoking inequality in favor of the rich ( $E = -0.084$ ) which is persistent over time.
- Main drivers are **education** and living in a **single-adult household**.

# Empirical Application: Kjellsson (2018)

	$V_k$	Static RE probit (Mundlak)			Dynamic RE probit (Mundlak)		
		PE	Contribution	%	PE	Contribution	%
<i>fath_white_high</i>	0.034 (.000)	-0.051 (.024)	-0.007 (.045)	8.4	-0.015 (.403)	-0.002 (.417)	2.5
<i>fath_white_low</i>	0.020 (.001)	-0.029 (.270)	-0.002 (.312)	2.8	-0.002 (.932)	-0.000 (.936)	0.2
<i>fath_farm</i>	-0.023 (.000)	-0.057 (.038)	0.005 (.082)	-6.2	-0.005 (.812)	0.000 (.817)	-0.6
<i>lm2</i>	-0.005 (.113)	0.057 (.198)	-0.001 (.356)	1.5	0.011 (.699)	-0.000 (.753)	0.3
<i>cohort40</i>	0.013 (.036)	0.074 (.030)	0.004 (.113)	-4.7	0.037 (.113)	0.002 (.202)	-2.4
<i>cohort50</i>	0.025 (.000)	0.135 (.025)	0.013 (.050)	-15.9	0.108 (.018)	0.011 (.038)	-12.7
<i>cohort60</i>	-0.015 (.037)	0.100 (.253)	-0.006 (.326)	7.3	0.071 (.290)	-0.004 (.391)	5.2
<i>age</i>	-0.104 (.611)	-0.007 (.048)	0.003 (.653)	-3.4	-0.002 (.368)	0.001 (.745)	-1.2
<i>yrschool</i>	0.540 (.000)	-0.026 (.000)	-0.056 (.000)	66.6	-0.015 (.000)	-0.033 (.000)	39.0
<i>child1</i>	0.004 (.224)	-0.019 (.116)	-0.000 (.392)	0.4	-0.015 (.353)	-0.000 (.533)	0.3
<i>child2plus</i>	-0.002 (.749)	-0.041 (.006)	0.000 (.753)	-0.4	-0.035 (.040)	0.000 (.761)	-0.3
<i>single</i>	-0.068 (.000)	0.032 (.032)	-0.009 (.034)	10.3	0.050 (.018)	-0.014 (.020)	16.3
<i>hinc</i>	0.156 (.000)	0.007 (.572)	0.004 (.571)	-4.9	0.009 (.547)	0.006 (.546)	-6.6
<i>ln_LIFEinc</i>	0.172 (.000)	-0.004 (.905)	-0.003 (.904)	3.4	-0.019 (.470)	-0.013 (.467)	15.7
<i>m_child2plus</i>	0.004 (.224)	-0.076 (.141)	-0.001 (.411)	1.5	-0.036 (.347)	-0.001 (.524)	0.7
<i>m_child1</i>	-0.002 (.749)	-0.195 (.000)	0.001 (.766)	-1.7	-0.074 (.064)	0.001 (.791)	-0.7
<i>m_single</i>	-0.068 (.000)	0.052 (.079)	-0.014 (.082)	16.9	-0.012 (.674)	0.003 (.676)	-3.8
$Y_0$	-0.018 (.006)				0.450 (.000)	-0.032 (.008)	37.8
$Y_{it-1}$	-0.021 (.000)				0.092 (.001)	-0.008 (.014)	9.1
Residual			-0.012	14.7		0.000	0.2
Erreygers' index			-0.084	100		-0.084	100

# Level-Dependent Indices

# Introduction

- Literature on decomposition of health inequality mainly focuses on **rank-dependent indices**.

# Introduction

- Literature on decomposition of health inequality mainly focuses on **rank-dependent indices**.
- However, if one is interested in decomposing an inequality index  $I$  into

# Introduction

- Literature on decomposition of health inequality mainly focuses on **rank-dependent indices**.
- However, if one is interested in decomposing an inequality index  $I$  into
  - within-subgroup inequality  $I_W$

# Introduction

- Literature on decomposition of health inequality mainly focuses on **rank-dependent indices**.
- However, if one is interested in decomposing an inequality index  $I$  into
  - within-subgroup inequality  $I_W$
  - and the between-subgroup inequality  $I_B$

rank-dependent indices usually produce a non-zero residual term  
 $I_X = I - I_W - I_B$ .

# Introduction

- Literature on decomposition of health inequality mainly focuses on **rank-dependent indices**.
- However, if one is interested in decomposing an inequality index  $I$  into
  - within-subgroup inequality  $I_W$
  - and the between-subgroup inequality  $I_B$

rank-dependent indices usually produce a non-zero residual term  $I_X = I - I_W - I_B$ .

- Erreygers et al. (2018) developed a level-dependent index with the property of **subgroup decomposability**.



# Introduction

Within-subgroup inequality  $I_W$ :

$$I_j = \frac{1}{n_j} \sum_{i \in G_j} w_i(\mathbf{y}_j) h_i \quad (5)$$

$$I_W = \sum_{j=1}^k s_j I_j \quad (6)$$

where

$I_j$  inequality index for subgroup  $j$

$n_j$  number of individuals in subgroup  $j$

$G_j$  individuals in subgroup  $j$

$w_i(\mathbf{y}_j)$  weight of individual  $i$  for position within group  $j$

$h_i$  health status of individual  $i$

$s_j$  weight of subgroup  $j$ .

# Introduction

Between-subgroup inequality  $I_B$ :

$$I_B = \frac{1}{n} \sum_{j=1}^k n_j w_j (\mu_y) \mu_{h_j} \quad (7)$$

where

$n$  total number of individuals

$w_j (\mu_y)$  weight reflecting average situation of individuals in subgroup  $j$

$\mu_{h_j}$  mean health status in subgroup  $j$ .

# Rank-Dependent Measures

The standard concentration index  $I^R$  can be written as

$$I^R = \frac{1}{n} \sum_{i=1}^n w_i(\mathbf{y}) h_i \quad (8)$$

with

$$w_i(\mathbf{y}) = \frac{2r_i(\mathbf{y}) - n - 1}{n} \quad (9)$$

where

$w_i(\mathbf{y})$  weight reflecting situation of individual  $i$  in total population.

# Rank-Dependent Measures

## Defining

$$w_i(\mathbf{y}_j) = \frac{2r_i(\mathbf{y}_j) - n_j - 1}{n_j}$$

$$w_j(\boldsymbol{\mu}_y) = \frac{2r_j(\boldsymbol{\mu}_y) - n - 1}{n}$$

with

$$r_j(\boldsymbol{\mu}_y) = \frac{n_j + 1}{2} + \sum_{l=0}^{j-1} n_l, \quad (10)$$

$\mu_{y_1} < \dots < \mu_{y_k}$ ,  $s_j = \frac{n_j}{n}$  and  $n_0 = 0$ , the residual term

$$I_X^R = \frac{1}{n} \sum_{j=1}^k \sum_{i \in G_j} \left[ w_i(\mathbf{y}) h_i - \frac{n_j}{n} w_i(\mathbf{y}_j) h_i - w_j(\boldsymbol{\mu}_y) \mu_{h_j} \right] \quad (11)$$

only equals zero if the subgroup income ranges do not overlap.

# Level-Dependent Measures

If we instead consider **level-dependent** weights

$$\begin{aligned}w_i(\mathbf{y}) &= \frac{y_i - \mu_y}{\mu_y} \\w_i(\mathbf{y}_j) &= \frac{y_i - \mu_{y_j}}{\mu_{y_j}} \\w_j(\boldsymbol{\mu}_y) &= \frac{\mu_{y_j} - \mu_y}{\mu_y},\end{aligned}$$

we obtain the total population inequality  $I^L$  as the sum of the within- ( $I_W^L$ ) and the between-subgroup inequality ( $I_B^L$ ):

$$\begin{aligned}I^L &= I_W^L + I_B^L \\&= \frac{1}{n\mu_y} \sum_{i=1}^n y_i h_i - \mu_h\end{aligned}$$

# Comparing Decompositions: An Example

- Erreygers et al. (2018) estimate inequality in health – measured as SF-6D health score – due to equivalised income for Australian population aged 15+.

Source: Erreygers et al. (2018).

# Comparing Decompositions: An Example

- Erreygers et al. (2018) estimate inequality in health – measured as SF-6D health score – due to equivalised income for Australian population aged 15+.
- Decompose inequality according to sex and compare rank- and level-dependent indices.

Table 3. Decomposition of health inequality by sex.

	$I^R$		$I^L$	
	Values	%	Values	%
Within	0.0325	49.47	0.0141	98.30
Between	0.0210	31.92	0.0002	1.70
Residual	0.0122	18.61	-	-
Total	0.0657	100.00	0.0144	100.00

Source: Erreygers et al. (2018).

# Two-Dimensional Decomposition



# Introduction

- The previously introduced CI decomposition relies on specifying the determinants of health  $h_i$ .

# Introduction

- The previously introduced CI decomposition relies on specifying the determinants of health  $h_i$ .
- However, **determinants of socioeconomic status** are ignored by this approach.

# Introduction

- The previously introduced CI decomposition relies on specifying the determinants of health  $h_i$ .
- However, **determinants of socioeconomic status** are ignored by this approach.
- Kessels and Erreygers (2019) propose an approach which takes determinants of both health and SES into account.

# Introduction

- The previously introduced CI decomposition relies on specifying the determinants of health  $h_i$ .
- However, **determinants of socioeconomic status** are ignored by this approach.
- Kessels and Erreygers (2019) propose an approach which takes determinants of both health and SES into account.
- This direct regression approach is easy to implement and applicable for rank- and level-dependent indices.

# Implementation

- We can rewrite the specific index  $I$  as

$$I = \frac{1}{n} \sum_{i=1}^n w_i h_i = \frac{1}{n} \sum_{i=1}^n u_i = \mu_u \quad (12)$$

where  $w_i$  are the specific index's weights.

# Implementation

- We can rewrite the specific index  $I$  as

$$I = \frac{1}{n} \sum_{i=1}^n w_i h_i = \frac{1}{n} \sum_{i=1}^n u_i = \mu_u \quad (12)$$

where  $w_i$  are the specific index's weights.

- The  $u_i$  values combines both health and socioeconomic performance and can be written as

$$u_i = \beta_0 + \sum_{j=1}^J \beta_j x_{ij} + \eta_i \quad (13)$$

where  $x_{ij}$  are determinants of health and/or SES.

# Implementation

- We can rewrite the specific index  $I$  as

$$I = \frac{1}{n} \sum_{i=1}^n w_i h_i = \frac{1}{n} \sum_{i=1}^n u_i = \mu_u \quad (12)$$

where  $w_i$  are the specific index's weights.

- The  $u_i$  values combines both health and socioeconomic performance and can be written as

$$u_i = \beta_0 + \sum_{j=1}^J \beta_j x_{ij} + \eta_i \quad (13)$$

where  $x_{ij}$  are determinants of health and/or SES.

- Estimating 13 via OLS and inserting into 12, we obtain

$$I = \hat{\beta}_0 + \sum_{j=1}^J \hat{\beta}_j \mu_{x_j} \quad (14)$$

# Interpretation

- For marginal changes  $\Delta\mu_{x_j}$ , the effect on  $I$  is equal to  $\hat{\beta}_j\Delta\mu_{x_j}$ .



# Interpretation

- For marginal changes  $\Delta\mu_{x_j}$ , the effect on  $I$  is equal to  $\hat{\beta}_j\Delta\mu_{x_j}$ .
- $\hat{\beta}_j\mu_{x_j}$  is **not** the contribution of  $x_j$  to the level of inequality.

# Interpretation

- For marginal changes  $\Delta\mu_{x_j}$ , the effect on  $I$  is equal to  $\hat{\beta}_j\Delta\mu_{x_j}$ .
- $\hat{\beta}_j\mu_{x_j}$  is **not** the contribution of  $x_j$  to the level of inequality.
- The importance of specific variables can be evaluated via the logworth statistic defined as

$$\text{Logworth} = -\log_{10}(p) \quad (15)$$

where  $p$  is the  $p$  value of the  $F$  test conducted on a certain set of variables.

# Interpretation

- For marginal changes  $\Delta\mu_{x_j}$ , the effect on  $I$  is equal to  $\hat{\beta}_j\Delta\mu_{x_j}$ .
- $\hat{\beta}_j\mu_{x_j}$  is **not** the contribution of  $x_j$  to the level of inequality.
- The importance of specific variables can be evaluated via the logworth statistic defined as

$$\text{Logworth} = -\log_{10}(p) \quad (15)$$

where  $p$  is the  $p$  value of the  $F$  test conducted on a certain set of variables.

- Kessels and Erreygers (2019) apply this approach to Australian data with SF-6D as health indicator and equivalent income to denote SES.

# Empirical Application - Kessels and Erreygers (2019)

Variable	$\hat{\chi}_j$	Prob >  t	Prob > F	Logworth
Male	0.0260	0.0033		2.488
Indigenous	-0.0967	0.0004		3.418
Age	-0.0015	< 0.0001		7.027
Not married	-0.1145	< 0.0001		30.566
Children 0-4	-0.0653	< 0.0001		12.526
Children 5-14	-0.0450	< 0.0001		12.480
Semi-detached house	0.0065	0.6999		
Flat	-0.0007	0.9611		
Non-private dwelling	-0.2691	0.0016		
Other dwelling	-0.1340	0.0249	0.0044	2.361
Managers & professionals	0.2266	< 0.0001		
Manual workers	-0.0893	< 0.0001		
Unemployed	-0.1702	< 0.0001		
Not in labour force	-0.2250	< 0.0001	< 0.0001	279.915
Living poorly	-0.2272	< 0.0001		
Just getting along	-0.2163	< 0.0001	< 0.0001	103.965
Smoking	-0.0481	< 0.0001		4.521

Very good sleep quality	0.0232	0.0398		
Fairly bad sleep quality	-0.0283	0.0113		
Very bad sleep quality	-0.0547	0.0155		
Not reported	-0.0505	0.1951	0.0006	3.213
Almost always stressed	-0.0197	0.2252		
Often stressed	-0.0049	0.6489		
Rarely stressed	-0.0302	0.0090		
Never stressed	-0.0276	0.2919	0.0895	1.048
Life satisfaction	0.0301	< 0.0001		18.983
Very satisfied with weight	0.0252	0.1268		
Satisfied with weight	-0.0072	0.5408		
Dissatisfied with weight	0.0342	0.0028		
Very dissatisfied with weight	-0.0022	0.8994	0.0012	2.910
No physical activity	-0.0886	< 0.0001		
Some physical activity	-0.0470	< 0.0001	< 0.0001	10.845
Constant	0.0368	0.2773		
$R^2$	0.2111			

Figure 1. Results for Level-Dependent Index

## Summary and Conclusions

# Summary and Conclusions

- The properties of the concentration index depend on the measurement **characteristics** of the health variable of interest.

# Summary and Conclusions

- The properties of the concentration index depend on the measurement **characteristics** of the health variable of interest.
- When the health variable is cardinal and has finite upper and lower bounds, the **Erreygers index**  $E(h)$  and the **Wagstaff index**  $W(h)$  satisfy the desired properties (sign, scale invariance, mirror property) and are superior to the CI.

# Summary and Conclusions

- The properties of the concentration index depend on the measurement **characteristics** of the health variable of interest.
- When the health variable is cardinal and has finite upper and lower bounds, the **Erreygers index**  $E(h)$  and the **Wagstaff index**  $W(h)$  satisfy the desired properties (sign, scale invariance, mirror property) and are superior to the CI.
- Level-dependent indices allow for decomposition into within- and between-subgroup inequality.



# Summary and Conclusions

- The properties of the concentration index depend on the measurement **characteristics** of the health variable of interest.
- When the health variable is cardinal and has finite upper and lower bounds, the **Erreygers index**  $E(h)$  and the **Wagstaff index**  $W(h)$  satisfy the desired properties (sign, scale invariance, mirror property) and are superior to the CI.
- Level-dependent indices allow for decomposition into within- and between-subgroup inequality.
- Applying decomposition methods to inequality indicators (like the CI) allows to analyse income-related inequalities in health across the **entire income distribution** (income proxying for SES).

# Summary and Conclusions

- The properties of the concentration index depend on the measurement **characteristics** of the health variable of interest.
- When the health variable is cardinal and has finite upper and lower bounds, the **Erreygers index**  $E(h)$  and the **Wagstaff index**  $W(h)$  satisfy the desired properties (sign, scale invariance, mirror property) and are superior to the CI.
- Level-dependent indices allow for decomposition into within- and between-subgroup inequality.
- Applying decomposition methods to inequality indicators (like the CI) allows to analyse income-related inequalities in health across the **entire** income **distribution** (income proxying for SES).
- In this case, each source of inequality is quantified – and not just the difference between the two groups.

# Literature I

- ERREYERS, G. (2009): "Correcting the concentration index," *Journal of health economics*, 28, 504–515.
- ERREYERS, G., R. KESSELS, L. CHEN, AND P. CLARKE (2018): "Subgroup Decomposability of Income-Related Inequality of Health, with an Application to Australia," *Economic Record*, 94, 39–50.
- ERREYERS, G. AND T. VAN OURTI (2011): "Measuring socioeconomic inequality in health, health care and health financing by means of rank-dependent indices: a recipe for good practice," *Journal of health economics*, 30, 685–694.
- KESSELS, R. AND G. ERREYERS (2019): "A direct regression approach to decomposing socioeconomic inequality of health," *Health economics*, 28, 884–905.
- KJELLSSON, G. (2018): "Extending decomposition analysis to account for unobserved heterogeneity and persistence in health behavior: Income-related smoking inequality among Swedish women," *Health Economics*, 27, 440–447.
- WAGSTAFF, A. (2005): "The bounds of the concentration index when the variable of interest is binary, with an application to immunization inequality," *Health economics*, 14, 429–432.

## Change in the CI: Oaxaca-Style Approach

- Suppose we want to compare and decompose the **differences** between concentration indices of two populations, for example to analyse how inequality changes in one country over time  $t$ .

# Change in the CI: Oaxaca-Style Approach

- Suppose we want to compare and decompose the **differences** between concentration indices of two populations, for example to analyse how inequality changes in one country over time  $t$ .
- Consider the decomposition of the CI:

$$C = \sum_k \eta_k C_k + \frac{GC_\epsilon}{\mu} \quad (16)$$

# Change in the CI: Oaxaca-Style Approach

- Suppose we want to compare and decompose the **differences** between concentration indices of two populations, for example to analyse how inequality changes in one country over time  $t$ .
- Consider the decomposition of the CI:

$$C = \sum_k \eta_k C_k + \frac{GC_\epsilon}{\mu} \quad (16)$$

- We may apply a Oaxaca-style decomposition:

$$\Delta C = \sum_k \eta_{kt} (C_{kt} - C_{k,t-1}) + \sum_k C_{k,t-1} (\eta_{kt} - \eta_{k,t-1}) + \Delta \left( \frac{GC_{\eta t}}{\mu_t} \right) \quad (17)$$

# Change in the CI: Oaxaca-Style Approach

- Suppose we want to compare and decompose the **differences** between concentration indices of two populations, for example to analyse how inequality changes in one country over time  $t$ .
- Consider the decomposition of the CI:

$$C = \sum_k \eta_k C_k + \frac{GC_\epsilon}{\mu} \quad (16)$$

- We may apply a Oaxaca-style decomposition:

$$\Delta C = \sum_k \eta_{kt} (C_{kt} - C_{k,t-1}) + \sum_k C_{k,t-1} (\eta_{kt} - \eta_{k,t-1}) + \Delta \left( \frac{GC_{\eta t}}{\mu_t} \right) \quad (17)$$

- It gives us differences in inequality at different  $t$  as a sum of:

# Change in the CI: Oaxaca-Style Approach

- Suppose we want to compare and decompose the **differences** between concentration indices of two populations, for example to analyse how inequality changes in one country over time  $t$ .
- Consider the decomposition of the CI:

$$C = \sum_k \eta_k C_k + \frac{GC_\epsilon}{\mu} \quad (16)$$

- We may apply a Oaxaca-style decomposition:

$$\Delta C = \sum_k \eta_{kt} (C_{kt} - C_{k,t-1}) + \sum_k C_{k,t-1} (\eta_{kt} - \eta_{k,t-1}) + \Delta \left( \frac{GC_{\eta t}}{\mu_t} \right) \quad (17)$$

- It gives us differences in inequality at different  $t$  as a sum of:
  - 1 CIs for determinants  $k$  weighted by their elasticities.



# Change in the CI: Oaxaca-Style Approach

- Suppose we want to compare and decompose the **differences** between concentration indices of two populations, for example to analyse how inequality changes in one country over time  $t$ .
- Consider the decomposition of the CI:

$$C = \sum_k \eta_k C_k + \frac{GC_\epsilon}{\mu} \quad (16)$$

- We may apply a Oaxaca-style decomposition:

$$\Delta C = \sum_k \eta_{kt} (C_{kt} - C_{k,t-1}) + \sum_k C_{k,t-1} (\eta_{kt} - \eta_{k,t-1}) + \Delta \left( \frac{GC_{\eta t}}{\mu_t} \right) \quad (17)$$

- It gives us differences in inequality at different  $t$  as a sum of:
  - 1 Cls for determinants  $k$  weighted by their elasticities.
  - 2 Elasticities weighted by the respective Cls.

# Change in the CI: Oaxaca-Style Approach

- Suppose we want to compare and decompose the **differences** between concentration indices of two populations, for example to analyse how inequality changes in one country over time  $t$ .
- Consider the decomposition of the CI:

$$C = \sum_k \eta_k C_k + \frac{GC_\epsilon}{\mu} \quad (16)$$

- We may apply a Oaxaca-style decomposition:

$$\Delta C = \sum_k \eta_{kt} (C_{kt} - C_{k,t-1}) + \sum_k C_{k,t-1} (\eta_{kt} - \eta_{k,t-1}) + \Delta \left( \frac{GC_{\eta t}}{\mu_t} \right) \quad (17)$$

- It gives us differences in inequality at different  $t$  as a sum of:
  - 1 Cls for determinants  $k$  weighted by their elasticities.
  - 2 Elasticities weighted by the respective Cls.
  - 3 Generalised Cls of the residuals.

# Example: Change in the CI, Different Components




Figure 2. Decomposition of change in CI for HAZ-scores of children < 10 y.o. in Vietnam, 1993-98.

# Change in the CI: Total Differential Approach

- However decomposition (17) does not allow to distinguish how changes within the elasticities  $\eta_k$  affect changes in socioeconomic-related inequality.

# Change in the CI: Total Differential Approach

- However decomposition (17) does not allow to distinguish how changes within the elasticities  $\eta_k$  affect changes in socioeconomic-related inequality.
- Consider the derivatives w.r.t. changes in  $\beta_k$  and  $\bar{x}_k$ :

$$\begin{aligned}\frac{dC}{d\beta_k} &= \frac{\partial C}{\partial \beta_k} + \frac{\partial C}{\partial \mu} \frac{d\mu}{d\beta_k} = \frac{\bar{x}_k C_k}{\mu} - \frac{\bar{x}_k}{\mu} C \\ \frac{dC}{d\bar{x}_k} &= \frac{\beta_k}{\mu} (C_k - C)\end{aligned}$$

# Change in the CI: Total Differential Approach

- Get the **total differential** of Eq. (16):

$$\begin{aligned}
 dC = & -\frac{C}{\mu}d\alpha + \sum_k \frac{\bar{x}_k}{\mu} (C_k - C) d\beta_k + \sum_k \frac{\beta_k}{\mu} (C_k - C) d\bar{x}_k \\
 & + \sum_k \frac{\beta_k \bar{x}_k}{\mu} dC_k + d\frac{GC_\epsilon}{\mu}
 \end{aligned} \tag{18}$$

# Change in the CI: Total Differential Approach

- Get the **total differential** of Eq. (16):

$$\begin{aligned} dC = & -\frac{C}{\mu}d\alpha + \sum_k \frac{\bar{x}_k}{\mu} (C_k - C) d\beta_k + \sum_k \frac{\beta_k}{\mu} (C_k - C) d\bar{x}_k \\ & + \sum_k \frac{\beta_k \bar{x}_k}{\mu} dC_k + d\frac{GC_\epsilon}{\mu} \end{aligned} \quad (18)$$

- Changes in  $\beta_k$  and  $\bar{x}_k$  have a **direct** effect on changes in  $C$ .

# Change in the CI: Total Differential Approach

- Get the **total differential** of Eq. (16):

$$\begin{aligned}
 dC = & -\frac{C}{\mu}d\alpha + \sum_k \frac{\bar{x}_k}{\mu} (C_k - C) d\beta_k + \sum_k \frac{\beta_k}{\mu} (C_k - C) d\bar{x}_k \\
 & + \sum_k \frac{\beta_k \bar{x}_k}{\mu} dC_k + d\frac{GC_\epsilon}{\mu}
 \end{aligned} \tag{18}$$

- Changes in  $\beta_k$  and  $\bar{x}_k$  have a **direct** effect on changes in  $C$ .
- They also have an **indirect** effect through  $\mu$ : an increase in inequality in  $\bar{x}_k$  increases the degree of inequality in  $h$ .



# Change in the CI: Total Differential Approach

- Get the **total differential** of Eq. (16):

$$\begin{aligned}
 dC = & -\frac{C}{\mu}d\alpha + \sum_k \frac{\bar{x}_k}{\mu} (C_k - C) d\beta_k + \sum_k \frac{\beta_k}{\mu} (C_k - C) d\bar{x}_k \\
 & + \sum_k \frac{\beta_k \bar{x}_k}{\mu} dC_k + d\frac{GC_\epsilon}{\mu}
 \end{aligned} \tag{18}$$

- Changes in  $\beta_k$  and  $\bar{x}_k$  have a **direct** effect on changes in  $C$ .
- They also have an **indirect** effect through  $\mu$ : an increase in inequality in  $\bar{x}_k$  increases the degree of inequality in  $h$ .
- $C$  increases for increases in  $\beta_k$  and  $\bar{x}_k$ ;  $C$  decreases for increases in  $\mu$ .

# Example: Change in HAZ-Scores

Table 4. Decomposition of changes in the CI for HAZ-scores: Comparison between total differential and Oaxaca-style approach.

Variable	Total differential approach (Eq. ??)					Oaxaca-style approach (Eq. 17)	
	$\beta$	$\bar{x}$	CI	Total	Percent	Total	Percent
Child's age (in months)	0.003	0.011	-0.002	0.012	-57	0.007	-30
Child's age squared	0.003	-0.010	0.001	-0.006	29	-0.006	26
Male	0.001	0.000	0.000	0.001	-5	0.001	-3
Household consumption	-0.005	-0.005	-0.002	-0.011	52	-0.016	74
Safe drinking water	-0.002	0.000	0.000	-0.003	14	-0.003	16
Satisfactory sanitation	0.003	-0.002	0.000	0.001	-5	0.001	-5
Years schooling household head	0.001	0.000	-0.001	0.000	0	0.000	1
Years schooling mother	0.005	0.000	-0.001	0.004	-19	0.003	-11
Commune (fixed effects)	0.000	-0.014	-0.010	-0.025	119	-0.012	55
Residual				0.005	-24	0.005	-24
Total	0.010	-0.021	-0.016	-0.021	100	-0.022	100