An aerial photograph of a coastal region, likely in the Mediterranean. The top half shows a coastline with a river or estuary flowing into the sea. The land is a patchwork of agricultural fields in various shades of green and brown. The bottom half shows a dense urban or agricultural area with a grid-like pattern of fields. The text is overlaid on a semi-transparent white box in the center.

# Using geospatial data to improve targeting of agricultural projects in the context of climate change and resource scarcity

Darius A. Görgen

2023-09-11



# Who am I?

- 🙋 Darius A. Görgen
- 🎓 M.Sc. Geography, B.Sc. Political Sciences
- 🌍 Focus on Climate Change and Agriculture
- 🌐 Part of the MAPME Initiative since 2020
- 🔬 Advocating for OpenSource and OpenScience



E-mail: [info@dariusgoergen.com](mailto:info@dariusgoergen.com)

Website: [www.dariusgoergen.com](http://www.dariusgoergen.com)

Source (title slide): [ESA](#), [CC BY-SA 3.0 IGO](#)


Inception Workshop of the Geo4Impact Program - September 11th, 2023, Paris

# Content

- What is geospatial data?
- From visual interpretation to automated analysis
  - i. Counting trees by hand
  - ii. Deep Learning for field boundary delineation
  - iii. Satellite time series for crop type identification
  - iv. Data fusion for crop biophysical monitoring
- Targeting in the context of agricultural projects
  - i. Mapping flood areas
  - ii. Analyzing climatological drought
  - iii. Water accounting
- Wrap-Up

# Geospatial data

# Geospatial data



Conceptualization of space in the dominant digital formats.

# From visual interpretation to automated analysis

# Visual interpretation



A tree plantation near the [Jordan EcoPark](#).

# Field boundary delineation

- very often our area is too large for manual interpretation
- we need tools that automate the interpretation of satellite imagery
- Meta AI's [Segment Anything](#) Network is already used in the agricultural sector
- field boundaries can be used to inform about the area distribution of farms ...
- ... but also they might be required for later analysis stages



Screenshot of agricultural boundaries produced by SAM.




# Crop type identification

Satellite imagery timeseries ...




Animation of a Sentinel-2 timeseries over an agricultural area.

... reveal distinct signatures of crops over time.



Temporal NDVI profiles of different crop types.

# Data fusion



Visualisation of an data fusion approach from Sentinel 1 and 2 for crop biophysical monitoring.

# Targeting




# Flood areas




Animation of 2022 monsoon floods in Bangladesh.

# Climatological drought

- SPI/SPEI to quantify intensity and duration of meteorological droughts
- SPEI is preferable when temperature or  $ET_0$  data is available
- gridded datasets allow drought analysis even in data scarce regions
- [CHELSA](#) has good performance for complex terrains and data-scarce regions
- includes climate projections for different CMIP6 scenarios





Comparison of SPI and SPEI to characterize meteorological drought.



[CHELSA](#) animation of average precipitation between 1981-2010.

# Water accounting

- Water accounting study in the Jordan River Basin by [FAO](#)
- Uses remote-sensing based variables by FAO's [WAPOR](#)
- Differentiates between water generating ( $P > ET_a$ ) and consuming ( $P < ET_a$ ) land cover classes



Difference between Precipitation ( $P$ ) and Actual Evapotranspiration and Interception ( $ET_a$ ).

Figure 1: Contribution of landcover classes to  $ET_a$  (a) and precipitation (b) in the Jordan River Basin.



# Wrap Up

- 🎯 geospatial data can help to better target areas to maximize benefit
- 🛰️ remote sensing can deliver valuable insights in data scarce regions
- 💬 *All models are wrong, but some are useful.* ([George Box](#))
- 🍏 evaluate the low-hanging fruits first
- 💰 gold standards require large amounts of high-quality and thus expensive data
- 💡 SSL might be a game-changer, but the training of foundation models is expensive

# Thank you for your attention!

# References

- FAO, and IHE Delft. 2020. “Water Accounting in the Jordan River Basin.” *FAO WaPOR Water Accounting Reports*.  
<https://doi.org/https://doi.org/10.4060/ca9181en>.
- Karger, Dirk Nikolaus, Olaf Conrad, Jürgen Böhner, Tobias Kawohl, Holger Kreft, Rodrigo Wilber Soria-Auza, Niklaus E. Zimmermann, H. Peter Linder, and Michael Kessler. 2021. “Climatologies at High Resolution for the Earth’s Land Surface Areas.” *EnviDat*. <https://doi.org/http://dx.doi.org/10.16904/envidat.228>.
- Saab, David. 2003. “Conceptualizing Space: Mapping Schemas as Meaningful Representations.” PhD thesis.  
<https://doi.org/10.13140/2.1.3030.1767>.
- Silva-Perez, Cristian, Armando Marino, and Iain Cameron. 2022. “Learning-Based Tracking of Crop Biophysical Variables and Key Dates Estimation from Fusion of SAR and Optical Data.” *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 15: 7444–57. <https://doi.org/10.1109/JSTARS.2022.3203248>.
- Tariq, Aqil, Jianguo Yan, Alexandre S. Gagnon, Mobushir Riaz Khan, and Faisal Mumtaz. 2022. “Mapping of Cropland, Cropping Patterns and Crop Types by Combining Optical Remote Sensing Images with Decision Tree Classifier and Random Forest.” *Geo-Spatial Information Science* 0 (0): 1–19. <https://doi.org/10.1080/10095020.2022.2100287>.
- Vicente-Serrano, Sergio M., Santiago Beguería, and Juan I. López-Moreno. 2010. “A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index.” *Journal of Climate* 23 (7): 1696–1718.  
<https://doi.org/https://doi.org/10.1175/2009JCLI2909.1>.



0 Comments - powered by utteranc.es

Write Preview

Sign in to comment

 Styling with Markdown is supported

Sign in with GitHub

