

HW2 Solution

①

1) Using a shift cipher to encrypt a 1 letter message is similar to using a one time pad - any plaintext could encrypt to any ciphertext and there is not enough information revealed from the ciphertext to allow further analysis (ie: frequency analysis). In this case,

$H(P) = H(P|C) \Rightarrow$ perfect secrecy is maintained

Let's suppose all 26 english letters are equal in frequency

$$H(P) = -26 \left(\frac{1}{26} \log_2 \frac{1}{26} \right)$$

\uparrow 26 total letters \uparrow $p(x) = \frac{1}{26}$ for ea. letter

$$H(P) = -\log_2 \frac{1}{26}$$

$$H(P|C) = -\sum p(c) \sum p(p|c) \log_2 p(p|c)$$

$$P(c) = \left(\frac{1}{26} \right) \left(\frac{1}{26} \right) + \left(\frac{1}{26} \right) \left(\frac{1}{26} \right) + \dots$$

\uparrow $p(PT=a)$ \uparrow $p(k \text{ that encrypts } a)$ \nwarrow repeated for ea. of the 26 plaintext

$$= 26 \left(\frac{1}{26} \right) \left(\frac{1}{26} \right) = \frac{1}{26}$$

$$P(P|C) = \frac{P(P,C)}{P(C)} = \frac{P(P)P(K)}{P(C)} = \frac{\left(\frac{1}{26} \right) \left(\frac{1}{26} \right)}{\frac{1}{26}} = \frac{1}{26}$$

\uparrow for each combo of P/C

$$H(P|C) = -26 \left[\frac{1}{26} \left(26 \left(\frac{1}{26} \log_2 \frac{1}{26} \right) \right) \right]$$

\uparrow repeated for all cipher letters \uparrow $p(c)$ for ea. letter \uparrow 26 letter combos \uparrow $p(p|c)$

$$H(P|C) = -\log_2 \frac{1}{26}$$

$$\therefore H(P) = H(P|C)$$

2)	<u>a</u>	<u>b</u>	<u>c</u>
	C	A	B
K_1			
K_2	A	B	C
K_3	B	C	A

$$p(a) = 0.8$$

$$p(b) = 0.15$$

$$p(c) = 0.05$$

$$p(K_1) = \frac{1}{3}$$

$$p(K_2) = \frac{1}{3}$$

$$p(K_3) = \frac{1}{3}$$

$$a) H(p) = - \sum_{x \in P} p(x) \log_2 p(x)$$

$$= - [p(a) \log_2 p(a) + p(b) \log_2 p(b) + p(c) \log_2 p(c)]$$

$$= - [0.8 \log_2 0.8 + 0.15 \log_2 0.15 + 0.05 \log_2 0.05]$$

$$= 0.884$$

b) see next page

(2)

$$b) H(P/C) = - \sum p(C) \sum p(P/C) \log_2 p(P/C)$$

* determine the necessary components:

$p(C)$ for ea. cipher-text value:

$$p(A) = p(b)p(k_1) + p(a)p(k_2) + p(c)p(k_3) \\ = 0.333 \text{ (or } \frac{1}{3})$$

$$p(B) = p(a)p(k_1) + p(b)p(k_2) + p(c)p(k_3) \\ = 0.333 \text{ (or } \frac{1}{3})$$

$$p(C) = p(a)p(k_1) + p(c)p(k_2) + p(b)p(k_3) \\ = 0.333 \text{ (or } \frac{1}{3})$$

$p(P/C)$ for ea. PT and CT combo:

$$p(a/A) = \frac{p(a)p(k_2)}{p(A)} = \frac{(0.8)(\frac{1}{3})}{(\frac{1}{3})} = 0.8$$

$$p(b/A) = 0.15$$

$$p(c/A) = 0.05$$

$$p(a/B) = 0.8$$

$$p(b/B) = 0.15$$

$$p(c/B) = 0.05$$

} same as above

$$p(a/C) = 0.8$$

$$p(b/C) = 0.15$$

$$p(c/C) = 0.05$$

} same as above

$$H(P/C) = - \left[\left(\frac{1}{3} \right) (0.8 \log_2 0.8 + 0.15 \log_2 0.15 + 0.05 \log_2 0.05) + \dots \right] \\ = \left[3 \left(\frac{1}{3} \right) (0.8 \log_2 0.8 + 0.15 \log_2 0.15 + 0.05 \log_2 0.05) \right] \\ = \underline{\underline{0.884}}$$

③
c) No information about the plaintext is revealed by the ciphertext because $H(p) = H(p|c)$.

If, however $E_{k_2}(a) = C$ and $E_{k_2}(b) = B$, this would not be the case. In that scenario $H(p|c) < H(p)$ because it is more likely that B decrypts to b than any other plaintext.

3) problem #1a pg 107

$$101 = 5 \cdot 17 + 16$$

$$17 = 16 \cdot 1 + 1 \leftarrow \gcd(17, 101) = 1 \quad \checkmark$$

$$1 = 17 \cdot 1 - 16 \cdot 1 \quad 1 \text{ as linear combination of } 16, 17$$

$$16 = 101 \cdot 1 + 17(-5) \quad \text{substitute ...}$$

$$1 = 17 \cdot 1 - (101 \cdot 1 + 17(-5))$$

$$= 17 \cdot 6 + 101(-1)$$

$$\boxed{x=6 \quad y=-1}$$

$$④) \quad 101 = (6)15 + 11$$

$$15 = (1)11 + 4$$

$$11 = (2)4 + 3$$

$$4 = (1)3 + 1 \quad \gcd(101, 15) = 1 \text{ therefore } 15^{-1}(\text{mod } 101) \text{ does in fact exist!}$$

$$3 = (3)1 + 0$$

$$1 = 4 - (1)3$$

$$= 4 - (1)(11 - (2)4)$$

$$= 4 - (1)11 + (2)4$$

$$= (3)4 - (1)11$$

$$= (3)(15 - (1)11) - (1)11$$

$$= (3)15 - (3)11 - (1)11$$

$$= (3)15 - (4)11$$

$$= (3)15 - (4)(101 - (6)15)$$

$$= (3)15 - (4)101 + (24)15$$

$$= (27)15 - (4)101$$

$$\boxed{15^{-1}(\text{mod } 101) \equiv 27}$$

$$\boxed{101^{-1}(\text{mod } 15) \equiv -4 \equiv 11(\text{mod } 15)}$$

5) prob #4 pg 104

$$30030 \bmod 257 = 218$$

$$257 \bmod 218 = 39$$

$$218 \bmod 39 = 23$$

$$39 \bmod 23 = 16$$

$$23 \bmod 16 = 7$$

$$16 \bmod 7 = 2$$

$$7 \bmod 2 = 1 \quad \checkmark$$

Just use Euclidean Algorithm (not extended), since all you need is gcd.

$$\gcd(257, 30030) = 1$$

b) $901^{-1} \pmod{2968}$

$$2968 = (3)901 + 265$$

$$901 = (3)265 + 106$$

$$265 = (2)106 + \boxed{53}$$

$$106 = (2)53 + 0$$

$$\gcd(901, 2968) = 53 \quad \ddots$$

No inverse exists!

7) a) $12x \equiv 28 \pmod{42}$

$$6x \equiv 14 \pmod{21}$$

↑

3 & 14 therefore

No solution

$\gcd(c, r, n) = 2$, so reduce eqn

$\gcd(c, n) = 3$ therefore:

1) cycle repeats $\frac{21}{3} = 7$

2) all r should be multiple of 3

b) $12x \equiv 30 \pmod{42}$ $\gcd(c, n) = 6$ so reduce ^⑥
(expect 6 answers)

$2x \equiv 5 \pmod{7}$ $\gcd(c, n) = 1$

1) cycle repeats every
 $n = 7$

2) all values of r are possible

$2^{-1} \pmod{7} = 4$ because $2 \cdot 4 = 8 \equiv 1 \pmod{7}$ and $3 \cdot 7 = 21$

$2^{-1} \cdot 2x \equiv 5 \cdot 2^{-1} \pmod{7}$

$x \equiv 55 \pmod{7}$

$x \equiv 6 \pmod{7}$

$x \equiv \{6, 13, 20, 27, 34, 41\}$

8 a) $x^2 \equiv 10 \pmod{31}$ 31 is prime, $31 \equiv 3 \pmod{4}$

check $x \equiv \pm 10^{(31+1)/4} \pmod{31}$ $x \equiv \pm 10^8 \pmod{31} = \{14, 17\}$ try these

$14^2 \equiv 10 \pmod{31} \checkmark$ so $x = \{14, 17\}$

b) $x^2 \equiv 9 \pmod{24}$ 24 composite, split into $3 \cdot 8$

$x^2 \equiv 0 \pmod{3}$ (1 soln) $x^2 \equiv 1 \pmod{8}$ $x = \{1, 3, 5, 7\}$ (4 solns)

$x \equiv 0 \pmod{3}$	$x \equiv 0 \pmod{8}$	$x \equiv 0 \pmod{3}$	$x \equiv 0 \pmod{8}$
$x \equiv 1 \pmod{3}$	$x \equiv 3 \pmod{8}$	$x \equiv 5 \pmod{3}$	$x \equiv 7 \pmod{8}$

$x = 9$

$x = 3$

$x = 21$

$x = 15$

find with trial error or spreadsheet or Gauss

$x = \{3, 9, 15, 21\}$

c) $x^2 \equiv 5 \pmod{17}$ 17 prime, $5 \neq 1$, $17 \not\equiv 3 \pmod{4}$ trial error or Excel

no solution

THE UNIVERSITY OF CHICAGO

DEPARTMENT OF CHEMISTRY

PHYSICAL CHEMISTRY

PROFESSOR OF CHEMISTRY

CHICAGO, ILL.

1911

TO THE EDITOR OF THE JOURNAL OF CHEMICAL PHYSICS

DEAR SIR

I have the honor to acknowledge the receipt of your letter of the 10th inst.

and in reply to inform you that