

On multiple infections by parasites with complex life cycles

1 ~~Manuscript elements: Figure 1, figure 2, figure 3, figure 4, figure 5, figure 6, figure 7,~~
2 ~~table 1~~

3 ~~Keywords: host manipulation, multiple infections, parasite pool, trophically transmitted~~
4 ~~parasites~~

5 ~~Manuscript type: e-article.~~

6 **Abstract**

7 Host manipulation is a common strategy of parasites of different complexity.
8 Host manipulation directly affects predator-prey dynamics in trophically transmit-
9 ted parasites, where parasite transmission requires predation. Theoretical studies
10 suggest that manipulation that enhances predation often results in a heavy bur-
11 den on the prey population. Consequently, the system is often destabilised,
12 making parasites prone to extinction. Host manipulation, however, can also sup-
13 press predation. Such suppression is possible if multiple parasites coinfect a host
14 with conflicting interests in manipulation. The interests could be misaligned for
15 various reasons, such as limited carrying capacity or parasitoid developmental
16 stage. Multiple infections are a norm in parasite ecology but are often neglected
17 in the theoretical assessment of host-parasite dynamics. We tease apart the
18 effect of host manipulation of coinfecting parasites and manipulation interests
19 via a mathematical model of a trophically transmitted parasite with a complex
20 life cycle. The life cycle comprises a free-living state, an intermediate and a
21 definitive host. With coinfection, we show that host manipulation that enhances
22 predation need not permanently destabilise the predator-prey system. However,
23 ~~cooperation between coinfecting parasites leading to increased predation and can~~

~~lead to sabotage in manipulation can induce~~ bistability such that a slight disturbance in the system drives the parasite population to extinction. ~~On the other hand, when coinfecting parasites sabotage the manipulative ability of one another, the stability of the predator-prey system is always guaranteed~~ Intriguingly, cooperation in both aspects, host manipulation and reproduction, might ensure system stability. In some cases, a lack of cooperation in reproduction may actually prevent the dynamical system from bistability. Our study highlights the necessity and means of incorporating the reality of multiple parasites and their multi-trophic life cycles in a single system in studying parasite ecology.

Introduction

Parasites infect life on earth ubiquitously, and many of these parasites have complex life cycles (Zimmer, 2001). While a complex lifecycle can be defined as abrupt ontogenic changes in morphology and ecology (Benesh, 2016), a complex parasitic lifecycle typically involves numerous hosts that a parasite needs to traverse to complete its life cycle. This complex lifecycle results in the evolution of various strategies that enable the success of parasite transmission from one host to another. One famous strategy that inspires many science fiction movies and novels is host manipulation, where a parasite can alter the morphology and/or behaviour of its host to enhance its transmission to the next host (Hughes et al., 2012). Host manipulation has been shown in many host-parasite systems, from parasites with simple life-cycle to those with complex life-cycle that involves more than one host (Hughes et al., 2012; Molyneux and Jefferies, 1986). For instance, sand flies infected by *Leishmania* parasites bite more and take more time for a blood meal from mammals (the definitive host of *Leishmania*) compared to their uninfected counterparts (Rogers and Bates, 2007). Copepods infected by cestode parasites are more active and accessible to sticklebacks (the definitive hosts of the cestodes) compared to uninfected copepods (Wedekind and Milinski, 1996).

Theoretical studies have long attempted to understand the ecological and evolutionary consequences of host manipulation. Roosien et al. (2013) and Hosack et al. (2008) showed that manipulative parasites could increase the disease prevalence in an epidemic. Gandon (2018) studied the evolution of the manipulative ability of infectious disease parasites, showing different evolutionary outcomes depending on whether the pathogen can control its vector or host. Haderl and Freedman (1989); Fenton and Rands (2006) and Rogawa et al. (2018) showed that host manipulation could stabilise or destabilise the predator-prey dynamics depending on how manipulation affects the predation response function and the assumption on the fertility of the definitive infected host. Seppälä and Jokela (2008) showed that host

manipulation could evolve even when it increases the risk of the intermediate host being eaten by a non-host predator, given that the initial predation risk is sufficiently low. These models, however, lack a crucial aspect of parasite dynamics, multiple infections (Kalbe et al., 2002)

Typical studies do not consider multiple infections, a phenomenon that is the norm rather than an exception in parasitism. Multiple infections result in the coinfection of more than one parasite inside a host, which may alter the manipulative outcomes (figure 1). An alignment of interest between coinfecting parasites may enhance manipulation, while a conflict of interest may reduce the manipulative effect. Indeed, Hafer and Milinski (2015) showed that copepods infected by two cestode parasites reduce the activity of copepods when both parasites are at the same noninfectious stage, i.e. both parasites are not ready to transmit. Thus the reduction in mobility is suggested to reduce the predation rate by the definitive hosts. When two infectious parasites infect the copepods, the copepods' activity increases, and so does the predation risk for the copepod. However, when the copepods are infected by one infectious and one noninfectious parasite, their interests clash, and one parasite wins over the other.

Theoretical work that considers multiple infections often focuses on the evolution of virulence (van Baalen and Sabelis, 1995; Alizon et al., 2013; Alizon and van Baalen, 2008; Choisy and de Roode, 2010; Alizon, 2012). They show that multiple infections can increase virulence (van Baalen and Sabelis, 1995; Choisy and de Roode, 2010). Evolutionary branching of a less virulent and a hypervirulent parasite can occur when within-host dynamics are considered (Alizon and van Baalen, 2008), and a reduction in virulence is possible if parasites are co-transmitted (Alizon, 2012). ~~In epidemiological models, higher virulence often assumes a link with a higher transmission rate; virulence is therefore associated with host manipulation in such cases~~As host manipulation is suggested to affect transmission rate, these studies also involve host manipulation to a certain extent even though they do not explicitly consider the trait. Host manipulation in trophically transmitted parasite receive less attention. Although manipulation is correlated with transmission rate in both trophically transmitted parasites and infectious disease, there are subtlely difference. Host manipulation influences the predation rate in trophically transmitted parasites, predominantly affecting predator-prey dynamics. Theoretical studies on host manipulation in trophically transmitted parasites with multiple infections are rare (Parker et al., 2003; Vickery and Poulin, 2009). Moreover, they do not consider the prey-predator dynamics, which will likely have important feedback on the evolution of host manipulation. A few studies considering the prey-predator dynamics do not incorporate multiple infections (Rogawa et al., 2018; Iritani and Sato, 2018; Haderler and Freedman, 1989; Fenton and Rands, 2006). More importantly, they assume that transmission from definitive hosts to intermediate hosts is due to direct contact between the



Figure 1: **Who is in control?.** Schistocephalus eggs, which overwinter at the bottom of bodies of water, hatch into microscopically small swimming larvae. These larvae are eaten by copepods (also known as Cyclops due to its single eye), where they develop to the second larval stage. However, the copepod is only the first intermediate host. The larvae are then eaten by sticklebacks, where they reach the third larval stage and grow significantly in size and weight. For the parasite to successfully reach its final host, a warm-blooded animal like a bird, it manipulates its intermediate hosts. The timing is crucial as the chances of success are greatest if the larvae develop in the copepod for 13 to 15 days before entering the stickleback. The presence of multiple parasites in the same host can lead to competition and strategic decision pertaining to investment in manipulation and growth. And indeed a stickleback can be infected by numerous tapeworms as shown above by Martin Kalbe.

94 two types of hosts. This is often not the case, as parasites are released from the defini-
 95 tive hosts into the environment. Transmission happens only when intermediate hosts have
 96 contact with this free-living parasite pool.

97 Our study addresses the gap in the theoretical work on host manipulation in trophically
 98 transmitted parasites. We include multiple infections and consider the dynamics of the free-
 99 living parasite pool. Our compartment model helps illustrate a parasite's complex lifecycle
 100 with two hosts: an intermediate host preyed upon by a definitive host. Transmission from
 101 the intermediate host to the definitive host occurs when predation on infected intermediate
 102 hosts happens. Reproduction only happens in the definitive hosts. New parasites are then

103 released into the environment, where they again have contact with the intermediate hosts
 104 to complete their lifecycle. We focus on the manipulation of the intermediate hosts, such
 105 that the parasite increases the predation rate on the intermediate host by the definitive host
 106 to increase its transmission rate. We then analyse the effect of host manipulation on the
 107 ecological dynamics in the prey-predator-parasite system. In contrast to the examples men-
 108 tioned above our model consists of a single intermediate host as it already provides enough
 109 complexity to discuss between transmission and manipulation. We found that ~~cooperation~~
 110 ~~sabotage~~ in host manipulation ~~leads to bistability in the predator-prey system, given that~~
 111 ~~reproduction from multiple infections is sufficiently high. This finding almost always pushes~~
 112 ~~the dynamical system toward bistability, provided that the reproduction in single infection~~
 113 ~~is sufficiently small. The bistable nature~~ suggests that the predator-prey parasite system is
 114 finely balanced and susceptible to extinction via ecological disturbances. Initially ~~surprising,~~
 115 ~~we show how sabotage in host manipulation guarantees a unique stable equilibrium in the~~
 116 ~~system~~ ~~surprising, we showed that cooperation in both host manipulation and reproduction~~
 117 ~~is not always beneficial and might expose the parasite population to the risk of extinction.~~

118 Model and Results

119 Our model concerns the complex lifecycle of a trophically transmitted parasite that requires
 120 two hosts: an intermediate host and a definitive host. Reproduction only happens inside the
 121 definitive hosts, releasing new parasitic progeny in the environment. An intermediate host
 122 can be infected if it encounters this free-living parasite pool. Finally, when a definitive host
 123 consumes an infected intermediate host, the definitive host gets infected, and the parasite
 124 completes its lifecycle.

For simplicity, we assume that hosts can be infected by one (single infection) or, at most,
 two parasites (double infections). Our model is, therefore, more relevant to the macropara-
 sitic system. Given that infection occurs, the probability that two parasites from the parasite
 pool co-transmit to an intermediate host is denoted by p . Thus $1 - p$ is the probability that a
 single parasite enters an intermediate host. When a definitive host consumes an intermediate
 host infected by two parasites, there is a probability q that the parasites co-transmit to the
 definitive host. With probability $1 - q$, only one parasite successfully transmits. This formu-
 lation assumes that infection always happens when hosts encounter parasites. The dynamics
 of a complex lifecycle parasite that requires two hosts is described by the following system

of equations, firstly for the intermediate host as,

$$\begin{aligned}\frac{dI_s}{dt} &= R(I_s, I_w, I_{ww}) - dI_s - P_s(D_s, D_w, D_{ww})I_s - \eta I_s \\ \frac{dI_w}{dt} &= (1-p)\eta I_s - (d + \alpha_w)I_w - P_w(D_s, D_w, D_{ww}, \beta_w)I_w \\ \frac{dI_{ww}}{dt} &= p\eta I_s - (d + \alpha_{ww})I_{ww} - P_{ww}(D_s, D_w, D_{ww}, \beta_{ww})I_{ww}\end{aligned}\quad (1)$$

where $R(I_s, I_w, I_{ww})$ represents the birth rate of the intermediate hosts, a function of both infected and uninfected individuals. P_s , P_w , P_{ww} are the predation functions of definitive hosts on susceptible, singly infected and doubly infected intermediate hosts. The predation function depends on the density of the definitive hosts and the manipulative strategies of parasites in the intermediate hosts. In particular, if a single parasite infects an intermediate host, the manipulation strategy is β_w . However, if the intermediate host is co-infected, the manipulation strategy is β_{ww} . In the scope of this model, we assume no specific relationship between β_w and β_{ww} to explore all possible ecological outcomes of the system. The force of infection by parasites in the environment is denoted by $\eta = \gamma W$. Since parasites can manipulate intermediate and definitive hosts, here, whenever we mention host manipulation, it specifically refers to the manipulation in intermediate hosts, which correlates to the predation rate.

For the definitive hosts we have,

$$\begin{aligned}\frac{dD_s}{dt} &= B(D_s, D_w, D_{ww}, I_s, I_w, I_{ww}) - \mu D_s - (2\lambda_{ww} + \lambda_w)D_s \\ \frac{dD_w}{dt} &= (\lambda_w + 2(1-q)\lambda_{ww})D_s - (\mu + \sigma_w)D_w - (2(1-q)\lambda_{ww} + \lambda_w)D_w \\ \frac{dD_{ww}}{dt} &= 2q\lambda_{ww}D_s + (2(1-q)\lambda_{ww} + \lambda_w)D_w - (\mu + \sigma_{ww})D_{ww}\end{aligned}\quad (2)$$

where $B(D_s, D_w, D_{ww}, I_s, I_w, I_{ww})$ represents the birth rate of definitive hosts. The birth rates depend on the density of both intermediate and definitive hosts, infected or uninfected. The force of infection that corresponds respectively to singly infected intermediate host (I_w) and doubly infected intermediate hosts (I_{ww}) is denoted respectively by $\lambda_w = \beta_w I_w$ and $\lambda_{ww} = \beta_{ww} I_{ww}$. $\lambda_w = h_1(\rho + \beta_w)I_w$ and $\lambda_{ww} = h_2(\rho + \beta_{ww})I_{ww}$, where ρ is the based line predation rate and h_1 and h_2 are the probability that the parasite successfully established inside the host. If there is no manipulation, that is, $\beta_w = \beta_{ww} = 0$, the parasite is still transmitted via the based line predation. The dynamics of the free-living parasites in the

environment are then given by,

$$\frac{dW}{dt} = f_w D_w + f_{ww} D_{ww} - \delta W - \eta I_s. \quad (3)$$

137 Definitions of different parameters can be found in Table 1.

138 Here, we focus on manipulation that enhances transmission from intermediate hosts to
 139 definitive hosts; we thus simplify the transmission from the parasite pool to intermediate
 140 hosts such that no sequential infection occurs at this transmission state. This assumption
 141 may not be implausible, given that the prey' lifecycle is often shorter than that of the
 142 predator. A prey likely encounters the free-living parasite pool once and then either die due
 143 to predation, making sequential transmission less likely at this state. Sequential infection
 144 can happen when parasites transmit from intermediate hosts to definitive hosts. Therefore,
 145 a singly infected definitive host can be further infected by another parasite if it consumes
 infected intermediate hosts. The system's dynamics are illustrated in figure (2).




Figure 2: Schematic of the model. Blue ovals represent intermediate host compartment, green diamonds represent definitive host compartment, and the transparent hexagon represents the parasite pool compartment with red circles illustrating individual parasites.

146

147 Basic reproduction ratio R_0 of the parasites

The basic reproduction ratio R_0 (or basic reproduction number as often used in epidemiology) indicates parasite fitness. It can be understood as the expected number of offspring a parasite produces during its lifetime when introduced to a susceptible host population. We calculate the basic reproduction ratio R_0 using the next-generation method (Diekmann et al., 1990, 2009; Hurford et al., 2010) (See SI1 for details).

$$R_0 = \overbrace{\gamma I_s^* \frac{2pqh(\rho + \beta_{ww})}{\alpha_{ww} + d + P_{ww}} \frac{D_s^*}{\mu + \sigma_{ww}} \frac{f_{ww}}{\delta + \gamma I_s^*}}^{\text{Double infections}} + \underbrace{\gamma I_s^* \left(\frac{(1-p)h(\rho + \beta_w)}{\alpha_w + d + P_w} + \frac{2p(1-q)h(\rho + \beta_{ww})}{\alpha_{ww} + d + P_{ww}} \right) \frac{D_s^*}{\mu + \sigma_w} \frac{f_w}{\delta + \gamma I_s^*}}_{\text{Single infection}} \quad (4)$$

148 where I_s^* and D_s^* are the densities of susceptible intermediate and definitive hosts at the
 149 disease-free equilibrium. Here, the expression of R_0 contains the possible reproduction routes
 150 of a parasite, which can be via double or single infections. The first component corresponds
 151 to the double infections route, in which the focal parasite co-transmits with another par-
 152 asite into a susceptible intermediate host, then co-transmits into a susceptible definitive
 153 host and reproduces. Here, parasites are so rare that only co-transmission matters and the
 154 compartments with sequential infections are therefore neglected. The second component
 155 corresponds to the single infection route, wherein the focal parasite infects a susceptible
 156 intermediate host via single or double infections. The parasite then transmits alone into the
 157 susceptible definitive host and eventually reproduces.

If $R_0 > 1$, a parasite spreads when introduced into the disease-free equilibrium of prey and predator. Intuitively, the higher the density of susceptible intermediate and definitive hosts, the larger the value of R_0 as the infection reservoir is more extensive. In contrast, regardless of the explicit form of the predation function, the higher the predation rate P_w and P_{ww} , the lower the value of R_0 given the smaller reservoir of intermediate hosts. The effect of host manipulation on the value of R_0 is not so straightforward; as host manipulation becomes efficient, the transmission rate from the intermediate host to the definitive host increases, but so does the predation rate. A higher predation rate results in a smaller intermediate host reservoir available for the parasites to infect. To understand the effect of manipulation on parasites' fitness and the system's ecological dynamics, we next specify the predation

functions. We consider linear functions for predation to begin with,

$$\begin{aligned} P_s(D_s, D_w, D_{ww}) &= \rho D_{total} \\ P_w(D_s, D_w, D_{ww}, \beta_w) &= (\rho + \beta_w) D_{total} \\ P_{ww}(D_s, D_w, D_{ww}, \beta_{ww}) &= (\rho + \beta_{ww}) D_{total} \end{aligned}$$

158 where $D_{total} = D_s + D_w + D_{ww}$ is the total density of the definitive hosts, and ρ is the
159 baseline capture rate of the predator on the prey. If an intermediate host is infected, it is
160 captured by the definitive hosts with rate $\rho + \beta_w$ if it is singly infected and with rate $\rho + \beta_{ww}$
161 if it is doubly infected. Zero values for β_w and β_{ww} suggest no manipulation, and predation
162 is at the baseline value ρ .

For simplicity, we also consider a linear function of the birth of definitive hosts

$$B(D_s, D_w, D_{ww}, I_s, I_w, I_{ww}) = \rho c D_{total} I_{total}$$

163 where c is the efficiency of converting prey into predator's offspring, and $I_{total} = I_s + I_w + I_{ww}$
164 is the total density of the intermediate hosts. It is important to note that host manipulation
165 affects the population dynamics via its influence on predation rate but not the physiological
166 aspect of the definitive host, i.e. the predator. The birth rate of the predators thus depends
167 on the capture rate, but it is not affected by host manipulation, as to our best knowledge,
168 there is no supporting evidence to necessarily consider otherwise.

169 The explicit form of I_s^* and D_s^* , capturing the predator-prey dynamics, depends on the
170 precise form of all birth and predation functions B, R, P_s, P_w and P_{ww} . But, it does not
171 depend on the manipulation ability or any other parameter of the parasite. Given that the
172 birth rate of the predator and the predation rate are linear functions in prey and predator
173 density, the form of the birth rate R of the prey has a significant effect on the susceptible
174 intermediate and definitive host dynamics. ~~Hence we explore the exact structure of these~~
175 ~~birth functions next.~~

176 ~~Linear birth~~ Birth function of intermediate hosts

~~Here, we consider the system when the birth function R of the intermediate host is linear,~~
~~specifically, $R(I_s, I_w, I_{ww}) = r I_{total}$. The equilibrium of intermediate and definitive hosts in~~

the disease-free state are,

$$I_s^* = \frac{\mu}{c\rho}$$

$$D_s^* = \frac{r - d}{\rho}$$

177 ~~This~~
 178 The simplest form of the prey's birth rate is a linear function, in which case the disease
 179 free equilibrium is always unstable. In particular, it has a cyclic behaviour because, at this
 180 equilibrium, the jacobian matrix of the system (1, 2, 3) always has two pure imaginary
 181 eigenvalues (see SI2). This follows from the Lotka-Volterra system using linear functions
 182 for prey birth and predation (Lotka, 1920). Since the disease-free dynamics is cyclic, it
 183 is difficult to analyse the spread of a parasite (~~often using the basic reproduction ratio,~~
 184 which is evaluated when the disease-free state is stable). Here, $R_0 > 1$ happens when
 185 γ , the transmission rate from the environment to intermediate hosts, and the reproduction
 186 rates f_w, f_{ww} are significantly large (the specific mathematical conditions can be found in
 187 SI2SI3). However, even when this condition is satisfied, the parasite may not be able to
 188 spread and persist in cyclic susceptible host dynamics (Figure 3). This result agrees with the
 189 conclusion in (Ripa and Dieckmann, 2013), which suggests that it is difficult for a mutant
 190 to invade a cyclic resident population. In our case, it is not the invasion of a mutant in a
 191 resident population but the invasion of a parasite in a cyclic disease-free host population;
 192 the argument, however, remains valid in both cases. This issue deserves a more thorough
 193 investigation, which is out of the scope of this article. ~~We, therefore,~~ Here, we choose a
 194 non-linear birth function of the intermediate hosts to obtain a stable disease circulation state
 195 and focus on the effect of host manipulation on the ecological dynamics.

196 **Non-linear birth function of intermediate hosts**

~~We chose the~~ The logistic growth for the non-linear birth function ~~, that is follows by~~

$$R(I_w, I_s, I_{ww}) = rI_{total}(1 - kI_{total})$$

where k is the intraspecific competition coefficient. The disease-free equilibrium is as follows

$$I_s^* = \frac{\mu}{c\rho} ; D_s^* = \frac{c\rho(r - d) - k\mu r}{c\rho^2}$$




Figure 3: Disease-free equilibrium using linear birth function, where parasite goes extinct (left panel), and susceptible hosts demonstrate cyclic dynamics (right panel). Solid gray line indicate the density of free-living parasites, blue lines indicate infected intermediate hosts while red lines indicate infected definitive hosts. Dashed lines indicate singly infected hosts while dot-dashed lines indicate doubly infected hosts. Parameter values $\rho = 1.2$, $d = 0.9$, $r = 2.5$, $\gamma = 2.9$, $\alpha_w = \alpha_{ww} = 0$, $\beta_w = 1.5$, $\beta_{ww} = 1.5$, $p = 0.1$, $c = 1.4$, $\mu = 0.9$, $\sigma_w = \sigma_{ww} = 0$, $q = 0.01$, $f_w = 6.5$, $f_{ww} = 7.5$, $\delta = 0.9$, $R_0 = 2.233$, $h_1 = h_2 = 0.8$, $R_0 = 4.997$

This equilibrium is ~~stable if the following three conditions are met,~~

$$\begin{aligned}
 & \text{(i) } r > d \\
 & \text{(ii) } \frac{2c\rho \left(\sqrt{\frac{-d+\mu+r}{\mu}} - 1 \right)}{r} \leq k < \frac{c\rho(r-d)}{\mu r} \\
 & \text{(iii) } \mu > \frac{4c^2\rho^2r - 4c^2d\rho^2}{4ck\rho r + k^2r^2}.
 \end{aligned}$$

197 The above conditions suggest that (i) the intrinsic reproduction of intermediate hosts r needs
 198 to be greater than their natural mortality rate d . More importantly, (ii) the intraspecific
 199 competition coefficient has to be within a range allowing the population to survive. Finally,
 200 (iii) the definitive host's natural mortality rate must be sufficiently large. Satisfying such

201 ~~conditions, we obtain a stable disease-free equilibrium~~ (positive and stable if components
 202 ~~of the parasite, such as reproduction and transmission are sufficiently small, details of the~~
 203 ~~condition can be found in SI4 (Figure 4B).~~




Figure 4: A, B) Disease free equilibrium where parasite cannot persist. C, D) Disease stable equilibrium. Annotations are the same in figure 3. Parameters for disease free equilibrium $\rho = 1.2, d = 0.9, r = 2.5, \gamma = 2.9, \alpha_w = \alpha_{ww} = 0, \beta_w = \beta_{ww} = 1.5, p = 0.1, e = 1.4, \mu = 3.9, \sigma$. Disease stable equilibrium have the same parameter values except for higher host manipulation $\beta_w = \beta_{ww} = 4.5$ and parasite reproduction $f_w = f_{ww} = 45$

204 When a parasite appears in the disease-free equilibrium, it spreads if its reproduction ratio

205 $R_0 > 1$. Since the expression is complicated, we could not obtain analytical solutions for
 206 this inequality without assumptions. We assume the same parasite virulence, $\alpha_w = \alpha_{ww}$,
 207 $\sigma_w = \sigma_{ww}$, and reproduction in double infection as a linear function concerning reproduction
 208 in single infections, $f_{ww} = \epsilon f_w$. When $\epsilon > 1$, reproduction in double infections is greater
 209 than reproduction in a single infection, whereas $\epsilon \leq 1$, reproduction in double infections is
 210 lower or equal to reproduction in a single infection. We found that the parasite can establish
 211 if its reproduction value in a single infection f_w is more significant than a threshold (Figure
 212 5, see SI3SI5).

213 Our numerical results show that the parasite reproduction is substantial compared to other
 214 parameters (its value is nearly 40 times greater than other parameters). This observation
 215 suggests that trophically transmitted parasites must release many offspring into the environ-
 216 ment to persist. Interestingly, bistability occurs if the reproduction rate of the parasite in
 217 double infections is greater than in the single infection state (Figure 5A, B). In the bistable
 218 region, the parasite population can reach a stable equilibrium if the initial density is large
 219 enough. In contrast, with sufficient disturbance, the parasite population could go extinct.

220 The effect of host manipulation on ecological dynamics

221 Host manipulation can be cooperative; two parasites increase the predation rate on interme-
 222 diate hosts, or $\beta_{ww} > \beta_w$. However, it can also be uncooperative; that is, the predation rate
 223 on doubly-infected intermediate hosts lower than that on singly-infected ones, or $\beta_{ww} < \beta_w$.
 224 Cooperation in parasite manipulation does increases the parasite's basic reproduction ratio
 225 (R_0) but the manipulation in single infection has a stronger effect on the value of R_0
 226 (Figure 6). Intuitively, if the manipulation in single infection is small, there is not enough
 227 transmission and the parasite goes extinct. However, suppose the ability to manipulate the
 228 host in a single infection is ~~not strong enough~~ just enough for the parasite population to
 229 escape extinction but not strong. In that case, cooperation in host manipulation leads to a
 230 bistable state of the system. Within the bistable region, the basic reproduction ratio ~~is~~ can
 231 be less than one, suggesting that the parasite cannot spread when its manipulative values
 232 are within this area of weak manipulation when coinfecting. ~~Parasites that can persist in~~
 233 ~~the population may have weak manipulative activity in a single infection but become much~~
 234 ~~more manipulative in coinfection. Likewise, parasites can persist if uncooperative but can~~
 235 ~~manipulate the intermediate hosts effectively when alone.~~

236 Cooperation between parasites need not be limited to host manipulation. Parasites can
 237 cooperate to have a higher reproduction rate in co-infections, i.e. $f_{ww} > f_w$. Likewise,
 238 they can compete for resources, so reproduction in double infection is smaller than in
 239 single infection. Without any assumption on the relationship between manipulative abil-




Figure 5: Effect of parasite reproduction on the ecological dynamics. A) Increasing reproduction in doubly infection leads to bistability, B, C) When Density of singly infected host at equilibrium when reproduction of parasites are the same in singly and doubly infected hosts $f_{ww} = f_w$, C, D) When and when reproduction of parasites in singly doubly infected hosts is four times greater than those in doubly singly infected hosts $f_{ww} = 4f_w$. Filled circles indicate stable equilibrium and open circles indicate unstable equilibrium. Parameter $\rho = 1.2, d = 0.9, r = 2.5, \gamma = 2.9, \alpha_w = 0, \alpha_{ww} = 0, \beta_w = 1.5, \beta_{ww} = 1.5, p = 0.1, c = 1.4, \mu$
 $\rho = 1.2, d = 0.9, r = 2.5, \gamma = 2.9, \alpha_w = 0, \alpha_{ww} = 0, \beta_w = 1.5, \beta_{ww} = 1.5, p = 0.05, c = 1.4,$

ity and reproduction, we explore all possible combinations of cooperation and sabotage in both manipulation and reproduction. Interestingly, higher cooperation in manipulation and reproduction enlarges the area of bistability even though it also shrinks the extinction space




Figure 6: R_0 values increase with more efficient manipulation in both single and double infection. The parasite goes extinct if its manipulative ability is insufficient (dotted-white area). Bistability region occurs when cooperation in manipulation is intermediate (hatched area). As manipulation in single infection increases, the system only has one stable equilibrium. On the black line, manipulation is indifference between single infection and double infection ($\beta_w = \beta_{ww}$). $R_0 < 1$ in Other parameters are the hatched area indicates that the parasite cannot establish same as in a disease-free prey-predator population Figure 6. $f_w = 30$

If parasites are uncooperative in both reproduction and manipulation, they can not persist (Figure 7). In contrast, if they are highly cooperative in both aspects (i.e. $f_{ww}/f_w \rightarrow \infty$ and $\beta_{ww}/\beta_w \rightarrow \infty$), there is one guaranteed single equilibrium for parasite existence.

For intermediate cooperation of both reproduction and manipulation, the bistable area could occur. However, the size of this area is highly sensitive to the value of reproduction and manipulation in single infection. In particular, higher values of these two parameters reduce the bistability area, whereas larger values increase the bistability area (Figure 7, Figure SI1). If the parasites sabotage in host manipulation, the system is highly prone to bistability, and only has a single equilibrium when cooperation in reproduction is extremely high. Interestingly, sufficiently high cooperation in reproduction leads to bistability (i.e. f_{ww} is at least 4 times f_w), and noncooperation in reproduction always lead to single equilibrium of the system (Figure 7). While a single equilibrium guarantees existence of parasite population, bistability indicates that a disturbance of the system may likely lead to extinction of parasite. This suggests that whether or not cooperation benefits the population is context dependent. Clearly, if there is no limitation to cooperation, and if reproduction or manipulation in single infection is large enough, cooperation always hold an advantage. On the other hand, cooperation in one aspect or insufficient cooperation in both aspects may make the parasite population more vulnerable to extinction than systems with less cooperative parasites. This suggests that systems in which parasites have much higher manipulative ability and reproduction rate when co-infected than when singly infected are more prone to instability than systems with less cooperative parasites or systems with parasites that sabotage each other in co-infection. In other words, having the best of both worlds, effective manipulation and reproduction, at the individual level may not benefit the population as a whole.

Increasing the co-transmission probability p from the parasite pool to intermediate hosts reduces the extinction area. When p is high, doubly infected intermediate hosts are more abundant. Cooperation in host manipulation then need not be too high to bring the population out of the bi-stability state. However, it also means that the singly infected intermediate hosts are few and parasites in a single infection must make more manipulative effort to successfully transmit (Figure 7B). On the other hand, increasing the has opposite effect on the bistable area compared to co-transmission probability q from intermediate hosts to definitive hosts broadens the extinction area. When from intermediate hosts to intermediate hosts (Figure 8). In particular, when parasite sabotage the manipulation, increasing p enlarge the bistable area whereas increasing q is high, successful transmission to definitive hosts relies on the predation of susceptible definitive hosts on doubly infected intermediate hosts. Cooperation in manipulation, therefore, needs




Figure 7: Changes of the bistability area (shaded areas) with respect to different reproduction rates in single ~~and double~~ infection (different boundary styles), ~~and varying cotransmission probability~~. Manipulation ~~and reproduction~~ is indifference between single infection and double infection on the ~~black line~~ vertical and horizontal lines respectively. Common parameter: $p = 1.2$, $d = 0.9$, $r = 2.5$, $\gamma = 2.9$, $\alpha_w = 0$, $\alpha_{ww} = 0$, $p = 0.1$, $c = 1.4$, $\mu = 3.9$, $\sigma_w = 0$, $\sigma_{ww} = 0$ for the thick boundary $\epsilon = 0.5$, $f_w = 36$, the dashed boundary $\epsilon = 1$, $f_w = 36$, and the dot-dashed boundary $\epsilon = 2$, $f_w = 35$.

279 to be sufficiently high to avoid bi-stability. Sequential transmission is also rarer because
 280 the probability of a single infection $1 - q$ is low. Suppose the number of doubly infected
 281 intermediate hosts is low. In that case, general transmission from intermediate hosts to
 282 definitive hosts is limited, which explains the wide extinction area reduce it. In contrast, when
 283 parasites cooperate in manipulation, reducing p decreases the bistable area while reducing
 284 q widen it. Evidently, if cooperation in manipulation is extremely high then regardless of
 285 the co-transmission value, the population will always exist with one single stable equilibrium.
 286 However, as there are always limitation and trade-off in nature, extremely high values may
 287 not possible. Considering bistability indicates vulnerability to disturbance, this suggests that
 288 cooperation in manipulation may be beneficial when the co-transmission from the pool to
 289 intermediate host increases. However, cooperation in manipulation may be harmful to the

290 population when the co-transmission from intermediate host to definitive host increases.




Figure 8: Common parameter: $\rho = 1.2$, $d = 0.9$, $r = 2.5$, $\gamma = 2.9$, $\alpha_w = 0$, $\alpha_{ww} = 0$, $p = 0.05$, $c = 1.4$, μ

291 Discussion & Conclusion

292 Host manipulation is a ubiquitous phenomenon suggested to affect the prey-predator dynam-
 293 ics in trophically transmitted parasites. In particular, manipulation of infected intermediate
 294 hosts to increase the predation rate of definitive hosts may result in a heavy burden of preda-
 295 tors on the intermediate host population. This pressure can make parasites more vulnerable
 296 to extinction (Haderler and Freedman, 1989; Fenton and Rands, 2006).

297 Our model shows that parasites cannot spread quickly in a cyclic predator-prey system.
 298 This delay is an expected result since even though the parasite's basic reproduction ratio R_0
 299 is larger than one, it is estimated at the predator and prey's unstable equilibrium (or cyclic
 300 equilibrium). Thus when the density of the prey and predator is at the minimum value of the
 301 cycle, the "effective" R_0 of the parasite can be smaller than one. Another interesting result
 302 is that the reproduction value is much larger than other parameter values. This result is
 303 likely due to the introduction of a free-living parasitic pool. Our model shows that in making
 304 the system more realistic, we also obtain a more realistic quantitative value for parasitic
 305 reproduction.

306 In the study by Rogawa et al. (2018), a non-manipulative parasite can invade a susceptible
 307 prey-predator population and causes the system to cycle. The system is stabilised when the
 308 parasite becomes manipulative, and the stability increases with the manipulative ability. In
 309 our model, non-manipulative parasites cannot persist in the system. ~~The parasite does~~
 310 ~~not necessarily destabilise the predator-prey system, which may contradict the result of~~
 311 Rogawa et al. (2018), and the parasite never leads the system into cyclic dynamics. These

312 results may contradict with Rogawa et al. (2018), where non-manipulative parasites lead to
313 a stable system, while increasing manipulation induces the cyclic behaviour of the population.
314 We suggest that the different results may be due to our introduction of a parasite pool and
315 multiple infections, unlike the model of Rogawa et al. (2018). In their system, transmission
316 from the definitive host to the intermediate host was assumed to result from direct contact
317 between the two hosts. Such immediate transmission could directly accelerate the feedback
318 loop between prey and predator. Hence, faster predator-prey dynamics occur, which may
319 lead to cyclic dynamics when parasites are introduced.

320 ~~However, in~~ In our study, ~~host manipulation can destabilise the predator-prey system under~~
321 ~~particular circumstances and in a different way than the models of Rogawa et al. (2018).~~ In
322 ~~particular, the destabilisation of the system is possible~~ population dynamics exhibit bistability
323 under certain circumstances. This is very likely due to the ~~occurrence of bistability when~~
324 ~~parasite reproduction in coinfection is boosted~~ introduction of co-transmission, which has
325 been shown to result in bistable population dynamics in plant virus Allen et al. (2019) and
326 infectious disease Gao et al. (2016). In this bistability region, if the system is disturbed (e.g.
327 migration of the intermediate or definitive hosts or predation of intermediate hosts by other
328 predators), then the density of the infected hosts may crash, leading to parasite extinction.
329 In other word, it is a way of destabilising the predator-prey system that is different from the
330 result of Rogawa et al. (2018) (where destabilising means cyclic behaviour). In particular,
331 the destabilisation of the system is possible due to the occurrence of bistability when parasite
332 reproduction in coinfection is boosted. The bistability region widens as ~~the manipulation~~
333 ~~in double infection increases, and manipulation in a single infection is insufficient~~ parasites
334 cooperate in reproduction but sabotage in manipulation. This extension is because the
335 density of the doubly infected hosts is always much smaller than the singly infected host
336 density, limited by sequential transmission and a small probability of co-transmission. Sup-
337 pose manipulation in a single infection is not sufficient. In that case, the transmission of
338 the parasites depends mainly on the double infection hosts, which is rare. So extinction is
339 possible if manipulation in double infection is not sufficiently high.

340 Iritani and Sato (2018) show that manipulative parasites can persist if they can alternate
341 manipulation between enhancing and suppressing predation rate. In our model, the parasite
342 cannot switch its manipulative strategy. However, we show that sabotage in manipulation
343 when parasites are coinfecting ~~almost~~ always leads to a single stable equilibrium scenario when
344 there is no cooperation in reproduction. This result suggests that manipulation suppression,
345 either by alternating manipulative strategy or sabotaging, can be crucial in maintaining the
346 parasite population.

347 Finally, our study focuses on the ecological dynamics of the trophically transmitted para-

348 site. However, investigating the evolution of host manipulation is a natural extension beyond
349 the scope of a single manuscript, given the complexities that arise in the ecological dynamics
350 itself~~(we tried). The~~. Studying evolution of host manipulation taking into account free-living
351 parasite pool calls for thorough analyses, which itself could be a standalone study. In addition,
352 the occurrence of bistability in our model suggests that the evolution of host manipulation
353 may drive the parasite population to extinction simply because of the scarcity of the mutant
354 and the Allee effect in the population dynamics. ~~Moreover, the~~ The parasite can enhance
355 both values if there is no tradeoff between manipulation and reproduction. Nevertheless,
356 our model shows that this strategy, which seems to make the best of both worlds, can make
357 the system even more unstable. Evolutionary dynamics here depend mainly on the tradeoff
358 between host manipulation and other traits of the parasites, such as reproduction, virulence,
359 and survivorship in the parasite pool, to list a few. This extension deserves thorough analysis,
360 and we will treat it as a separate matter.

361 ~~Acknowledgments~~Acknowledgements

362 Removed for review

363 **Statement of Authorship**

364 Removed for review

365 **Data and Code Availability**

366 All data and simulation codes for generating figures are available on [https://anonymous.](https://anonymous.4open.science/r/multipleinfections)
367 [4open.science/r/multipleinfections](https://anonymous.4open.science/r/multipleinfections)

368 **References**

- 369 Alizon, S., 2012. Parasite co-transmission and the evolutionary epidemiology of virulence.
370 Evolution 67:921–933. URL <https://doi.org/10.1111/j.1558-5646.2012.01827.x>.
- 371 Alizon, S. and M. van Baalen, 2008. Multiple infections, immune dynamics, and the evolution
372 of virulence. The American Naturalist 172:E150–E168. URL [https://doi.org/10.](https://doi.org/10.1086/590958)
373 [1086/590958](https://doi.org/10.1086/590958).

374 Alizon, S., J. C. de Roode, and Y. Michalakis, 2013. Multiple infections and the evolution of
375 virulence. *Ecology Letters* 16:556–567. URL <https://doi.org/10.1111/ele.12076>.

376 Allen, L. J. S., V. A. Bokil, N. J. Cuniffe, F. M. Hamelin, F. M. Hilker, and M. J. Jeger, 2019.
377 Modelling Vector Transmission and Epidemiology of Co-Infecting Plant Viruses. *Viruses*
378 11:1153. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6950130/>.

379 van Baalen, M. and M. W. Sabelis, 1995. The dynamics of multiple infection and the
380 evolution of virulence. *The American Naturalist* 146:881–910. URL [https://doi.org/](https://doi.org/10.1086/285830)
381 [10.1086/285830](https://doi.org/10.1086/285830).

382 Benesh, D. P., 2016. Autonomy and integration in complex parasite life cycles. *Parasitology*
383 143:1824 – 1846.

384 Choisy, M. and J. C. de Roode, 2010. Mixed infections and the evolution of virulence: Effects
385 of resource competition, parasite plasticity, and impaired host immunity. *The American*
386 *Naturalist* 175:E105–E118. URL <https://doi.org/10.1086/651587>.

387 Diekmann, O., J. Heesterbeek, and J. Metz, 1990. On the definition and the computation
388 of the basic reproduction ratio r_0 in models for infectious diseases in heterogeneous
389 populations. *Journal of Mathematical Biology* 28. URL [https://doi.org/10.1007/](https://doi.org/10.1007/bf00178324)
390 [bf00178324](https://doi.org/10.1007/bf00178324).

391 Diekmann, O., J. A. P. Heesterbeek, and M. G. Roberts, 2009. The construction of next-
392 generation matrices for compartmental epidemic models. *Journal of The Royal Society*
393 *Interface* 7:873–885. URL <https://doi.org/10.1098/rsif.2009.0386>.

394 Fenton, A. and S. A. Rands, 2006. The impact of parasite manipulation and predator
395 foraging behavior on predator - prey communities. *Ecology* 87:2832–2841. URL [https:](https://doi.org/10.1890/0012-9658(2006)87[2832:tiopma]2.0.co;2)
396 [//doi.org/10.1890/0012-9658\(2006\)87\[2832:tiopma\]2.0.co;2](https://doi.org/10.1890/0012-9658(2006)87[2832:tiopma]2.0.co;2).

397 Gandon, S., 2018. Evolution and manipulation of vector host choice. *The American Naturalist*
398 192:23–34. URL <https://doi.org/10.1086/697575>.

399 Gao, D., T. C. Porco, and S. Ruan, 2016. Coinfection dynamics of two diseases in a single
400 host population. *Journal of Mathematical Analysis and Applications* 442:171–188. URL
401 <https://www.sciencedirect.com/science/article/pii/S0022247X16300841>.

402 Hader, K. P. and H. I. Freedman, 1989. Predator-prey populations with parasitic infec-
403 tion. *Journal of Mathematical Biology* 27:609–631. URL [https://doi.org/10.1007/](https://doi.org/10.1007/bf00276947)
404 [bf00276947](https://doi.org/10.1007/bf00276947).

- 405 Hafer, N. and M. Milinski, 2015. When parasites disagree: evidence for parasite-induced
406 sabotage of host manipulation. *Evolution* 69:611 – 620.
- 407 Hosack, G. R., P. A. Rossignol, and P. van den Driessche, 2008. The control of vector-borne
408 disease epidemics. *Journal of Theoretical Biology* 255:16–25. URL [https://doi.org/](https://doi.org/10.1016/j.jtbi.2008.07.033)
409 10.1016/j.jtbi.2008.07.033.
- 410 Hughes, D. P., J. Brodeur, and F. Thomas, 2012. *Host Manipulation by Parasites*. Oxford
411 University Press, London, England.
- 412 Hurford, A., D. Cownden, and T. Day, 2010. Next-generation tools for evolutionary invasion
413 analyses. *Journal of The Royal Society Interface* 7:561–571.
- 414 Iritani, R. and T. Sato, 2018. Host-manipulation by trophically transmitted parasites: The
415 switcher-paradigm. *Trends in Parasitology* 34:934–944. URL [https://doi.org/10.](https://doi.org/10.1016/j.pt.2018.08.005)
416 1016/j.pt.2018.08.005.
- 417 Kalbe, M., K. M. Wegner, and T. B. H. Reusch, 2002. Dispersion patterns of parasites in
418 0+ year three-spined sticklebacks: a cross population comparison. *Journal of Fish Biology*
419 60:1529–1542.
- 420 Lotka, A. J., 1920. Analytical note on certain rhythmic relations in organic systems. *Pro-*
421 *ceedings of the National Academy of Sciences* 6:410–415. URL [https://doi.org/10.](https://doi.org/10.1073/pnas.6.7.410)
422 1073/pnas.6.7.410.
- 423 Molyneux, D. H. and D. Jefferies, 1986. Feeding behaviour of pathogen-infected vectors.
424 *Parasitology* 92:721–736.
- 425 Parker, G. A., J. C. Chubb, G. N. Roberts, M. Michaud, and M. Milinski, 2003. Optimal
426 growth strategies of larval helminths in their intermediate hosts. *Journal of Evolutionary*
427 *Biology* 16:47–54. URL <https://doi.org/10.1046/j.1420-9101.2003.00504.x>.
- 428 Ripa, J. and U. Dieckmann, 2013. Mutant invasions and adaptive dynamics in variable
429 environments. *Evolution* 67:1279–1290. URL [https://onlinelibrary.wiley.com/](https://onlinelibrary.wiley.com/doi/abs/10.1111/evo.12046)
430 doi/abs/10.1111/evo.12046.
- 431 Rogawa, A., S. Ogata, and A. Mougi, 2018. Parasite transmission between trophic levels
432 stabilizes predator–prey interaction. *Scientific Reports* 8. URL [https://doi.org/10.](https://doi.org/10.1038/s41598-018-30818-7)
433 1038/s41598-018-30818-7.

- 434 Rogers, M. E. and P. A. Bates, 2007. Leishmania manipulation of sand fly feeding behavior
435 results in enhanced transmission. PLoS Pathogens 3:e91. URL <https://doi.org/10.1371/journal.ppat.0030091>.
- 437 Roosien, B. K., R. Gomulkiewicz, L. L. Ingwell, N. A. Bosque-Pérez, D. Rajabaskar, and
438 S. D. Eigenbrode, 2013. Conditional vector preference aids the spread of plant pathogens:
439 Results from a model. Environmental Entomology 42:1299–1308. URL <https://doi.org/10.1603/en13062>.
- 441 Seppälä, O. and J. Jokela, 2008. Host manipulation as a parasite transmission strategy
442 when manipulation is exploited by non-host predators. Biology Letters 4:663–666. URL
443 <https://doi.org/10.1098/rsbl.2008.0335>.
- 444 Vickery, W. L. and R. Poulin, 2009. The evolution of host manipulation by parasites: a
445 game theory analysis. Evolutionary Ecology 24:773–788. URL <https://doi.org/10.1007/s10682-009-9334-0>.
- 447 Wedekind, C. and M. Milinski, 1996. Do three-spined sticklebacks avoid consuming cope-
448 pods, the first intermediate host of *Schistocephalus solidus*? - an experimental analysis
449 of behavioural resistance. Parasitology 112:371–383. URL <https://doi.org/10.1017/s0031182000066609>.
- 451 Zimmer, C., 2001. Parasite Rex: Inside the Bizarre World of Nature's Most Dangerous
452 Creatures. Atria Books.

453 Tables

454 ~~Figure legends~~

Table 1: Description of variables and parameters

Parameters and Variables	Description
I_i	Density of intermediate hosts that are susceptible $i = s$, singly infected $i = w$, or doubly infected $i = ww$
D_i	Density of definitive hosts that are susceptible $i = s$, singly infected $i = w$, or doubly infected $i = ww$
W	Density of parasites released from definitive hosts into the environment
d	Natural death rate of intermediate hosts
α_i	Additional death rate of intermediate hosts due to infection by a single parasite ($i = w$) or two parasites ($i = ww$)
p	Probability that two parasites cotransmit from the environment to an intermediate host
γ	Transmission rate of parasites in the environment to intermediate hosts
μ	Natural death rate of definitive hosts
σ_i	Additional death rate of definitive hosts due to infection by a single parasite ($i = w$) or two parasites ($i = ww$)
σ_i	Additional death rate of the hosts due to being infected by a singly parasite ($i = w$) or two parasites ($i = ww$)
q	Probability that two parasites cotransmit from intermediate hosts to definitive hosts
β_i	Transmission rate of parasites from intermediate hosts to definitive hosts
f_i	Reproduction rate of parasites in singly infected definitive hosts ($i = w$) or doubly infected hosts ($i = ww$)
δ	Natural death rate of parasites in the environment
\underline{h}	<u>Probability that the parasites successfully established inside the definitive host</u>