



Example 1: Table below gives the activities of a project, their predecessor(s), durations and costs.

- Draw the network for the project.
- Determine the critical path.
- Construct the Time Schedule.
- Obtain programs with minimum costs.

Activities	Predecessor(s)	Durations		Costs	
		Normal	Crashed	Normal	Crashed
A		4	2	100	160
B		5	5	200	200
C		7	4	250	310
D	A	7	5	180	210
E	B, C	6	6	120	120
F	C	8	4	150	250
G	F	9	6	200	320
H	E, F	4	3	160	220
I	D, E	6	5	240	300
J	F, G	10	5	150	350
K	H, I	5	5	250	250

UCC
30
20
15
25
40
60
60
40
—

a)




Program Summary
Critical Path: C-F-J
Total Duration: 25
Total Cost: 2000

- The crashing amount is determined by two factors: (1) maximal crashing amount of selected critical activities (2) Free floats of alternative noncritical paths.

Crashing amount of C = $\min \{7-4, 0+4, 0+4\} = 3$
FF values of AD, BE

Alternative noncritical paths


- Start from the same or backward event with the selected critical activities
- Lead to the critical path(s).



Program Summary
Critical Path: C-F-J
Total Duration: 22
Total Cost: 2000 + 3 * 20 = 2060


UCCs of Critical activities: C-F-J
X 25 40
should be crashed?

Crashing amount of F = $\min \{8-4, 0+1, 0+1, 3+1\} = 1$
FF values of AD, BE, GK, it starts from FD, it should have started earlier?



Program Summary
Critical Paths: CFT, BEJ, ADJ
Total Duration: 21
Total Cost: 2060 + 1 * 25 = 2085


Crashing options:
X FJ
X BJ
X ADJ
Crashing amount = $\min \{10-5, 4, 1, 3\} = 1$
of J, I, HK, GK



Program Summary
Critical Paths: CFJ, ADJ, BEJ
CFHK, BEHK, ADHK
Total Duration: 20
Total Cost: 2085 + 1 * 40 = 2125

Crashing amount of J and H simultaneously = $\min \{9-5, 4-3, 2, 3\} = 1$
J H G, I

Crashing options:
X FJ
X ADJ
X BJ
X FHK
X BEHK
X ADHK
J and H should be crashed simultaneously



Program Summary:
CP: CFJ, ADJ, BEJ, CFHK, BEHK
TD: 19
TC: 2125 + 1 * 40 + 1 * 60 = 2225

Crashing options:
X FJ
X ADJ
X BJ
X FHK
X BEHK
X ADHK

