

Synthetic Pelvic Floor Disorder CT Dataset

Radiologist Review — 4 Representative Cases

Study: PFD Prevalence Study — South India Cohort

Method: Real CT source + computational PFD deformation

HU range: -1000 to +1500 (full diagnostic range)

Reference: Africa PFD Epidemiological Study

Source data: CTPelvic1K (de-identified)

Total dataset planned: 350 patients (175 Plain + 175 Hilly)



This package shows one representative Grade 2 (moderate) case per PFD pattern: combined multi-compartment, cystocele, rectocele, and uterine prolapse. Each case shows the original CT, the deformed CT, and a difference heatmap, followed by axial survey slices and sagittal/coronal MPR reconstructions.


CASE 1 Multi-Compartment Pelvic Floor Disorder — Combined PFD

Grade 2 — Moderate | Plain pelvis

Simultaneous anterior (bladder), posterior (rectal), and apical (uterine) compartment involvement. Grade 2 across all compartments. The most common symptomatic presentation. Assess coordinated multi-compartment descent and levator ani hiatal changes.

ORIGINAL CT / PFD-DEFORMED / DIFFERENCE HEATMAP (3 HIGHEST-DEFORMATION SLICES)



**AXIAL SURVEY — SOFT-TISSUE WINDOW (INFERIOR TO SUPERIOR)****SAGITTAL MPR****CORONAL MPR**





Patient ID: REAL_001_PLAIN_COMBINED_PFD_G2

CASE 2 Anterior Compartment Prolapse — Cystocele Grade 2 — Moderate | Plain pelvis

Bladder base descends through the anterior vaginal wall. Grade 2: descent to the level of the vaginal introitus. Assess bladder base position relative to the pubococcygeal line, urethrovesical junction descent, and anterior vaginal wall support.



ORIGINAL CT / PFD-DEFORMED / DIFFERENCE HEATMAP (3 HIGHEST-DEFORMATION SLICES)**AXIAL SURVEY — SOFT-TISSUE WINDOW (INFERIOR TO SUPERIOR)****SAGITTAL MPR****CORONAL MPR**




Patient ID: REAL_003_PLAIN_CYSTOCELE_G2

CASE 3 Posterior Compartment Prolapse — Rectocele Grade 2 — Moderate | Plain pelvis

Anterior herniation of the rectal ampulla through the posterior vaginal wall. Grade 2: moderate anterior bulging. Assess anterior rectal wall contour, rectovaginal septal integrity, and extent of herniation.

ORIGINAL CT / PFD-DEFORMED / DIFFERENCE HEATMAP (3 HIGHEST-DEFORMATION SLICES)**AXIAL SURVEY — SOFT-TISSUE WINDOW (INFERIOR TO SUPERIOR)****SAGITTAL MPR****CORONAL MPR**







Patient ID: REAL_005_PLAIN_RECTOCELE_G2

CASE 4 Apical Compartment Prolapse — Uterine Prolapse

Grade 2 — Moderate | Plain pelvis

Inferior displacement of the uterine cervix toward or through the vaginal introitus. Grade 2: cervix at the level of the hymen (POP-Q C point). Assess uterine axis, cervical position relative to ischial spines, and associated cul-de-sac changes.

ORIGINAL CT / PFD-DEFORMED / DIFFERENCE HEATMAP (3 HIGHEST-DEFORMATION SLICES)**AXIAL SURVEY — SOFT-TISSUE WINDOW (INFERIOR TO SUPERIOR)**

SAGITTAL MPR**CORONAL MPR**

Patient ID: REAL_022_PLAIN_UTERINE_PROLAPSE_G2
