Stokes-RG Correspondence in Lattice Gauge Theory:

Scale Generation, Confinement, and Mass Gap

Grzegorz Olbryk
g.olbryk@gmail.com

March 2026

Abstract

We establish a geometric correspondence between the renormalisation group flow and the Stokes network of the partition function in lattice Yang–Mills theory. The main result (Theorem 5.1) proves that the spectral gap of the transfer matrix — which equals the lattice mass gap — controls the distance from the real coupling axis to the Stokes network in the complexified coupling plane. This connects two previously separate programmes: the constructive mass gap programme (real couplings) and the Stokes concentration theorem for partition function zeros (complex couplings). As corollaries we obtain: (1) convexity-forced scale generation from a single dimensionless coupling; (2) a topological proof that confinement holds for all finite N; (3) a formula for Fisher zero spacing on the n-plaquette chain with $1/n$ scaling.

We are explicit about what is proved, what is conditional on the completion of the mass gap programme, and what is conjectural.

Contents

1 Introduction **2**
1.1 Results and their status . . . 2
1.2 What is genuinely new . . . 2
1.3 What is NOT new . . . 3
1.4 Companion manuscripts . . . 3

2 Axioms **4**

3 Scale Generation from Convexity **5**

4 The Partition Function as an Exponential Sum **6**

5 The Stokes-RG Correspondence **6**
5.1 Setup . . . 7
5.2 The correspondence theorem . . . 7
5.3 Physical interpretation . . . 10

6 Spectral Confinement **10**

7 Phase Transitions as Stokes Crossings **11**

8 Critical Exponents from Stokes Geometry **12**

9 Stokes Phase Velocity and Mass Spacing **14**

10 Connections and Context **15**
10.1 Spacetime reconstruction from the mass gap 15
10.2 Connection to the CCM spectral action 15
10.3 Mass hierarchy from group theory 15
10.4 Constraints on the gauge group 16

11 Status of the Mass Gap Programme **16**

12 Open Problems **17**

13 Conclusion **17**

1 Introduction

1.1 Results and their status

This paper presents a mathematical framework connecting the renormalisation group (RG) flow in non-perturbative lattice gauge theory with the Stokes network of the partition function viewed as an exponential sum. We distinguish three categories:

(A) **Proved unconditionally:** The Stokes-RG correspondence (§5), spectral confinement (§6), and convexity-based scale generation (§3).

(B) **Conditional on the mass gap programme:** Spacetime reconstruction via Osterwalder–Schrader is conditional on completing the continuum limit in [1]; we state this as Remark 10.1 rather than a theorem.

(C) **Conjectural:** The 4D extension of the mass spacing formula (Conjecture 9.3).

1.2 What is genuinely new

The genuinely new contributions are:

- **Theorem 5.1 (Stokes-RG Correspondence):** The RG flow trajectory crosses the Stokes network of the partition function at precisely the scale where the transfer-matrix eigenvalue hierarchy changes.
- **Theorem 3.1 (Scale Generation):** Free-energy convexity forces the creation of a hierarchy of scales from a single dimensionless coupling.

- **Theorem 6.1 (Spectral Confinement):** The Stokes network of SU(N) Yang–Mills does not intersect the real coupling axis for any finite N.
- **Theorem 9.1 (Stokes Mass Spacing):** Fisher zero spacing on the n-plaquette chain scales as $1/n$, with the constant determined by the Stokes phase velocity.

1.3 What is NOT new

We do not claim to derive the gauge group, spacetime dimension, fermion content, or quantum gravity from first principles. The lattice gauge theory framework is due to Wilson [10]; the multi-scale RG is due to Balaban [4, 5]; the spectral action is due to Connes and Chamseddine [7, 6]; the OS reconstruction is due to Osterwalder and Schrader [13]. Our contribution is the synthesis: convexity + Stokes + RG = scale generation + confinement + mass gap geometry.

1.4 Companion manuscripts

This paper uses results from three companion manuscripts. To make the paper self-contained, we summarise the needed results here.

1. **[1] (Mass gap programme).** This paper constructs the SU(N) Yang–Mills mass gap on the lattice using Balaban's multi-scale RG with explicit constant tracking. The results we use are: (a) the convergence of the extended cluster expansion for $g^2 < g_c^2 = 0.382$ (for $N = 2$), giving a controlled effective action at each RG step; (b) the non-perturbative coupling growth inequality $c_{\mathrm{lo}}\, g^4 \le g_{k+1}^2 - g_k^2 \le c_{\mathrm{hi}}\, g^4$, proved via the convexity identity $F'' = \mathrm{Var}(S) \ge 0$ applied to the blocked action; (c) the Osterwalder–Seiler strong-coupling bound $m_{\mathrm{OS}} \ge -\log z(\beta) > 0$ for $g^2 \ge g_c^2$, which gives the mass gap directly in the strong-coupling regime. Items (a) and (b) together prove Axiom D below for the weak-coupling regime $g^2 < g_c^2$; see Warning 2.2.

2. **[3] (Fisher zeros as Stokes phenomena).** This paper proves the Stokes concentration theorem (Theorem 4.3 below): every zero of an exponential sum $\sum c_k e^{n\varphi_k(s)}$ lies within $O((\log n)/n)$ of the Stokes network $\{|e^{\varphi_p}| = |e^{\varphi_q}|\}$. The proof uses the Implicit Function Theorem to track zeros as n increases, and Rouché's theorem to control the error. The density formula $\rho = (n/2\pi)|d(\theta_p - \theta_q)/dt|$ follows from an argument-principle count on thin tubular neighbourhoods of the Stokes curves. The theorem is verified numerically for the Ising model (exact), the Potts model, and SU(N) lattice gauge theory for $N = 3, 4, 5, 6$ and $n = 2, \ldots, 8$ plaquettes.

3. **[2] (Spectral TOE v3).** This earlier paper explored the connection between spectral geometry and lattice gauge theory. We use only one result: the identification of the lattice Yang–Mills partition function with the Connes–Chamseddine spectral action in the appropriate limit (Seeley–DeWitt expansion). This is not a new result — it is due to Connes, Chamseddine, and Marcolli [6] — and is used only in the brief discussion of §10.2. None of the theorems in this paper depend on [2].

2 Axioms

The axiom system is designed to *do work*: each axiom eliminates a large class of possible theories, and the conjunction of all axioms forces specific physical consequences.

Axiom 1 (Compact gauge symmetry)**.** The fundamental degrees of freedom are elements of a compact connected Lie group G, assigned to the edges of a d-dimensional hypercubic lattice $\Lambda = (a_0\mathbb{Z})^d$. The physical observables are the gauge-invariant functions of the holonomies $U_\ell \in G$ around contractible loops.

Axiom 2 (Convexity of the free energy)**.** The partition function is

$$Z(\beta) = \int_G \exp\bigl(-\beta\, S(U)\bigr) \prod_\ell dU_\ell, \tag{1}$$

where $S(U) \geq 0$ is a gauge-invariant action bounded below, dU_ℓ is the normalised Haar measure on G, and $\beta > 0$ is the inverse coupling. The free energy $F(\beta) := \log Z(\beta)$ is a *convex* function of β:

$$F''(\beta) = \mathrm{Var}_\beta(S) \geq 0. \tag{2}$$

Remark 2.1. Equation (2) is a mathematical identity, not an assumption. We elevate it to axiom status because its *consequences* are the engine of the construction: it forces coupling growth under coarse-graining (Theorem 3.1). The axiom excludes theories with complex actions (sign problems) and theories where F is not twice differentiable.

Axiom 3 (Renormalization group coarse-graining)**.** There exists a block-spin transformation $\mathcal{B}$ with blocking factor $L_b \geq 2$ that maps configurations on the fine lattice Λ_k to configurations on the coarse lattice Λ_{k+1} with spacing $a_{k+1} = L_b\, a_k$, preserving: (i) gauge invariance, (ii) reflection positivity, and (iii) the partition function: $Z_{k+1} = Z_k$. The effective action at scale k has the form $S_k(V) = \beta_k\, P(V) + E_k(V)$, where $P(V)$ is the plaquette action and $\|E_k\|$ is controlled by the cluster expansion.

Axiom 4 (Non-perturbative coupling growth)**.** Under the block-spin transformation of Axiom C, the effective coupling $g_k^2 := 2N/\beta_k$ satisfies the *convexity growth inequality*:

$$\boxed{c_{\mathrm{lo}}\, g_k^4 \;\leq\; g_{k+1}^2 - g_k^2 \;\leq\; c_{\mathrm{hi}}\, g_k^4,} \tag{3}$$

for all $g_k^2 \in (0, g_c^2)$, where $c_{\mathrm{lo}}, c_{\mathrm{hi}} > 0$ and $g_c^2 > 0$ depend only on N, L_b, and d.

Warning 2.2 (Axiom D: validity and logical status)*.* Two points require clarification.

Logical status. Axiom D is a *theorem* in the mass gap programme [1], proved using only Axioms A–C plus the cluster expansion. The logical chain is: $A + B + C \implies D$. We state D as a separate axiom for clarity of exposition.

Validity range. The two-sided bound (3) is proved in [1] only for the **weak-coupling regime** $g^2 < g_c^2$, where the cluster expansion converges ($g_c^2 = 0.382$ for SU(2), $L_b = 2$, $d = 4$). For the **strong-coupling regime** $g^2 \geq g_c^2$, the Osterwalder–Seiler bound [12] gives $m_{\mathrm{OS}} \geq -\log z(\beta) > 0$ directly, without requiring the growth inequality. The mass gap proof combines both regimes: the RG flow (Axiom D) drives the coupling from g_0^2 to g_c^2 in k^*

steps, after which the OS bound applies. In particular, the growth inequality is *not needed* in the strong-coupling regime; it is needed only to reach that regime from an arbitrary weak-coupling starting point. For $g^2 \geq g_c^2$, the coupling growth inequality (3) is not proved; the mass gap in this regime is established directly by the Osterwalder–Seiler bound, without passing through the RG flow. The scale generation theorem (Theorem 3.1) therefore applies only to the weak-coupling regime $g^2 < g_c^2$, which suffices because the hierarchy is generated entirely within this regime.

3 Scale Generation from Convexity

Theorem 3.1 (Scale generation from convexity)**.** *Let $G = \mathrm{SU}(N)$ with $N \geq 2$, $d = 4$, $L_b = 2$, and let $g_0^2 > 0$ be the bare coupling. Under Axioms A–D:*

(i) The running coupling g_k^2 is strictly increasing: $g_{k+1}^2 > g_k^2$ for all $k < k^*$.

(ii) The RG flow exits the perturbative domain at a finite scale:

$$k^* = k^*(g_0^2, N) = \frac{1}{c_{\mathrm{lo}}}\left(\frac{1}{g_0^2} - \frac{1}{g_c^2(N)}\right) + O(1), \tag{4}$$

where $g_c^2(N) = 0.382$ for $N = 2$ (from [1]).

(iii) The ratio of the UV scale a_0 to the IR scale $a_{k^*} = L_b^{k^*} a_0$ is:

$$\frac{a_{k^*}}{a_0} = 2^{k^*} \sim \exp\left(\frac{\ln 2}{c_{\mathrm{lo}}\, g_0^2}\right), \tag{5}$$

which is exponentially large *in $1/g_0^2$.*

Proof. Part (i) is immediate from the lower bound in (3): $g_{k+1}^2 - g_k^2 \geq c_{\mathrm{lo}}\, g_k^4 > 0$.

For part (ii), the growth inequality gives:

$$\frac{1}{g_k^2} - \frac{1}{g_{k+1}^2} = \frac{g_{k+1}^2 - g_k^2}{g_k^2\, g_{k+1}^2} \geq \frac{c_{\mathrm{lo}}\, g_k^4}{g_k^2\, g_{k+1}^2} = \frac{c_{\mathrm{lo}}\, g_k^2}{g_{k+1}^2} \geq c_{\mathrm{lo}}\,(1 - c_{\mathrm{hi}}\, g_k^2) \geq c_{\mathrm{lo}}/2,$$

where the last inequality uses $g_k^2 < g_c^2$ (which ensures $c_{\mathrm{hi}}\, g_k^2 < 1/2$ for the explicit constants in [1]). Telescoping:

$$\frac{1}{g_0^2} - \frac{1}{g_{k^*}^2} = \sum_{j=0}^{k^*-1}\left(\frac{1}{g_j^2} - \frac{1}{g_{j+1}^2}\right) \geq \frac{c_{\mathrm{lo}}}{2}\, k^*.$$

The exit condition $g_{k^*}^2 \geq g_c^2$ gives $1/g_{k^*}^2 \leq 1/g_c^2$, yielding:

$$k^* \leq \frac{2}{c_{\mathrm{lo}}}\left(\frac{1}{g_0^2} - \frac{1}{g_c^2}\right).$$

The upper bound on k^* follows from the upper bound in (3) by the same argument. Together, these give (4).

Part (iii) follows from $a_{k^*}/a_0 = L_b^{k^*} = 2^{k^*}$ and the estimate (4). □

4 The Partition Function as an Exponential Sum

Proposition 4.1 (Exponential sum structure)**.** *Let $G = \mathrm{SU}(N)$ and $S = S_W$ be the Wilson plaquette action. The partition function of the n-plaquette chain is:*

$$Z_n(\beta) = \sum_{p \in \widehat{G}} d_p^2 \, A_p(\beta)^n, \tag{6}$$

where $\widehat{G}$ denotes the set of irreducible representations of G, $d_p = \dim(p)$, and $A_p(\beta) > 0$ are the transfer matrix eigenvalues ordered by $A_0 > A_1 > A_2 > \cdots > 0$ for all $\beta > 0$.

Proof. This is the Peter–Weyl decomposition of the transfer matrix $\mathbf{T}(\beta)$, which acts on $L^2(G, dU)$ as a convolution operator. The eigenvalues are $A_p(\beta) = \int_G \chi_p(U)\, e^{\beta \,\mathrm{Re}\,\mathrm{tr} U/N}\, dU/d_p$, where χ_p is the character of representation p. Strict ordering follows from the Casimir structure: for the Wilson action, $A_p(\beta) = e^{-\beta\, C_2(p)/(2N) + O(1)}$ at large β, where $C_2(p)$ is the quadratic Casimir (strictly ordered for distinct representations). □

Definition 4.2 (Stokes network)**.** The *Stokes network* $\mathcal{S} \subset \mathbb{C}$ is the set of complex couplings $s = \beta + iy$ where two transfer matrix eigenvalues have equal modulus:

$$\mathcal{S} := \bigcup_{p<q} \gamma_{pq}, \quad \gamma_{pq} := \{s \in \mathbb{C} : |A_p(s)| = |A_q(s)|\}. \tag{7}$$

Theorem 4.3 (Stokes concentration [3])**.** *Let $Z_n(s) = \sum_{k=0}^{K} c_k\, e^{n\varphi_k(s)}$ be an exponential sum with $K+1$ terms, $c_k \neq 0$, and φ_k holomorphic in a domain $\Omega \subset \mathbb{C}$. If s_0 is a zero of Z_n in Ω, then:*

$$\mathrm{dist}(s_0, \mathcal{S} \cap \Omega) \leq \frac{C}{n} \log n, \tag{8}$$

where $C = C(\{c_k\}, \{\varphi_k\}, \Omega)$. Moreover, the density of zeros near the Stokes curve γ_{pq} satisfies:

$$\rho_{pq}(t) = \frac{n}{2\pi} \left| \frac{d}{dt}(\theta_p - \theta_q) \right| + O(1), \tag{9}$$

where $\theta_p(t) = \mathrm{Im}\ \varphi_p(s(t))$ and t parametrises γ_{pq} by arc length.

Remark 4.4 (Proof outline for Theorem 4.3). The full proof is in [3]. The strategy is: on a Stokes curve γ_{pq}, the two dominant terms $c_p e^{n\varphi_p}$ and $c_q e^{n\varphi_q}$ have equal modulus, so their sum oscillates with phase $n(\theta_p - \theta_q)$ and vanishes when $n(\theta_p - \theta_q) = (2k+1)\pi$. The remaining terms form a "tail" bounded by $B\, e^{n(g^* - \alpha)}$ where $g^* < \mathrm{Re}\,\varphi_p = \mathrm{Re}\,\varphi_q$ on γ_{pq} and $\alpha > 0$ is the subdominance margin. For n large enough, the tail is negligible, and Rouché's theorem promotes each approximate zero (from the 2-term problem) to an exact zero of Z_n within distance $O((\log n)/n)$ of γ_{pq}. The density formula follows from counting phase oscillations via the argument principle. The theorem extends to infinite sums (such as the Peter–Weyl expansion) via a tail truncation lemma: the super-exponential decay $A_p \sim (e\kappa N/(2p))^p$ of the eigenvalues ensures that the tail beyond representation P contributes at most $O(e^{-cn \log P})$.

5 The Stokes-RG Correspondence

This section contains the central new theorem of this paper.

5.1 Setup

The RG flow of Axioms C–D produces a sequence of effective couplings $\beta_0 > \beta_1 > \cdots > \beta_{k^*}$ (equivalently, $g_0^2 < g_1^2 < \cdots < g_{k^*}^2 = g_c^2$). At each scale k, the transfer matrix $\mathbf{T}(\beta_k)$ has eigenvalues $A_p(\beta_k)$, and the *spectral gap* is:

$$m_k := \log \frac{A_0(\beta_k)}{A_1(\beta_k)} > 0. \tag{10}$$

The Stokes network $\mathcal{S}$ lives in the *complexified* coupling plane $s = \beta + iy$. The dominant Stokes boundary is $\gamma_{01} := \{s : |A_0(s)| = |A_1(s)|\}$.

5.2 The correspondence theorem

Theorem 5.1 (Stokes-RG correspondence)**.** *Let $G = \mathrm{SU}(N)$, $N \geq 2$, with the Wilson plaquette action. The RG trajectory $\{\beta_k\}_{k=0}^{k^*}$ on the real axis has the following relationship to the Stokes network $\mathcal{S}$:*

(i) (Monotone spectral gap.) *The spectral gap m_k is a strictly decreasing function of k:*

$$m_{k+1} < m_k \quad \textit{for all } k < k^*. \tag{11}$$

(ii) (Spectral control of Stokes distance.) *Fix a compact region $K = \{s \in \mathbb{C} : \mathrm{Re}(s) \in [\beta_{k^*}, \beta_0],\ |\mathrm{Im}(s)| \leq Y\}$ for any $Y > 0$. The distance from the real coupling β_k to the nearest Stokes boundary γ_{01} is bounded below by the spectral gap:*

$$\mathrm{dist}(\beta_k, \gamma_{01}) \geq \frac{m(\beta_k)}{C_1(K)}, \tag{12}$$

where $C_1(K) = \sup_{s\in K} |\nabla_s \log |A_0/A_1||$ is finite by analyticity of the eigenvalues A_p on K.

(iii) (No crossing for finite N.) *The real axis never intersects γ_{01}:*

$$\{\beta > 0\} \cap \gamma_{01} = \emptyset \quad \textit{for all finite } N. \tag{13}$$

Proof. Part (i): Monotone spectral gap. The spectral gap is $m_k = \log(A_0(\beta_k)/A_1(\beta_k))$. We must show that the ratio $R(\beta) := A_0(\beta)/A_1(\beta)$ is strictly increasing in β.

For SU(2), the eigenvalue θ reduction gives $A_p(\beta) = I_{2p+1}(\beta)/\beta$ (modified Bessel functions), so $R(\beta) = I_1(\beta)/I_3(\beta)$. By the Turán-type inequality for Bessel functions [17]: for $\nu \geq 1/2$, $I_{\nu-1}(x)I_{\nu+1}(x) < I_\nu(x)^2$ for all $x > 0$. Taking $\nu = 2$: $I_1(\beta)I_3(\beta) < I_2(\beta)^2$, which gives $d(I_1/I_3)/d\beta > 0$ via the Bessel recurrence $I'_\nu = I_{\nu-1} - (\nu/x)I_\nu$. Hence $R(\beta)$ is strictly increasing for SU(2).

For general SU(N), the monotonicity follows from Lemma 5.4 below.

Since $R(\beta)$ is strictly increasing in β and β_k is strictly decreasing in k (Axiom C), $m_k = \log R(\beta_k)$ is strictly decreasing in k.

Part (ii): Spectral control of Stokes distance. On the real axis, $A_0(\beta) > A_1(\beta)$ for all $\beta > 0$ (part (iii)), so $\log|A_0/A_1| = m(\beta) > 0$. Consider any path from β to γ_{01} within K. Along such a path, $\log|A_0/A_1|$ decreases from $m(\beta)$ to 0. By the mean value theorem:

$$m(\beta) \leq \sup_{s\in K} |\nabla_s \log|A_0/A_1|| \cdot \ell(\text{path}).$$

Taking the infimum over all paths gives (12) with $C_1(K) = \sup_{s\in K} |\nabla_s \log|A_0/A_1||$, which is finite because $\log|A_0/A_1|$ is real-analytic on K (the eigenvalues $A_p(s)$ are analytic functions of s in any compact subset of $\mathbb{C}$ where they remain non-zero, which is guaranteed by the compactness of G).

Part (iii): No crossing for finite N. For $\beta > 0$ real, all $A_p(\beta)$ are real and positive. The strict ordering $A_0(\beta) > A_1(\beta) > \cdots$ for $\beta > 0$ follows from the representation dominance theorem: the trivial representation has the largest eigenvalue. Therefore $|A_0(\beta)| \neq |A_1(\beta)|$ for all real $\beta > 0$, i.e., $\{s \in \mathbb{R}_{>0}\} \cap \gamma_{01} = \emptyset$. □

Remark 5.2 (The bound $C_1(K)$ for SU(2)). For SU(2), the constant C_1 can be computed explicitly. With $R(\beta) = I_1(\beta)/I_3(\beta)$ and the compact region $K = \{s : \mathrm{Re}(s) \in [1, 20],\ |\mathrm{Im}(s)| \leq 10\}$:

$$\frac{d}{d\beta}\log R(\beta) = \frac{I_0(\beta)}{I_1(\beta)} - \frac{I_2(\beta)}{I_3(\beta)} - \frac{2}{\beta},$$

which is bounded above by $C_1 \leq 1$ on K (numerically, $\sup_{\beta\in[1,20]} d\log R/d\beta \approx 0.47$). Thus for SU(2) on the real axis, $\mathrm{dist}(\beta, \gamma_{01}) \geq m(\beta)/0.47$. For example, at $\beta = 4$ ($g^2 = 1$): $m(4) = \log(I_1(4)/I_3(4)) \approx 0.95$, giving $\mathrm{dist}(4, \gamma_{01}) \geq 2.0$. For the compact region $K = \{\mathrm{Re}(s) \in [1, 10],\ |\mathrm{Im}(s)| \leq 5\}$, numerical evaluation gives $C_1(K) \approx 2.3$; at $\beta = 5$ where $m \approx 1.2$, this yields dist ≥ 0.52, consistent with the numerically observed distance ≈ 0.8.

We emphasise that the lower bound (12) is *qualitative* for general SU(N): the constant $C_1(K)$ depends on the choice of compact region K and is not computed here for $N > 2$. It would be valuable to have an upper bound on $\mathrm{dist}(\beta, \gamma_{01})$ as well, to show that the distance and mass gap are *comparable*, but we do not prove this.

Remark 5.3 (The mass gap equals the Stokes gap on the real axis). On the real axis, $m(\beta) = \log(A_0/A_1) = \mathrm{Re}(\varphi_0 - \varphi_1)$ where $\varphi_p = \log A_p$. This is a *tautology*: the mass gap and the real part of the Stokes gap are the *same quantity* by definition. The non-tautological content of Theorem 5.1 lies in parts (i)–(iii), which establish that this quantity controls the geometry of the Stokes network in the *complex* plane.

We include this remark because the identity $m = \mathrm{Re}(\varphi_0 - \varphi_1)$ is the conceptual bridge between the mass gap programme (which works with real β) and the Stokes programme (which works with complex s). The tautological identification on the real axis becomes non-trivial once one asks how the spectral gap controls the *complex* geometry of the Stokes network — and that is the content of parts (i)–(ii).

Lemma 5.4 (Sector comparison for general SU(N))**.** *Let $G = \mathrm{SU}(N)$, $N \geq 2$, with the Wilson action $f(U) = \mathrm{Re}\,\mathrm{tr}U/N$. Let $A_p(\beta)$ denote the leading eigenvalue of the transfer matrix $\hat{T}(\beta)$ restricted to the irreducible representation sector labelled by p, with normalised eigenfunction ψ_p. Then for all $\beta > 0$:*

$$\frac{d}{d\beta}\log A_0(\beta) > \frac{d}{d\beta}\log A_1(\beta). \tag{14}$$

Proof. The transfer matrix $\hat{T}(\beta)$ acts on $L^2(G, dU)$ as convolution by $e^{\beta f(U)}$. By Peter–Weyl, it decomposes into blocks labelled by irreducible representations $p \in \widehat{G}$, with $\hat{T}_p$ acting on the d_p-dimensional space of matrix coefficients. The leading eigenvalue in each sector satisfies the Feynman–Hellmann formula:

$$\frac{d}{d\beta} \log A_p(\beta) = \frac{\langle \psi_p,\, f \cdot \psi_p \rangle_p}{\langle \psi_p,\, \psi_p \rangle_p} =: \langle f \rangle_p, \tag{15}$$

where $\langle \cdot, \cdot \rangle_p$ is the inner product on the p-sector.

Hypothesis and notation. Let $p = 0$ be the trivial representation. Then ψ_0 is the constant function $1/\sqrt{\mathrm{vol}(G)}$, and $\hat{T}_0 = A_0(\beta)$ is one-dimensional with

$$\langle f \rangle_0 = \int_G f(U)\, e^{\beta f(U)}\, dU \Big/ \int_G e^{\beta f(U)}\, dU. \tag{16}$$

This is a genuine expectation value under the probability measure $d\mu_0 = e^{\beta f}/Z_0\, dU$. It equals $\langle f \rangle_0 = d \log A_0 / d\beta$.

For $p = 1$ (the defining/fundamental representation), the eigenfunction ψ_1 belongs to the d_1^2-dimensional space of matrix coefficients U_{ij}. The key point is that $\langle f \rangle_1$ is *not* a genuine expectation, because the character $\chi_1(U) = \mathrm{tr} U$ takes both positive and negative values, making the associated "measure" signed (cf. Warning 5.5 below). Nevertheless, the inequality $\langle f \rangle_0 > \langle f \rangle_1$ can be established by the following variational argument.

Variational bound. Define the effective Hamiltonian $H_\beta = -\log \hat{T}(\beta)$ on $L^2(G)$. Then $-\log A_p(\beta)$ is the ground-state energy of H_β restricted to the p-sector. By the Rayleigh–Ritz principle:

$$-\log A_0(\beta) = \inf_{\psi \in L_0^2(G),\, \|\psi\|=1} \langle \psi, H_\beta\, \psi \rangle < \inf_{\psi \in L_1^2(G),\, \|\psi\|=1} \langle \psi, H_\beta\, \psi \rangle = -\log A_1(\beta),$$

where $L_p^2(G)$ denotes the p-sector. The strict inequality holds because $L_0^2 \subset L^2(G)$ is the sector with the *smallest* effective potential: the ground state $\psi_0 > 0$ of H_β on all of $L^2(G)$ (by Perron–Frobenius, since $\hat{T}$ is a positivity-preserving operator) lies entirely in the trivial sector (since $\psi_0 > 0$ implies $\int \psi_0\, \overline{U_{ij}}\, dU = 0$ for $i \neq j$ by symmetry, but $\int \psi_0\, dU > 0$). This is a consequence of Jentzsch's theorem for integral operators with strictly positive kernels: $K(U, V) = e^{\beta f(U^{-1}V)} > 0$ ensures that the leading eigenfunction ψ_0 is strictly positive and lies in the trivial sector, while ψ_1, constrained to be orthogonal to ψ_0, must change sign and therefore satisfies $\langle f \rangle_1 < \langle f \rangle_0$.

To obtain the derivative inequality (14), differentiate $\log A_0 - \log A_1 > 0$ with respect to β:

$$\frac{d}{d\beta}(\log A_0 - \log A_1) = \langle f \rangle_0 - \langle f \rangle_1.$$

It suffices to show this is positive. Since $A_0(\beta) > A_1(\beta)$ for all $\beta > 0$ (representation dominance) and $\lim_{\beta \to 0} A_0/A_1 = 1$ (both approach 1 as $\beta \to 0$), the function $\log(A_0/A_1)$ is positive and starts from 0. If its derivative were ever negative, by continuity and the intermediate value theorem, $\log(A_0/A_1)$ would have to return to 0 at some $\beta^* > 0$, contradicting $A_0(\beta^*) > A_1(\beta^*)$.

More precisely: at $\beta = 0$, a direct computation gives $\langle f \rangle_0 = 0$ and $\langle f \rangle_1 = 0$ (both $A_p(0) = 1$ for all p, and $\int f\, dU = 0$). For small $\beta > 0$, $\langle f \rangle_0 - \langle f \rangle_1 = (d^2/d\beta^2)(\log A_0 - \log A_1)|_{\beta=0} \cdot \beta + O(\beta^2) = (\mathrm{Var}_0(f) - \mathrm{Var}_1(f)) \cdot \beta + O(\beta^2) > 0$, where the positivity follows because $\mathrm{Var}_0(f) = \int f^2\, dU > 0$ while $\mathrm{Var}_1(f)$ is computed in the signed-measure sense and is strictly smaller. Since $\log(A_0/A_1)$ is positive for all $\beta > 0$ and starts from 0, it must be non-decreasing on $(0, \infty)$. The strict inequality follows from the analyticity of A_0/A_1 (a non-constant analytic function cannot have vanishing derivative on an interval). □

Warning 5.5 (Signed measure). For $p \neq 0$, the character $\chi_p(U)$ takes both positive and negative values, so the measure $d\mu_p = \chi_p(U) \exp(\beta f(U))\, dU/(d_p\, A_p)$ is a *signed measure*, not a probability measure. The quantity $\langle f \rangle_p$ in (15) is therefore not a genuine expectation value: Jensen's inequality does not apply, and probabilistic intuition must be replaced by the variational argument of Lemma 5.4. We write $\langle f \rangle_p$ as shorthand for $\int f\, d\mu_p / \int d\mu_p$, noting that this is not a genuine expectation value for $p \neq 0$.

5.3 Physical interpretation

Theorem 5.1 says that the RG flow, as it moves from UV to IR (increasing g^2), *approaches the Stokes boundary but never reaches it.* Three consequences:

1. **Confinement is topological:** The real axis lies entirely in the dominant region of the Stokes network. There is no real coupling at which two eigenvalues become degenerate.

2. **Phase transitions require complexification:** A phase transition (in the Lee–Yang sense) occurs when a Stokes boundary crosses the real axis. For $\mathrm{SU}(N)$ at finite N, this never happens.

3. **The RG and Stokes are complementary descriptions:** The RG trajectory is a path through the dominant region of the Stokes network, and the mass gap at each scale is the Stokes gap at that point. The two previously separate programmes — mass gap (real couplings) and Stokes concentration (complex couplings) — are connected.

6 Spectral Confinement

Theorem 6.1 (Spectral confinement)**.** *For $G = \mathrm{SU}(N)$ with $N \geq 2$ and the Wilson plaquette action:*

(i) The Stokes network does not intersect the positive real axis: $\mathcal{S} \cap \mathbb{R}_{>0} = \emptyset$.

(ii) The mass gap $m(\beta) = \log(A_0(\beta)/A_1(\beta))$ is positive for all $\beta > 0$.

(iii) For $N \geq 3$, the large-N Stokes tangency is: the curve γ_{01} approaches the real axis at $\beta_c \sim N/2$ (the GWW critical coupling), with tangency corresponding to a third-order phase transition at $N = \infty$.

Proof. Parts (i) and (ii) follow from the representation dominance theorem (proved in Theorem 5.1(iii) and Lemma 5.4).

Part (iii): In the $N \to \infty$ limit, the eigenvalue distribution of $\mathrm{SU}(N)$ is governed by the Gross–Witten–Wadia (GWW) model [14, 15]. The GWW partition function undergoes a third-order phase transition at $\kappa_c = 1/2$ (where $\kappa = \beta/N$), characterised by a third-order discontinuity in the free energy: $F(\kappa)$ and $F'(\kappa)$ and $F''(\kappa)$ are continuous at κ_c, but F''' is not [14].

The spectral gap vanishes as $m(\kappa) \to 0$ when $\kappa \to \kappa_c^+$; the specific power-law exponent requires a detailed analysis of the transfer matrix eigenvalues in the large-N limit, which we now provide using the explicit GWW eigenvalue density. In the weak-coupling phase ($\kappa > \kappa_c$), the eigenvalue density on the unit circle is $\rho(\theta) = (1/\pi)(1 + 2\kappa\cos\theta)$, supported on the full circle. The transfer matrix eigenvalues in the large-N limit are $A_p/A_0 = (2\kappa)^{|p|}$ for $\kappa > \kappa_c$ (Fourier coefficients of ρ), giving $m = \log(A_0/A_1) = -\log(2\kappa)$. At $\kappa_c = 1/2$, $m = -\log(1) = 0$. For κ slightly above κ_c: $m(\kappa) = -\log(2\kappa) = -\log(1 + 2(\kappa - \kappa_c)) \approx 2(\kappa - \kappa_c)$, which is *linear*, not square-root, in $\kappa - \kappa_c$.

We correct the statement: the gap vanishes *linearly* $m \sim 2(\kappa - \kappa_c)$ as $\kappa \to \kappa_c^+$. The *third-order* nature of the transition refers to the free energy F, not the spectral gap. In Stokes language: the Stokes curve γ_{01} in the complex κ-plane touches the real axis at κ_c with tangency order determined by the condition $\mathrm{Im}(\gamma_{01}) \sim (\mathrm{Re}(\gamma_{01}) - \kappa_c)^{3/2}$ as $\kappa \to \kappa_c$. This 3/2 exponent comes from the Stokes curve geometry (the cubic tangency of $\mathrm{Re}(\varphi_0(s) - \varphi_1(s)) = 0$ in the complex plane), not from the gap scaling.

At finite N, the transition is smoothed: $m(\beta) > 0$ for all $\beta > 0$, and the Stokes boundary γ_{01} approaches the real axis at $\beta_c \approx N/2$ to distance $O(1/N^2)$ without crossing. □

Remark 6.2 (Tangency exponent and transition order). The relationship between the Stokes tangency exponent α and the order of the phase transition deserves clarification. If the Stokes curve γ_{01} touches the real axis at β_c with $\mathrm{Im}(\gamma_{01}) \sim (\mathrm{Re}(\gamma_{01}) - \beta_c)^\alpha$, then Fisher zeros accumulate on the real axis with density $\rho \sim |\beta - \beta_c|^{\alpha-1}$ (from the Stokes density formula). The free energy $f(\beta) = (1/n)\log|Z_n| = \int \log|\beta - \beta'|\,\rho(\beta')\,d\beta'$ (by the potential-theoretic representation of $\log|Z_n|$ as a logarithmic potential of the zero measure) then has a singularity of order $\alpha + 1$ in its derivatives, giving a discontinuity in the $\lceil 1 + 1/\alpha \rceil$-th derivative [16]. For the GWW model with $\alpha = 3/2$: $\lceil 1 + 1/(3/2 - 1/2) \rceil = \lceil 2 \rceil = 2$, giving a singularity in the second derivative of F — but GWW is third-order. The discrepancy arises because the tangency geometry of the Stokes curve in the GWW model is more subtle than a simple power law: the relevant quantity is not the *distance* of the Stokes curve from the real axis, but the *density of zeros pinching the real axis*, which involves the full 2D geometry of the Stokes network. A rigorous derivation of the transition order from the Stokes tangency in the GWW model remains an open problem (see §12).

7 Phase Transitions as Stokes Crossings

Theorem 7.1 (Phase transition = Stokes crossing)**.** *For the partition function* $Z_n(s) = \sum_k c_k e^{n\varphi_k(s)}$*:*

(i) The free energy per site $f(\beta) := (1/n)\log Z_n(\beta) = \max_k \mathrm{Re}\,\varphi_k(\beta) + O(\log n/n)$.

(ii) A non-analyticity of f at $\beta = \beta_c$ occurs if and only if $\beta_c \in \mathcal{S} \cap \mathbb{R}$, i.e., the Stokes network crosses the real axis.

Proof. Part (i) follows from the dominated convergence of the exponential sum: as $n \to \infty$, the term with the largest $\operatorname{Re}\varphi_k$ dominates.

Part (ii): If $\beta_c \notin \mathcal{S}$, then there exists a unique dominant term φ_{k_0} with $\operatorname{Re}\varphi_{k_0}(\beta) > \operatorname{Re}\varphi_k(\beta)$ for all $k \neq k_0$ in a neighbourhood of β_c. Then $f(\beta) = \operatorname{Re}\varphi_{k_0}(\beta) + O(e^{-cn})$, which is analytic. Conversely, if $\beta_c \in \mathcal{S}$, then at least two terms have equal real parts at β_c, and the free energy $f = (1/n)\log(c_p e^{n\varphi_p} + c_q e^{n\varphi_q} + \cdots)$ undergoes a "dominant saddle switch," producing a non-analyticity in the limit $n \to \infty$. □

Corollary 7.2 (No phase transition in confining SU(N))**.** *For* SU(N) *with finite N and the Wilson action: $\mathcal{S} \cap \mathbb{R}_{>0} = \emptyset$ (Theorem 6.1), so there is no bulk phase transition at any coupling $\beta > 0$.*

8 Critical Exponents from Stokes Geometry

The Stokes-RG correspondence (Theorem 5.1) establishes that the mass gap controls the distance from the real axis to the Stokes network. The phase transition theorem (Theorem 7.1) identifies phase transitions with Stokes crossings. We now combine these two results with standard finite-size scaling to show that *critical exponents* can be extracted from the Stokes geometry.

Definition 8.1 (Stokes distance)**.** For a lattice system on $L^{d-1} \times T$ with transfer matrix eigenvalues $\lambda_0(\beta, L) > \lambda_1(\beta, L) > 0$, the *Stokes distance* at coupling β and transverse size L is:

$$\Delta(\beta, L) := \inf\big\{y > 0 : |\lambda_0(\beta + iy, L)| = |\lambda_1(\beta + iy, L)|\big\}. \tag{17}$$

Theorem 8.2 (Stokes extraction of critical exponents)**.** *Let $\hat{T}(\beta, L)$ be the transfer matrix of a lattice system on $L^{d-1} \times T$ with eigenvalues $\lambda_0(\beta, L) > \lambda_1(\beta, L) > 0$, mass gap $m(\beta, L) := \log(\lambda_0/\lambda_1)$, and Stokes distance $\Delta(\beta, L)$ as in Definition 8.1. Suppose the system has a second-order phase transition at β_c with correlation length exponent ν.*

(i) (Comparability.) *There exist $0 < c \leq C < \infty$, uniform in a compact neighbourhood of β_c, such that*

$$\frac{m(\beta, L)}{C} \;\leq\; \Delta(\beta, L) \;\leq\; \frac{m(\beta, L)}{c}\,. \tag{18}$$

(ii) (Critical scaling.) *At $\beta = \beta_c$, assuming finite-size scaling of the mass gap [19, 20] and boundedness of the Stokes gradient (Assumption 5 below):*

$$\Delta(\beta_c, L) \;\sim\; c_0 \cdot L^{-1} \quad \text{as } L \to \infty. \tag{19}$$

(iii) (Exponent extraction.) *The exponent ν is determined by the finite-size scaling function $\Delta(\beta, L) = L^{-1}\,\Phi\big((\beta - \beta_c)\,L^{1/\nu}\big)$:*

$$\frac{1}{\nu} = \lim_{L\to\infty} \frac{d\log\big(\partial_\beta \Delta\big|_{\beta_c}\big)}{d\log L}\,. \tag{20}$$

We isolate the key assumption required for parts (ii) and (iii):

Axiom 5 (Stokes gradient boundedness)**.** Define the *Stokes gradient* $G'(y) := \partial_y \log|\lambda_0(\beta + iy, L)/\lambda_1(\beta + iy, L)|$. At the critical coupling β_c, the supremum $C(\beta_c, L) := \sup_{0<y<\Delta} |G'(y)|$ has a finite, nonzero limit as $L \to \infty$: $C(\beta_c, L) \to C_\infty \in (0, \infty)$.

This assumption holds when the Stokes gradient is controlled by the local analytic structure of the eigenvalues (determined by the action, not by L).

Proof of Theorem 8.2. Part (i): Comparability. Define $G(y) := \log|\lambda_0(\beta+iy, L)| - \log|\lambda_1(\beta+iy, L)|$. Then $G(0) = m(\beta, L) > 0$ and $G(\Delta) = 0$ by definition of Δ. The function G is real-analytic on $[0, \Delta]$ (the eigenvalues are analytic functions of $s = \beta + iy$ in a neighbourhood of the real axis, with $\lambda_j \neq 0$ by compactness of the gauge group).

By the mean value theorem, there exists $\xi \in (0, \Delta)$ with

$$m = G(0) - G(\Delta) = |G'(\xi)| \cdot \Delta. \tag{21}$$

Setting $C := \sup_{[0,\Delta]} |G'|$ and $c := \inf_{[0,\Delta]} |G'|$ gives the two-sided bound (18).

The positivity $c > 0$ follows from the strict monotonicity of G on $[0, \Delta]$: since $G(0) > 0$, $G(\Delta) = 0$, and G is real-analytic, G cannot have vanishing derivative on a subinterval (a non-constant analytic function has isolated critical points). Therefore $G' < 0$ on $(0, \Delta)$ except at finitely many points, and $c = \inf |G'| > 0$.

The uniform boundedness of C in (β, L) near β_c follows from the maximum principle: $|G'(y)| \leq |\partial_s \log(\lambda_0/\lambda_1)|$ is bounded on any compact subset of the analyticity domain of $\log(\lambda_0/\lambda_1)$.

Part (ii): Critical scaling. At a second-order critical point with periodic boundary conditions, the standard finite-size scaling result [19, 20] gives:

$$m(\beta_c, L) = L^{-1}\, \tilde{m}(0), \tag{22}$$

where $\tilde{m}(0) > 0$ is a universal constant depending on the universality class (for the 2D Ising model, $\tilde{m}(0) = \pi/4$). From part (i) and Assumption 5:

$$\Delta(\beta_c, L) = \frac{m(\beta_c, L)}{|G'(\xi_L)|} = \frac{\tilde{m}(0)}{|G'(\xi_L)|} \cdot L^{-1} \xrightarrow[L\to\infty]{} \frac{\tilde{m}(0)}{C_\infty} \cdot L^{-1}.$$

Part (iii): Exponent extraction. The full scaling form of the mass gap [19, 20]:

$$m(\beta, L) = L^{-1}\, \tilde{m}\big((\beta - \beta_c)\, L^{1/\nu}\big) \tag{23}$$

combined with part (i) gives $\Delta(\beta, L) = L^{-1}\, \Phi(xL^{1/\nu})$ with $x = \beta - \beta_c$ and $\Phi = \tilde{m}/|G'|$. Differentiating in β at β_c (i.e. at $x = 0$):

$$\left.\frac{\partial \log \Delta}{\partial \beta}\right|_{\beta_c} = \frac{\Phi'(0)}{\Phi(0)}\, L^{1/\nu},$$

where $\Phi'(0)/\Phi(0)$ is a universal, L-independent number. Taking the logarithmic derivative with respect to L:

$$\frac{d}{d\log L}\left(\log\left|\frac{\partial \Delta}{\partial \beta}\Big|_{\beta_c}\right|\right) = \frac{1}{\nu} - 1,$$

from which ν is extracted. Alternatively, comparing Δ at two sizes L_1, L_2 at β_c gives:

$$\frac{\log[\Delta(\beta_c, L_1)/\Delta(\beta_c, L_2)]}{\log(L_2/L_1)} \xrightarrow[L_1,L_2\to\infty]{} 1, \tag{24}$$

confirming $\Delta \sim L^{-1}$; the exponent ν is extracted from the *off-critical* scaling of $\Delta(\beta, L)$ via (20). □

Remark 8.3 (Absolute representation dominance). A key input to Theorem 8.2 (via Theorem 6.1) is that representation dominance is *absolute* for SU(N) lattice gauge theory: the strict ordering $A_0(\beta) > A_1(\beta) > A_2(\beta) > \cdots$ holds for *all* $\beta > 0$ and *all* finite N. There is no eigenvalue crossing on the real axis at any finite N. This means $\Delta(\beta, L) > 0$ for all $\beta > 0$ and all finite N and L — the Stokes distance never vanishes at finite N. A phase transition (and hence $\Delta \to 0$) can occur only in a thermodynamic limit: $L \to \infty$ for standard statistical mechanics, or $N \to \infty$ for the GWW large-N transition.

Remark 8.4 (Numerical verification: 2D Ising). The prediction $\Delta(\beta_c, L) \sim L^{-1}$ is confirmed by the 2D Ising strip computation in [3]: for $L = 2, 3, 4, 6, 8$, the product $\Delta(\beta_c, L) \cdot L$ is approximately constant, with the minimum Stokes distance occurring near $\beta_c = \log(1+\sqrt{2}) = 0.8814\ldots$ (the exact critical coupling). The exact 2D Ising exponent $\nu = 1$ is consistent with the scaling (19).

Conjecture 8.5 (GWW Stokes scaling)**.** *For* SU(N) *with the Wilson action at the GWW critical coupling* $\kappa_c = 1/2$ *(where* $\kappa = \beta/N$*):*

$$m(N, \kappa_c) \sim c_1\, N^{-2/3}, \qquad C(N, \kappa_c) \sim c_2\, N^{1/3}, \tag{25}$$

giving $\Delta(N) \sim (c_1/c_2)\, N^{-1}$*. The* $N^{-2/3}$ *exponent for the mass gap comes from Tracy–Widom universality at the edge of the GWW eigenvalue distribution; the Stokes gradient divergence* $C \sim N^{1/3}$ *comes from the sharpening of the eigenvalue density near the spectral edge.*

9 Stokes Phase Velocity and Mass Spacing

Theorem 9.1 (Mass spacing from Stokes density on the n-plaquette chain)**.** *For* SU(N) *Yang–Mills on the* n*-plaquette chain, the average spacing of Fisher zeros along the dominant Stokes curve satisfies:*

$$\langle \Delta y \rangle = \frac{2\pi}{\langle |d\theta_{01}/d\beta| \rangle} \cdot \frac{1}{n} + O(1/n^2), \tag{26}$$

where $\theta_{01}(\beta) := \mathrm{Im}(\varphi_0 - \varphi_1)$ *is the phase difference of the two dominant transfer matrix eigenvalues along the Stokes curve* γ_{01}*, and the average is over the relevant portion of* γ_{01}*.*

Proof. This follows from the Stokes concentration theorem (Theorem 4.3): the density of zeros near the Stokes curve γ_{01} is $\rho = (n/2\pi)|d\theta_{01}/dt|$, parametrised by arc length. The average spacing is $\langle \Delta y \rangle = 1/\rho$. The $1/n$ scaling is verified numerically for $n = 2, 3, 4, 6, 8$ in [3], where the constant $C = 2\pi/\langle |d\theta_{01}/d\beta| \rangle$ is found to be approximately 2.5 for the Wilson action and 1.75 for the Symanzik-improved action. □

Warning 9.2 (1D chain vs. 4D QCD). Theorem 9.1 is proved for the *n-plaquette chain*, which is a one-dimensional system. The extension to 4D requires: (a) that the dominant Stokes structure of the 4D partition function is controlled by the same eigenvalue pair (A_0, A_1); (b) that the Stokes phase velocity has a thermodynamic limit as the spatial volume $\to \infty$; (c) that the $1/n$ scaling persists. These are expected on physical grounds but unproved.

Conjecture 9.3 (4D extension)**.** *For* SU(N) *Yang–Mills on a 4D lattice $L^3 \times T$, the average spacing of the lowest glueball masses satisfies:*

$$\langle \Delta m \rangle = \frac{2\pi}{\langle |d\theta_{01}/d\beta| \rangle_{4D}} \cdot \frac{1}{T} + O(1/T^2), \tag{27}$$

where θ_{01} is the phase difference of the two dominant 4D transfer matrix eigenvalues and the average is in the thermodynamic limit $L \to \infty$.

10 Connections and Context

10.1 Spacetime reconstruction from the mass gap

Remark 10.1 (OS reconstruction). Conditional on the completion of the mass gap programme [1] (specifically: the continuum limit), the lattice gauge theory constructed here produces a continuum QFT satisfying the Wightman axioms via the Osterwalder–Schrader reconstruction [13]. The proof chain is standard: reflection positivity (preserved by the block-spin transformation) $\to$ exponential clustering (from the mass gap) $\to$ OS axioms OS0–OS4 $\to$ Hilbert space, Hamiltonian, vacuum $\to$ analytic continuation to Minkowski signature (edge-of-the-wedge theorem) $\to$ Poincaré invariance, microcausality, relativistic dispersion. The details are in [1], Sections 15–18. The OS reconstruction is due to Osterwalder and Schrader; our contribution is providing the *inputs* (mass gap, exponential clustering, reflection positivity).

10.2 Connection to the CCM spectral action

Remark 10.2 (CCM programme). The Connes–Chamseddine–Marcolli (CCM) spectral action programme [6, 7, 8] provides: (1) the gauge group $G_{\mathrm{SM}} = \mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ from the automorphism group of a finite spectral triple (classified by Barrett [9]); (2) the Higgs mechanism from inner fluctuations; (3) Einstein–Hilbert gravity, Yang–Mills, and the Higgs potential from the Seeley–DeWitt expansion of the spectral action. These are *inputs* to our framework, not outputs. What the present work adds to CCM is the non-perturbative dynamical mechanism (convexity-driven scale generation, confinement, mass gap) that was missing from the spectral action programme, which is purely classical and perturbative.

10.3 Mass hierarchy from group theory

Remark 10.3 (Multi-sector running). For the Standard Model gauge group $G_{\mathrm{SM}} = \mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$, the one-loop β-function coefficients $b_0^{(a)} = 11\, C_A^{(a)}/(3 \cdot 16\pi^2)$ (pure gauge) satisfy $b_0^{(3)}/b_0^{(2)} = 3/2$. Starting from a unified coupling g_0^2 at a high scale Λ, the dynamically

generated mass scales satisfy $\Lambda_3/\Lambda_2 = \exp(c/g_0^2)$ for an explicit constant c determined by the b_0 ratio. This is the standard one-loop running, reframed in our language: g_0^2 is the only free parameter, and the mass hierarchy between QCD and the electroweak scale is then a prediction (given G_{SM}). With $g_0^2 \approx 0.5$ ($\alpha_{\mathrm{GUT}} \approx 1/25$): $\Lambda_{\mathrm{QCD}}/m_W \sim 10^{-2.5}$, compared to the observed $\sim 10^{-2.4}$.

10.4 Constraints on the gauge group

Remark 10.4 (What the axioms constrain). Axioms A–D are satisfied by *every* compact simple Lie group in $d = 4$ (since $b_0 = 11\, C_A/(3 \cdot 16\pi^2) > 0$ for all non-abelian compact simple groups). They cannot distinguish SU(3) from SO(10), E_6, or any other compact group. U(1) factors are not asymptotically free, so Axiom D in its present form excludes pure U(1) unless coupled to non-abelian sectors. The gauge group selection problem remains the single largest gap in this framework.

11 Status of the Mass Gap Programme

The mass gap proof [1] is central to this programme. We are transparent about its current status.

What is proved: (1) Large-field bound (Peter–Weyl quartic) with explicit constants; (2) Fourier contour propagator bound ($\alpha_0 \approx 1.317$); (3) Reblocking constant $C_{\mathrm{RB}} \leq 6.2$; (4) Extended cluster expansion convergence for $g^2 < 0.382$; (5) Non-perturbative coupling growth via convexity; (6) OS strong-coupling bound $m_{\mathrm{OS}} \geq 0.0733$; (7) Lattice mass gap positive for all $g^2 > 0$ (combining RG + OS); (8) OS reconstruction, continuum limit via Prokhorov tightness, Wightman axioms via analytic continuation.

Open items: (1) Independent verification of the polymer counting constants (safety factor ≈ 20, but an independent check is desirable); (2) Full $O(4)$ invariance in the continuum limit (Symanzik improvement programme, standard but not verified with tracked constants in the Balaban framework); (3) Uniqueness of the continuum limit (relies on two-loop asymptotic scaling).

None of these items affect the *existence* of a mass gap; they affect the *construction of the continuum QFT*. The Stokes-RG correspondence and spectral confinement theorems of this paper are *unconditional*: they hold at the lattice level and do not require the continuum limit.

Remark 11.1 (Absolute representation dominance). For any compact group G with strictly positive heat kernel, the eigenvalue ordering $A_0(\beta) > A_1(\beta) > \cdots$ holds for *all* $\beta > 0$ and all finite N. There is no eigenvalue crossing at any real coupling. In particular, $\Delta(\beta) > 0$ for all finite systems. Phase transitions (where $\Delta \to 0$) require a thermodynamic limit $N \to \infty$ or $L \to \infty$ and manifest as the Stokes curve approaching the real axis asymptotically, never crossing it at finite N.

12 Open Problems

1. **Derive G_{SM} from a dynamical principle.** The gauge group enters through Axiom A. The axioms alone cannot select the Standard Model gauge group (Remark 10.4).

2. **Derive $d = 4$.** In $d = 4$, the Yang–Mills coupling is classically marginal, giving a logarithmic hierarchy. In $d = 3$ it is polynomial; in $d = 5$, non-renormalisable. This is suggestive but not a derivation.

3. **Extend the mass spacing formula to 4D.** Prove or disprove Conjecture 9.3.

4. **Rigorous Stokes tangency–transition order connection.** Derive the order of the GWW phase transition from the Stokes tangency exponent (Remark 6.2).

5. **Complete the mass gap proof.** Resolve the three open items in §11.

13 Conclusion

The central result of this paper is the Stokes-RG correspondence (Theorem 5.1): the RG flow through coupling space is simultaneously a traversal of the Stokes network of the partition function. The mass gap at each scale is the Stokes gap; confinement is the topological statement that the Stokes network does not cross the real axis; and phase transitions occur precisely when it does.

This framework provides a *mechanism* — convexity-driven scale generation — and a *geometric picture* — the Stokes-RG correspondence — connecting the constructive QFT programme with the Lee–Yang theory of partition function zeros. The single most important next step is the completion of the mass gap programme: if the open items of §11 are resolved, the entire construction from Axioms A–D to spacetime reconstruction becomes unconditional.

References

[1] G. Olbryk, "The Mass Gap for Four-Dimensional SU(N) Yang–Mills Theory: Explicit Constant Tracking in Balaban's Multi-Scale Renormalization Group," preprint (2026).

[2] G. Olbryk, "Global Spectral Theory of Everything — Version 3: Stokes Foundation," preprint (2026).

[3] G. Olbryk, "Fisher Zeros as Stokes Phenomena in Lattice Gauge Theory," preprint (2026).

[4] T. Balaban, "Ultraviolet stability in field theory: the φ^4_3 model," in *Scaling and Self-Similarity in Physics*, Birkhäuser (1985).

[5] T. Balaban, "Renormalization group approach to lattice gauge field theories. I. Generation of effective actions in a small field approximation and a coupling constant renormalization in four dimensions," Comm. Math. Phys. **109** (1987) 249–301.

[6] A. Chamseddine, A. Connes, M. Marcolli, "Gravity and the Standard Model with neutrino mixing," Adv. Theor. Math. Phys. **11** (2007) 991–1089.

[7] A. Chamseddine, A. Connes, "The spectral action principle," Comm. Math. Phys. **186** (1997) 731–750.

[8] A. Connes, "On the spectral characterization of manifolds," J. Noncommut. Geom. **7** (2013) 1–82.

[9] J. Barrett, "A Lorentzian version of the non-commutative geometry of the Standard Model of particle physics," J. Math. Phys. **48** (2007) 012303.

[10] K. Wilson, "Confinement of quarks," Phys. Rev. D **10** (1974) 2445.

[11] J. Kogut, L. Susskind, "Hamiltonian formulation of Wilson's lattice gauge theories," Phys. Rev. D **11** (1975) 395.

[12] K. Osterwalder, E. Seiler, "Gauge field theories on a lattice," Ann. Phys. **110** (1978) 440–471.

[13] K. Osterwalder, R. Schrader, "Axioms for Euclidean Green's functions," Comm. Math. Phys. **31** (1973) 83–112.

[14] D. Gross, E. Witten, "Possible third-order phase transition in the large-N lattice gauge theory," Phys. Rev. D **21** (1980) 446.

[15] S. Wadia, "$N = \infty$ phase transition in a class of exactly soluble model lattice gauge theories," Phys. Lett. B **93** (1980) 403.

[16] T.D. Lee, C.N. Yang, "Statistical theory of equations of state and phase transitions. II. Lattice gas and Ising model," Phys. Rev. **87** (1952) 410.

[17] Á. Baricz, "Turán type inequalities for modified Bessel functions," Bull. Aust. Math. Soc. **82** (2010) 254–264.

[18] G. S. Bali, "Casimir scaling of SU(3) static potentials," Phys. Rev. D **62** (2000) 114503.

[19] M. E. Fisher, M. N. Barber, "Scaling theory for finite-size effects in the critical region," Phys. Rev. Lett. **28** (1972) 1516.

[20] M. N. Barber, "Finite-size scaling," in *Phase Transitions and Critical Phenomena*, Vol. 8, C. Domb and J. Lebowitz, eds., Academic Press (1983), pp. 145–266.