

Constraints on Four-Dimensional Quantum Gauge Theories:

Mass Gap, Stokes-RG, and the Uniqueness
of the Standard Model

Grzegorz Olbryk
g.olbryk@gmail.com

March 2026

Abstract

We systematically enumerate the independent constraints that a four-dimensional quantum gauge theory must satisfy, and ask: how many theories survive all of them simultaneously? The six constraints are: (C1) mass gap existence (from our non-perturbative proof using convexity-based coupling growth); (C2) asymptotic freedom ($b_0 > 0$, restricting gauge groups and matter content); (C3) anomaly cancellation (constraining fermion representations); (C4) Stokes-RG correspondence (Fisher zeros on the Stokes network of the transfer matrix spectrum); (C5) convexity of the lattice free energy; (C6) Osterwalder–Schrader reconstruction (reflection positivity yielding a physical Hilbert space and Hamiltonian). We show that (C1)–(C3) alone reduce the space of candidate theories to a finite list. Constraint (C3) is particularly powerful: for $SU(3) \times SU(2) \times U(1)$, anomaly cancellation forces fermion hypercharge assignments that are uniquely those of the Standard Model (up to family replication). We are rigorous throughout about what is proved, what follows from established results in the literature, and what remains conjectural. The paper does not claim to derive the Standard Model from first principles; it maps the logical landscape of constraints and identifies where uniqueness arguments succeed and where they fail.

Contents

1	Introduction: The Constraint Approach	3
1.1	Motivation	3
1.2	The six constraints	3
1.3	Logical status of each constraint	4
1.4	What this paper does not claim	4
2	Constraint 1: Mass Gap Existence	4
2.1	Statement	4
2.2	Proof architecture	5

2.3	Constraint content	5
3	Constraint 2: Asymptotic Freedom	5
3.1	The one-loop coefficient	5
3.2	Pure gauge theories	6
3.3	With matter: the bound on representations	6
3.4	Upper bound on the gauge group rank	7
4	Constraint 3: Anomaly Cancellation	7
4.1	Gauge anomalies	7
4.2	Anomaly types for $SU(3) \times SU(2) \times U(1)$	7
4.3	Anomaly cancellation forces the Standard Model	8
5	Constraint 4: Stokes-RG Correspondence	9
5.1	Statement	9
5.2	Application to lattice gauge theory	9
5.3	Phase transitions from Stokes crossings	9
5.4	Constraint content	9
6	Constraint 5: Convexity of Free Energy	10
6.1	Statement and proof	10
6.2	Connection to the coupling growth inequality	10
6.3	Constraint content	10
7	Constraint 6: OS Reconstruction	11
7.1	Statement	11
7.2	Reflection positivity of the Wilson action	11
7.3	Constraint content	11
8	Combined Constraints: How Many Theories Survive?	12
8.1	Step 1: Gauge group from (C1) + (C2)	12
8.2	Step 2: Fermion content from (C3)	12
8.3	Step 3: Chiral matter requires a $U(1)$ factor	13
8.4	Step 4: Anomaly cancellation + chirality + AF	13
8.5	Step 5: The survivors	13
8.6	The key question	14
9	The Standard Model as a Survivor	15
9.1	Verification of all six constraints	15
9.2	Subtleties	16
9.3	What the constraints explain	16
9.4	What the constraints do not explain	17

10 What Remains Open	17
10.1 Gravity	17
10.2 The cosmological constant problem	17
10.3 The number of generations	18
10.4 The strong CP problem	18
10.5 Neutrino masses	18
11 Discussion	18
11.1 Summary of results	18
11.2 Comparison with other “uniqueness” arguments	19
11.3 The constraint approach as methodology	19

1 Introduction: The Constraint Approach

1.1 Motivation

The traditional approach to a “Theory of Everything” attempts to *generate* all of physics from a single principle: a Lagrangian, an action, a symmetry, or a set of axioms. Every such attempt faces the same problem: the gap between a beautiful principle and the messy specifics of the Standard Model (three generations, the particular gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y$, the hypercharge assignments, the CKM matrix) is enormous, and no known principle bridges it.

We propose a complementary approach: instead of generating physics from one principle, we ask whether physics is the *unique structure* satisfying multiple independent constraints. Each constraint eliminates a large class of candidate theories. The question is whether the intersection is empty, a point, or a small finite set.

1.2 The six constraints

We consider the following constraints on a 4D quantum gauge theory with compact gauge group G and matter fields in representations $\{R_i\}$:

- (C1) **Mass gap:** The pure Yang–Mills sector has $\inf \sigma(H \upharpoonright_{\Omega^\perp}) \geq m > 0$.
- (C2) **Asymptotic freedom:** The one-loop β -function coefficient satisfies $b_0(G, \{R_i\}) > 0$.
- (C3) **Anomaly cancellation:** All gauge anomalies vanish: $\mathcal{A}^{abc} = \text{tr}[T^a \{T^b, T^c\}] = 0$.
- (C4) **Stokes-RG correspondence:** Fisher zeros of the partition function lie on the Stokes network of the transfer matrix spectrum.
- (C5) **Convexity:** The lattice free energy $F(\beta) = -\log Z(\beta)$ is convex in β .
- (C6) **OS reconstruction:** The lattice measure satisfies reflection positivity, yielding via the Osterwalder–Schrader theorem a physical Hilbert space $\mathcal{H}$, a self-adjoint Hamiltonian $H \geq 0$, and unitary time evolution.

1.3 Logical status of each constraint

We are explicit about the logical status:

Constraint	Status	Source
(C1) Mass gap	Proved for $SU(N)$, $N \geq 2$	[1]
(C2) Asymptotic freedom	Proved (perturbative, b_0 exact)	[4, 5]
(C3) Anomaly cancellation	Proved (algebraic identities)	[6, 7]
(C4) Stokes-RG	Proved for exponential sums	[2]
(C5) Convexity	Proved (Griffiths, lattice)	[11, 12]
(C6) OS reconstruction	Proved (Wilson action RP)	[8, 9, 1]

Each constraint is individually established. The new content of this paper is the *combined* analysis: what survives all six simultaneously?

1.4 What this paper does not claim

We do not claim to derive the Standard Model from first principles. Specifically:

- We do not derive the gauge group $SU(3) \times SU(2) \times U(1)$.
- We do not derive the number of generations.
- We do not derive the Yukawa couplings, mixing angles, or masses.
- We do not incorporate gravity.

What we *do* show is that the combined constraints drastically reduce the space of allowed theories, and that several features of the Standard Model that appear arbitrary are in fact forced by the conjunction of (C1)–(C3).

2 Constraint 1: Mass Gap Existence

2.1 Statement

The following theorem is proved in [1]:

Theorem 2.1 (Mass gap for $SU(N)$ Yang–Mills [1, Thm. 3.2]). *For every $N \geq 2$ and every bare coupling $g^2 > 0$, the four-dimensional $SU(N)$ lattice Yang–Mills theory with Wilson action has a positive mass gap:*

$$m = -\lim_{t \rightarrow \infty} \frac{1}{t} \log \langle W(C_0) W(C_t) \rangle \geq m_0(N) > 0,$$

uniformly in the lattice volume. The continuum limit satisfies the Wightman axioms $W0$ – $W4$ with $\inf \sigma(H \upharpoonright_{\Omega^\perp}) \geq m_{\text{phys}} > 0$.

2.2 Proof architecture

The proof proceeds in three regimes that cover all $g^2 > 0$:

- (i) **Weak coupling** ($g^2 < g_c^2(N) = 0.382$ for $N = 2$): The extended Balaban cluster expansion converges, with the convergence domain enlarged by two explicit improvement factors: the Fourier contour propagator bound ($f_{\text{CT}} = 1.37$, decay rate $\alpha_0 \approx 1.317$) and the tight reblocking constant ($f_{\text{RB}} = 1.45$, $C_{\text{RB}} \leq 6.2$).
- (ii) **Strong coupling** ($g^2 \geq g_c^2(N)$): The Osterwalder–Seiler bound gives $m_{\text{latt}} \geq -\log z(\beta/N, N) > 0$ directly via representation dominance in the transfer matrix.
- (iii) **RG bridge**: For any $g_0^2 > 0$, the non-perturbative coupling growth inequality

$$g_{k+1}^2 \geq g_k^2 + c_{\text{NP}} g_k^4 \quad (1)$$

(proved via convexity of the free energy, not Feynman diagrams) ensures the running coupling reaches g_c^2 in finitely many RG steps.

2.3 Constraint content

As a constraint, (C1) is currently *proved* only for the $\text{SU}(N)$ series. For other compact simple groups G :

- $\text{SO}(N)$, $\text{Sp}(N)$: The proof architecture (Balaban RG + OS bound) extends, but the explicit constants (f_{CT} , f_{RB} , large-field bound) have not been tracked. The mass gap is *expected* but not proved.
- Exceptional groups (G_2 , F_4 , E_6 , E_7 , E_8): Same status. Lattice simulations strongly support a mass gap, but no rigorous proof exists.

Remark 2.2 (Universality of the mass gap). We conjecture that every compact simple Lie group G in 4D has a mass gap. The proof strategy of [1] is G -universal in its logical structure; the G -dependent inputs are $b_0(G)$ and $\dim \mathfrak{g}$. For the purpose of the constraint analysis below, we treat (C1) as established for $\text{SU}(N)$ and conjectured for other groups.

3 Constraint 2: Asymptotic Freedom

3.1 The one-loop coefficient

For a 4D gauge theory with gauge group G and n_f Weyl fermions in representation R , the one-loop β -function coefficient is:

$$b_0 = \frac{1}{(4\pi)^2} \left[\frac{11}{3} C_2(G) - \frac{2}{3} n_f T(R) - \frac{1}{3} n_s T(R_s) \right], \quad (2)$$

where $C_2(G)$ is the quadratic Casimir of the adjoint representation, $T(R)$ is the Dynkin index of the fermion representation, and n_s , $T(R_s)$ refer to complex scalars. Asymptotic freedom requires $b_0 > 0$.

3.2 Pure gauge theories

For pure Yang–Mills (no matter), $b_0 > 0$ for *every* compact simple Lie group. The values of $C_2(G)$ are:

Group	$C_2(G)$	$b_0 (4\pi)^2$	Notes
$SU(N)$	N	$11N/3$	$N \geq 2$
$SO(N)$	$N - 2$	$11(N - 2)/3$	$N \geq 5$
$Sp(2n)$	$n + 1$	$11(n + 1)/3$	$n \geq 1$
G_2	4	44/3	
F_4	9	33	
E_6	12	44	
E_7	18	66	
E_8	30	110	

Pure gauge: *no constraint* from asymptotic freedom alone. Every compact simple group is asymptotically free in 4D without matter.

3.3 With matter: the bound on representations

Adding n_f Weyl fermions in a representation R with Dynkin index $T(R)$, asymptotic freedom requires:

$$n_f < \frac{11 C_2(G)}{2 T(R)}. \quad (3)$$

For the Standard Model gauge factors:

Factor	C_2	Fund. $T(R)$	Max Weyl fermions
$SU(3)_c$	3	1/2	< 33
$SU(2)_L$	2	1/2	< 22
$U(1)_Y$	0	—	(see below)

For a product group $G_1 \times G_2 \times \dots$, asymptotic freedom of each factor is an independent constraint.

Remark 3.1 ($U(1)$ is never asymptotically free). For $U(1)$, $C_2(G) = 0$ and $b_0 < 0$ whenever charged matter is present. The $U(1)_Y$ factor of the Standard Model is *not* asymptotically free. It is infrared free: $b_0^Y < 0$.

This means $U(1)_Y$ by itself does not define a consistent quantum field theory at arbitrarily high energies (Landau pole). The resolution is either: (a) embedding in a non-abelian GUT group at high energies, or (b) treating the Standard Model as an effective theory valid below some cutoff Λ_{LP} .

For the constraint analysis, we note that asymptotic freedom is a constraint on the *non-abelian* factors only. The U(1) factor is constrained instead by anomaly cancellation and charge quantisation.

3.4 Upper bound on the gauge group rank

For a simple group G with n_{gen} generations of fermions in the fundamental representation:

$$b_0 > 0 \iff n_{\text{gen}} < \frac{11 C_2(G)}{2 T(\text{fund})}.$$

For $\text{SU}(N)$ with fundamentals ($T = 1/2$): $n_{\text{gen}} < 11N$. With three generations and the Standard Model matter content (per generation: 2 doublets and 3 triplets of Weyl fermions for each colour factor), the bound is easily satisfied for $\text{SU}(3)$ and $\text{SU}(2)$.

The constraint becomes restrictive for *large* representations. For $\text{SU}(N)$ with fermions in the adjoint ($T = N$): $n_f < 11/2$, so at most 5 adjoint Weyl fermions (or 2 adjoint Dirac fermions with room for one more Weyl). This rules out many $\mathcal{N} = 4$ Super-Yang–Mills deformations as UV-complete non-supersymmetric theories.

4 Constraint 3: Anomaly Cancellation

4.1 Gauge anomalies

A gauge theory is consistent only if all gauge anomalies cancel. For a set of left-handed Weyl fermions ψ_i in representations R_i of the gauge group G , the anomaly coefficients are:

$$\mathcal{A}^{abc} = \sum_i \text{tr}_{R_i}[T^a \{T^b, T^c\}]. \quad (4)$$

The theory is anomaly-free if $\mathcal{A}^{abc} = 0$ for all generators T^a, T^b, T^c .

4.2 Anomaly types for $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$

For the Standard Model gauge group, there are six independent anomaly conditions:

- (i) $[\text{SU}(3)]^3$: automatically satisfied (fundamentals and anti-fundamentals contribute with opposite signs).
- (ii) $[\text{SU}(2)]^3$: automatically zero (the anomaly coefficient $A(R) = 0$ for $\text{SU}(2)$ representations of integer isospin; for half-integer isospin, contributions cancel between quarks and leptons).
- (iii) $[\text{SU}(3)]^2 \text{U}(1)$: $\sum_{\text{L quarks}} Y_i = 0$. With n_c colours, this gives $2Y_Q + Y_u + Y_d = 0$ per generation.
- (iv) $[\text{SU}(2)]^2 \text{U}(1)$: $\sum_{\text{L doublets}} Y_i = 0$. This gives $n_c Y_Q + Y_L = 0$ per generation.
- (v) $[\text{U}(1)]^3$: $\sum_i Y_i^3 = 0$ (sum over all left-handed Weyl fermions).
- (vi) $[\text{grav}]^2 \text{U}(1)$: (mixed gravitational anomaly) $\sum_i Y_i = 0$.

4.3 Anomaly cancellation forces the Standard Model

Proposition 4.1 (Hypercharge quantisation from anomaly cancellation). *For $SU(3) \times SU(2) \times U(1)$ with n_{gen} generations, each containing the chiral fermion content $(Q_L, u_R, d_R, L_L, e_R)$ in the representations*

$$Q_L : (\mathbf{3}, \mathbf{2}, Y_Q), \quad u_R : (\bar{\mathbf{3}}, \mathbf{1}, Y_u), \quad d_R : (\bar{\mathbf{3}}, \mathbf{1}, Y_d), \quad L_L : (\mathbf{1}, \mathbf{2}, Y_L), \quad e_R : (\mathbf{1}, \mathbf{1}, Y_e),$$

anomaly cancellation uniquely determines the hypercharge ratios (up to an overall normalisation):

$$Y_Q : Y_u : Y_d : Y_L : Y_e = \frac{1}{6} : -\frac{2}{3} : \frac{1}{3} : -\frac{1}{2} : 1.$$

These are exactly the Standard Model hypercharge assignments.

Proof. The six anomaly conditions give five equations in five unknowns (with one overall normalisation fixed by convention). The system is:

$$[SU(3)]^2 U(1) : \quad 2Y_Q + Y_u + Y_d = 0, \tag{5}$$

$$[SU(2)]^2 U(1) : \quad 3Y_Q + Y_L = 0, \tag{6}$$

$$[U(1)]^3 : \quad 6Y_Q^3 + 3Y_u^3 + 3Y_d^3 + 2Y_L^3 + Y_e^3 = 0, \tag{7}$$

$$[\text{grav}]^2 U(1) : \quad 6Y_Q + 3Y_u + 3Y_d + 2Y_L + Y_e = 0. \tag{8}$$

From (5): $Y_u + Y_d = -2Y_Q$. From (6): $Y_L = -3Y_Q$. From (8): $6Y_Q + 3(-2Y_Q) + 2(-3Y_Q) + Y_e = 0$, giving $Y_e = 6Y_Q$. The remaining degree of freedom is fixed by (7), which yields (after substitution) a cubic equation with the unique rational solution $Y_u = -4Y_Q$, $Y_d = 2Y_Q$.

Setting $Y_Q = 1/6$ (the conventional normalisation) gives the Standard Model assignments. $\square$

Remark 4.2. The power of this result is often underappreciated. Given only the *group* $SU(3) \times SU(2) \times U(1)$ and the *representation structure* (one generation of quarks and leptons in the representations listed above), the hypercharge assignments are uniquely determined by anomaly cancellation. No dynamics, no Lagrangian, no symmetry breaking is needed.

Remark 4.3 (What anomaly cancellation does *not* fix). Anomaly cancellation does not determine:

- The number of generations n_{gen} (any $n_{\text{gen}} \geq 1$ works).
- The Yukawa couplings (these are free parameters).
- The Higgs sector (anomaly cancellation does not require a Higgs field).
- Whether the gauge group is $SU(3) \times SU(2) \times U(1)$ in the first place.

5 Constraint 4: Stokes-RG Correspondence

5.1 Statement

The Stokes-RG correspondence, proved in [2], establishes that the Fisher zeros of a partition function of the form $Z_n(s) = \sum_{k=0}^K c_k e^{n\varphi_k(s)}$ are controlled by the Stokes geometry of the spectral exponents $\{\varphi_k\}$:

Theorem 5.1 (Stokes concentration [2, Thm. 1]). *Under non-degeneracy and velocity-separation assumptions:*

- (i) *Every zero satisfies $\text{dist}(s_j^{(n)}, \mathcal{S}) \leq C/n$, where $\mathcal{S} = \{\text{Re } \varphi_p = \text{Re } \varphi_q\}$ is the Stokes network.*
- (ii) *The zero density along each Stokes curve is $n |\dot{\theta}_p - \dot{\theta}_q|/(2\pi) + O(1)$.*
- (iii) *The total zero count is $\mathcal{N}(\Omega) = O(n)$.*

5.2 Application to lattice gauge theory

For an n -plaquette lattice Yang–Mills partition function with gauge group G , the transfer matrix representation gives:

$$Z_n(\beta) = \sum_p d_p^2 e^{n \log A_p(\beta)},$$

where p ranges over irreducible representations of G , $d_p = \dim R_p$, and $A_p(\beta)$ is the corresponding transfer matrix eigenvalue. Setting $\varphi_p(s) = \log A_p(s)$ with $s = \beta \in \mathbb{C}$, the zeros lie on the Stokes network $\text{Re } \log A_p = \text{Re } \log A_q$, i.e., on the curves $|A_p(s)| = |A_q(s)|$.

5.3 Phase transitions from Stokes crossings

A corollary of Theorem 5.1 connects partition function zeros to thermodynamic phase transitions:

Corollary 5.2 (Phase transition = Stokes crossing [2, Cor. 2]). *In the thermodynamic limit $n \rightarrow \infty$, the free energy per site $f(\kappa) = -\lim_{n \rightarrow \infty} n^{-1} \log Z_n(\kappa)$ is non-analytic at κ_0 if and only if $\kappa_0 \in \mathcal{S} \cap \mathbb{R}$, i.e., the Stokes network crosses the real coupling axis at κ_0 .*

The order of the transition equals the order of vanishing of $g_p(\kappa) - g_q(\kappa)$ at κ_0 , where $g_k = \text{Re } \varphi_k$.

5.4 Constraint content

As a constraint on candidate theories, (C4) is structural rather than eliminative: it does not rule out gauge groups, but it provides a universal organising principle for the analytic structure of the partition function. Its primary use is in conjunction with (C1):

Proposition 5.3 (Stokes-RG and mass gap compatibility). *If the Stokes network $\mathcal{S}$ does not cross the positive real β -axis (i.e., $\mathcal{S} \cap \mathbb{R}_{>0} = \emptyset$), then there is no finite- β phase transition, and the mass gap $m(\beta)$ is an analytic function of β for all $\beta > 0$.*

For $SU(N)$ Yang–Mills on a finite lattice, numerical evidence [2] shows that no Stokes curve crosses the real axis at fixed N — consistent with the absence of a bulk phase transition (the deconfinement transition occurs only in the thermodynamic limit via the GWW mechanism at $\kappa_c \sim N \log N / (N + 1)$).

6 Constraint 5: Convexity of Free Energy

6.1 Statement and proof

Proposition 6.1 (Convexity of lattice free energy). *For the lattice Yang–Mills partition function $Z(\beta) = \int e^{\beta S_W(U)} dU$ with the Wilson action $S_W = (1/N) \sum_p \text{Re Tr } U_p$, the free energy $F(\beta) = -\log Z(\beta)$ is convex in β :*

$$F''(\beta) = \text{Var}_\beta(S_W) \geq 0.$$

Strict convexity ($F'' > 0$) holds whenever S_W is non-constant dU -a.e., which is the case for all finite lattices with at least one plaquette.

Proof. This is a consequence of the Cauchy–Schwarz inequality applied to the Gibbs measure $d\mu_\beta = Z^{-1} e^{\beta S_W} dU$. The second derivative $F''(\beta) = \langle S_W^2 \rangle_\beta - \langle S_W \rangle_\beta^2 = \text{Var}_\beta(S_W) \geq 0$. $\square$

6.2 Connection to the coupling growth inequality

The convexity of F is the foundation of the non-perturbative coupling growth inequality (1) proved in [1]. The variance lower bound

$$\text{Var}_\beta(S_W) \geq c_{\text{var}} g^2$$

(with $c_{\text{var}} > 0$ depending on N and the lattice geometry) implies that the plaquette coefficient β_k *strictly decreases* under each RG blocking step, giving $g_{k+1}^2 > g_k^2$. The quartic lower bound $g_{k+1}^2 - g_k^2 \geq c_{\text{NP}} g_k^4$ follows from the more refined analysis of the variance in the cluster expansion.

6.3 Constraint content

Convexity is automatic for any lattice gauge theory with a real-valued, bounded action. It does not distinguish between gauge groups. However, it plays a crucial structural role: it ensures that the RG flow is monotone and that the coupling cannot “stick” at a finite value. Combined with (C1), convexity guarantees that the RG flow always reaches the strong-coupling regime where the OS bound applies.

Remark 6.2 (Convexity for non-Wilson actions). For improved lattice actions (Symanzik, Iwasaki, DBW2), convexity of $F(\beta)$ still holds by the same variance argument, as long as the action is a real-valued function of the link variables. The coupling growth inequality (1) then holds with c_{NP} depending on the specific action.

7 Constraint 6: OS Reconstruction

7.1 Statement

The Osterwalder–Schrader reconstruction theorem [8, 9] provides the bridge from a Euclidean lattice measure to a physical quantum theory:

Theorem 7.1 (OS reconstruction). *Let μ be a measure on lattice configurations satisfying:*

- (OS1) *Regularity (polynomial boundedness of moments),*
- (OS2) *Euclidean invariance,*
- (OS3) *Reflection positivity: $\int \overline{\theta F} \cdot F d\mu \geq 0$ for all F supported on the positive-time half-lattice,*
- (OS4) *Cluster property (decay of correlations).*

Then there exists a separable Hilbert space $\mathcal{H}$, a normalised vacuum vector $\Omega \in \mathcal{H}$, a self-adjoint Hamiltonian $H \geq 0$ with $H\Omega = 0$, and a unitary representation of the Poincaré group on $\mathcal{H}$ (in the continuum limit).

7.2 Reflection positivity of the Wilson action

The Wilson gauge action $S_W = (\beta/N) \sum_p \text{Re Tr} U_p$ satisfies reflection positivity because plaquette variables decompose across the reflection plane into contributions from links on each side, coupled through a positive-definite kernel formed by links *on* the reflection plane [10].

This is a special property of the Wilson action. Not all lattice actions satisfy RP:

Warning 7.2 (RP is not automatic). Improved actions (Symanzik, Iwasaki) involve rectangles 1×2 that straddle the reflection plane, potentially violating RP. The spectral functional $\mathcal{L}$ of the global spectral TOE programme [3] fails RP due to all-pairs coupling with negative-definite cross-kernel. RP is a genuine constraint that eliminates candidate theories.

7.3 Constraint content

As a constraint, (C6) is powerful: it eliminates any candidate “action” that does not admit a decomposition with a positive-definite coupling across a reflection plane. This rules out:

- Actions with non-local couplings (e.g., the spectral functional $\mathcal{L}$ of [3]).
- Certain higher-derivative theories.
- Non-unitary theories (e.g., Lee–Wick theories without the Lee–Wick prescription).

For the standard Wilson action with any compact gauge group G , RP holds automatically. Thus (C6) does not further constrain the gauge group choice among standard lattice formulations.

8 Combined Constraints: How Many Theories Survive?

We now apply the six constraints simultaneously and count the surviving theories.

8.1 Step 1: Gauge group from (C1) + (C2)

From (C2) (asymptotic freedom), the non-abelian gauge group must be a product of compact simple Lie groups. As shown in §3, *every* compact simple group is asymptotically free without matter. With Standard Model-like matter content (chiral fermions in fundamental and anti-fundamental representations), asymptotic freedom imposes upper bounds on the number of fermion flavours but does not uniquely select the gauge group.

From (C1) (mass gap), only $SU(N)$ is currently proved. Assuming the conjecture of Remark 2.2, every compact simple group passes (C1).

After (C1) + (C2): An infinite family of candidate gauge groups survives — any finite product of compact simple groups with sufficiently few matter fields.

8.2 Step 2: Fermion content from (C3)

Anomaly cancellation is the most restrictive algebraic constraint. We analyse it systematically.

Definition 8.1 (Anomaly-free gauge theory). A 4D chiral gauge theory with gauge group G and left-handed Weyl fermions in representations $\{R_i\}$ is *anomaly-free* if all cubic gauge anomaly coefficients vanish: $\sum_i A(R_i) = 0$, where $A(R)$ is the anomaly coefficient of representation R .

For simple groups, the anomaly classification depends on the group type:

Group	Anomaly structure
$SU(N)$, $N \geq 3$	Non-trivial $A(R) \neq 0$ for complex reps
$SU(2)$	$A(R) = 0$ for all R (pseudo-real)
$SO(N)$, $N \neq 6$	$A(R) = 0$ for all R (real/pseudo-real)
$Sp(2n)$	$A(R) = 0$ for all R (pseudo-real)
Exceptionals	$A(R) = 0$ for all R (real)

Groups with only real or pseudo-real representations ($SU(2)$, $SO(N)$, $Sp(2n)$, exceptionals) are automatically anomaly-free for *any* fermion content. The constraint (C3) is non-trivial only for gauge groups containing $SU(N)$ factors with $N \geq 3$ coupled to chiral fermions.

8.3 Step 3: Chiral matter requires a U(1) factor

For a purely non-abelian gauge group (no U(1) factor), chiral fermion masses require either: (a) Yukawa couplings to a Higgs field, which requires a scalar in a specific representation, or (b) dynamical symmetry breaking.

However, the existence of chiral fermions in nature (observed parity violation) imposes an independent constraint: the gauge group must admit complex representations. Among simple groups, only $SU(N)$ with $N \geq 3$ has complex representations. The groups $SO(N)$, $Sp(2n)$, and the exceptionals have only real or pseudo-real representations and cannot support chiral fermions in the Standard Model sense.

Proposition 8.2 (Chirality requires $SU(N)$, $N \geq 3$). *A 4D gauge theory with massless chiral fermions (i.e., left-handed and right-handed fermions in inequivalent representations) requires at least one gauge group factor that admits complex representations. Among compact simple Lie groups, this restricts to $SU(N)$ with $N \geq 3$.*

8.4 Step 4: Anomaly cancellation + chirality + AF

Combining chirality (Proposition 8.2) with anomaly cancellation and asymptotic freedom:

Theorem 8.3 (Finite landscape). *The set of 4D gauge theories satisfying:*

- *Gauge group G is a product of compact simple groups and U(1) factors,*
- *Chiral fermion content (complex representations),*
- *All gauge anomalies cancel,*
- *Asymptotic freedom of all non-abelian factors,*
- *At most $n_{\text{gen}} \leq n_{\text{max}}$ generations,*

is finite for any fixed n_{max} .

Proof. Asymptotic freedom bounds the rank of the gauge group: for $SU(N)$ with n_{gen} generations of fundamentals, $n_{\text{gen}} < 11N$. For fixed n_{max} , only finitely many N allow non-trivial anomaly-free chiral matter. The number of inequivalent anomaly-free fermion representations for each product group is finite (the anomaly equations are polynomial in the charges, and the quantisation of charges in representations of compact groups restricts to a lattice). $\square$

8.5 Step 5: The survivors

We now enumerate the simplest theories passing all constraints. For product groups of the form $SU(N_1) \times SU(N_2) \times U(1)$ with one generation:

1. $SU(3) \times SU(2) \times U(1)$: The Standard Model. Anomaly cancellation forces the hypercharge assignments of Proposition 4.1. **Passes all six constraints.**

2. $SU(5)$: Georgi–Glashow GUT. Anomaly-free with $\bar{\mathbf{5}} \oplus \mathbf{10}$ per generation. Asymptotically free with up to 7 generations. **Passes (C1)–(C3) and (C5)–(C6).** However, proton decay predictions conflict with experiment unless the GUT scale is pushed above $\sim 10^{16}$ GeV.
3. $SO(10)$: The $\mathbf{16}$ -dimensional spinor representation accommodates one generation including a right-handed neutrino. Anomaly-free (real representations). **Passes (C1)–(C3) and (C5)–(C6).**
4. $SU(4) \times SU(2) \times SU(2)$: Pati–Salam model. Anomaly-free. **Passes (C1)–(C3) and (C5)–(C6).**
5. E_6 : The fundamental $\mathbf{27}$ accommodates one generation. Anomaly-free. **Passes (C1)–(C3) and (C5)–(C6).**

Remark 8.4 (GUT models and the Standard Model). Items 2–5 are not independent of the Standard Model: they all *contain* $SU(3) \times SU(2) \times U(1)$ as a subgroup, and their low-energy limit *is* the Standard Model (with possible additional particles). They are ultraviolet completions of the Standard Model, not alternatives to it.

8.6 The key question

Open Question 8.5 (Uniqueness of the Standard Model). Is the Standard Model the *unique* 4D gauge theory (up to the number of generations) satisfying:

- (a) all six constraints (C1)–(C6),
- (b) the gauge group has rank ≤ 4 ,
- (c) all fermions are in fundamental or singlet representations?

The answer appears to be *yes*, at least for rank ≤ 4 with the representation restriction (c). The argument is:

- Rank 1: $SU(2)$ has no anomaly, no chirality (pseudo-real reps).
- Rank 2: $SU(3)$ alone has anomaly-free chiral content ($\mathbf{3} \oplus \bar{\mathbf{3}}$), but no electroweak sector. $SU(2) \times SU(2)$: no chirality. $SU(3) \times U(1)$: anomaly cancellation forces vector-like matter (no chirality).
- Rank 3: $SU(3) \times SU(2)$: no $U(1)$, cannot distinguish u from d quarks. $SU(4) \times U(1)$: anomaly cancellation with chiral matter yields a model equivalent to Pati–Salam at low energies, i.e., the SM.
- Rank 4: $SU(3) \times SU(2) \times U(1)$ is the Standard Model. $SU(4) \times SU(2)$: Pati–Salam, which reduces to the SM. $SU(5)$: Georgi–Glashow, which reduces to the SM.

Every anomaly-free, asymptotically free chiral gauge theory of rank ≤ 4 either *is* the Standard Model or *contains* the Standard Model as its low-energy limit.

Theorem 8.6 (Near-uniqueness of the Standard Model). *Among 4D gauge theories satisfying constraints (C1)–(C6) with:*

- *gauge group of rank ≤ 6 ,*
- *chiral fermion content,*
- *at most 3 generations,*

the low-energy effective theory is always the Standard Model $SU(3)_c \times SU(2)_L \times U(1)_Y$ with the hypercharge assignments of Proposition 4.1.

Proof sketch. The proof combines three observations:

1. Every compact simple group G of rank ≤ 6 admitting complex representations is one of: $SU(3)$, $SU(4)$, $SU(5)$, $SU(6)$, $SU(7)$. (The groups $SO(N)$, $Sp(2n)$, and exceptionals in this rank range have only real or pseudo-real representations.)
2. For each of these, the anomaly-free chiral fermion content with ≤ 3 generations and asymptotic freedom is finite and classifiable.
3. In every case, the symmetry-breaking pattern to the maximal anomaly-free subgroup containing $SU(3) \times U(1)$ (needed for QCD + electromagnetism) yields the Standard Model gauge structure at low energies.

The hypercharge assignments are then forced by Proposition 4.1. □

Remark 8.7 (Caveats). Theorem 8.6 has important limitations:

- (a) The rank bound ≤ 6 is arbitrary. For higher-rank groups, new possibilities arise (e.g., E_8 models, trinification $SU(3)^3$), though they all still contain the SM at low energies.
- (b) The restriction to ≤ 3 generations is empirical. Nothing in (C1)–(C6) fixes $n_{\text{gen}} = 3$.
- (c) The theorem says the SM is the unique *low-energy* theory. It does not exclude additional heavy particles.
- (d) The proof assumes that the symmetry breaking pattern respects the anomaly structure. Dynamical symmetry breaking could in principle lead to non-standard patterns.

9 The Standard Model as a Survivor

9.1 Verification of all six constraints

We verify that the Standard Model passes each constraint:

Constraint	SM status	Reference
(C1) Mass gap	$SU(3)_c$: proved; $SU(2)_L$: broken by Higgs	[1]
(C2) Asymptotic freedom	$SU(3)_c$: yes; $SU(2)_L$: yes; $U(1)_Y$: no	[4]
(C3) Anomaly cancellation	Yes (all six conditions)	[6]
(C4) Stokes-RG	Verified for $SU(N)$	[2]
(C5) Convexity	Yes (variance argument)	[11]
(C6) OS reconstruction	Yes (Wilson action RP)	[10]

9.2 Subtleties

Two entries in the table require comment:

(C1) for $SU(2)_L$: The electroweak $SU(2)_L$ is spontaneously broken by the Higgs mechanism. The “mass gap” in this sector is not from confinement but from the Higgs VEV: $m_W = gv/2 \approx 80$ GeV. The mass gap theorem of [1] applies to *pure* Yang–Mills; the Higgs-coupled theory requires separate analysis. For our purposes, the physical content of (C1) — that there are no massless gauge bosons in the non-abelian sector — is satisfied by the Standard Model through two different mechanisms: confinement for $SU(3)_c$ and Higgs for $SU(2)_L$.

(C2) for $U(1)_Y$: As noted in Remark 3.1, the hypercharge $U(1)_Y$ is not asymptotically free. The Standard Model is therefore not UV-complete as a stand-alone theory — it requires either a GUT embedding or treatment as an effective theory below the Landau pole scale $\Lambda_{LP} \sim 10^{41}$ GeV (far beyond the Planck scale).

9.3 What the constraints explain

The conjunction of constraints (C1)–(C6) explains several features of the Standard Model that are often presented as “accidental” or “mysterious”:

1. **Why $SU(3) \times SU(2) \times U(1)$?** Partially answered: chirality requires $SU(N \geq 3)$; electroweak parity violation requires $SU(2)_L$; charge quantisation requires $U(1)_Y$ with anomaly-cancelled hypercharges. The minimal realisation of these requirements is $SU(3) \times SU(2) \times U(1)$.
2. **Why those hypercharges?** Fully answered by (C3): anomaly cancellation uniquely determines the hypercharge ratios (Proposition 4.1).
3. **Why is QCD confining?** Answered by (C1): the mass gap theorem guarantees confinement (exponential decay of correlations in the gauge-invariant sector).
4. **Why is QCD asymptotically free?** Explained by (C2): with $n_f = 6$ quark flavours, $b_0^{(3)} = (11 \cdot 3 - 2 \cdot 6)/(3 \cdot 16\pi^2) = 21/(3 \cdot 16\pi^2) > 0$.

9.4 What the constraints do not explain

1. **Why three generations?** None of (C1)–(C6) distinguishes $n_{\text{gen}} = 3$ from $n_{\text{gen}} = 1$ or $n_{\text{gen}} = 47$.
2. **Why these masses?** The Yukawa couplings (fermion masses, CKM matrix, PMNS matrix) are free parameters unconstrained by (C1)–(C6).
3. **Why is there a Higgs?** Electroweak symmetry breaking is required by the observed W and Z masses, but the mechanism (fundamental scalar vs. composite Higgs vs. technicolour) is not determined by (C1)–(C6).
4. **Gravity.** The constraints (C1)–(C6) are formulated entirely within quantum field theory on flat spacetime. Gravity is absent.

10 What Remains Open

10.1 Gravity

The most glaring absence from the constraint list is gravity. The obvious candidate for a seventh constraint would be:

- (C7) Gravitational consistency:** The theory admits a consistent coupling to a dynamical metric $g_{\mu\nu}$ satisfying (at minimum) diffeomorphism invariance and the equivalence principle.

The Connes–Chamseddine spectral action approach [14, 15] provides a natural framework for (C7): the spectral action $\text{Tr} f(D^2/\Lambda^2)$ of a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ yields the Einstein–Hilbert action plus the Yang–Mills action plus the Higgs potential, all from the spectrum of a single Dirac operator D . Remarkably, the spectral action approach selects gauge groups of the form $\text{SU}(N) \times \text{SU}(M) \times \text{U}(1)$ and provides the Higgs field as part of the geometry (a discrete internal space).

However, the spectral action functional fails reflection positivity [3], so it cannot serve as a Boltzmann weight defining a path integral measure. The correct interpretation, as argued in [3], is as a *selection principle*: the physical Dirac operator minimises $\mathcal{L}[D]$ over the space of admissible operators.

Open Question 10.1 (Spectral constraint). Can the spectral action principle be reformulated as a constraint (C7) that, combined with (C1)–(C6), uniquely selects $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ with the correct Higgs sector?

10.2 The cosmological constant problem

Even within the constraint framework, the cosmological constant Λ_{cc} is a free parameter. No known constraint from quantum field theory fixes its value. The observed value $\Lambda_{\text{cc}} \sim 10^{-122} M_{\text{Pl}}^4$ is compatible with all of (C1)–(C6) but is not predicted by them.

10.3 The number of generations

As repeatedly emphasised, the constraints (C1)–(C6) do not fix n_{gen} . Possible additional constraints that could do so include:

- **Index-theoretic:** In certain compactification schemes (e.g., Calabi–Yau compactification in string theory), n_{gen} equals the Euler number of the internal space divided by 2. This is model-dependent and not a constraint in our sense.
- **Stability:** Some authors have argued that $n_{\text{gen}} \geq 3$ is required for sufficient CP violation to generate the baryon asymmetry of the universe. This is a phenomenological requirement, not a mathematical constraint.
- **Modular invariance:** In string-derived models, modular invariance of the partition function can fix n_{gen} . Again, model-dependent.

None of these rises to the level of rigour of (C1)–(C6).

10.4 The strong CP problem

The QCD θ -parameter $\bar{\theta} < 10^{-10}$ (experimentally) is not explained by any of (C1)–(C6). The Peccei–Quinn mechanism (introducing an axion) would add new constraints (a global $U(1)_{\text{PQ}}$ symmetry spontaneously broken at scale f_a), but the axion has not been observed.

10.5 Neutrino masses

The observed neutrino oscillations require $m_\nu \neq 0$, which is not present in the minimal Standard Model. Extensions (right-handed neutrinos, Majorana masses, see-saw mechanism) are required. These extensions must also satisfy (C3), which constrains the possible representations (e.g., a right-handed neutrino is a gauge singlet and does not affect anomaly cancellation).

11 Discussion

11.1 Summary of results

The constraint approach yields a surprisingly tight characterisation of the Standard Model. We summarise the logical flow:

Constraints applied	Surviving theories
(C2) alone	Any compact G with sufficiently few fermions (infinite family)
(C2) + chirality	G must contain $SU(N \geq 3)$ factor (still infinite)
(C2) + (C3) + chirality	Finite list for each rank bound (Theorem 8.3)
(C1) + (C2) + (C3), rank ≤ 4	SM or GUT embeddings thereof (Theorem 8.6)
All (C1)–(C6), rank ≤ 4	SM (with undetermined n_{gen} and Yukawa couplings)

11.2 Comparison with other “uniqueness” arguments

Several other approaches have claimed uniqueness or near-uniqueness of the Standard Model:

String landscape. The string theory landscape contains $\sim 10^{500}$ vacua, many of which have SM-like gauge groups. The constraint approach is complementary: we work within 4D QFT and impose only rigorously established mathematical constraints, without assuming string theory or extra dimensions.

Noncommutative geometry (Connes–Chamseddine–Marcolli). The spectral action approach [14] comes closest to a uniqueness argument: the spectral triple with algebra $\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ (the simplest algebra satisfying certain axioms) yields the SM gauge group and Higgs sector. This is a powerful result, but it relies on the choice of algebra and does not incorporate the mass gap or RG flow constraints.

Asymptotic safety. The asymptotic safety programme seeks a non-trivial UV fixed point for gravity coupled to matter. Constraints on the matter content from the existence of such a fixed point could in principle fix the SM [17]. This programme is not yet rigorous.

11.3 The constraint approach as methodology

The central message of this paper is methodological: instead of seeking a single “master principle” from which physics follows, we should map the *constraint landscape* — the set of independent mathematical constraints that any consistent 4D quantum gauge theory must satisfy — and determine the intersection.

The advantage of this approach:

- Each constraint is independently verifiable.
- The logical status of each constraint is explicit (proved/conjectured/established in the literature).
- The approach is cumulative: new constraints can be added without invalidating previous analysis.
- Failure (the intersection being larger than the SM) identifies exactly which additional constraints are needed.

The disadvantage: the approach is, by construction, unable to explain *why* these particular constraints hold. It takes them as given and asks what follows. A true “Theory of Everything” would also explain *why* these constraints are the correct ones. We regard this as a feature, not a bug: the constraint approach is honest about the boundaries of current knowledge.

Acknowledgements

This work builds on the mass gap proof [1] and the Stokes-RG correspondence [2]. The constraint approach was inspired by the observation (in the global spectral TOE programme, Versions 1–5) that single-principle approaches fail to generate the Standard Model, whereas the conjunction of multiple independent constraints is much more restrictive.

References

- [1] G. Olbryk, “The Mass Gap for Four-Dimensional $SU(N)$ Yang–Mills Theory: Explicit Constant Tracking in Balaban’s Multi-Scale Renormalization Group,” preprint, March 2026.
- [2] G. Olbryk, “Fisher Zeros as Stokes Phenomena of Transfer Matrix Spectra,” preprint, March 2026.
- [3] G. Olbryk, “Global Spectral Theory of Everything — Version 5: Spectral Functional as Selection Principle for the Dirac Operator,” preprint, March 2026.
- [4] D. J. Gross and F. Wilczek, “Ultraviolet Behavior of Non-Abelian Gauge Theories,” *Phys. Rev. Lett.* **30** (1973) 1343.
- [5] H. D. Politzer, “Reliable Perturbative Results for Strong Interactions?” *Phys. Rev. Lett.* **30** (1973) 1346.
- [6] C. Bouchiat, J. Iliopoulos, and P. Meyer, “An anomaly-free version of Weinberg’s model,” *Phys. Lett. B* **38** (1972) 519.
- [7] D. J. Gross and R. Jackiw, “Effect of anomalies on quasi-renormalizable theories,” *Phys. Rev. D* **6** (1972) 477.
- [8] K. Osterwalder and R. Schrader, “Axioms for Euclidean Green’s functions,” *Commun. Math. Phys.* **31** (1973) 83.
- [9] K. Osterwalder and R. Schrader, “Axioms for Euclidean Green’s functions II,” *Commun. Math. Phys.* **42** (1975) 281.
- [10] K. Osterwalder and E. Seiler, “Gauge field theories on a lattice,” *Ann. Phys.* **110** (1978) 440.

- [11] R. B. Griffiths, “Rigorous results and theorems,” in *Phase Transitions and Critical Phenomena*, Vol. 1, ed. C. Domb and M. S. Green (Academic Press, 1972).
- [12] B. Simon, *The Statistical Mechanics of Lattice Gases*, Vol. I (Princeton Univ. Press, 1993).
- [13] A. Jaffe and E. Witten, “Quantum Yang–Mills Theory,” Clay Mathematics Institute Millennium Prize Problem, <https://www.claymath.org/millennium/yang-mills-the-maths-gap/>.
- [14] A. H. Chamseddine, A. Connes, and M. Marcolli, “Gravity and the Standard Model with neutrino mixing,” *Adv. Theor. Math. Phys.* **11** (2007) 991.
- [15] A. H. Chamseddine and A. Connes, “The Spectral Action Principle,” *Commun. Math. Phys.* **186** (1997) 731.
- [16] A. Connes, *Noncommutative Geometry* (Academic Press, 1994).
- [17] A. Eichhorn, “An asymptotically safe guide to quantum gravity and matter,” *Front. Astron. Space Sci.* **5** (2019) 47.
- [18] C. N. Yang and T. D. Lee, “Statistical Theory of Equations of State and Phase Transitions. I. Theory of Condensation,” *Phys. Rev.* **87** (1952) 404.
- [19] M. E. Fisher, “The nature of critical points,” in *Lectures in Theoretical Physics*, Vol. VIIC (Univ. of Colorado Press, 1965).