Micro: Problem Set 1.

Victor H. Aguiar

This version: 2024

Exercise 1. Consider a choice set X, and a consumer with preference defined over X, $\succeq \subseteq X \times X$, such that the preferences are a preorder (but not necessarily complete).

- Assume in this literal that X is finite, prove that the preferences $\succeq \subseteq X \times X$ defined above, in general, cannot be represented by a utility function $u: X \mapsto \mathbb{R}$. Formally, find a counterexample to the statement: There exists a $u: X \mapsto \mathbb{R}$ such that for any $a, b \in X, a \succeq b \iff u(a) \geq u(b)$.
- Assume in this literal that X is finite, prove that the preferences $\succeq \subseteq X \times X$ defined above, have a multiple utility representation, i.e., there is a set of utility functions \mathcal{U} such that for any $a, b \in X$ with $a \succeq b \iff u(a) \geq u(b)$ for all $u \in \mathcal{U}$. (Hint: the set of utilities \mathcal{U} can be finite. Also you could use the idea of a binary relation closure).
- c) Maintain the assumptions in b). Let $C^{\succeq}(A) = \{a \in A | \text{there is no } b \in A, b \succeq a\}$. Consider the data set $\{C^{\succeq}(A)\}_{A \in \mathcal{A}}$, where $\mathcal{A} \equiv 2^X \setminus \emptyset$, show (by means of a counterexample) that the data set may fail GARP.
- d) Maintain the assumptions in b), and consider the data set $\{C(A)\}_{A\in\mathcal{A}}$, where $\mathcal{A}\equiv 2^X\setminus\emptyset$. Prove that the following axiom is a necessary condition for the data set $\{C(A)\}_{A\in\mathcal{A}}$ to be generated by the preferences in b) (i.e., $C(A)\equiv\{a\in A|\text{there is no }b\in A,b\succ a\}\}$.
- Weak Axiom of Revealed Non-Inferiority (WARNI). For any $A \in \mathcal{A}$ and $y \in A$, if for every $x \in C(A)$ there exists a $B \in \mathcal{A}$ with $y \in C(B)$ and $x \in B$, then $y \in C(A)$.