The Call Option and The Black-Scholes Formula

Anton Yang

May 4, 2023

Suppose I tell you that I will give you \$10 tomorrow if it rains, and if it doesn't, nothing happens.

Suppose I tell you that I will give you \$10 tomorrow if it rains, and if it doesn't, nothing happens.

Would you accept the offer?



You would definitely take the offer because there's no risk to you.

You would definitely take the offer because there's no risk to you.

So what actually is a call option?


A call option is a financial derivative where the trader buys a contract to have the right, not the obligation, to purchase a share of stock at the strike price.

A call option is a financial derivative where the trader buys a contract to have the right, not the obligation, to purchase a share of stock at the strike price.

Ex: Suppose you decided to buy a call option at the strike price of \$165 with a premium price of \$5, and Tesla's current stock price is \$160. If the next day, Tesla's stock price increases to \$180. You can exercise the option and get a payoff of \$15 and a profit of \$10 per share.

Payoff = $\max[(S_t-K),0]$

Profit=max[$(S_t$ -K),0]-C



$$C(S_0, K, \sigma, r, T, \delta) = \frac{1}{\sqrt{2\pi}} \left(Se^{-\delta T} \int_{-\infty}^{\frac{\ln \frac{S}{K} + (r-\delta + \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}} e^{-\frac{1}{2}x^2} dx - Ke^{-rT} \int_{-\infty}^{\frac{\ln \frac{S}{K} + (r-\delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}} e^{-\frac{1}{2}x^2} dx \right)$$

 $S_0 = \text{Stock Price}$

K =Strike Price

 $\sigma = Volatility$

r =Interest Rate

T = Time to Expiree (in years)

 δ = Dividend Yield



$$C(S_0, K, \sigma, r, T, \delta) = Se^{-\delta T}N(d_1) - Ke^{-rT}N(d_2)$$

$$C(S_0, K, \sigma, r, T, \delta) = Se^{-\delta T}N(d_1) - Ke^{-rT}N(d_2)$$

$$d_{1,2} = \frac{\ln S - \ln K + (r - \delta \pm \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}$$

$$C(S_0, K, \sigma, r, T, \delta) = Se^{-\delta T}N(d_1) - Ke^{-rT}N(d_2)$$

$$d_{1,2} = \frac{\ln S - \ln K + (r - \delta \pm \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}$$

$$N(d) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{d} e^{-\frac{1}{2}x^{2}} dx$$



$$C(S, K, \sigma, r, T, \delta) = e^{-rT} [\mathbb{E}(S_T - K | S_T > K)] P(S_T > K)$$

$$C(S, K, \sigma, r, T, \delta) = e^{-rT} [\mathbb{E}(S_T - K | S_T > K)] P(S_T > K)$$

$$= e^{-rT} \mathbb{E}(S_T | S_T > K) P(S_T > K) - e^{-rT} \mathbb{E}(K | S_T > K) P(S_T > K)$$

$$C(S, K, \sigma, r, T, \delta) = e^{-rT} [\mathbb{E}(S_T - K | S_T > K)] P(S_T > K)$$

$$= e^{-rT} \mathbb{E}(S_T | S_T > K) P(S_T > K) - e^{-rT} \mathbb{E}(K | S_T > K) P(S_T > K)$$

$$= e^{-rT} \mathbb{E}(S_T | S_T > K) P(S_T > K) - Ke^{-rT} P(S_T > K)$$



Assume that continuously compounded returns, on the stock underlying the call option, from t=0 to t=T, are log-normal distributed with a mean and a variance $((r - \delta - \frac{1}{2}\sigma^2)T, \sigma^2T)$

Assume that continuously compounded returns, on the stock underlying the call option, from t=0 to t=T, are log-normal distributed with a mean and a variance $((r - \delta - \frac{1}{2}\sigma^2)T, \sigma^2T)$

$$\ln{(\frac{S_T}{S})} \sim \mathcal{N}((r - \delta - \frac{1}{2}\sigma^2)T, \sigma^2T)$$



Apply the identity $\ln \left(\frac{a}{b} \right) = \ln a - \ln b$ and shift the mean:

$$\ln S_T - \ln S \sim \mathcal{N}((r - \delta - \frac{1}{2}\sigma^2)T, \sigma^2 T)$$

Apply the identity $\ln \left(\frac{a}{b} \right) = \ln a - \ln b$ and shift the mean:

$$\ln S_T - \ln S \sim \mathcal{N}((r - \delta - \frac{1}{2}\sigma^2)T, \sigma^2 T)$$

$$\ln S_T \sim \mathcal{N}(\ln S + (r - \delta - \frac{1}{2}\sigma^2)T, \sigma^2 T)$$



$$Z = \frac{X - \mu}{\sigma}$$

$$Z = \frac{X - \mu}{\sigma}$$

$$=\frac{\ln S_T - (\ln S + (r-\delta - \frac{1}{2}\sigma^2))T}{\sigma\sqrt{T}}$$

$$Z = \frac{X - \mu}{\sigma}$$

$$=\frac{\ln S_T - (\ln S + (r-\delta - \frac{1}{2}\sigma^2))T}{\sigma\sqrt{T}}$$

$$=\frac{\ln S_T - \ln S - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}$$



If $X_1 < X_2$, then $\ln X_1 < \ln X_2$, so $P(S_T < K) = P(\ln S_T < K)$. Subtract $\ln S + (r - \delta - \frac{1}{2}\sigma^2)T$ from both sides and dividing both sides by $\sigma\sqrt{T}$:

If $X_1 < X_2$, then $\ln X_1 < \ln X_2$, so $\mathrm{P}(S_T < K) = \mathrm{P}(\ln S_T < K)$. Subtract $\ln S + (r - \delta - \frac{1}{2}\sigma^2)T$ from both sides and dividing both sides by $\sigma\sqrt{T}$:

$$P\left(\frac{\ln S_T - \ln S - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}} < \frac{\ln K - \ln S - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}\right)$$

If $X_1 < X_2$, then $\ln X_1 < \ln X_2$, so $P(S_T < K) = P(\ln S_T < K)$. Subtract $\ln S + (r - \delta - \frac{1}{2}\sigma^2)T$ from both sides and dividing both sides by $\sigma\sqrt{T}$:

$$P\left(\frac{\ln S_T - \ln S - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}} < \frac{\ln K - \ln S - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}\right)$$

The left side of the inequality is Z. The right side:



If $X_1 < X_2$, then $\ln X_1 < \ln X_2$, so $\mathrm{P}(S_T < K) = \mathrm{P}(\ln S_T < K)$. Subtract $\ln S + (r - \delta - \frac{1}{2}\sigma^2)T$ from both sides and dividing both sides by $\sigma\sqrt{T}$:

$$P\left(\frac{\ln S_T - \ln S - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}} < \frac{\ln K - \ln S - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}\right)$$

The left side of the inequality is Z. The right side:

$$-d_2 = -\frac{\ln S - \ln K + (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma\sqrt{T}}$$



$$P(S_T > K) = 1 - P(S_T < K)$$

$$P(S_T > K) = 1 - P(S_T < K)$$

= 1 - N(-d₂)

$$P(S_T > K) = 1 - P(S_T < K)$$

= 1 - N(-d₂)
= N(-(-d₂))

$$P(S_T > K) = 1 - P(S_T < K)$$

$$= 1 - N(-d_2)$$

$$= N(-(-d_2))$$

$$= N(d_2)$$

Present Value of Expected Call Payoff

 $C(S, K, \sigma, r, T, \delta)$

Present Value of Expected Call Payoff

$$C(S, K, \sigma, r, T, \delta)$$

$$= e^{-rT} \mathbb{E}(S_T|S_T > K) P(S_T > K) - Ke^{-rT} P(S_T > K)$$

Present Value of Expected Call Payoff

$$C(S, K, \sigma, r, T, \delta)$$

$$= e^{-rT} \mathbb{E}(S_T|S_T > K) P(S_T > K) - Ke^{-rT} P(S_T > K)$$

and we already proved that $P(S_T > K) = N(d_2)$



Suppose $X \sim \mathcal{N}(\mu, \sigma^2)$ is a normally distributed random variable.

Suppose $X \sim \mathcal{N}(\mu, \sigma^2)$ is a normally distributed random variable.

$$\mathbb{E}(X) = \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} x e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2} dx$$

Suppose $X \sim \mathcal{N}(\mu, \sigma^2)$ is a normally distributed random variable.

$$\mathbb{E}(X) = \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} x e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2} dx$$

Then we can calculate the expected value over a limited range of values:

Suppose $X \sim \mathcal{N}(\mu, \sigma^2)$ is a normally distributed random variable.

$$\mathbb{E}(X) = \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} x e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2} dx$$

Then we can calculate the expected value over a limited range of values:

$$\mathbb{E}_{x>d}(X) = \frac{1}{\sigma\sqrt{2\pi}} \int_{d}^{\infty} x e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2}$$



Let S be a log-normal distributed random variable with $\ln S \sim \mathcal{N}(m, v^2)$, and let K be a positive number:

Let S be a log-normal distributed random variable with $\ln S \sim \mathcal{N}(m, v^2)$, and let K be a positive number:

$$C = \begin{cases} S & \text{for } S > K \\ 0 & \text{otherwise} \end{cases}$$

Let S be a log-normal distributed random variable with $\ln S \sim \mathcal{N}(m, v^2)$, and let K be a positive number:

$$C = \begin{cases} S & \text{for } S > K \\ 0 & \text{otherwise} \end{cases}$$

Then:

$$\mathbb{E}_{S>K}(C) = e^{m + \frac{1}{2}v^2} N\left(v - \frac{\ln K - m}{v}\right)$$



Calculating the partial expectation of ln *S*:

Calculating the partial expectation of ln *S*:

$$\mathbb{E}_{S>K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\ln K}^{\infty} Se^{-\frac{1}{2}(\frac{\ln S - m}{v})^2} d\ln S$$

Calculating the partial expectation of ln *S*:

$$\mathbb{E}_{S>K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\ln K}^{\infty} Se^{-\frac{1}{2}(\frac{\ln S - m}{v})^2} d\ln S$$

Let $x = \ln S$ and $S = e^x$:

Calculating the partial expectation of ln *S*:

$$\mathbb{E}_{S>K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\ln K}^{\infty} Se^{-\frac{1}{2}(\frac{\ln S - m}{v})^2} d\ln S$$

Let $x = \ln S$ and $S = e^x$:

$$\mathbb{E}_{S > K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\ln K}^{\infty} e^{x} e^{-\frac{1}{2}(\frac{x-m}{v})^{2}} dx$$

Calculating the partial expectation of ln *S*:

$$\mathbb{E}_{S>K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\ln K}^{\infty} Se^{-\frac{1}{2}(\frac{\ln S - m}{v})^2} d\ln S$$

Let $x = \ln S$ and $S = e^x$:

$$\mathbb{E}_{S > K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\ln K}^{\infty} e^{x} e^{-\frac{1}{2}(\frac{x-m}{v})^{2}} dx$$

Then let
$$z = \frac{x-m}{v}$$
 with $dz = \frac{1}{v}dx$:

Then let
$$z = \frac{x-m}{v}$$
 with $dz = \frac{1}{v}dx$:

$$\mathbb{E}_{S>K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\frac{\ln K - m}{v}}^{\infty} e^{m + vz} e^{-\frac{1}{2}z^2} dz$$

Then let
$$z = \frac{x-m}{v}$$
 with $dz = \frac{1}{v}dx$:
$$\mathbb{E}_{S>K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\frac{\ln K - m}{v}}^{\infty} e^{m+vz} e^{-\frac{1}{2}z^2} dz$$

$$= \frac{e^{m + \frac{1}{2}v^2}}{\sqrt{2\pi}} \int_{\frac{\ln K - m}{v}}^{\infty} e^{-\frac{1}{2}z^2 + vz - \frac{1}{2}v^2} dz$$

Then let
$$z = \frac{x-m}{v}$$
 with $dz = \frac{1}{v}dx$:

$$\mathbb{E}_{S>K}(C) = \frac{1}{v\sqrt{2\pi}} \int_{\frac{\ln K - m}{v}}^{\infty} e^{m + vz} e^{-\frac{1}{2}z^2} dz$$

$$= \frac{e^{m + \frac{1}{2}v^2}}{\sqrt{2\pi}} \int_{\frac{\ln K - m}{v}}^{\infty} e^{-\frac{1}{2}z^2 + vz - \frac{1}{2}v^2} dz$$

$$= \frac{e^{m + \frac{1}{2}v^2}}{\sqrt{2\pi}} \int_{\frac{\ln K - m}{v}}^{\infty} e^{-\frac{1}{2}(z - v)^2} dz$$



$$\mathbb{E}_{S > K}(C) = \frac{e^{m + \frac{1}{2}v^2}}{\sqrt{2\pi}} \int_{\frac{\ln K - m}{v} - v}^{\infty} e^{-\frac{1}{2}u^2} du$$

$$\mathbb{E}_{S>K}(C) = \frac{e^{m+\frac{1}{2}v^2}}{\sqrt{2\pi}} \int_{\frac{\ln K - m}{v} - v}^{\infty} e^{-\frac{1}{2}u^2} du$$

$$= e^{m + \frac{1}{2}v^2} \left[1 - N(\frac{\ln K - m}{v} - v) \right]$$

$$\mathbb{E}_{S>K}(C) = \frac{e^{m + \frac{1}{2}v^2}}{\sqrt{2\pi}} \int_{\frac{\ln K - m}{v} - v}^{\infty} e^{-\frac{1}{2}u^2} du$$

$$= e^{m + \frac{1}{2}v^2} \left[1 - N(\frac{\ln K - m}{v} - v) \right]$$

$$= e^{m + \frac{1}{2}v^2} N\left(v - \frac{\ln K - m}{v}\right)$$

$$e^{m+\frac{1}{2}v^2}$$

$$e^{m+\frac{1}{2}v^2} = e^{\ln S}e^{(r-\delta)T}e^{-\frac{1}{2}\sigma^2T}e^{\frac{1}{2}\sigma^2T}$$

$$e^{m+\frac{1}{2}v^2} = e^{\ln S}e^{(r-\delta)T}e^{-\frac{1}{2}\sigma^2T}e^{\frac{1}{2}\sigma^2T}$$

$$= Se^{(r-\delta)T}$$

$$e^{m+\frac{1}{2}v^2} = e^{\ln S}e^{(r-\delta)T}e^{-\frac{1}{2}\sigma^2T}e^{\frac{1}{2}\sigma^2T}$$

$$= Se^{(r-\delta)T}$$

$$N\left(v - \frac{lnK - m}{v}\right)$$

$$e^{m+\frac{1}{2}v^2} = e^{\ln S}e^{(r-\delta)T}e^{-\frac{1}{2}\sigma^2T}e^{\frac{1}{2}\sigma^2T}$$

$$= Se^{(r-\delta)T}$$

$$N\left(v-\frac{lnK-m}{v}\right)$$

$$= N \left(\sigma \sqrt{T} - \frac{lnK - lnS_T - (r - \delta - \frac{1}{2}\sigma^2)T}{\sigma \sqrt{T}} \right)$$



$$= N(\sigma\sqrt{T} - (-d_2))$$

$$= N(\sigma\sqrt{T} - (-d_2))$$
$$= N(d_1)$$

$$= N(\sigma\sqrt{T} - (-d_2))$$
$$= N(d_1)$$

Combine them:

$$= N(\sigma\sqrt{T} - (-d_2))$$
$$= N(d_1)$$

Combine them:

$$\mathbb{E}_{S_T > K}(S_T) = Se^{(r-\delta)T}N(d_1)$$

Let *X* be a discrete random variable representing the outcome of a single roll of a six-sided die:

Let *X* be a discrete random variable representing the outcome of a single roll of a six-sided die:

$$\mathbb{E} = \frac{1}{6}(1) + \frac{1}{6}(2) + \frac{1}{6}(3) + \frac{1}{6}(4) + \frac{1}{6}(5) + \frac{1}{6}(6) = \frac{21}{6} = 3.5$$

Let *X* be a discrete random variable representing the outcome of a single roll of a six-sided die:

$$\mathbb{E} = \frac{1}{6}(1) + \frac{1}{6}(2) + \frac{1}{6}(3) + \frac{1}{6}(4) + \frac{1}{6}(5) + \frac{1}{6}(6) = \frac{21}{6} = 3.5$$

 $\mathbb{E}(X|X\geqslant 4)=5$. However, if we calculate the partial expectation of $\mathbb{E}_{X\geqslant 4}(X)$:



Let *X* be a discrete random variable representing the outcome of a single roll of a six-sided die:

$$\mathbb{E} = \frac{1}{6}(1) + \frac{1}{6}(2) + \frac{1}{6}(3) + \frac{1}{6}(4) + \frac{1}{6}(5) + \frac{1}{6}(6) = \frac{21}{6} = 3.5$$

 $\mathbb{E}(X|X\geqslant 4)=5$. However, if we calculate the partial expectation of $\mathbb{E}_{X\geqslant 4}(X)$:

$$\mathbb{E}_{X \geqslant 4}(X) = \frac{1}{6}(4) + \frac{1}{6}(5) + \frac{1}{6}(6) = \frac{15}{6} = 2.5$$



Therefore to get the $\mathbb{E}(X|X \ge 4) = 5$, we have to divide the expected value by the probability:

Therefore to get the $\mathbb{E}(X|X \ge 4) = 5$, we have to divide the expected value by the probability:

$$\mathbb{E}(X|X \ge 4) = \frac{\mathbb{E}_{X \ge 4}(X)}{P(X \ge 4)} = \frac{2.5}{0.5} = 5$$

Therefore to get the $\mathbb{E}(X|X \ge 4) = 5$, we have to divide the expected value by the probability:

$$\mathbb{E}(X|X \ge 4) = \frac{\mathbb{E}_{X \ge 4}(X)}{P(X \ge 4)} = \frac{2.5}{0.5} = 5$$

So for The Black-Scholes formula:

Therefore to get the $\mathbb{E}(X|X \ge 4) = 5$, we have to divide the expected value by the probability:

$$\mathbb{E}(X|X \ge 4) = \frac{\mathbb{E}_{X \ge 4}(X)}{P(X \ge 4)} = \frac{2.5}{0.5} = 5$$

So for The Black-Scholes formula:

$$\mathbb{E}(S_T|S_T > K) = \frac{\mathbb{E}_{S_T > K}(S_T)}{P(S_T > K)}$$



$$C(S, K, \sigma, r, T, \delta) =$$

$$C(S, K, \sigma, r, T, \delta) =$$

$$e^{-rT} \mathbb{E}(S_T | S_T > K) P(S_T > K) - K e^{-rT} P(S_T > K)$$

$$C(S, K, \sigma, r, T, \delta) =$$

$$e^{-rT} \mathbb{E}(S_T | S_T > K) P(S_T > K) - K e^{-rT} P(S_T > K)$$

$$= e^{-rT} \frac{\mathbb{E}_{S_T > K}(S_T)}{P(S_T > K)} P(S_T > K) - K e^{-rT} P(S_T > K)$$

$$C(S, K, \sigma, r, T, \delta) =$$

$$e^{-rT} \mathbb{E}(S_T | S_T > K) P(S_T > K) - Ke^{-rT} P(S_T > K)$$

$$= e^{-rT} \frac{\mathbb{E}_{S_T > K}(S_T)}{P(S_T > K)} P(S_T > K) - Ke^{-rT} P(S_T > K)$$

$$= e^{-rT} Se^{(r-\delta)T} N(d_1) - Ke^{-rT} N(d_2)$$

$$C(S, K, \sigma, r, T, \delta) =$$

$$e^{-rT} \mathbb{E}(S_T | S_T > K) P(S_T > K) - K e^{-rT} P(S_T > K)$$

$$= e^{-rT} \frac{\mathbb{E}_{S_T > K}(S_T)}{P(S_T > K)} P(S_T > K) - K e^{-rT} P(S_T > K)$$

$$= e^{-rT} S e^{(r-\delta)T} N(d_1) - K e^{-rT} N(d_2)$$

$$= S e^{-\delta T} N(d_1) - K e^{-rT} N(d_2)$$