

Notes on Measurement of Brownian Motion for Two SmA Islands

Joe Becker

February 27, 2013

1 2 Island Interaction

We measured the relative Brownian motion of a Smectic A island interacting with another Smectic A island on a freely suspended film of 8CB liquid crystal. To achieve this we measure the relative motion of the islands parallel to the separation, d , and perpendicular to the separation, τ . The geometry is shown in figure 1.

Note this operates under the assumption that the island A's movement is not large enough to affect the direction of d and τ .

Note that the data collection is done through video of island motion through a $40\times$ objective. The video is decomposed into individual frames which allows the islands to be tracked. The tracking is done by an IDL algorithm written by Joe MacLennan which fits an ellipse to the islands and outputs the islands' center coordinates and the major and minor axes. A single tracked frame is shown in figure 2.

From this tracking data we convert to the $d - \tau$ basis through the calculations

$$d_n = d_{n-1} + (x_n - x_{n-1}) \cos(\theta_{n-1}) + (y_n - y_{n-1}) \sin(\theta_{n-1}) \quad (1)$$

$$\tau_n = \tau_{n-1} + (y_n - y_{n-1}) \cos(\theta_{n-1}) - (x_n - x_{n-1}) \sin(\theta_{n-1}) \quad (2)$$

where θ_n is the angle the separation line makes with the x axis. It is calculated through the arctangent of the rectangular coordinates. We then analyze the tracked data by calculating the time step change of d and τ over 8 to 12 time steps. This allows us to calculate the average change $\overline{\Delta d}_m$ and $\overline{\Delta \tau}_m$ for each timestep m . We then can find the variance from these means, σ_m^τ and σ_m^d . We obtain the diffusion constant by using the fact that the variance is proportional to elapsed time where the constant of proportionality is the diffusion constant. An example plot is shown in figure

3




Figure 1: *The geometry of d and τ .*




Figure 2: An example tracking frame. The red ring is the ellipse fitted to the island.




Figure 3: An example plot of σ^d and σ^τ versus time. Note that the plot labeled x corresponds to d and the plot labeled y corresponds to τ .




Figure 4: Predicted and measured reduced mobility versus the ratio of separation and island radius for island island interaction. Note that this is in the direction along the island separation.

From here we wish to compare our measured results with the results generated by the model created by Tatiana. To do this we must convert our diffusion constant to a reduced mobility. This is done through the *Einstein Relation*

$$D_T = k_B T b_T \quad (3)$$

where b is the mobility. We then find the reduced mobility by dividing out μ_0 which is given by

$$\mu_0 = \frac{1}{4\pi\eta h}$$

where η is the liquid crystal viscosity and h is the film thickness. Note that the mobility is related to the reduced mobility, μ_T , by

$$b_T = \mu_0 \mu_T$$

So we calculate our reduced mobility by

$$\mu_T = \frac{D_T}{\mu_0 k_B T} = \frac{4\pi\eta h}{k_B T} D_T$$

. Finally, we must account for the confinement which is not accounted for in the model. To do this we use Saffman's theory to get the reduced mobility due to the confinement effect as

$$\mu_{\text{conf}} = \ln \left(\frac{R}{a} - \frac{1}{2} \right)$$

where R is the radius of the film and a is the radius of the island. We then correct the model's prediction by

$$\mu_{\text{corr}} = \frac{\mu_{\text{pred}} \mu_{\text{conf}}}{\mu_{\text{pred}} + \mu_{\text{conf}}}$$

Figures 4, 5, and 6 show the comparison between the model and measured results.




Figure 5: *Predicted and measured reduced mobility versus the ratio of separation and island radius for island island interaction. Note that this is in the direction perpendicular to the island separation.*



Figure 6: *Ratio of the reduced mobilities for both along and perpendicular to the island separation.*




Figure 7: *Predicted and measured reduced mobility versus the ratio of separation and island radius for island wall interactions. Note that this is in the direction along the island separation.*

2 Island Wall Interaction

A study of mobilities of smecticA islands created in 8CB was conducted by a now graduated group member, Aaron Goldfain. The island wall diffusion data was collected the fall and winter of 2011. We took his data and compared it to the predicted results of Tatiana's model. These comparisons are shown in figures 7, 8, and 9.

Note that the data analysis was preformed by Arron Goldfain, using similar methods as described in the 2 island interaction section, and by Zhi Yuan in his notes about island wall interaction. We neglected to account for the confinement effects with this data so that the ratios can remain consistent with the "Ghost Particle" island wall interaction.




Figure 8: *Predicted and measured reduced mobility versus the ratio of separation and island radius for island wall interactions. Note that this is in the direction perpendicular to the island wall separation.*




Figure 9: *Ratio of the reduced mobilities for both along and perpendicular to the island wall separation.*