

Physics 601
Analytical Mechanics
Professor Siu Chin

Homework #9

Joe Becker
UID: 125-00-4128
November 11th, 2015

1 Problem #1

For a one-dimensional harmonic oscillator of mass, m , and spring constant k we can construct a Hamiltonian by finding the Lagrangian as

$$L = \frac{1}{2}m\dot{q}^2 - \frac{1}{2}m\omega^2 q^2$$

where $\omega = \sqrt{k/m}$. We find the canonical momentum of the system as

$$p = \frac{\partial L}{\partial \dot{q}} = m\dot{q}$$

which allows us to construct the Hamiltonian by

$$\begin{aligned} H = p\dot{q} - L &= \frac{p^2}{m} - \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2 \\ &= \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2 \\ &= \frac{1}{2}m \left(\left(\frac{p}{m} \right)^2 + (\omega q)^2 \right) \end{aligned}$$

For the *Canonical Transformation* of a one-dimensional harmonic oscillator given by

$$Q = C(p + im\omega q), \quad P = C(p - im\omega q)$$

We can find the constant C by using the fact that the determinant of the Jacobian is unity. This implies that

$$\begin{aligned} 1 &= \frac{\partial Q}{\partial q} \frac{\partial P}{\partial p} - \frac{\partial Q}{\partial p} \frac{\partial P}{\partial q} \\ &= (Cim\omega)C - C(-Cim\omega) \\ &= 2C^2im\omega \\ &\Downarrow \\ C &= \frac{1}{\sqrt{2im\omega}} \end{aligned}$$

So our transformations become

$$Q = \frac{1}{\sqrt{2im\omega}}(p + im\omega q), \quad P = \frac{1}{\sqrt{2im\omega}}(p - im\omega q)$$

Now we can find the generating function $F_2(q, P)$ for this transformation by noting that

$$\frac{\partial F_2}{\partial q} = p, \quad \frac{\partial F_2}{\partial P} = Q$$

Where we can solve the transformations such that

$$p(q, P) = \sqrt{2im\omega}P + im\omega q$$

and

$$Q(q, P) = \frac{1}{\sqrt{2im\omega}}(p(q, P) + im\omega q) = P + \sqrt{2im\omega}q$$

Which allows us to find

$$\begin{aligned}\frac{\partial F_2}{\partial P} &= Q = P + \sqrt{2im\omega}q \\ &\Downarrow \\ F_2(q, P) &= \frac{1}{2}P^2 + \sqrt{2im\omega}qP + f(q)\end{aligned}$$

Now we can find the function of q by

$$\begin{aligned}\frac{\partial F_2}{\partial q} &= P = \sqrt{2im\omega}P + f'(q) \\ &\Downarrow \\ \sqrt{2im\omega}P + im\omega q &= \sqrt{2im\omega}P + f'(q) \\ &\Downarrow \\ f(q) &= \frac{1}{2}im\omega q^2\end{aligned}$$

So our generator function is

$$F_2(q, P) = \frac{1}{2}P^2 + \sqrt{2im\omega}qP + \frac{1}{2}im\omega q^2$$

which allows us see that

$$\begin{aligned}QP &= \frac{1}{2im\omega}(p + im\omega q)(p - im\omega q) \\ &= -\frac{i}{2m\omega}(p^2 + (m\omega q)^2) \\ &= -\frac{i}{2}m\left(\frac{1}{\omega}\left(\frac{p}{m}\right)^2 + \omega q^2\right)\end{aligned}$$

Which implies that the Hamiltonian is

$$H = i\omega QP$$

so the equations of motion are given by

$$\dot{P} = -\frac{\partial H}{\partial Q} = -i\omega P, \quad \dot{Q} = \frac{\partial H}{\partial P} = i\omega Q$$

Which easily allows us to solve the equations of motion as

$$P(t) = Ae^{-i\omega t}, \quad Q(t) = Be^{i\omega t}$$

2 Problem #2

We note that the generator $F(q, Q) = F_1(q, Q)$ follows from the Legendre transform that states

$$\delta \int_{t_1}^{t_2} (p\dot{q} - H) - (P\dot{Q} - K) dt = 0$$

which implies that the difference in the integrand can only differ by a total time derivative

$$(p\dot{q} - H) - (P\dot{Q} - K) = \frac{dF}{dt}$$

which implies that

$$dF = (K - H)dt + pdq - PdQ$$

We note that F is a function of q and Q . We can use this to find $F_3(p, Q)$ by noting that

$$d(pq) = pdq + qdp$$

which allows us to write

$$\begin{aligned} dF &= (K - H)dt + pdq - PdQ \\ &\Downarrow \\ dF &= (K - H)dt + d(pq) - qdp - PdQ \\ &\Downarrow \\ d(F - pq) &= (K - H)dt - qdp - PdQ \end{aligned}$$

So $F_3(p, Q) = F - pq$. Next we repeat the process to find $F_4(p, P)$ by noting that

$$d(PQ) = PdQ + QdP$$

which implies

$$\begin{aligned} dF &= (K - H)dt + pdq - PdQ \\ &\Downarrow \\ dF &= (K - H)dt + pdq - (d(PQ) - QdP) \\ &\Downarrow \\ d(F + PQ) &= (K - H)dt + pdq + QdP \end{aligned}$$

which implies that $F_4(p, P) = F + PQ$.

3 Problem #3

(a) For the infinitesimal transformation with the generator given by

$$F_2(q, P) = qP + \epsilon H(q, P)$$

where H is the Hamiltonian given by

$$H(q, p) = \frac{p^2}{2m} + V(q)$$

we can find the canonical transformations associated with $F_2(q, P)$ by

$$\begin{aligned} Q = \frac{\partial F_2}{\partial P} &= q + \epsilon \frac{\partial}{\partial P} \left(\frac{P^2}{2m} + V(q) \right) \\ &= q + \epsilon \frac{P}{m} \end{aligned}$$

Then we can find p and invert by

$$\begin{aligned} p = \frac{\partial F_2}{\partial q} &= P + \epsilon \frac{dV}{dq} \\ &\Downarrow \\ P &= p - \epsilon \frac{dV}{dq} \end{aligned}$$

So replacing P in the Q term we get the canonical transformations to first order in ϵ

$$\begin{aligned} P &= p - \epsilon \frac{dV}{dq} \\ Q &= q + \epsilon \frac{p}{m} \end{aligned}$$

(b) If we take ϵ to be a small time step we can see that for $q = q(t)$ and $p = p(t)$ we have the transformations

$$\begin{aligned} q(t + \epsilon) = Q &= q(t) + \epsilon \frac{p(t)}{m} \\ p(t + \epsilon) = P &= p(t) - \epsilon \frac{dV}{dq} \end{aligned}$$

If we Taylor expand $q(t + \epsilon)$ about ϵ to first order we see that

$$\begin{aligned} q(t + \epsilon) &= q(t) + \dot{q}\epsilon \\ p(t + \epsilon) &= p(t) + \dot{p}\epsilon \end{aligned}$$

We note that the velocity \dot{q} is related to momentum p by $\dot{q} = p/m$ which implies that The q expansion is the same as the canonical transformation. We also note that $\dot{p} = F = -dV/dq$ so the expansion of p also is the same to first order.

(c) Using the results from above we can find the transformed Hamiltonian $K(P, Q)$ by noting

$$\begin{aligned}
 H(q, p) &= \frac{p^2}{2m} + V(q) \\
 &\Downarrow \\
 K(Q, P) &= \frac{1}{2m} \left(P + \epsilon \frac{dV}{dq} \right)^2 + V \left(Q - \epsilon \frac{P}{m} \right) \\
 &= \frac{1}{2m} \left(P^2 + P\epsilon \frac{dV}{dq} + \epsilon^2 \left(\frac{dV}{dq} \right)^2 \right) + V(Q) - \epsilon \frac{P}{m} \frac{dV}{dQ} + \epsilon^2 \frac{p^2}{m^2} \frac{d^2V}{dQ^2} \\
 &= \frac{P^2}{2m} + V(Q) + \epsilon \left(P \frac{dV}{dq} - \frac{P}{m} \frac{dV}{dQ} \right) + \epsilon^2 \left(\left(\frac{dV}{dq} \right)^2 + \frac{p^2}{m^2} \frac{d^2V}{dQ^2} \right)
 \end{aligned}$$

4 Problem #4

(a) For the generator

$$F_3(p, Q) = -pQ + \epsilon H(Q, p)$$

where H is the same Hamiltonian from problem 3 this allows up to find the transformations by

$$q = -\frac{\partial F_3}{\partial p} = Q - \epsilon \frac{p}{m}$$

and

$$P = -\frac{\partial F_3}{\partial Q} = p - \epsilon \frac{dV}{dQ}$$

So solving for Q we have the canonical transformations by noting that

$$\frac{\partial V}{\partial q} = \frac{\partial V}{\partial Q} \frac{\partial V}{\partial q}$$

but as we can see $dQ/dq = 1$ so

$$Q = q + \epsilon \frac{p}{m}$$

$$P = p - \epsilon \frac{dV}{dq}$$

which are the same canonical transformations as problem 3.

(b) For ϵ as a time step we can find the same result that

$$q(t + \epsilon) = Q = q(t) + \epsilon \frac{p(t)}{m}$$

$$p(t + \epsilon) = P = p(t) - \epsilon \frac{dV}{dq}$$

This again is the same as the Taylor expansion about ϵ to first order. Where $\dot{p} = -dV/dq$ and $\dot{q} = p/m$.

(c) We note that the canonical transformation found in part (a) is the same as the transformation from problem 3. Therefore the transformed Hamiltonian to second order is the same given by

$$K(Q, P) = \frac{P^2}{2m} + V(Q) + \epsilon \left(P \frac{dV}{dq} - \frac{P}{m} \frac{dV}{dQ} \right) + \epsilon^2 \left(\left(\frac{dV}{dq} \right)^2 + \frac{p^2}{m^2} \frac{d^2V}{dQ^2} \right)$$

(d)

5 Problem #5

For the Kepler problem with a three dimensional Hamiltonian given by

$$H = \frac{\mathbf{p}^2}{2m} - \frac{k}{|\mathbf{r}|}.$$

Using this we can show that

$$\mathbf{L} = \mathbf{r} \times \mathbf{p}$$

is a constant of motion by taking the Poisson bracket with H and noting that the l component of \mathbf{L} is

$$\left(\mathbf{r} \times \mathbf{p} \right)_l = \epsilon_{lmn} r_m p_n$$

$$\mathbf{L}, H =$$