Sharp quantitative stability of the Brunn-Minkowski inequality

Alessio Figalli, Peter van Hintum, Marius Tiba

November 1, 2023

Abstract

The Brunn-Minkowski inequality states that for bounded measurable sets A and B in \mathbb{R}^n , we have $|A+B|^{1/n} \geq |A|^{1/n} + |B|^{1/n}$. Also, equality holds if and only if A and B are convex and homothetic sets in \mathbb{R}^d . The stability of this statement is a well-known problem that has attracted much attention in recent years. This paper gives a conclusive answer by proving the sharp stability result for the Brunn-Minkowski inequality on arbitrary sets.

Contents

T	11111	roduction	3
	1.1	An alternative approach to Theorem 1.5	
	1.2	Notation and conventions	
	1.3	Overview of the proofs of the main results	
	1.4	Structure of the paper	8
2	Init	ial reductions of Theorem 1.4, Theorem 1.5, and Theorem 1.6	8
	2.1	Setup	8
	2.2	Proposition	9
	2.3	Auxiliary Lemmas	9
	2.4	Proof of Proposition 2.9	10
	2.5	Proofs of Auxiliary Lemmas	10
		2.5.1 Proof of Lemma 2.11	10
		2.5.2 Proof of Lemma 2.12	11
3	Out	tline of the proof of the Quadratic Theorem (Theorem 1.5)	12
	Inte	ermediate results for the Quadratic Theorem (Theorem 1.5): Part I	13
	Inte	ermediate results for the Quadratic Theorem (Theorem 1.5): Part I Setup	
			13
	4.1	Setup	13 13
	$4.1 \\ 4.2$	Setup	13 13 14
	4.1 4.2 4.3	Setup	13 13 14 14
	4.1 4.2 4.3 4.4	Setup	13 13 14 14 15
	4.1 4.2 4.3 4.4 4.5	Setup Theorem Propositions Auxiliary Lemmas Proof of Theorem 4.11	13 13 14 14 15 16
	4.1 4.2 4.3 4.4 4.5	Setup Theorem Propositions Auxiliary Lemmas Proof of Theorem 4.11 Proof of Propositions	13 13 14 14 15 16 16
	4.1 4.2 4.3 4.4 4.5	Setup Theorem Propositions Auxiliary Lemmas Proof of Theorem 4.11 Proof of Propositions 4.6.1 Proof of Proposition 4.12	13 13 14 14 15 16 16 17
	4.1 4.2 4.3 4.4 4.5 4.6	Setup Theorem Propositions Auxiliary Lemmas Proof of Theorem 4.11 Proof of Propositions 4.6.1 Proof of Proposition 4.12 4.6.2 Proof of Proposition 4.13	13 13 14 14 15 16 16 17 20
	4.1 4.2 4.3 4.4 4.5 4.6	Setup Theorem Propositions Auxiliary Lemmas Proof of Theorem 4.11 Proof of Propositions 4.6.1 Proof of Proposition 4.12 4.6.2 Proof of Proposition 4.13 Proof of Auxiliary Lemmas	13 13 14 14 15 16 16 17 20 20
	4.1 4.2 4.3 4.4 4.5 4.6	Setup Theorem Propositions Auxiliary Lemmas Proof of Theorem 4.11 Proof of Propositions 4.6.1 Proof of Proposition 4.12 4.6.2 Proof of Proposition 4.13 Proof of Auxiliary Lemmas 4.7.1 Proof of Lemma 4.14	13 13 14 14 15 16 16 17 20 20 21
	4.1 4.2 4.3 4.4 4.5 4.6	Setup Theorem Propositions Auxiliary Lemmas Proof of Theorem 4.11 Proof of Propositions 4.6.1 Proof of Proposition 4.12 4.6.2 Proof of Proposition 4.13 Proof of Auxiliary Lemmas 4.7.1 Proof of Lemma 4.14 4.7.2 Proof of Lemma 4.15	13 13 14 14 15 16 16 17 20 20 21 21

5	Intermediate results for Quadratic Theorem (Theorem 1.5): Part II	22
	5.1 Setup	22
	5.2 Theorem	22
	5.3 Propositions	22
	5.4 Proof of Theorem	24
	5.4.1 Proof of Theorem 5.3	24
	5.5 Proofs of Propositions	24
	5.5.1 Proof of Proposition 5.4	24
	5.5.2 Proof of Proposition 5.5	25
	5.5.3 Proof of Proposition 5.6	25
	5.5.4 Proof of Proposition 5.7	25
_		a=
6	Intermediate results for the Quadratic Theorem (Theorem 1.5): Part III	27
	6.1 Setup	27
	6.2 Propositions	27
	6.3 Proof of Propositions	28
	6.3.1 Proof of Quadratic Theorem: Proposition 6.6	
	6.3.2 Proof of Proposition 6.7	30
	6.3.3 Proof of Proposition 6.8	30
7	Proof of the Quadratic Symmetric Difference result (Theorem 1.5)	31
'	1 roof of the Quadratic Symmetric Difference result (Theorem 1.9)	31
8	Intermediate results for Linear Theorem with few vertices (Theorem 1.6)	33
	8.1 Outline of the proof of Theorem 8.1	33
	8.2 Auxiliary propositions	33
	8.3 Auxiliary lemmas	34
	8.4 Proof of Theorem 8.1	35
	8.5 Proof of Propositions	36
	8.5.1 Sublinearity of doubling in subsimplices; Proposition 8.2	36
	8.5.2 Proof of Proposition 8.4	37
	8.6 Proof of Auxiliary Lemmas	37
	8.6.1 Proof of Lemma 8.5	37
	8.6.2 Proof of Lemma 8.6	38
	8.6.3 Proof of Lemma 8.8	38
	8.6.4 Proof of Lemma 8.9	39
	8.6.5 Proof of Lemma 8.10	40
		10
9	Proof of the Linear Theorem for few vertices (Theorem 1.6)	40
10		1 (70)
10	O Intermediate results for the Linear Theorem: from few vertices (Theorem 1.6) to the general 1.1. Outline of the proof of Theorem 1.4.	
	10.1 Outline of the proof of Theorem 1.4	41 42
	10.2 Setup	
	10.3 Propositions	42
	10.4 Lemmas	42
	10.5 Proof of Propositions	43
	10.5.1 Proof of Proposition 10.3	43
	10.5.2 Proof of Proposition 10.4	44
	10.5.3 Proof of Proposition 10.5	44
	10.6 Proof of Lemmas	45
	10.6.1 Proof of Lemma 10.6	45
	10.6.2 Proof of Lemma 10.7	45
	10.6.3 Proof of Lemma 10.8	47
	10.6.4 Proof of Lemma 10.9	48
	10.6.5 Proof of Lemma 10.10	49
	10.6.6 Proof of Lemma 10.11	49
11	1 Proof of general Linear Theorem (Theorem 1.4)	50

12 .	Pro	of of Symmetric difference vs Common convex hull result (Theorem 1.7)	50	
	12.1	Propositions	50	
	12.2	Proofs of Propositions	51	
		12.2.1 Proof of Proposition 12.1	51	
		12.2.2 Proof of Proposition 12.2	51	
	12.3	Proof of Theorem 1.7	52	
13 Putting it all together: Proof of Theorem 1.3				
14 (Оре	en Problems	53	

1 Introduction

Given measurable sets $X, Y \subset \mathbb{R}^n$ with positive measure, the Brunn-Minkowski inequality says that

$$|X+Y|^{\frac{1}{n}} \ge |X|^{\frac{1}{n}} + |Y|^{\frac{1}{n}}.$$

Alternatively, for equal sized measurable sets $A, B \subset \mathbb{R}^n$ and a parameter $t \in (0,1)$, this is equivalent to

$$|tA + (1-t)B| \ge |A|,$$

with equality for equal convex sets A and B (less a measure zero set). Here, $A+B=\{a+b\mid a\in A, \text{ and } b\in B\}$ is the $Minkowski\ sum$, $tA:=\{ta:a\in A\}$, and $|\cdot|$ refers to the outer Lebesgue measure.

The Brunn-Minkowski inequality is part of a vast body of geometric inequalities, such as the isoperimetric inequality, the Prékopa-Leindler inequality, and the Borell-Brascamb-Lieb inequality. The famous isoperimetric inequality, which states that for a given volume the body minimizing its perimeter is the ball, follows from Brunn-Minkowski by taking A a ball and letting t tend to zero. The Prékopa-Leindler inequality asserts that for $t \in (0,1)$ and functions $f,g,h \colon \mathbb{R}^n \to \mathbb{R}_{\geq 0}$ with the property that $h(tx+(1-t)y) \geq f^t(x)g^{1-t}(y)$ for all $x,y \in \mathbb{R}^n$ and $\int f = \int g$, we have $\int h \geq \int f$ with equality if and only if $f(x) = ag(x-x_0)$ is a log-concave function for some $a \in \mathbb{R}_{>0}$ and $x_0 \in \mathbb{R}^n$. The Prékopa-Leindler inequality implies Brunn-Minkowski by taking f and g to be the indicator functions of f and f and f and f and f are prékopa-Leindler inequality in turn is subsumed by the Borell-Brascamb-Lieb inequality. Studying these inequalities and their stabilities has sparked a fruitful field of research in recent years.

The stability of the Brunn-Minkowski inequality says that if we are *close* to equality, then the sets are *close* to being equal and convex (up to translates), and the aim is to quantify the two notions of closeness (see e.g. [Fig14]). The major folklore conjecture concerning the stability of the Brunn-Minkowski inequality is that we are within a factor $1+\delta$ from equality, then the distance from A and B to a common convex set is $O_n(t^{-1/2}\delta^{1/2})$.

Conjecture 1.1. For $n \in \mathbb{N}$, $n \geq 2$ and $t \in (0, 1/2)$ there exist $c_n, d_{n,t} > 0$ such that the following holds. Let A and B be measurable sets of equal size with $|tA + (1-t)B| \leq (1+\delta)|A|$ and $\delta < d_{n,t}$. Then there exists a convex set K such that, up to translation, $K \supset A, B$ and

$$|K \setminus A| = |K \setminus B| \le c_n t^{-1/2} \delta^{1/2} |A|.$$

Another important conjecture regarding the stability of the Brunn-Minkowski inequality is that the distance from A and B to their individual convex hulls is linear $O_{n,t}(\delta)$. Figalli and Jerison [FJ21] formulated this conjecture in the case of equal sets, and van Hintum, Spink and Tiba [vHSTarb, Section 12] considered this conjecture for arbitrary sets in the plane.

Conjecture 1.2. For $n \in \mathbb{N}$ and $t \in (0, 1/2)$ there exist $c_n, d_{n,t} > 0$ such that the following holds. Let A and B be measurable sets of equal size with $|tA + (1-t)B| \le (1+\delta)|A|$ and $\delta < d_{n,t}$. Then

$$|\operatorname{co}(A) \setminus A| + |\operatorname{co}(B) \setminus B| \le c_{n,t} \delta |A|.$$

These conjectures have received a lot of attention becoming central problems in convex geometry (see e.g. [FMP09, FMP10a, Chr12b, Chr12a, EK14, FJ15, Fig15, FJ17, BJ17, CM17, FJ21, vHST22, vHK23a, vHK23b, vHSTara, vHSTarb]).

The first contribution to the study of stability was made by Freiman [Fre59] in dimension n=1. Freiman's celebrated 3k-4 Theorem [Fre59, LS95, Sta96] from additive combinatorics, implies the following strong version of Conjecture 1.2: If $t \in (0, 1/2]$ and $A, B \subset \mathbb{R}$ are measurable sets with equal volume such that

 $|tA + (1-t)B| \le (1+\delta)|A|$ with $\delta < t$, then $|\cos(A) \setminus A| \le t^{-1}\delta|A|$ and $|\cos(B) \setminus B| \le (1-t)^{-1}\delta|B|$. Simple examples show that this result is optimal.

Stability in higher dimensions is considerably more difficult; in [Chr12b, Chr12a] Christ showed a qualitative result: If $n \in \mathbb{N}$, $t \in (0, 1/2]$, and $A, B \subset \mathbb{R}^n$ are measurable sets with equal volume such that $|tA + (1-t)B| \le (1+\delta)|A|$ with δ sufficiently small in terms of t and n, then there exists a convex set K such that, up to translation, $K \supset A, B$ and $|K \setminus A| = |K \setminus B| = o_{n,t,\delta}(1)|A|$, where $o_{n,t,\delta}(1) \to 0$ as $\delta \to 0$ for fixed n and t. In a cornerstone result, Figalli and Jerison [FJ17] obtained the first quantitative bounds: $|K \setminus A| = |K \setminus B| \le \delta^{(t/|\log(t)|)^{\exp(O(n))}}|A|$. A similar result for the Prékopa-Leindler inequality was recently established by Böröcky, Figalli, and Ramos [BFRar].

The only instance of Conjecture 1.1 for arbitrary sets was established in two dimensions by van Hintum, Spink, and Tiba [vHSTarb]. In an independent direction, van Hintum and Keevash [vHK23b] (see also Proposition 8.3) determined the optimal value $d_{n,t} = t^n$ for all $n \in \mathbb{N}$ and $t \in (0, 1/2]$, with the same bound on the distance to a common convex set as in the result of Figalli and Jerison.

Even partial answers to Conjecture 1.1 for restricted classes of sets A and B have received much attention. These papers have focused on controlling the weaker "asymmetry" measure $\inf_x |A\triangle(B+x)|$, which a priori does not control $|K\setminus A|$. In [FMP09, FMP10b], Figalli, Maggi, and Pratelli established that given $n\in\mathbb{N}$, $t\in(0,1/2]$, and $A,B\subset\mathbb{R}^n$ convex sets with equal volume, if $|tA+(1-t)B|\leq(1+\delta)|A|$ with δ sufficiently small in terms of t and n, then, up to translation, $|A\triangle B|\leq O_d(t^{-1/2}\delta^{1/2})|A|$. Figalli, Maggi, and Mooney [FMM18] showed the analogous result when A is a ball and B is arbitrary. Note that this is closely related to the stability of the isoperimetric inequality. Barchiesi and Julin [BJ17] extended the previous results to A convex and B arbitrary. Despite all these results supporting Conjecture 1.1, a conclusive answer remained wide open and outside the scope of the available techniques.

The particular case of equal sets A=B in Conjecture 1.2 has been thoroughly investigated. Indeed, after establishing in [FJ15] some quantitative bounds for Conjecture 1.2 in all dimensions, Figalli and Jerison [FJ21] resolved Conjecture 1.2 for A=B in dimensions n=1,2,3, and subsequently Van Hintum, Spink, and Tiba [vHST22] resolved Conjecture 1.2 for A=B in all dimensions. Moreover, they determined the optimal dependency on t. Furthermore, van Hintum, Spink, and Tiba [vHSTara, Theorem 1.1] established the optimal dependency on t in dimensions t in dimensions t is the hypograph of a function over a convex domain. Another closely related result by van Hintum and Keevash [vHK23a] is that if t is that if t is the hypograph of t with t is the hypograph of t with t in the theorem t is the hypograph of t in the theorem t in the t in the hypograph of t is that if t is the hypograph of t in the hypograph of t in the hypograph of t is that if t is the hypograph of t in the hypograp

For distinct sets A and B, Conjecture 1.2 has proved much more difficult. Van Hintum, Spink, and Tiba in [vHSTara, Theorem 1.5], resolved Conjecture 1.2, when A and B are hypographs of functions over the same convex domain. The only instance of Conjecture 1.2 for arbitrary sets was established by van Hintum, Spink, and Tiba [vHSTarb, Section 12] in two dimensions. Despite these determined efforts, for arbitrary sets in higher dimensions Conjecture 1.2 remained open.

Our main results resolve the conjectured quadratic stability to a common convex hull and the conjectured linear stability to the individual convex hulls in the Brunn-Minkowski inequality, concluding a long line of research on these problems.

Theorem 1.3. For all $n \in \mathbb{N}$ and $t \in (0, 1/2]$, there are computable constants $c_n^{1,3}, d_{n,t}^{1,3} > 0$ such that the following holds. Assume $\delta \in [0, d_{n,t}^{1,3})$ and let $A, B \subset \mathbb{R}^n$ be measurable sets with equal volume satisfying

$$|tA + (1-t)B| = (1+\delta)|A|.$$

Then, up to translation¹, there is a convex set $K \supset A \cup B$ such that

$$|K \setminus A| + |K \setminus B| \le c_n^{1.3} t^{-1/2} \delta^{1/2} |A|.$$

Theorem 1.4. For $n \in \mathbb{N}$ and $0 < t \le \frac{1}{2}$, there are constants $c^{1.4}$, $d^{1.4}_{n,t} > 0$ such that the following hold. Assume $\delta \in [0, d^{1.4}_{n,t})$, and assume $A, B \subset \mathbb{R}^n$ are measurable sets of equal volume so that $|tA + (1-t)B| \le (1+\delta)|A|$. Then

$$|\operatorname{co}(A)\setminus A|+|\operatorname{co}(B)\setminus B|\leq t^{-c^{1}.4_{n^8}}\delta|A|.$$

The proofs of Theorem 1.3 and Theorem 1.4 are very involved and will be obtained by combining a series of intermediate results, all of which have their own interest. Moreover, Theorem 1.3 uses Theorem 1.4.

We prove Theorem 1.3 by first showing a sharp control of the symmetric difference between A and B.

 $^{^{1}\}text{That is, there exist }x,y\in\mathbb{R}^{n}\text{ so that }x+A,y+B\subset K\text{ and }|K\setminus(x+A)|+|K\setminus(y+B)|\leq t^{-c^{\textstyle 1.3}}n^{8}\delta^{\textstyle \frac{1}{2}}|A|.$

Theorem 1.5. For all $n \in \mathbb{N}$ and $t \in (0, 1/2]$, there are computable constants $c_n^{1.5}, d_{n,t}^{1.5} > 0$ such that the following holds. Assume $\delta \in [0, d_{n,t}^{1.5})$ and assume $A, B \subset \mathbb{R}^n$ are measurable sets with equal volume so that $|tA + (1-t)B| = (1+\delta)|A|$. Then, up to translation,

$$|A\triangle B| \le c_n^{1.5} t^{-1/2} \delta^{1/2} |A|.$$

The exponents of δ and t are optimal as shown by the example $A = [0, 1 + \sqrt{\delta/t}] \times [0, 1]^{d-1}$ and $B = [0, 1] \times [0, 1 + \sqrt{\delta/t}] \times [0, 1]^{d-2}$. In this case, we find $tA + (1-t)B = [0, 1 + t\sqrt{\delta/t}] \times [0, 1 + (1-t)\sqrt{\delta/t}] \times [0, 1]^{d-2}$, so that $|tA + (1-t)B| \le (1+\delta)|A|$, while $|A\triangle B| \ge 2\sqrt{\delta/t}|A|$.

An important step in proving Theorem 1.4 is to establish the case where co(A) and co(B) have a bounded number of vertices.

Theorem 1.6. For all $n, v \in \mathbb{N}$ and $t \in (0,1)$, there are computable constants $k_n^{1.6}(v), c^{1.6}, d_{n,t}^{1.6} > 0$ such that the following holds. Assume $\delta \in [0, d_{n,t}^{1.6}]$, and assume $A, B \subset \mathbb{R}^n$ are measurable sets of equal volume, so that co(A) has at most v vertices and $|tA + (1-t)B| = (1+\delta)|A|$. Then

$$|\operatorname{co}(A) \setminus A| \le k_n^{1.6}(v) \min\{t, 1-t\}^{-c^{1.6}n^8} \delta |A|.$$

As mentioned above, the volume of the symmetric difference $|A\triangle B|$ is a commonly used parameter to measure stability in geometric inequalities. Here, instead of controlling $|A\triangle B|$, we want to improve this notion of closeness by finding a common convex set K that contains both A and B. The key step to achieve this is contained in the next general theorem about convex sets, which is of independent interest.

Theorem 1.7. There exists a constant c_n such that given convex sets $X, Y \subset \mathbb{R}^n$, we have

$$\frac{|\operatorname{co}(X \cup Y)|}{\min\{|X|,|Y|\}} - 1 \le c_n \frac{|X \triangle Y|}{|X \cap Y|}.$$

Remark 1.8. In Theorem 1.3, Theorem 1.4 and Theorem 1.5, we can assume that, for fixed $n \in \mathbb{N}$, the function $d_{n,t} : (0,1/2] \to \mathbb{R}_+$ is increasing. Similarly, in Theorem 1.6, we can assume that for fixed $n \in \mathbb{N}$, the function $d_{n,t}$ increases in (0,1/2] and decreases in [1/2,1). This follows from the proofs.

1.1 An alternative approach to Theorem 1.5.

While working on this project, we proved the following result.

Theorem 1.9. For all $n \in \mathbb{N}$ and $t \in (0, 1/2]$, there are computable constants $c_n^{1,9}, d_{n,t}^{1,9}, \Gamma_{n,t}^{1,9} > 0$ such that the following holds. Assume $\delta \in [0, d_{n,t}^{1,9}]$, $\gamma \in [0, \Gamma_{n,t}^{1,9}]$, and assume that $A, B \subset \mathbb{R}^n$, are measurable sets with equal volume so that

$$|tA + (1-t)B| \le (1+\delta)|A|$$
 and $|\operatorname{co}(A) \setminus A| + |\operatorname{co}(B) \setminus B| \le \gamma |A|$.

Then, up to translation,

$$|A\triangle B| \le c_n^{1.9} \sqrt{\frac{\delta + \gamma}{t}} |A|.$$

This theorem, combined with Theorem 1.4, has the following important corollary.

Corollary 1.10. For all $n \in \mathbb{N}$ and $t \in (0, 1/2]$, there are computable constants $c_{n,t}^{1.10}, d_{n,t}^{1.10} > 0$ such that the following holds. Assume $\delta \in [0, d_{n,t}^{1.10}]$ and assume that $A, B \subset \mathbb{R}^n$, are measurable sets with equal volume so that $|tA + (1-t)B| \le (1+\delta)|A|$. Then, up to translation,

$$|A\triangle B| \le c_{n,t}^{1.10} \delta^{1/2} |A|.$$

Although this corollary provides a sharp stability bound in terms of δ , it is weaker than Theorem 1.5 in that the dependency on t is suboptimal. Actually, even combining Theorem 1.9 with the optimal result contained in Conjecture 14.1 would not obtain the optimal t-dependence provided by Theorem 1.5. For this reason, in this paper, we develop a completely different approach to prove Theorem 1.5 that bypasses the use of Theorem 1.9.

Still, we believe that the proof of Theorem 1.9 brings a lot of value in studying the stability of the Brunn-Minkowski inequality, as it uses a mass transport approach in a new original way, and we defer its proof to a forthcoming paper [FvHT].

1.2 Notation and conventions.

Before starting our proofs, it is convenient to briefly explain the notation that we will use throughout the paper. With c>0, we shall denote a universal constant independent of the dimension, while $c_n>0$ (and analogous notations) denote dimensional constants. Saying that the quantity a is controlled by $O_n(b)$ means that $|a| \le c_n b$, while notation $a = \Omega_n(b)$ means that $a \ge c_n |b|$. When a constant also depends on t, we write $c_{n,t}$. To distinguish the constants that appear in the different statements, $c^{\ell,m}$ means that the constant c is the one appearing in Theorem/Proposition/Lemma $\ell.m$.

Throughout the paper, we fix $n \in \mathbb{N}$ with $n \geq 3$ and either $t \in (0, 1/2]$ or $t \in (0, 1)$; unless otherwise specified, we assume the former. We use $|\cdot|$ to denote the outer Lebesgue measure in \mathbb{R}^n .

Given $s \in \mathbb{R}$ and sets X and Y in \mathbb{R}^n , we define $sX = \{sx \colon x \in X\}$ and $X + Y = \{x + y \colon x \in X, y \in Y\}$. A set X in \mathbb{R}^n is convex if for all $t \in [0,1]$ we have $tX + (1-t)X \subset X$. The convex hull co(X) of a set X in \mathbb{R}^n is the intersection of all convex sets containing X. In particular, co(X) is a convex set. Two sets X and Y of \mathbb{R}^n are homothetic if there exist a point z in \mathbb{R}^n and a scalar s > 0 such that X = sY + z.

Given a bounded convex set X in \mathbb{R}^n , we define \overline{X} as the closure of X, which is also a convex set. The vertices of X, denoted by V(X), represent the set $V(X) = \{x \in \overline{X} : \operatorname{co}(\overline{X} \setminus \{x\}) \neq \operatorname{co}(\overline{X})\}$. It follows that $\overline{X} = \operatorname{co}(V(X))$.

Measureable sets X_1, \ldots, X_k in \mathbb{R}^n are said to form an essential partition of \mathbb{R}^n if $|\cap_i X_i^c| = 0$ and for $j_1 \neq j_2$, we have $|X_{j_1} \cap X_{j_2}| = 0$. By a basis e_1, \ldots, e_n in \mathbb{R}^n , we mean an orthogonal set of vectors with unit length. In light of Proposition 2.9, we can assume that the sets A and B (as well as all parts into which we subdivide A and B) are compact.

1.3 Overview of the proofs of the main results

We first prove the linear stability Theorem 1.4 which is a crucial tool at several steps in the proof of the quadratic stability Theorem 1.3. The proof of Theorem 1.4 breaks up into two parts: first we show a linear result for sets A so that co(A) has few vertices (Theorem 1.6), then we use this to prove the result for an arbitrary number of vertices.

First consider Theorem 1.6, i.e., the linear stability to the convex hull for sets with few vertices. The first step is to reduce to the case where $\operatorname{co}(A)$ is a simplex (see Theorem 8.1). Assume that A and B are a finite union of points and boxes, $V(\operatorname{co}(A)) = \{x_0, \dots, x_n\}$, and assume that |A| = |B| and $|tA + (1-t)B| \le (1+\delta)|A|$. Our move now is to pick a point x in A and construct cones C_0, \dots, C_n where C_i has a vertex at x and is generated by rays $xx_0, xx_1, \dots, xx_{i-1}, xx_{i+1}, \dots, xx_{d-1}, xx_n$. The cones C_0, \dots, C_n partition A into subsets A_0, \dots, A_n with the property that $\operatorname{co}(A_i)$ is again a simplex. We find a translation y such that the cones $y + C_0, \dots, y + C_n$ partition B into subsets B_0, \dots, B_n where $|A_i| = |B_i|$ (see Proposition 5.4). Repeating this move in each part, we create a partition of A and B into sets A_1, \dots, A_m and B_1, \dots, B_m with the property that $|A_i| = |B_i|$, the sets $\{tA_i + (1-t)B_i\}_{1 \le i \le m}$ are all disjoint, and (crucially) $\operatorname{co}(A) = \sqcup_i \operatorname{co}(A_i)$. Our aim then becomes to show Theorem 1.6 for essentially all parts A_i and B_i , that is, $|\operatorname{co}(A_i) \setminus A_i| \le O_{n,t}(\delta_i)|A_i|$. Then, we can combine all the pieces to get

$$|\operatorname{co}(A)\setminus A|=\sum_{i}|\operatorname{co}(A_{i})\setminus A_{i}|\leq \sum_{i}O_{n,t}(\delta_{i})|A_{i}|=O_{n,t}(\delta)|A|.$$

In this process, we stop further subdividing a part A_i when either $A_i = \operatorname{co}(A_i)$ or $|A_i| \leq 0.01 |\operatorname{co}(A_i)|$. In Proposition 8.4, we show that we can pick the points in each part sufficiently centrally in such a way as to guarantee the following two facts. First, provided that some part A_i satisfies $|A_i| \geq 0.01 |\operatorname{co}(A_i)|$, then all the n+1 parts $A_{i,0},\ldots,A_{i,n}$ in which we subdivide A_i satisfy $|A_{i,j}| \geq \Omega_n(1) |\operatorname{co}(A_{i,j})|$. Second, for any parameter $\varepsilon > 0$, there exists $k \in \mathbb{N}$ such that essentially all parts A_i constructed in generation k are small, in the sense that the total volumes of all large parts A_i with $\operatorname{diam}(A_i) \geq \varepsilon$ in generation ℓ are at most ε .

To conclude, we choose $\varepsilon > 0$ smaller than $\delta^2 |A|$, and sufficiently small so that the boundary of A (recall that A is a finite union of boxes and points) thickened by ε has size at most $\delta^2 |A|$. We run the process up to generation ℓ and note that all parts fall into four categories. First, there are the parts A_i where we stopped further subdividing because $|A_i| \leq 0.01 |\cos(A_i)|$; for such A_i we also get $|A_i| \geq \Omega_n(1) |\cos(A_i)|$. Second, there are the parts A_i where we stopped further subdividing because $A_i = \cos(A_i)$. Third are the small parts A_i in generation ℓ with diam $(A_i) \leq \varepsilon$ and $0.01 |\cos(A_i)| < |A_i| < |\cos(A_i)|$. As these are neither empty nor full, $\cos(A_i)$ must intersect the boundary of A so that the combined volume of these $\cos(A_i)$'s is at most $\delta^2 |A|$. Finally, there are the big parts in generation ℓ with diam $(A_i) > \varepsilon$, the total volume of which is at most $\delta^2 |A|$.

Now, neglecting the parts in the third and fourth categories, as they contribute very little, it is easy to check Theorem 1.6 for the parts in the first and second categories. Indeed, in the first category we can use a

qualitative stability result (see Proposition 8.3), and in the second category there is nothing to prove as the set is already convex.

Having proved Theorem 1.6, we turn our attention to generalizing it to an arbitrary number of vertices of co(A).

The first step (Proposition 10.3) is to identify a collection of disjoint convex regions $X_i \subset co(A)$ of small diameter ϵ that contain a positive proportion of the missing volume of $co(A) \setminus A$, i.e., $\sum |co(A \cap X_i) \setminus (A \cap X_i)| = \Omega_n(|co(A) \setminus A|)$. To find the regions X_i of small diameter, we induct akin to the proof for few vertices. First, we find as follows a triangulation of co(A) so that all simplices are pretty full. We triangulate $\partial co(A)$ and consider the n-simplices formed by a (n-1)-simplex in $\partial co(A)$ together with the origin o. Within each of these simplices, we find a central point in A which partitions the simplex into smaller simplices all of which preserve the convex hull. We iterate until we find a simplex with low density (between 98% and 99%) or small diameter. For simplices with low density, finding a convex subregion with smaller diameter containing a positive proportion of the missing region is a lot simpler (Lemma 10.8).

A standard reduction (Proposition 2.9) allows us to assume that

$$B^n(o, \Omega_n(1)) \subset K \subset A, B \subset (1+\eta)K \subset B^n(o, O_n(1))$$

for some convex set K and η small in terms of n and t. We consider a simplicial tube U i.e., a set of the form $T \times \mathbb{R}^+$ with $T \subset \mathbb{R}^{n-1}$ a regular simplex centered at the origin. We insist that the diameter of U is small in terms of n but much larger than the diameter ϵ of the regions X_i . We take a random rotation of U. As each X_i is completely contained inside U with probability $\Omega(1)$, we get that

$$|\operatorname{co}(A) \setminus A| = O(1) \sum |\operatorname{co}(A \cap X_i) \setminus (A \cap X_i)| = O(1) \mathbb{E}|\operatorname{co}(A \cap U) \setminus (A \cap U)|.$$

So it suffices to show that for every rotation of U we have $|\cos(A \cap U) \setminus (A \cap U)| = O_{n,t}(\delta)|A|$. After some reductions, we may assume $|A \cap U| = |B \cap U|$ and $|t(A \cap U) + (1-t)(B \cap U)| - |A \cap U| \le |tA + (1-t)B| - |A|$. Partitioning U into smaller parallel tubes U_i , we find that

$$\sum_{i} \left(|t(A \cap U_i) + (1-t)(B \cap U_i)| - t|A \cap U_i| - (1-t)|B \cap U_i| \right) \le |t(A \cap U) + (1-t)(B \cap U)| - |A \cap U|,$$

although we might have $|A \cap U_i| \neq |B \cap U_i|$. In the direction of the tubes, every fibre of $A \cap U_i$ and $B \cap U_i$ starts with a long interval, so that extending one of the sets at the bottom does not affect $|t(A \cap U_i) + (1-t)(B \cap U_i)| - t|A \cap U_i| - (1-t)|B \cap U_i|$ (cf. Lemma 10.11). Extending appropriately (which is always quite little), we retrieve $|A \cap U_i| = |B \cap U_i|$, and by choosing U_i appropriately we find that $co(A \cap U_i)$ has few vertices. Applying Theorem 1.6 concludes.

We turn to proof of Theorem 1.3. The first step is to reduce Theorem 1.3 to Theorem 1.5, that is, the first step is to switch from showing that A+x and B+y are contained in a common convex set K with almost the same volume, to show that $|(A+z)\triangle B|$ is small. This reduction follows quickly from Theorem 1.7 and Theorem 1.4.

The starting point for the proof of Theorem 1.5 is inspired by a classical proof of the Brunn-Minkowski inequality. Assume that the sets A and B are finite unions of boxes and assume that |A| = |B| and $|tA + (1 - t)B| \le (1 + \delta)|A|$. Our move is to choose a hyperplane B that partitions A into A_1 and A_2 , and then to find a parallel hyperplane B that partitions B into B_1 and B_2 such that $|A_i| = |B_i|$. Repetition of this move creates a partition of A and B into sets A_1, \ldots, A_m and B_1, \ldots, B_m with the property that $|A_i| = |B_i|$ and the sets $\{tA_i + (1 - t)B_i\}_{1 \le i \le m}$ are all disjoint. Hence, if $|tA_i + (1 - t)B_i| = (1 + \delta_i)|A_i|$, then $\sum_i \delta_i |A_i| \le \delta |A|$. Our aim is to show Theorem 1.5 for each pair of sets A_i and B_i , that is, $|(A_i + x_i) \triangle B_i| \le O_{n,t}(\delta_i^{1/2})|A_i|$. This is motivated by the fact that, by cutting with sufficiently many hyperplanes, the sets A_i and A_i become simple enough (e.g. boxes). Now, under the much too optimistic assumption that all translates x_i coincide, we can put all the pieces together to get

$$|(A+x)\triangle B| = \sum_{i} |(A_i + x_i)\triangle B_i| \le \sum_{i} O_{n,t}(\delta_i^{1/2})|A_i| \le O_{n,t}(\delta^{1/2})|A|.$$

The main problem with the above plan is that there is no reason to believe that all the translates x_i coincide. In general, given sets Y and Z, it is difficult to grasp the optimal translation x that minimizes $|(Y+x)\triangle Z|$. However, for certain classes of sets, which we call cone-like sets, this is possible.

A cone C is the intersection of a finite family of half-spaces generated by hyperplanes through the origin o. In particular, the vertex of the cone is at the origin. We always assume that the cone is not too wide, that is, all angles at o are less than 179°. We say that $Y \subset C$ is C-like if

$$C \cap B^n(o, \Omega_n(1)) \subset Y \subset C \cap B^n(o, O_n(1))$$

Now, fix a cone C and let Y and Z be convex (or approximately convex) subsets of C that are C-like. For such sets, if for a translate x we have $|(Y+x)\triangle Z| \leq \sqrt{\delta}|Y|$ then $||x||_2$ must be small so that also $|Y\triangle Z| \leq O_n(\sqrt{\delta})|Y|$. So, in effect, the essentially optimal translation for cone-like sets is x = o.

Motivated by this idea, the plan is to partition the whole space into cones $C_1, \ldots C_m$ using hyperplane cuts, which induce the partitions of A and B into subsets A_1, \ldots, A_m and B_1, \ldots, B_m , in order to reduce Theorem 1.5 from A and B to A_i and B_i , which we hope to be simpler to deal with. For this to work, we must impose that the subsets A_i and B_i of C_i are C_i -like and have the same size. The first condition turns out to be easy to satisfy, as after an affine transformation we can assume without loss of generality that $B^n(o, \Omega_n(1)) \subset A, B \subset B^n(o, O_n(1))$ (see Proposition 2.9). The second condition is also rather easy to satisfy.

In Theorem 4.11, we show that we can always find such a partition into cones, each of which is the convex hull of a set of bounded size of rays through the origin. Moreover, all the rays are clustered arbitrarily close to a pair of rays. Thus, each cone C_i has a bounded number of faces and is arbitrarily narrow in all but one direction. (In general, one cannot hope for all cones to be arbitrarily narrow in all directions, as one can see already in the two-dimensional case when $A = [0, 1 - \varepsilon] \times [0, 1 + \varepsilon]$ and $B = [0, 1 + \varepsilon] \times [0, 1 - \varepsilon]$.)

Now, recalling that A and B are finite unions of boxes and insisting that the narrow directions of the cones are much smaller than the sides of the boxes, we can assume without loss of generality that the subsets A_i and B_i of C_i are such that each sectional cut in the short directions is either completely full or completely empty.

So far, we have reduced Theorem 1.5 to rather simple sets A_i and B_i . So, given such sets A_i and B_i with $|A_i| = |B_i|$ and $|tA_i| + (1-t)B_i| \le (1+\delta_i)|A_i|$, we need to show that $|A_i \triangle B_i| \le O_n(\delta_i^{1/2}/t^{1/2})|A_i|$.

We proceed akin to the final step for the linear result. It is enough to show that for each large tube U loosely oriented in the same direction as C_i we have $|(A_i \cap U) \triangle (B_i \cap U)| \leq O_n(\delta_i^{1/2}/t^{1/2})|A_i|$. After some reductions, we may assume $|A_i \cap U| = |B_i \cap U|$. Now by the linear result we know that $|\cos(A_i \cap U) \setminus (A_i \cap U)| \leq O_{n,t}(\delta)|A_i|$. Recall that in A_i and B_i each sectional cut of C_i in the short directions is either completely full or completely empty. Combining the two facts we get that $A_i \cap U$ and $B_i \cap U$ are essentially one-codimension compressed sets. For such sets we can prove a sharp quadratic stability and thus conclude.

1.4 Structure of the paper

Given the complexity of our proofs, we kindly encourage the reader to refer to the table of contents as a guide for navigation and orientation within the paper. We note that:

- The proof of Theorem 1.3 uses Theorem 1.4, Theorem 1.5, and Theorem 1.7;
- The proof of Theorem 1.5 uses Theorem 1.4;
- The proof of Theorem 1.4 uses Theorem 1.6;
- Theorem 1.7 and Theorem 1.6 are proved directly.

Acknowledgements. AF acknowledges the support of the ERC Grant No.721675 "Regularity and Stability in Partial Differential Equations (RSPDE)" and of the Lagrange Mathematics and Computation Research Center.

2 Initial reductions of Theorem 1.4, Theorem 1.5, and Theorem 1.6

We shall deduce some basic allowed additional assumptions for Theorem 1.5, Theorem 1.4, and Theorem 1.6, but before we do so, we need a few definitions.

2.1 Setup

Definition 2.1. A set $X \subset \mathbb{R}^n$ is called simple if

$$X = \bigsqcup_{i \le k} x_i + [0, 1]^n$$

for some $k \in \mathbb{N}$, that is, X is a finite disjoint union of translates of the unit cube.

Definition 2.2. A convex set $C \subset \mathbb{R}^n$ is called a cone if there exists a hyperplane H not containing the origin and a bounded convex set $P \subset H$ such that

$$C = \bigsqcup_{t \ge 0} tP.$$

Definition 2.3. We write S^{v_0,\dots,v_n} for the simplex with vertices v_0,\dots,v_n . Assuming that S^{v_0,\dots,v_n} contains the origin in the interior, construct the family of cones $\mathfrak{C}^{v_0,\dots,v_n} := \{C_i : 0 \le i \le n\}$, where

$$C_i = \bigsqcup_{t>0} t \operatorname{co}(v_0, \dots v_{i-1}, v_{i+1}, \dots, v_n).$$

Note that the cones in $\mathfrak{C}^{v_0,\dots,v_n}$ form an essential partition of \mathbb{R}^n .

Definition 2.4. Fix vectors $e_0, \ldots, e_n \in \mathbb{R}^n$ such that S^{e_0, \ldots, e_n} is a regular unit volume simplex centered at the origin o. Denote $S = S^{e_0, \ldots, e_n}$ and $\mathfrak{C} = \mathfrak{C}^{e_0, \ldots, e_n}$.

Definition 2.5. A pair of sets $X, Y \subset \mathbb{R}^n$ is λ -bounded if there exists an r > 0 so that

$$rS \subset X, Y \subset \lambda rS$$
.

Observation 2.6. If $A, B \subset \mathbb{R}^n$ are λ -bounded and |A| = |B| = 1, then $B(o, (2\lambda n)^{-1}) \subset A, B \subset B(o, 2\lambda n)$.

Definition 2.7. Given a cone $F \subset C' \in \mathfrak{C}$, a pair of sets $X, Y \subset \mathbb{R}^n$ is (λ, F) -bounded if there exists an r > 0 so that

$$r(F \cap S) \subset X, Y \subset \lambda r(F \cap S).$$

Definition 2.8. A pair of sets $X,Y \subset \mathbb{R}^k$ is called a η -sandwich if there exists a convex set P such that $o \in P \subset X, Y \subset (1+\eta)P$.

Given a cone F and an η -sandwich $X,Y \subset \mathbb{R}^n$, the pair $X \cap F, Y \cap F$ is also an η -sandwich.

2.2 Proposition

Proposition 2.9. Fix $n \in \mathbb{N}$ and $t \in (0, 1/2]$.

- Assume that there exist constants $\lambda_n^{2.9} > 2n^2$ and $\eta_{n,t}^{2.9} > 0$ such that Theorem 1.5 holds for all simple $\lambda_n^{2.9}$ -bounded $\eta_{n,t}^{2.9}$ -sandwiches $A, B \subset \mathbb{R}^n$. Then it holds for all measurable sets $A, B \subset \mathbb{R}^n$.
- Assume that there exist constants $\lambda_n^{2.9} > 2n^2$ and $\eta_{n,t}^{2.9} > 0$ such that Theorem 1.4 holds for all $\lambda_n^{2.9}$ -bounded $\eta_{n,t}^{2.9}$ -sandwiches $A, B \subset \mathbb{R}^n$ so that A is a simple set intersected with co(A) and analogously for B. Then it holds for all measurable sets $A, B \subset \mathbb{R}^n$.
- Similarly, fix $n \in \mathbb{N}$ and $t \in (0,1)$. If Theorem 1.6 holds for all $\lambda_n^{2.9}$ -bounded $\eta_{n,t}^{2.9}$ -sandwiches $A, B \subset \mathbb{R}^n$ so that A is a simple set intersected with $\operatorname{co}(A)$, then it holds for all measurable sets $A, B \subset \mathbb{R}^n$.

2.3 Auxiliary Lemmas

We first collect some auxiliary results that will be used to prove Proposition 2.9. The proof of such results will be given in Section 2.5 below.

We recall the following result by Michael Christ.

Theorem 2.10 (Christ 2012, [Chr12a]). For all $n \in \mathbb{N}$, $t \in (0,1)$ and $\eta > 0$, there exist constants $\Delta_n^{2.10} > 0$, so that for all measurable $X, Y \subset \mathbb{R}^n$ of equal volume with the property that $|tX + (1-t)Y| \le (1 + \Delta_n^{2.10})|X|$, then

$$\min_{v \in \mathbb{R}^n} |\operatorname{co}(X \cup (v+Y))| \le (1+\eta)|X|.$$

We also need two lemmas.

Lemma 2.11. For $n \in \mathbb{N}$, $t \in (0,1/2]$ and $\eta > 0$, there exist constants $c^{2.11}$ and $\Delta^{2.11}_{n,t}(\eta) > 0$ so that the following holds. If $X,Y \subset \mathbb{R}^n$ are measurable sets with |X| = |Y| and $|tX + (1-t)Y| = (1+\delta)|X|$ with $\delta \in [0,\Delta^{2.11}_{n,t}]$, then, up to translation, there exist measurable sets $X',Y' \subset \mathbb{R}^n$ so that

1. X', Y' is an η -sandwich,

- 2. |X'| = |Y'| = |X|,
- 3. $\operatorname{co}(X') = \operatorname{co}(X)$ and $\operatorname{co}(Y') = \operatorname{co}(Y)$
- 4. $|X' \triangle X| + |Y' \triangle Y| \le c^{2.11} t^{-1} \delta |X|$,
- 5. $|tX' + (1-t)Y'| \le (1+\delta)|X|$.

Moreover, if $X \subset Y$, we additionally find $X' \subset Y'$.

Lemma 2.12. For $n \in \mathbb{N}$, and $\eta > 0$ the following holds. If $X, Y \subset \mathbb{R}^n$ is an η -sandwich, then there exists $v \in \mathbb{R}^n$ and there exists a linear transformation $\theta \colon \mathbb{R}^n \to \mathbb{R}^n$ such that $\theta(v+X), \theta(v+Y)$ is a $(n^2+n^3\eta)$ -bounded $n\eta$ -sandwich.

2.4 Proof of Proposition 2.9

Proof of Proposition 2.9. Choose η sufficiently small in terms of n, $\eta_n^{2.9}$, and $\lambda_n^{2.9}$. Choose ζ small in terms of η , and choose ξ sufficiently small in terms of ζ and η .

Apply Lemma 2.11 with parameter η to A,B to find A_1,B_1 . Apply Lemma 2.12 to A_1,B_1 to find A_2,B_2 which is a $(n^2+n^3\eta)$ -bounded $n\eta$ -sandwich. To additionally get that the sets are simple we use a standard approximation (see e.g. [vHK23a, Lemma 3.13]). Find compact subset $A_3\subset A_2$, so that $|A_2\setminus A_3|\to 0$ as $\zeta\to 0$. We can ensure $|\operatorname{co}(A_3)|\to |\operatorname{co}(A_2)|$ as $\zeta\to 0$, by requiring A_3 to contain a large finite subset of the vertices of $\operatorname{co}(A_2)$ (or all of them if $V(\operatorname{co}(A_2))$ is finite). Analogously define B_3 . Note that we may additionally ask that A_3,B_3 have the same size and form a $(n^2+n^3\eta)$ -bounded $n\eta$ -sandwich.

From here we consider Theorem 1.5, Theorem 1.4, and Theorem 1.6 separately. First, consider Theorem 1.5. Let

$$A_4 := \{x \in (\xi \mathbb{Z})^n : (x + [0, \xi]^n) \cap A_3 \neq \emptyset\} + [0, \xi]^n \supset A_3$$

and B_4 analogously. Since A_3 is compact, $|A_4 \setminus A_3| \to 0$ as $\xi \to 0$. Similarly, as $tA_3 + (1-t)B_3$ is compact, $|(tA_4 + (1-t)B_4) \setminus (tA_3 + (1-t)B_3)| \to 0$ as $\xi \to 0$. We can construct subsets A_5 and B_5 of A_4 and B_4 , respectively such that $A_5 = X + [0, \xi]^n$ and $B_5 = Y + [0, \xi]^n$ with $X, Y \subset (\xi \mathbb{Z})^n$, $|A_5| = |B_5| = \min\{|A_4|, |B_4|\}$ and A_5, B_5 is a simple $(n^2 + n^3\eta)$ -bounded $2n\eta$ -sandwich as $\xi \to 0$. By the above,

$$\lim_{\xi \to 0} \lim_{\xi \to 0} |tA_5 + (1-t)B_5|/|A_5| \le |tA_2 + (1-t)B_2|/|A_2| \le 1 + \delta.$$

Choosing η sufficiently small, we can apply Theorem 1.5 to find $\lim_{\zeta \to 0} |A_5 \triangle B_5|/|A_5| \le c_n^{1.5} t^{-1/2} \delta^{\frac{1}{2}}$ (up to a translation). This implies that, up to a translation, $|A_2 \triangle B_2|/|A_2| \le 2c_n^{1.5} t^{-1/2} \delta^{\frac{1}{2}}$, hence $|A \triangle B|/|A| \le 3c_n^{1.5} t^{-1/2} \delta^{\frac{1}{2}}$, as desired.

For Theorem 1.4 and Theorem 1.6, proceed analogously to the previous paragraph but with

$$A_4 := (\{x \in (\xi \mathbb{Z})^n : (x + [0, \xi]^n) \cap A_3 \neq \emptyset\} + [0, \xi]^n) \cap \operatorname{co}(A_3) \supset A_3$$

and B_4 similarly.

2.5 Proofs of Auxiliary Lemmas

2.5.1 Proof of Lemma 2.11

The idea will be to show that a large homothetic copy of $co(X \cup Y)$ is contained in tX + (1-t)Y, so that adding a slightly smaller homothetic copy of $co(X \cup Y)$ to X and Y will not change tX + (1-t)Y.

Proof of Lemma 2.11. Translate X and Y so that $|\cos(X \cup Y)|$ is minimal. Additionally, taking an affine transformation if necessary, we may assume that the John ellipsoid $E \subset \cos(X \cup Y)$ is a ball centered at the origin. Note that $|E| \ge n^{-n} |\cos(X \cup Y)| \ge n^{-n} |X|$.

Let $\eta' = t\eta$ and $\xi = \min\left\{\frac{1}{3}\left(\frac{t\eta'}{(1-t)n}\right)^n, 2^{-n}\right\}$. Apply Theorem 2.10 with parameters t, n, and $\eta^{2.10} = \xi$ to produce $\Delta^{2.10}$, and choose $\Delta_n = \Delta_n^{2.10}$, so that $|\cos(X \cup Y)| \le (1+\xi)|X|$.

Claim 2.13.
$$(1 - \eta') \cos(X \cup Y) \subset tX + (1 - t)Y$$

Proof of claim. Consider a point $p \in (1 - \eta') \operatorname{co}(X \cup Y)$. Find the point $p' \in \partial \operatorname{co}(X \cup Y)$ so that $p = \lambda p'$ for some $\lambda \in [0, 1 - \eta']$. Let $H : \mathbb{R}^n \to \mathbb{R}^n$ be the homothety of ratio $1 - \lambda > \eta'$ centered at p'. By convexity $H(E) \subset H(\operatorname{co}(X \cup Y)) \subset \operatorname{co}(X \cup Y)$ and H(o) = p', so that $\operatorname{co}(X \cup Y)$ contains a set of size

$$|H(E)| \ge (1 - \lambda)^n |E| \ge \eta'^n |E| \ge (\eta'/n)^n |X|$$

symmetric around p. Note that

$$|H(E) \setminus Y| \le |\operatorname{co}(X \cup Y) \setminus Y| \le \xi |Y| \le \frac{1}{3} \left(\frac{t\eta'}{(1-t)n}\right)^n |Y| \le \frac{1}{3} |H(E)|,$$

hence $|H(E) \cap Y| \ge \frac{2}{3}|H(E)|$. Now consider the homothety $H' : \mathbb{R}^n \to \mathbb{R}^n$ of ratio -t/(1-t) centered at p. Consider the set $H'(H(E) \cap Y) \subset H(E)$ and note that $|H'(H(E) \cap Y)| \ge \frac{2}{3}|H'(H(E))|$. Similarly as before, we find

$$|H'(H(E)\cap Y)\setminus X|\leq |\operatorname{co}(X\cup Y)\setminus X|\leq \frac{1}{3}\left(\frac{t\eta'}{(1-t)n}\right)^n|X|\leq \frac{1}{3}|H'(H(E))|\leq \frac{1}{2}|H'(H(E)\cap Y)|.$$

Hence, there exists a point $y \in H(E) \cap Y$, so that $H'(y) \in X \cap H'(H(E))$. Note that this implies that tx + (1-t)y = p, so that $p \in tX + (1-t)Y$. This concludes the claim.

Consider the sets $X'' := X \cup (1 - t^{-1}\eta')\operatorname{co}(X \cup Y)$ and $Y'' := Y \cup (1 - t^{-1}\eta')\operatorname{co}(X \cup Y)$. Note that we get

$$tX'' + (1-t)Y'' \subset (tX + (1-t)Y) \cup (t(1-t^{-1}\eta')\operatorname{co}(X \cup Y) + (1-t)\operatorname{co}(X \cup Y))$$

= $(tX + (1-t)Y) \cup (1-\eta')\operatorname{co}(X \cup Y)) = tX + (1-t)Y,$

where in the last equality we used the claim. We find $\frac{|X''\setminus X|}{|X|} \leq \frac{|\cos(X\cup Y)\setminus X|}{|X|} \leq \xi \leq 2^{-n}$. Now using the Brunn-Minkowski inequality, we find $(t|X''|^{\frac{1}{n}}+(1-t)|Y''|^{\frac{1}{n}})^n \leq |tX''+(1-t)Y''| = |tX+(1-t)Y| \leq (1+\delta)|X|$, which yields

$$\left(1 + t \frac{|X'' \setminus X|}{2n|X|} + (1-t) \frac{|Y'' \setminus Y|}{2n|Y|}\right)^n \le \left(t \left(1 + \frac{|X'' \setminus X|}{|X|}\right)^{\frac{1}{n}} + (1-t) \left(1 + \frac{|Y'' \setminus Y|}{|Y|}\right)^{\frac{1}{n}}\right)^n \le 1 + \delta,$$

so that $|X'' \setminus X| + |Y'' \setminus Y| \le 2\delta t^{-1}|X|$.

Having gained control over the size of X'' and Y'', we remove some of it to get the equality in sizes. To this end choose subsets $X' \subset X''$ so that the vertices of co(X) = co(X'') are in X' (so that co(X') = co(X)), $(1 - t^{-1}\eta') co(X \cup Y) \subset X'$ and |X'| = |X|. Similarly, choose $Y' \subset Y''$ and ensure |X'| = |Y'|. Now note that

$$|X' \triangle X| + |Y' \triangle Y| = |X'' \setminus X| + |X'' \setminus X'| + |Y'' \setminus Y| + |Y'' \setminus Y'| = 2(|X'' \setminus X| + |Y'' \setminus Y|) \le 4\delta t^{-1}|X|,$$

and $|tX' + (1-t)Y'| \le |tX'' + (1-t)Y''| = |tX + (1-t)Y| \le (1+\delta)|X|$.

Finally, if $X \subset Y$, note that $|Y \setminus X| = 0$. Constructing X' as before and setting $Y' = X' \cup (Y \setminus X)$ concludes the proof.

2.5.2 Proof of Lemma 2.12

Proof of Lemma 2.12. We begin the proof with the following two claims.

Claim 2.14. Let $\alpha, \eta \in (0,1)$, $P \subset \mathbb{R}^n$ convex and X, Y, so that $o \in P \subset X, Y \subset (1+\eta)P$ and let $o' \in (1-\alpha)P$. Let P' = P - o', X' = X - o' and Y' = Y - o'. Then $o \in P' \subset X', Y' \subset (1+\eta/\alpha)P'$.

Proof. The first and second containment is trivial so we focus on the last containment. By hypothesis, it is enough to show $(1+\eta)P - o' \subset (1+\eta/\alpha)(P-o')$. This is equivalent to $(1+\eta)P + (\eta/\alpha)o' \subset (1+\eta/\alpha)P$. Because $o' \in (1-\alpha)P$, it is enough to note that $(1+\eta)P + (\eta/\alpha)(1-\alpha)P = (1+\eta/\alpha)P$. This concludes the proof of the claim.

Claim 2.15. Assume that $o \in P$ is a convex set. Let o' be the center of John ellipsoid E of P. Then $o' \in (1 - \frac{1}{n})P$.

Proof. John ellipsoid has the following property. $(E-o')\subset (P-o')\subset n(E-o')$. In particular, by symmetry of E, we have $o-o'\in n(E-o')=-n(E-o')$, which, is equivalent to $o'\in \frac{1}{n+1}o+\frac{n}{n+1}E$. As $o\in P$ and $E\subset P$, this implies $o'\in \frac{n}{n+1}P$.

We now return to the proof of Lemma 2.12. Let $P \subset \mathbb{R}^n$ convex $o \in P \subset X, Y \subset (1+\eta)P$ using that X, Y is an η sandwich. Let o' be the center of the John ellipsoid E in P. Let P' = P - o', E' = E - o', X' = X - o' and Y' = Y - o'.

By Claim 2.15, we get $o' \in (1 - \frac{1}{n})P$. By Claim 2.14 we get $o \in P' \subset X', Y' \subset (1 + n\eta)P'$. In particular, we deduce that X', Y' is a $n\eta$ -sandwich.

By construction, we get that E' is the John ellipsoid of P' and it is centered at o. It has the property that $E' \subset P' \subset nE'$. It follows that $E' \subset X', Y' \subset (n+n^2\eta)E'$.

As E' is an ellipsoid centered at the origin o, there exists a linear transformation $\theta \colon \mathbb{R}^n \to \mathbb{R}^n$ such that $\theta(E')$ is a ball centered at the origin. In particular, we get that $\theta(E') \subset \theta(X'), \theta(Y') \subset (n+n^2\eta)\theta(E')$.

By observing that the John ellipsoid of the regular simplex S is a ball centered at the origin, we immediately get that $rS \subset \theta(E') \subset nrS$ for some r > 0. Putting together all of the above, we get that

$$rS \subset \theta(X'), \theta(Y') \subset (n^2 + n^3 \eta)rS.$$

We conclude that $\theta(X - o')$, $\theta(Y - o')$ is $(n^2 + n^3\eta)$ -bounded. As sandwiches are preserved under linear transformations, we also conclude that $\theta(X - o')$, $\theta(Y - o')$ is a $n\eta$ -sandwich.

3 Outline of the proof of the Quadratic Theorem (Theorem 1.5)

In this section, we give an outline of the results contained in the next four sections. We shall assume that Theorem 1.4 has been proved.

Thanks to the simple reduction performed in the previous section, see Section 2 and specifically Proposition 2.9, we can assume that A and B are already pretty convex, are sandwiched between two balls of comparable sizes, and are the finite union of axis aligned cubes. The strategy is now the following.

- 1. Moving A and B slightly we can partition \mathbb{R}^n into n+1 reasonably shaped (not too large or small) convex cones $C \in \mathcal{C}_1$ so that $|A \cap C| = |B \cap C|$ (see Proposition 5.4).
- 2. We will refine this partition into cones using the following procedure. Given a cone C and a codimension-two subspace S, we find a hyperplane $H \supset S$ so that $|C \cap H^{\pm} \cap A| = |C \cap H^{\pm} \cap B|$ (see Proposition 5.5).
- 3. Choosing the codimension-two subspaces carefully, we obtain a partition into convex cones $C \in \mathcal{C}_2$ essentially all of which satisfy the following properties (see Proposition 5.6, the engine of which is Theorem 4.11):
 - $\bullet |A \cap C| = |B \cap C|,$
 - C is the convex hull of few half-lines through the origin,
 - C is very narrow in all but one direction of an orthogonal basis depending on C.

These imply
$$\sum_{C \in \mathcal{C}_2} |t(A \cap C) + (1-t)(B \cap C)| \le |tA + (1-t)B|$$
.

- 4. Removing a negligible part of A and B, we may additionally assume that the sections of C in the narrow directions are completely contained in or disjoint from $A \cap C$ and $B \cap C$ for $C \in \mathcal{C}_2$ (see Proposition 5.7). In some sense this reduces $A \cap C$ and $B \cap C$ to two-dimensional sets, as all information about the sets is captured by $\pi(A \cap C)$ and $\pi(B \cap C)$, where π is the projection along the narrow directions.
- 5. In Proposition 6.8 we construct a bounded family \mathcal{U} of cylinders. All cylinders have the same simplex base, which is contained in a face of T and inside C. Moreover, the cylinders cover a big ball intersected with C (so, in particular, $A \cap C$ and $B \cap C$). Furthermore, for $U \in \mathcal{U}$, we have $|U \cap A \cap C| = \Omega_n(1)|A \cap C|$.
- 6. In Proposition 6.7, for $U \in \mathcal{U}$, we find a matching cylinder V parallel to U such that $|U \cap A \cap C| = |V \cap B \cap C|$ and $|t(U \cap A \cap C) + (1-t)(V \cap A \cap C)| |U \cap A \cap C| \le |t(A \cap C) + (1-t)(B \cap C)| |A \cap C|$. Moreover, the distance between U and V is small, namely $|(U \triangle V) \cap S \cap C| \le O_n(1)\delta^{1/2}t^{-1/2}|S \cap C|$. This allows us to assume U = V and further reduce the problem to the sets $A \cap U \cap C$ and $B \cap V \cap C$ which satisfy $|t(U \cap A \cap C) + (1-t)(V \cap A \cap C)| = (1+\delta')|U \cap A \cap C|$.
- 7. By Theorem 1.4, we find $|\operatorname{co}(U \cap A \cap C) \setminus (U \cap A \cap C)| = O_{n,t}(\delta')|(U \cap A \cap C)|$. We also know that that the sections of C in the narrow directions are completely contained in or disjoint from $A \cap C$ and $B \cap C$. Combining these, we can deduce that $U \cap A \cap C$ and $V \cap B \cap C$ are essentially one-codimensional compressed.
- 8. In Proposition 6.6 we resolve the problem for one-codimensional (in the long direction) compressed sets.

4 Intermediate results for the Quadratic Theorem (Theorem 1.5): Part I

4.1 Setup

Definition 4.1. Let K^n be the family of convex sets in \mathbb{R}^n with a finite number of vertices and let S_k^n be the set of codimension k affine subspaces of \mathbb{R}^n .

Definition 4.2. Say a function $f: \mathcal{K}^n \times \mathcal{S}_2^n \to \mathcal{S}_1^n$ is a respectful function if $L \subset f(K, L)$. A respectful function f induces functions $f^-, f^+: \mathcal{K}^n \times \mathcal{S}_2^n \to \mathcal{K}^n$, where $f^-(K, L), f^+(K, L)$ are the convex sets the affine hyperplane f(K, L) essentially partitions K into.

Definition 4.3. Given a respectful function $f: \mathcal{K}^n \times \mathcal{S}_2^n \to \mathcal{S}_1^n$ and a convex set P, we say \mathcal{F} is a valid partition of P into convex subsets if there exists a sequence of families $\{P\} = \mathcal{G}_0, \ldots, \mathcal{G}_j = \mathcal{F}$ such that if $\mathcal{G}_i = \{P_1, P_2, \ldots, P_k\}$, then there exists codimension-two affine subspaces L_1, \ldots, L_k such that $\mathcal{G}_{i+1} = \bigcup_{j=1}^k R_j$, where $R_j = \{f^+(P_j, L_j), f^-(P_j, L_j)\}$ or $R_j = \{P_j\}$.

We now consider the analogous definitions for cones (cf. Definition 2.2).

Definition 4.4. Let C^n be the family of cones in \mathbb{R}^n and let \mathcal{T}^n_k be the set of codimension k subspaces of \mathbb{R}^n .

Definition 4.5. Say a function $f: \mathcal{C}^n \times \mathcal{T}_2^n \to \mathcal{T}_1^n$ is a respectful function if $L \subset f(C, L)$. A respectful function f induces functions $f^-, f^+: \mathcal{C}^n \times \mathcal{T}_2^n \to \mathcal{C}^n$, where $f^-(C, L), f^+(C, L)$ are the cones the hyperplane f(C, L) partitions C into.

Definition 4.6. Given a respectful function $f: \mathcal{C}^n \times \mathcal{T}_2^n \to \mathcal{T}_1^n$ and a cone C, we say \mathcal{F} is a valid partition of C into cones if there exists a sequence of families $\{C\} = \mathcal{G}_0, \ldots, \mathcal{G}_j = \mathcal{F}$ such that if $\mathcal{G}_i = \{C_1, C_2, \ldots, C_k\}$, then there exists codimension-two subspaces L_1, \ldots, L_k such that $\mathcal{G}_{i+1} = \bigcup_{j=1}^k R_j$, where $R_j = \{f^+(C_j, L_j), f^-(C_j, L_j)\}$ or $R_j = \{C_j\}$.

Definition 4.7. Given a cone $C \subset \mathbb{R}^n$, define

$$\mu_n(C) := |C \cap S|,$$

where S is the unit volume simplex from Definition 2.4.

Definition 4.8. A function $f: C \to \mathcal{K}^n$ from some convex domain $C \subset \mathbb{R}^m$ is linear if f(tx + (1-t)y) = tf(x) + (1-t)f(y) for all $x, y \in C$.

For example, the function $f: \mathbb{R}^2_{\geq 0} \to \mathcal{K}^2$ given by $f(x,y) = [x/2,3x/2] \times [y,3y]$ is linear. Note that given a linear function $f: C \to \mathcal{K}^n$ with $C \in \mathcal{C}^m$ and $f(o) = \{o\}$, the set $\bigcup_{x \in C} f(x) \times \{x\}$ is a cone in \mathcal{C}^{m+n} .

Definition 4.9. A cone $C \in \mathcal{C}^n$ is (i, ℓ, ε) -good if there exists a basis e_1^C, \ldots, e_n^C and there exists a cone $C' \in \mathcal{C}^i$ and a linear function $f: C' \to \mathcal{K}^{n-i}$ such that f(x) has at most ℓ vertices for all $x \in C'$,

$$\sup_{y,z\in f(x)}||y-z||_2\leq \varepsilon||x||_2, \qquad and \qquad C=\cup_{x\in C'}\{f(x)\times\{x\}\}.$$

Definition 4.10. When a basis is established, let $\pi_i : \mathbb{R}^n \to \mathbb{R}$ be the projection onto the ith coordinate and let $\pi_{i,j} : \mathbb{R}^n \to \mathbb{R}^2$ be the projection onto the plane spanned by the ith and jth coordinate.

4.2 Theorem

In this section, we will prove the following theorem. Recall from Definition 2.4 that S is a regular simplex with unit volume. Let F_0, \ldots, F_n be the faces of S defined by $F_i = C_i \cap \partial S$ where $C_i \in \mathfrak{C}$.

Theorem 4.11. There exist constants $\ell_n^{4.11}(m)$ such that for every $\varepsilon > 0$ the following holds. Given a respectful $f: \mathcal{C}^n \times \mathcal{T}_2^n \to \mathcal{T}_1^n$ and a cone C defined by m lines which is a subcone of some $C_i \in \mathfrak{C}$, there exists a valid partition \mathcal{F} of C that can be written as $\mathcal{F} = \mathcal{F}_0 \sqcup \mathcal{F}_1 \sqcup \mathcal{F}_2$ such that

- 1. $\sum_{F \in \mathcal{F}_0} \mu_n(F) \leq \varepsilon$.
- 2. Every cone $F \in \mathcal{F}_1$ is $(1, \ell_n^{4.11}(m), \varepsilon)$ -good.

3. For every cone $F \in \mathcal{F}_2$ there exists a sub-cone F' of F with $\mu_n(F') \geq (1 - \varepsilon)\mu_n(F)$ such that F' is $(2, \ell_n^{4.11}(m), \varepsilon)$ -good.

Furthermore, given H + v the affine hyperplane containing F_i , where H is a hyperplane through the origin, we can insist that for all cones $F \in \mathcal{F}_1$ we have $e_n^F \in H^{\perp}$ and for all $F \in \mathcal{F}_2$ we have $e_n^{F'} \in H^{\perp}$.

We note that a slightly more general result holds where we drop S. The motivation to include S comes from the way we apply the theorem.

4.3 Propositions

To prove Theorem 4.11, we first state some propositions and lemmas that will be used in the proofs. All these results will be proved later below.

Proposition 4.12. There exists a constant $\ell_2^{4.12}$ such that for every $\varepsilon > 0$ the following holds. Given a respectful $f: \mathcal{K}^2 \times \mathcal{S}_2^2 \to \mathcal{S}_1^2$ and convex set P, there exists a valid partition \mathcal{F} of P that can be written as $\mathcal{F} = \mathcal{F}_0 \sqcup \mathcal{F}_1 \sqcup \mathcal{F}_2$ such that

- 1. $\sum_{F \in \mathcal{F}_0} |F| \leq \varepsilon$.
- 2. For every $F \in \mathcal{F}_1$ we have $|V(F)| \leq \ell_2^{4.12}$ and for a basis e_1, e_2 and for i = 1, 2 we have that $|\pi_i(F)| \leq \varepsilon$,
- 3. For every $F \in \mathcal{F}_2$ we have $|V(F)| \leq \ell_2^{4.12}$ there exists a basis (dependent on F) e_1, e_2 such that we have $|\pi_1(F)| \leq \varepsilon$ and $|\pi_2(F)| \geq \varepsilon$ and $\pi_2(V(F)) \subset V(\pi_2(F)) + (-\varepsilon^2, \varepsilon^2)$.

Here $\pi_i : \mathbb{R}^n \to \mathbb{R}$ is the projection onto the *i*-th coordinate (as in Definition 4.10).

Proposition 4.13. There exists a constant $\ell_n^{4.13}(m)$ such that for every $\varepsilon > 0$ the following holds. Given a respectful $f: \mathcal{K}^n \times \mathcal{S}_2^n \to \mathcal{S}_1^n$ and convex set P with $|V(P)| \leq m$, there exists a valid partition \mathcal{F} of P that can be written as $\mathcal{F} = \mathcal{F}_0 \sqcup \mathcal{F}_1 \sqcup \mathcal{F}_2$ such that

- 1. $\sum_{F \in \mathcal{F}_0} |F| \leq \varepsilon$.
- 2. For every $F \in \mathcal{F}_1$ we have $|V(F)| \leq \ell_n^{4.13}(m)$ and for a basis e_1, e_2, \ldots, e_n we have for every $i \mid \pi_i(F) \mid \leq \varepsilon$.
- 3. For every $F \in \mathcal{F}_2$ we have $|V(F)| \leq \ell_n^{4.13}(m)$ and there exists a basis (dependent on F) e_1, e_2, \ldots, e_n such that for every $i \neq n$ we have $|\pi_i(F)| \leq \varepsilon$ and $|\pi_n(F)| \geq \varepsilon$ and $\pi_n(V(F)) \subset V(\pi_n(F)) + (-\varepsilon^2, \varepsilon^2)$.

4.4 Auxiliary Lemmas

Lemma 4.14. There exist constants $\lambda_n^{4.14} > 0$ so that given a convex set $P \subset \mathbb{R}^n$ and an affine subspace $L' \in \mathcal{S}_2^n$, there exists a translate $L \in \mathcal{S}_2^n$ of L' so that any affine hyperplane $H \supset L$ essentially partitioning the space into two parts H^+ and H^- satisfies

$$\frac{|H^+\cap P|}{|H^-\cap P|}\in \left[\lambda_n^{4.14},(\lambda_n^{4.14})^{-1}\right].$$

Lemma 4.15. For any convex body $P \subset \mathbb{R}^2$ with at least 7 vertices, there exists a point $p \in P$, so that for any line $\ell \ni p$ essentially partitioning the plane into halfplanes ℓ^+ and ℓ^- , we have that both convex sets $\ell^+ \cap P$ and $\ell^- \cap P$ have fewer vertices than P.

Lemma 4.16. Let $P \subset \mathbb{R}^2$ be a convex set with at most C vertices and a respectful function $f: \mathcal{K}^2 \times \mathbb{R}^2 \to \mathcal{S}_1^2$. Then there exists a valid partition of P into at most 2^{C-6} parts so that all parts have at most 6 vertices.

Lemma 4.17. Let $P \subset \mathbb{R}^n$ be a convex set. For any $\xi > 0$, there exist $\zeta_0, \eta_0 > 0$ (depending on P) so that for any $\zeta < \zeta_0$ and $\eta < \eta_0$ the following holds. Say a line $L \subset \mathbb{R}^n$ is ζ permissible if for one of the points $x \in L \cap \partial P$, there exists a line L_x tangent to P at x with $\angle L, L_x \leq \zeta$. Let

$$Q_{\zeta,\eta} := P \cap \bigcup_{L:\zeta \ permissible} L + B(o,\eta),$$

then $Q_{\zeta,\eta} \subset \partial P + B(o,\xi)$

Lemma 4.18. There exists a constant $\alpha > 0$, so that for any $\varepsilon > 0$ the following holds. Given a basis e_1, e_2 in \mathbb{R}^2 , a convex set $P \subset \mathbb{R}^2$ with at most six vertices and a respectful $f: \mathcal{K}^2 \times \mathbb{R}^2 \to \mathcal{S}_1^2$, there exists a valid partition \mathcal{G} of P and an element $P' \in \mathcal{G}$ so that $|P'| \geq \alpha |P|$ and

- either $|\pi_2(P')| < |\pi_2(P)| \varepsilon^2$,
- or $\pi_2(V(P')) \subset V(\pi_2(P')) + (-\varepsilon^2, \varepsilon^2)$.

Furthermore, all convex sets in G have at most 10 vertices.

4.5Proof of Theorem 4.11

Theorem 4.11 can be considered as Proposition 4.13 coned off at the origin.

Proof of Theorem 4.11. Let $\ell_n^{4.11}(m) := (\ell_{n-1}^{4.13}(m))^2$. Let $c_n^{4.11}$ be 1/n+1 times the radius of the largest ball inside T. Choose $\eta > 0$ sufficiently small in terms of ε .

Let $P := (v + H) \cap C \in \mathcal{K}^{n-1}$. For any subset $P' \subset v + H$, we define $C(P') := \bigcup_{t \geq 0} tP'$. Note that if P'is a bounded convex set in H+v, then C(P') is a cone. In particular, C=C(P). Note moreover that for a codimension 2 affine subspace L of v + H, C(L) is a codimension 2 subspace of \mathbb{R}^n .

In the following we will slightly abuse notation to interpret v + H as a copy \mathbb{R}^{n-1} to aid the application of

Proposition 4.13. For instance, we will write \mathcal{K}^{n-1} to indicate the convex subsets of v+H.

We construct a respectful function $f': \mathcal{K}^{n-1} \times \mathcal{S}_2^{n-1} \to \mathcal{S}_1^{n-1}$ from $f: \mathcal{C}^n \times \mathcal{T}_2^n \to \mathcal{T}_1^n$ as follows. For any convex subset $P' \subset P$ and a codimension 2 affine subspace L of v+H, define $f'(P',L) = (v+H) \cap f(C(P'),C(L))$. Note that indeed

$$L = (v + H) \cap C(L) \subset (v + H) \cap f(C(P'), C(L)) = f'(P', L),$$

so that f' is respectful.

Now apply Proposition 4.13 to P with respectful function f' and parameter η (as ε), to find valid partition $\mathcal{F}' = \mathcal{F}'_0 \cup \mathcal{F}'_1 \cup \mathcal{F}'_2$ of P, where all $P' \in \mathcal{F}'_1 \cup \mathcal{F}'_2$ have at most $\ell_{n-1}^{4.13}(m)$ vertices. Define $\mathcal{F} = \mathcal{F}_0 \cup \mathcal{F}_1 \cup \mathcal{F}_2$ by $\mathcal{F}_i := \{C(P') : P' \in \mathcal{F}'_i\}$. Note that clearly \mathcal{F} is a valid partition by the construction of f', so it remains to check that it indeed satisfies the conditions on \mathcal{F}_0 , \mathcal{F}_1 , and \mathcal{F}_2 . By definition of H, we find that $\mu_n(C(P')) = c_n^{4.11}|P'|$, so that

$$\sum_{F \in \mathcal{F}_0} \mu_n(F) = c_n^{4.11} \sum_{P' \in \mathcal{F}'_0} |P'| \le c_n^{4.11} \eta \le \varepsilon,$$

as $c_n^{4.11}$ represents 1/(n+1) of the distance from o to F_i .

Consider $F \in \mathcal{F}_1$ and choose a basis e_1, \ldots, e_n with e_n perpendicular to H. Let $P' \in \mathcal{F}'_1$, so that F = C(P'). To show that F is $(1, \ell, \varepsilon)$ -good, let $C' = \mathbb{R}_{>0}e_n$ and $g: C' \to \mathcal{K}^{n-1}$, $te_n \mapsto (tv + H) \cap F = tP'$, which is clearly linear. Note that $g(te_n)$ is homothetic for all t, so always has the same number of vertices (as P'), in particular at most $\ell_{n-1}^{4.13}(m) \le \ell_n^{4.11}(m)$. Note

$$\sup_{x,y \in g(te_n)} ||x - y||_2 = \sup_{x,y \in tP'} ||x - y||_2 \le t\sqrt{n - 1}\eta \le \varepsilon ||te_n||_2.$$

This concludes that F is $(1, \ell, \varepsilon)$ -good.

Consider $F \in \mathcal{F}_2$ and the $P' \in \mathcal{F}'_2$, so that F = C(P'). Let e_n be the unit vector orthogonal to H, and let e_1, \ldots, e_{n-1} be the basis of H so that $|\pi_i(P')| \leq \eta$ for $i \leq n-2$ and $|\pi_{n-1}(P')| \geq \eta$ and $\pi_{n-1}(V(P')) \subset \eta$ $V(\pi_{n-1}(P')) + (-\eta^2, \eta^2)$. Let

$$P'' := P' \cap \pi_{n-1}^{-1} \left(\pi_{n-1}(P') \setminus \left[V(\pi_{n-1}(P')) + \left(-\eta^2, \eta^2 \right) \right] \right),$$

so that by construction $\pi_{n-1}(V(P'')) = V(\pi_{n-1}(P''))$. By convexity, we have $|P' \setminus P''| \leq O_n(\eta |P'|)$. Now let $F' := C(P'') \subset F$, and note that $\mu_n(F') \geq (1 - \varepsilon)\mu_n(F)$. Let $C' := C(\pi_{n-1,n}(F')) = \pi_{n-1,n}(F')$, where $\pi_{n-1,n} : \mathbb{R}^n \to \mathbb{R}^2$ is the projection onto the last two coordinates and let

$$g\colon C'\to \mathcal{K}^{n-2}, x\mapsto \pi_{n-1,n}^{-1}(x)\cap F'.$$

To see that g is linear, consider the auxiliary function

$$g' : \pi_{n-1}(P'') \to \mathcal{K}^{n-2}, x \mapsto \pi_{n-1}^{-1}(x) \cap P''$$

and note that g' is linear because $\pi_{n-1}(V(P'')) = V(\pi_{n-1}(P''))$. Let h > 0 be such that $\pi_n(v) = h$ and note that $\pi_n^{-1}(th) \cap F' = tP''$, so that for a point $(x,y) \in \pi_{n-1,n}(F'')$, we have $g(x,y) = \frac{y}{h}g'(\frac{hx}{y})$. Hence, we get for $(x,y), (x',y') \in C'$, that

$$hg(tx + (1-t)x', ty + (1-t)y') = (ty + (1-t)y')g'\left(\frac{tx + (1-t)x'}{ty + (1-t)y'}h\right)$$

$$= (ty + (1-t)y')\left(g'\left(\frac{ty}{ty + (1-t)y'} \cdot \frac{x}{y}h + \frac{(1-t)y'}{ty + (1-t)y'} \cdot \frac{x'}{y'}h\right)\right)$$

$$= (ty + (1-t)y')\left(\frac{ty}{ty + (1-t)y'}g'\left(\frac{x}{y}h\right) + \frac{(1-t)y'}{ty + (1-t)y'}g'\left(\frac{x'}{y'}h\right)\right)$$

$$= tyg'\left(\frac{x}{y}h\right) + (1-t)y'g'\left(\frac{x'}{y'}h\right)$$

$$= thg(x,y) + (1-t)hg(x',y'),$$

so that g is linear. Finally, to count the number of vertices of g(x,y), note that it is the intersection between a homothetic copy of P'' and a hyperplane, so that the number of vertices is bounded by the number of edges in P'' which is at most $\binom{\ell^4.13}{n-1} \binom{m}{2} \le \ell^4.11 \binom{m}{n}$.

4.6 Proof of Propositions

4.6.1 Proof of Proposition 4.12

Note that $S_2^2 = \mathbb{R}^2$.

Proof of Proposition 4.12. Choose $\ell_2^{4.12} = 10$. We first show that we may assume that $|\pi_1(P)| \leq \varepsilon$, per the following claim.

Claim 4.19. The result for convex sets P with $|\pi_1(P)| \le \varepsilon$ implies the result for general convex P.

Proof of claim. Iteratively produce a valid partition of P, starting with $\mathcal{G}_0 = \{P\}$. Given $\mathcal{G}_i = \{P_1, \dots, P_{2^i}\}$, by Lemma 4.14, find points $p_j \in P_j$ and construct

$$\mathcal{G}_{i+1} := \{ f^+(P_1, p_1), f^-(P_1, p_1), \dots, f^+(P_{2^i}, p_{2^i}), f^-(P_{2^i}, p_{2^i}) \}.$$

Now note that by construction $\max\{|P'|: P' \in \mathcal{G}_{i+1}\} \leq (1-\lambda_2) \max\{|P'|: P' \in \mathcal{G}_i\}$, where $\lambda_2 > 0$ is the constant from Lemma 4.14. Hence, for i_0 sufficiently large, we find that $\max\{|P'|: P' \in \mathcal{G}_{i_0}\} < \frac{\varepsilon^2}{8}$. This implies that for all $P' \in \mathcal{G}_{i_0}$ there exists a basis e_1, e_2 so that $|\pi_1(P')| \leq \varepsilon$. Indeed, let E' be the outer Lowner-John ellipsoid of P' and let e_1, e_2 be the short and long axis of E'. Hence, $|\pi_1(E')|^2/2 \leq |\pi_1(E')||\pi_2(E')|/2 \leq |E'| \leq 4|P| \leq \frac{\varepsilon^2}{2}$. Therefore, $|\pi_1(P')| \leq |\pi_1(E')| \leq \varepsilon$. Now applying the result in each $P' \in \mathcal{G}_{i_0}$, we deduce the result in P.

Henceforth, assume $\pi_1(P) \leq \varepsilon$. We iteratively construct a sequence of valid partitions \mathcal{G}_i and \mathcal{G}'_i of P, starting with $\mathcal{G}_0 = \{P\}$.

Given \mathcal{G}_i , apply Lemma 4.16 to all elements of \mathcal{G}_i to find refinement \mathcal{G}_i' in which all elements have at most 6 vertices. Then apply Lemma 4.18 to each of the elements of \mathcal{G}_i' to produce refinement \mathcal{G}_{i+1} . Given this sequence we will construct a valid partition of P with the desired properties. Consider the uniform probability measure \mathbb{P} on P, so that $\sum_{P' \in \mathcal{G}_i} \mathbb{P}(P') = 1$ for all i. we are going to analyse the change in expected value of the following parameter. Let $f_0(P) = |\pi_2(P)|$. Given $f_i : \mathcal{G}_i \to \mathbb{R}$, construct $f_{i+1} : \mathcal{G}_{i+1} \to \mathbb{R}$ as follows.

$$f_{i+1}(P') := \begin{cases} 0 & \text{if } P' \subset P'' \in \mathcal{G}_i \text{ with } f_i(P'') = 0, \text{ or if } \pi_2(V(P')) \subset V(\pi_2(P')) + (-\varepsilon^2, \varepsilon^2) \\ |\pi_2(P')| & \text{otherwise.} \end{cases}$$

Note that if $|\pi_2(P')| < 2\varepsilon^2$, then $f_i(P') = 0$ by the second clause, so if $f_i(P') \neq 0$, then $f_i(P') \geq 2\varepsilon^2$. In terms of this function, we construct our families $\mathcal{F}_0^i, \mathcal{F}_1^i, \mathcal{F}_2^i$, as follows; $\mathcal{F}_0^0 = \{P\}, \mathcal{F}_1^0 = \emptyset, \mathcal{F}_2^0 = \emptyset$, and

- $\mathcal{F}_0^{i+1} = \{ P' \in \mathcal{G}_{i+1} : f_{i+1}(P') > 0 \}$
- $\mathcal{F}_1^{i+1} = \mathcal{F}_1^i \cup \{P' \in \mathcal{G}_{i+1} : f(P') = 0, |\pi_2(P')| \le \varepsilon, \text{ and } \exists P'' \in \mathcal{F}_0^i, P' \subset P''\}$
- $\mathcal{F}_2^{i+1} = \mathcal{F}_2^i \cup \{P' \in \mathcal{G}_{i+1} : f(P') = 0, |\pi_2(P')| > \varepsilon, \text{ and } \exists P'' \in \mathcal{F}_0^i, P' \subset P''\}$

It is easy to see that $\mathcal{F}^i = \mathcal{F}^i_0 \sqcup \mathcal{F}^i_1 \sqcup \mathcal{F}^i_2$ is a valid partition of P. We will show that for i sufficiently large \mathcal{F}^i satisfies the desired the conditions. We use shorthand $\mathbb{P}\left(\mathcal{F}^i_0\right)$ for $\mathbb{P}\left(\bigcup_{P'\in\mathcal{F}^i_0}P'\right) = \sum_{P'\in\mathcal{F}^i_0}\mathbb{P}(P')$.

Claim 4.20. $\mathbb{P}\left(\mathcal{F}_0^i\right) \to 0 \text{ as } i \to \infty.$

Proof of claim. First note that f_i is non-negative and non-increasing, in the sense that if $P' \in \mathcal{G}_i$ and $P'' \in \mathcal{G}_j$ with $P' \subset P''$, then $f_j(P'') \geq f_i(P')$. Partition \mathcal{G}_i into two parts \mathcal{F}_0^i and $\mathcal{H}_i := \mathcal{G}_i \setminus \mathcal{F}_0^i$ (so that \mathcal{H}_i is a refinement of $\mathcal{F}_1^i \cup \mathcal{F}_2^i$). Consider some $P' \in \mathcal{F}_0^i$ (i.e., with $f_i(P') > 2\varepsilon^2$) and the valid partition $\{P_1, \ldots, P_j\} \subset \mathcal{G}_i'$ of P'. Note that by Lemma 4.18, for every $1 \leq k \leq j$, we can find $P_k \supset P_k' \in \mathcal{G}_{i+1}$ with $\mathbb{P}(P_k') \geq \alpha \mathbb{P}(P_k)$ and

either
$$|\pi_2(P_k')| \le |\pi_2(P_k)| - \varepsilon^2$$
, or $\pi_2(V(P_k')) \subset V(\pi_2(P_k')) + (-\varepsilon^2, \varepsilon^2)$.

Either way, we find $f_{i+1}(P'_k) \leq f_i(P') - \varepsilon^2$. Hence, if we let $\mathcal{G}_{i+1}(P') := \{P'' \in \mathcal{G}_{i+1} : P'' \subset P'\}$, then

$$\sum_{P'' \in \mathcal{G}_{i+1}(P')} \mathbb{P}(P'') f_{i+1}(P'') \le \mathbb{P}(P') f_i(P') - \alpha \mathbb{P}(P') \varepsilon^2.$$

Summing this over all $P' \in \mathcal{F}_0^i$ and using induction, we find

$$0 \leq \sum_{P'' \in \mathcal{G}_{i+1}} \mathbb{P}(P'') f_{i+1}(P'') \leq \sum_{P' \in \mathcal{G}_i} \mathbb{P}(P') f_i(P') - \alpha \varepsilon^2 \mathbb{P}\left(\mathcal{F}_0^i\right) \leq f_0(P) - \alpha \varepsilon^2 \sum_{j=0}^i \mathbb{P}\left(\mathcal{F}_0^i\right).$$

This implies $\mathbb{P}\left(\mathcal{F}_0^i\right) \to 0$ as $i \to 0$ and thus the conclusion follows.

By this claim, we can find i_0 so that $\mathbb{P}\left(\mathcal{F}_0^{i_0}\right) \leq \frac{\varepsilon}{|P|}$, thus $\sum_{P' \in \mathcal{F}_0^{i_0}} |P'| = \mathbb{P}\left(\mathcal{F}_0^{i_0}\right) |P| \leq \varepsilon$. Hence, let $\mathcal{F}_0 = \mathcal{F}_0^{i_0}, \mathcal{F}_1 = \mathcal{F}_1^{i_0}$, and $\mathcal{F}_2 = \mathcal{F}_2^{i_0}$. By construction, we have for every $F \in \mathcal{F}_1$, $|\pi_1(F)|, |\pi_2(F)| \leq \varepsilon$ and that, for $F \in \mathcal{F}_2$,

$$|\pi_1(F)| \le \varepsilon$$
, $|\pi_2(F)| > \varepsilon$, and $\pi_2(V(F)) \subset V(\pi_2(F)) + (-\varepsilon^2, \varepsilon^2)$.

Finally, by Lemma 4.18 all parts in \mathcal{G}_i have at most 10 vertices, so in particular so do the parts in $\mathcal{F}_1 \sqcup \mathcal{F}_2$. \square

4.6.2 Proof of Proposition 4.13

Proof of Proposition 4.13. Let $\xi \gg \zeta \gg \eta$ be chosen sufficiently small in terms of ε and n to make various statements throughout the proof.

We first find a valid partition so that all parts are small in all but one direction.

We iteratively construct a sequence of valid partitions \mathcal{G}^i for i = 0, ..., n, starting with $\mathcal{G}^0 = \{P\}$. Each of the \mathcal{G}^i 's can be partitioned into two parts \mathcal{G}_0^i , and \mathcal{G}_1^i , so that the following hold

- 1. $\sum_{F \in \mathcal{G}_0^i} |F| \leq i\eta$
- 2. For every $F \in \mathcal{G}_1^i$, there exists a basis e_1, \ldots, e_n so that $|\pi_i(F)| \leq \eta$ for all $1 < j \leq i$.
- 3. For every $F \in \mathcal{G}_0^i \cup \mathcal{G}_1^i$, we have that $F = P \cap F'$ where F' is the intersection of at most $i\ell_2^{4.12}$ halfspaces, each of which contains all but two of the basis vectors corresponding to F.

Assume that $\mathcal{G}_0^i, \mathcal{G}_1^i$ have been constructed. Fix a $F \in \mathcal{G}_1^i$ and the corresponding basis e_1, \ldots, e_n . Consider the plane H spanned by e_1 and e_{i+1} and the projection $\pi = \pi_{1,i+1}$ onto that plane. Now note that any translate S of the codimension-two subspace of \mathbb{R}^n spanned by $e_2, \ldots, e_i, e_{i+2}, \ldots, e_n$, has that $\pi(S)$ is a single point in \mathbb{R}^2 . Hence, the respectful function $f: \mathcal{K}^n \times \mathcal{S}_2^n \to \mathcal{S}_1^n$ corresponds to a respectful function $f': \mathcal{K}^2 \times \mathcal{S}_2^2 \to \mathcal{S}_1^2$. Indeed, for $X \in \mathcal{K}^2$ and $X \in \mathcal{S}_2^2 = \mathbb{R}^2$, let

$$f'(X,x) := \pi(f(\pi^{-1}(X) \cap F, \pi^{-1}(x))).$$

Now apply Proposition 4.12 to $\pi(F)$ with respectful function f' and parameter $\eta \cdot \min \left\{ 1, \frac{1}{\max_{x \in \mathbb{R}^2} |\pi^{-1}(x) \cap F|} \right\}$, to find a valid partition $\mathcal{H} = \mathcal{H}_0 \sqcup \mathcal{H}_1 \sqcup \mathcal{H}_2$ of $\pi(F)$ so that

- 1. $\sum_{Q \in \mathcal{H}_0} |Q| \le \frac{\eta}{\max_{x \in \mathbb{R}^2} |\pi^{-1}(x) \cap F|}.$
- 2. For every $Q \in \mathcal{H}_1$ there exists a basis e'_1, e'_2 such that for i = 1, 2 we have $|\pi'_i(Q)| \leq \eta$.

- 3. For every $Q \in \mathcal{H}_2$ there exists a basis e_1', e_2' such that we have $|\pi_1'(Q)| \leq \eta$, $|\pi_2'(Q)| \geq \eta$ and $\pi_2'(V(Q)) \subset V(\pi_2'(Q)) + (-\eta^2, \eta^2)$.
- 4. For every $Q \in \mathcal{H}_1 \cup \mathcal{H}_2$, we have $|V(Q)| \leq \ell_2^{4.12}$, i.e., Q is the intersection of at most ℓ_2 halfplanes.

Note that 2 and 3, imply the weaker statement that for every $Q \in \mathcal{H}_1 \cup \mathcal{H}_2$, there exists a basis e'_1, e'_2 such that we have $|\pi'_1(Q)| \leq \eta$. The partition \mathcal{H} naturally corresponds to a valid partition $\mathcal{H}' = \mathcal{H}'_0 \sqcup \mathcal{H}'_1 \sqcup \mathcal{H}'_2$ of F, where $\mathcal{H}'_i := \{\pi^{-1}(Q) \cap F : Q \in \mathcal{H}_i\}$. This is a valid partition by construction of f'. The properties of \mathcal{H} translate to the following properties of \mathcal{H}' .

- 1. $\sum_{Q \in \mathcal{H}_0'} |Q| \le \eta.$
- 2. For every $Q \in \mathcal{H}'_1 \cup \mathcal{H}'_2$ there exists a basis e'_1, e'_{i+1} of the plane spanned by e_1 and e_{i+1} such that we have $|\pi'_{i+1}(Q)| \leq \eta$.
- 3. For every $Q \in \mathcal{H}'_1 \cup \mathcal{H}'_2$, we have $Q = F \cap Q'$, where Q' is the intersection of at most $\ell_2^{4.12}$ halfspaces all of which contain $e_2, \ldots, e_i, e_{i+2}, \ldots, e_n$.

Now for Q, we choose the basis $e'_1, e_2, \ldots, e_i, e'_{i+1}, e_{i+2}, \ldots, e_n$. Given $F \in \mathcal{G}_1^i$, let $\mathcal{H}'_0(F)$, $\mathcal{H}'_1(F)$, and $\mathcal{H}'_2(F)$ be the sets produced here. We define

$$\mathcal{G}_0^{i+1} := \mathcal{G}_0^i \cup \bigcup_{F \in \mathcal{G}_1^i} \mathcal{H}_0'(F), \quad \text{ and } \quad \mathcal{G}_1^{i+1} := \bigcup_{F \in \mathcal{G}_1^i} \mathcal{H}_1'(F) \cup \mathcal{H}_2'(F).$$

Note that these satisfy the properties 1, 2 and 3 of \mathcal{G}_0^{i+1} , \mathcal{G}_1^{i+1} set out above.

Now consider the elements of \mathcal{G}_1^n . These can only be 'long' in at most one direction, viz e_1 . Now that we have established this direction, we will repeat essentially the same process to show that the sets are either short in the e_1 direction as well, or (most of) the vertices are close to the extremes in the e_1 direction.

We continue constructing a sequence of valid partitions \mathcal{G}^{n+i} for i = 1, ..., n, starting with $\mathcal{G}^0 = \{P\}$. Each of the \mathcal{G}^{n+i} 's can be partitioned into three parts $\mathcal{G}_0^{n+i}, \mathcal{G}_1^{n+i}, \mathcal{G}_2^{n+i}$, so that the following hold

- 1. $\sum_{F \in \mathcal{G}_0^{n+i}} |F| \le n\eta + i\zeta.$
- 2. For every $F \in \mathcal{G}_1^{n+i}$, there exists a basis e_1, \ldots, e_n so that $|\pi_j(F)| \leq \zeta$ for all $j = 1, \ldots n$.
- 3. For every $F \in \mathcal{G}_2^{n+i}$, there exists a basis e_1, \ldots, e_n so that $|\pi_j(F)| \leq \eta$ for all $j = 2, \ldots n$. Moreover, we have $\pi_1(V(\pi_{1,j}(F))) \subset V(\pi_1(F)) + (-2\zeta^2, 2\zeta^2)$ for all $j = 2, \ldots, i$.
- 4. For every $F \in \mathcal{G}_1^{n+i} \cup \mathcal{G}_2^{n+i}$, we have that $F = P \cap F'$ where F' is the intersection of at most $(n+i)\ell_2^{4.12}$ halfspaces, each of which contains all but two of the basis vectors corresponding to F.

Assume that \mathcal{G}_0^{n+i} , \mathcal{G}_1^{n+i} , \mathcal{G}_2^{n+i} have been constructed. Fix a $F \in \mathcal{G}_2^{n+i}$ and the corresponding basis e_1, \ldots, e_n . As before, consider the plane spanned by e_1 and e_{i+1} and the projection $\pi = \pi_{1,i+1}$ onto that plane. Now note that any translate L of the codimension-two subspace of \mathbb{R}^n spanned by $e_2, \ldots, e_i, e_{i+2}, \ldots, e_n$, has that $\pi(L)$ is a single point in \mathbb{R}^2 . Hence, the respectful function $f: \mathcal{K}^n \times \mathcal{S}_2^n \to \mathcal{S}_1^n$ corresponds to a respectful function $f': \mathcal{K}^2 \times \mathcal{S}_2^2 \to \mathcal{S}_1^2$. Indeed, for $X \in \mathcal{K}^2$ and $x \in \mathcal{S}_2^2 = \mathbb{R}^2$, let

$$f'(X,x) := \pi(f(\pi^{-1}(X) \cap F, \pi^{-1}(x))).$$

Now apply Proposition 4.12 to $\pi(F)$ with respectful function f' and parameter $\zeta \cdot \min \left\{ 1, \frac{1}{\max_{x \in \mathbb{R}^2} |\pi^{-1}(x) \cap F|} \right\}$, to find a valid partition $\mathcal{H} = \mathcal{H}_0 \sqcup \mathcal{H}_1 \sqcup \mathcal{H}_2$ of $\pi(F)$ so that

- 1. $\sum_{Q \in \mathcal{H}_0} |Q| \le \frac{\zeta}{\max_{x \in \mathbb{R}^2} |\pi^{-1}(x) \cap F|}.$
- 2. For every $Q \in \mathcal{H}_1$ there exists a basis e'_1, e'_2 such that for i = 1, 2 we have $|\pi'_i(Q)| \leq \zeta$.
- 3. For every $Q \in \mathcal{F}_2$ there exists a basis e_1', e_2' such that we have $|\pi_2'(Q)| \leq \zeta$, $|\pi_1'(Q)| \geq \zeta$ and $\pi_1'(V(Q)) \subset V(\pi_1'(Q)) + (-\zeta^2, \zeta^2)$.
- 4. For every $Q \in \mathcal{H}_1 \cup \mathcal{H}_2$, we have $|V(Q)| \leq \ell_2^{4.12}$, i.e., Q is the intersection of at most $\ell_2^{4.12}$ halfplanes.

Let Q satisfy property 3. Given the information we already have about Q (viz $|\pi_2(Q)| \leq \eta$), we will show that we can get essentially the same property 3, with basis e'_1, e'_2 replaced by e_1, e_2 .

Claim 4.21. $|\pi_2(Q)| \le \zeta/2$, $|\pi_1(Q)| \ge \zeta/2$ and $\pi_1(V(Q)) \subset V(\pi_1(Q)) + (-4\zeta^2, 4\zeta^2)$.

Proof of claim. First note that, up to translation, in the e_1, e_2 basis we have $Q \subset [0, |\pi_1(Q)|] \times [0, |\pi_2(Q)|]$. Hence, we get

$$\zeta \le |\pi_1'(Q)| \le |\pi_1(Q)| + |\pi_2(Q)| \le |\pi_1(Q)| + \eta,$$

so $|\pi_1(Q)| \ge \zeta - \eta \ge \zeta/2$. Also, $|\pi_2(Q)| \le \eta \le \zeta/2$.

For the last part, it is enough to show that there exists two vertices $x, y \in V(Q)$ such that for any other vertex $z \in V(Q)$ we have $\min(|xz|, |yz|) < 2\zeta^2$.

It is easy to see that there exist two vertices $x, y \in V(Q)$ such that $V(\pi_1'(Q)) = {\{\pi_1'(x), \pi_1'(y)\}}$. We have $|xy| \ge |\pi_1'(x)\pi_1'(y)| = \underline{|\pi_1'(Q)|} \ge \zeta$. Moreover, $|\pi_1'(x)\pi_1'(y)| \ge \zeta$ and $|\pi_2'(x)\pi_2'(y)| \le \zeta$ implies that $\left|\left\langle \frac{y-x}{|y-x|}, e_1' \right\rangle\right| \le \sin(45^\circ) = 1/\sqrt{2}.$

Fix $z \in V(Q)$ and assume that $\pi_1'(z) \in (\pi_1'(x) - \zeta^2, \pi_1'(x) + \zeta^2)$, i.e., $|\pi_1'(x)\pi_1'(z)| \le \zeta^2$. It is enough to show that $|zx| \le 2\zeta^2$. Assume for a contradiction that $|zx| \ge 2\zeta^2$. Combining the last two inequalities, we deduce that $\left|\left\langle \frac{z-x}{|z-x|}, e_1'\right\rangle\right| \ge \sin(60^\circ) = \sqrt{3}/2$.

As
$$\left|\left\langle \frac{y-x}{|y-x|}, e_1'\right\rangle\right| \leq \sin(45^\circ)$$
 and $\left|\left\langle \frac{z-x}{|z-x|}, e_1'\right\rangle\right| \geq \sin(60^\circ)$ we deduce that $\left|\left\langle \frac{y-x}{|y-x|}, \frac{z-x}{|z-x|}\right\rangle\right| \geq \sin(15^\circ)$.

As $|\langle \frac{y-x}{|y-x|}, e_1' \rangle| \leq \sin(45^\circ)$ and $|\langle \frac{z-x}{|z-x|}, e_1' \rangle| \geq \sin(60^\circ)$ we deduce that $|\langle \frac{y-x}{|y-x|}, \frac{z-x}{|z-x|} \rangle| \geq \sin(15^\circ)$. In the triangle $\cos(\{x, y, z\})$ the radius r of the inscribed circle has the formula $r = \frac{|\langle y-x, z-x \rangle|}{|xz|+|yz|+|xy|}$. Using the above, we deduce that

$$r \ge \frac{|xy||xz|\sin(15^\circ)}{2|xy| + 2|xz|} = \frac{\sin(15^\circ)}{\frac{2}{|xy|} + \frac{2}{|xz|}} \ge \frac{\sin(15^\circ)}{4} \min(|xy|, |xz|) \ge \frac{\zeta^2 \sin(15^\circ)}{2}.$$

Hence $|\pi_2(Q)| \ge |\pi_2(\operatorname{co}(\{x,y,z\}))| \ge \zeta^2 \sin(15^\circ)$. This gives the desired result as $|\pi_2(Q)| \le \eta < \zeta^2 \sin(15^\circ)$. The conclusion follows.

As before, we can translate back to a valid partition $\mathcal{H}' = \mathcal{H}'_0 \sqcup \mathcal{H}'_1 \sqcup \mathcal{H}'_2$ of F, where $\mathcal{H}'_i := \pi - 1(Q) \cap F : Q \in$ \mathcal{H}_i . This is a valid partition by construction of f'. The properties of \mathcal{H} translate to the following properties

- 1. $\sum_{Q \in \mathcal{H}'_{\alpha}} |Q| \le \zeta$
- 2. For every $Q \in \mathcal{H}'_1$, there exists a basis e'_1, e'_{i+1} of the plane spanned by e_1 and e_{i+1} such that we have $|\pi'_1(Q)|, |\pi'_{i+1}(Q)| \le \zeta.$
- 3. For every $Q \in \mathcal{H}'_2$, we have $\pi_1(V(\pi_{1,i+1}(Q))) \subset V(\pi_1(Q)) + (-4\zeta^2, 4\zeta^2)$.
- 4. For every $Q \in \mathcal{H}'_1 \cup \mathcal{H}'_2$, we have $Q = F \cap Q'$, where Q' is the intersection of at most $\ell_2^{4.12}$ halfspaces all of which contain $e_2, \ldots, e_i, e_{i+2}, \ldots, e_n$.

Given $F \in \mathcal{G}_2^{n+i}$, let $\mathcal{H}_0'(F)$, $\mathcal{H}_1'(F)$, and $\mathcal{H}_2'(F)$ be the sets produced here. We define $\mathcal{G}^{n+i+1} = \mathcal{G}_0^{n+i+1} \sqcup \mathcal{G}_1^{n+i+1}$ as follows:

$$\mathcal{G}_0^{n+i+1}:=\mathcal{G}_0^{n+i}\cup\bigcup_{F\in\mathcal{G}_2^{n+i}}\mathcal{H}_0'(F),\qquad \mathcal{G}_1^{n+i+1}:=\mathcal{G}_1^{n+i}\cup\bigcup_{F\in\mathcal{G}_2^{n+i}}\mathcal{H}_1'(F),\qquad \mathcal{G}_2^{n+i+1}:=\bigcup_{F\in\mathcal{G}_2^{n+i}}\mathcal{H}_2'(F)$$

Change the basis to $e'_1, e_2, \ldots, e_i, e'_{i+1}, e_{i+2}, \ldots, e_n$ for those parts $Q \in \mathcal{H}'_1(F)$ for some $F \in \mathcal{G}_2^{n+i}$. It is easy to see that this valid partition $\mathcal{G}^{n+i+1} = \mathcal{G}_0^{n+i+1} \sqcup \mathcal{G}_1^{n+i+1} \sqcup \mathcal{G}_2^{n+i+1}$ satisfies the properties as set out before. Consider the valid partition $\mathcal{G}^{2n} = \mathcal{G}_0^{2n} \sqcup \mathcal{G}_1^{2n} \sqcup \mathcal{G}_2^{2n}$ which has the following properties:

- 1. $\sum_{F \in \mathcal{G}_0^{2n}} |F| \leq 2n\zeta$
- 2. For every $F \in \mathcal{G}_1^{2n}$, there exists a basis e_1, \ldots, e_n so that $|\pi_i(F)| \leq \zeta$ for all $i = 1, \ldots, n$.
- 3. For every $F \in \mathcal{G}_2^{2n}$, there exists a basis e_1, \ldots, e_n so that $|\pi_i(F)| \leq \eta$ for all $i = 2, \ldots n$. Moreover, we have $\pi_1(V(\pi_{1,i}(F))) \subset V(\pi_1(F)) + (-4\zeta^2, 4\zeta^2)$ for all $i = 2, \ldots, n$.
- 4. For every $F \in \mathcal{G}_1^{2n} \cup \mathcal{G}_2^{2n}$, we have that $F = P \cap F'$ where F' is the intersection of at most $2n\ell_2^{4.12}$ halfspaces, each of which contains all but one of the basis vectors e_2, \ldots, e_n corresponding to F.

This last property shows that F is the intersection $2n\ell_2^{4.12} + \ell_P$ halfspaces, where $\ell_P \leq \binom{m}{k}$ is the number of halfspaces needed to construct P. Note that a vertex of F arises from n halfspaces, so that $|V(F)| \leq \binom{2n\ell_2^{4.12} + \ell_P}{n} \leq \ell_n^{4.13}(m)$. Hence, what remains is to strengthen property 3, to show that for $F \in \mathcal{G}_2^{2n}$, we have $\pi_1(V(F)) \subset V(\pi_1(F)) + (-4\zeta^2, 4\zeta^2)$. By the above we can partition V(F) into those vertices that are also a vertex of F' and those that are not.

First consider a vertex v of F that is also a vertex of F'. Note that as each of the defining hyperplanes of F' contains all but one of e_2, \ldots, e_n , and there are n hyperplanes needed to define a vertex of F', v must be defined by at least two planes containing the same n-2 elements from e_2, \ldots, e_n . Say v is defined by two planes containing $e_2, \ldots, e_{i-1}, e_{i+1}, e_n$, then we find that $\pi_{1,i}(v) \in V(\pi_{1,i}(F))$, so by property 3:

$$\pi_1(v) = \pi_1(\pi_{1,i}(v)) \in \pi_1(V(\pi_{1,i}(F))) \subset V(\pi_1(F)) + (-4\zeta^2, 4\zeta^2).$$

Hence, it remains to deal with vertices that lie on ∂P . we will show that if such a vertex lies far from $V(\pi_1(F))$, then F is contained very close to the boundary of P.

Note that as we have that $|\pi_i(F)| \leq \eta$ for all i = 2, ... n, we find that up to translation, we have $F \subset [0, |\pi_1(F)|] \times [0, \eta]^{n-1}$, so that if we let $C := \mathbb{R} \times [0, \eta]^{n-1}$, then $F \subset C \cap P$.

Claim 4.22. If $\exists u, v, w \in C \cap \partial P$, so that $|\pi_1(u) - \pi_1(v)|, |\pi_1(v) - \pi_1(w)|, |\pi_1(w) - \pi_1(u)| > 2\zeta^2$, then there exists a point x in $C \cap \partial P$ and a line L_x tangent to P at x so that $\angle L_x, e_1 \leq \zeta$

Proof. Note that by convexity of $P, C \cap \partial P$ can have at most two connected components, so we may assume u, v are in the same connected component. Let $\pi_1^c : \mathbb{R}^n \to \mathbb{R}^{n-1}$ be the projection away from the first coordinate. Note that $|\pi_1^c(u) - \pi_1^c(v)| \leq \sqrt{n-1}\eta$ as $\pi_1^c(u), \pi_1^c(v) \in [0, \eta]^{n-1}$. Hence,

$$\frac{1}{2}\angle(u-v), e_1 \le \sin(\angle(u-v), e_1) \le \frac{\sqrt{n-1}\eta}{\zeta^2} \le \frac{1}{2}\zeta,$$

using that η is sufficiently small in terms of ζ .

Consider the plane H containing u,v and direction e_1 . Note that again $H \cap C \cap \partial P$ has at most two connected components with u and v being in the same component. By the mean value theorem, there exists a point $x \in H \cap C \cap \partial P$ between u and v with a line L_x tangent to $H \cap P$ (and thus to P) parallel to u - v. Clearly, $\angle L_x, e_1 = \angle (u - v), e_1 \le \zeta$.

Note that because $F \subset C \cap P$, we have the intervals $\pi_1(F) \subset \pi_1(C \cap P)$. Moreover, because $C = \mathbb{R} \times [0,\eta]^{n-1}$, there exist $u,w \in C \cap \partial P$ such that $\pi_1(C \cap P) = [\pi_1(u),\pi_1(w)]$. Assume now v is a vertex of F that lies on ∂P with the property that $\pi_1(v) \notin V(\pi_1(F)) + (-2\zeta^2,2\zeta^2)$. It immediately follows that $\pi_1(v) \notin \{\pi_1(u),\pi_1(w)\} + (-2\zeta^2,2\zeta^2)$ i.e., $|\pi_1(u)-\pi_1(v)|,|\pi_1(v)-\pi_1(w)|,|\pi_1(w)-\pi_1(u)| > 2\zeta^2$. Hence, this claim shows that $F \subset L + B(0,\sqrt{n}\cdot\eta)$ for some line L which makes an angle at most ζ with a tangent line at a point in $L \cap \partial P$. By Lemma 4.17, this implies $F \subset \partial P + B(0,\xi)$.

With this setup, we are ready to conclude. Define \mathcal{F} as follows:

$$\mathcal{F}_0 := \mathcal{G}_0^{2n} \cup \left\{ F \in \mathcal{G}_2^{2n} : \exists v \in V(F) : \pi_1(v) \notin V(\pi_1(F)) + (-2\zeta^2, 2\zeta^2) \right\}$$

$$\mathcal{F}_1 := \mathcal{G}_1^{2n}$$

$$\mathcal{F}_2 := \left\{ F \in \mathcal{G}_2^{2n} : \forall v \in V(F) : \pi_1(v) \in V(\pi_1(F)) + (-2\zeta^2, 2\zeta^2) \right\}.$$

Indeed,

$$\sum_{F \in \mathcal{F}_0} |F| = 2n\zeta + |\partial P + B(0,\xi)| \le \varepsilon,$$

where we used that ξ is sufficiently small in terms of ε and P.

4.7 Proof of Auxiliary Lemmas

4.7.1 Proof of Lemma 4.14

Proof of Lemma 4.14. Let $\lambda_n = n^{-n}$ Translate P so that the John ellipsoid $E \subset P \subset nE$ is centred at the origin. Consider a translate L of L' that contains the origin. Let $H \supset L$ be any hyperplane containing L. Note that as H also contains the centre of E, we find

$$\lambda_n = n^{-n} = \frac{\frac{1}{2}|E|}{\frac{1}{2}|nE|} = \frac{|H^+ \cap E|}{|H^- \cap nE|} \le \frac{|H^+ \cap P|}{|H^- \cap P|} \le \frac{|H^+ \cap nE|}{|H^- \cap E|} = \frac{\frac{1}{2}|nE|}{\frac{1}{2}|E|} = n^n = \lambda_n^{-1},$$

which concludes the proof of the lemma.

4.7.2 Proof of Lemma 4.15

Proof of Lemma 4.15. Let v_1, \ldots, v_k where $k \geq 7$ be the vertices of P appearing around ∂P in that order. For every $i \in \{1, \ldots k\}$, consider two vertices v_i and v_{i+3} where the indices are considered mod k. The line through v_i and v_{i+3} partitions the plane into two halfplanes. Let H_i be the halfplane that contains all vertices except for v_{i+1} and v_{i+2} .

Claim 4.23.
$$\bigcap_{i=1}^k H_i \neq \emptyset$$

Proof of Claim. By Helly's theorem it suffices to show that any three of these halfplanes have non-empty intersection. Note that $|H_i^c \cap \{v_1, \ldots, v_k\}| = 2$, so for any i, i', i'' we find

$$|(H_i \cap H_{i'} \cap H_{i''}) \cap \{v_1, \dots, v_k\}| = |\{v_1, \dots, v_k\} \setminus (H_i^c \cup H_{i'}^c \cup H_{i''}^c)| \ge k - 6 \ge 1.$$

Hence, the intersection of all H_i is non-empty.

Choose some $p \in \bigcap_{i=1}^k H_i$ and consider any line ℓ containing p. Consider the two intersection points between ℓ and ∂P , say they lie on the line segment between v_i and v_{i+1} and on the line segment between v_j and v_{j+1} . Assume for a contradiction one of the sets $\ell^+ \cap P$ and $\ell^- \cap P$ has at least k vertices. Then we find that $|i-j| \leq 2$ (again mod k), say $j \in \{i+1, i+2\}$. However, this implies that $p \in \ell \cap P \subset H_i^c$, which is clearly a contradiction.

4.7.3 Proof of Lemma 4.16

Proof of Lemma 4.16. Iteratively construct a sequence of valid partitions \mathcal{G}_i of P, starting with $\mathcal{G}_0 = \{P\}$. Given $\mathcal{G}_i = \{P_1, \dots, P_j\}$, use Lemma 4.15 to find points $p_i \in P_i$ (in those P_i with at least 7 vertices) with the property that every line through them partitions P_i into polygons with fewer vertices than P_i . Now let

$$G_{i+1} = \{f^+(P_1, p_1), f^-(P_1, p_1), \dots, f^+(P_j, p_j), f^-(P_j, p_j)\}.$$

Note that if we let $v(\mathcal{G}_i)$ be the maximal number of vertices among all parts in \mathcal{G}_i , then $v(\mathcal{G}_{i+1}) \leq \max\{v(\mathcal{G}_i) - 1, 6\}$. Also, if we let $n(\mathcal{G}_i)$ be the number of parts of \mathcal{G}_i , then $n(\mathcal{G}_i) \leq 2^i$. Hence, we find $v(\mathcal{G}_{C-6}) \leq 6$ and $n(\mathcal{G}_i) \leq 2^{c-6}$. Clearly \mathcal{G}_{C-6} is a valid partition with the desired properties.

4.7.4 Proof of Lemma 4.17

Proof of Lemma 4.17. Let $D(P) := \sup_{x,y \in P} |x-y|$, we will show that $Q_{\zeta,\eta} \subset \partial P + B(0,\eta + \zeta D(P))$. Indeed, consider a ζ -permissible line L with the corresponding point $x \in L \cap \partial P$ and tangent line L_x . For any point $y \in L \cap P$, we have $|x-y| \leq D(P)$. By the definition of the sin, we have

$$\min_{y' \in L_x} |y - y'| = |x - y| \sin(\angle L, L_x) \le D(P)\zeta.$$

If we let $y' \in L_x$ the point realizing $\min_{y' \in L_x} |y - y'|$, then the line segment between y and y' intersects ∂P , so that $\min_{y' \in \partial P} |y - y'| \le \min_{y' \in L_x} |y - y'| \le \zeta D(P)$. Choosing ζ and η sufficiently small so that $\eta + \zeta D(P) \le \xi$, the lemma follows.

4.7.5 Proof of Lemma 4.18

Proof of Lemma 4.18. Let $\alpha = \lambda_2^4$, where λ_2 is the constant from Lemma 4.14.

Iteratively produce a valid partition of P, starting with $\mathcal{G}_0 = \{P\}$. Given $\mathcal{G}_i = \{P_1, \dots, P_{2^i}\}$, by Lemma 4.14, find points $p_j \in P_j$ and construct

$$\mathcal{G}_{i+1} := \{ f^+(P_1, p_1), f^-(P_1, p_1), \dots, f^+(P_{2^i}, p_{2^i}), f^-(P_{2^i}, p_{2^i}) \}.$$

Consider \mathcal{G}_4 , which has $2^4 = 16$ elements. Note that by construction, each $P' \in \mathcal{G}_4$ has $|P'| \geq \lambda_2^4 |P| = \alpha |P|$. Consider the set of newly created vertices $\bigcup_{P' \in \mathcal{G}_4} V(P') \setminus V(P)$. If there is a $v \in \bigcup_{P' \in \mathcal{G}_4} V(P') \setminus V(P)$ so that $\pi_2(v) \notin V(\pi_2(P)) + (-\varepsilon^2, \varepsilon^2)$ (or a line $f(P_j, p_j)$ through a vertex in V(P)), then consider the line $f(P_j, p_j)$ that created this vertex. Note that

$$\min\{|\pi_2(f^+(P_i, p_i))|, |\pi_2(f^-(P_i, p_i))|\} \le |\pi_2(P)| - \varepsilon^2,$$

so that at least one element P' of \mathcal{G}_4 has $|\pi_2(P')| \leq |\pi_2(P)| - \varepsilon^2$.

Alternatively, we find that

$$\pi_2\left(\bigcup_{P'\in G_*}V(P')\setminus V(P)\right)\subset V(\pi_2(P))+(-\varepsilon^2,\varepsilon^2).$$

As there are at most 4 vertices of P whose projection is not in $V(\pi_2(P)) + (-\varepsilon^2, \varepsilon^2)$, each of which is in at most element of \mathcal{G}_4 , and there are 16 sets in \mathcal{G}_4 , there is an set $P' \in \mathcal{G}_4$ with $V(P') \subset V(\pi_2(P)) + (-\varepsilon^2, \varepsilon^2)$. Note that

$$(V(\pi_2(P)) + (-\varepsilon^2, \varepsilon^2)) \cap \pi_2(P') \subset V(\pi_2(P')) + (-\varepsilon^2, \varepsilon^2).$$

Finally, note that every element of \mathcal{G}_4 has at most 10 vertices. The lemma follows.

5 Intermediate results for Quadratic Theorem (Theorem 1.5): Part II

5.1 Setup

Definition 5.1. Given a (i, ℓ, ε) -good cone $C \in \mathcal{C}^n$ (see Definition 4.9), we say a measurable subset $X \subset C$ is filled if for all $x \in C'$ we have

$$f(x) \times \{x\} \subset X \text{ or } (f(x) \times \{x\}) \cap X = \emptyset.$$

Definition 5.2. A pair of sets $X, Y \subset \mathbb{R}^n$ is a (η, γ) -approximate sandwich if there exists a convex set $P \subset \mathbb{R}^n$ containing the origin, so that $X, Y \subset (1 + \eta)P$ and $|P \setminus X| + |P \setminus Y| \le \gamma |P|$.

5.2 Theorem

Theorem 5.3. There exists an $\ell = \ell_n^{5.3}$ so that for every $\xi, \lambda, \eta, \gamma > 0$ the following holds. Given $A, B \subset \mathbb{R}^n$ a simple λ_n -bounded η_n -sandwich with the property that for all cones $C_i \in \mathfrak{C}$, we have $|A \cap C_i| = |B \cap C_i|$. Then there exist measurable subsets $A' \subset A$ and $B' \subset B$ and there exists a family of cones \mathcal{G} essentially partitioning \mathbb{R}^n refining \mathfrak{C} and a partition $\mathcal{G} = \mathcal{G}_0 \sqcup \mathcal{G}_1 \sqcup \mathcal{G}_2$ such that

- 1. $|A'| = |B'| \ge (1 \xi)|A|$
- 2. Every cone $F \in \mathcal{G}_0$ has $|A' \cap F| = |B' \cap F| = 0$.
- 3. Every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$ has $|A' \cap F| = |B' \cap F|$.
- 4. Every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$ has $A' \cap F$ and $B' \cap F$ are $(2\lambda, F)$ -bounded (η, γ) -approximate sandwiches inside F.
- 5. Every cone $F \in \mathcal{G}_1$ is $(1, \ell, \infty)$ -good and every cone $F \in \mathcal{G}_2$ is $(2, \ell, \infty)$ -good.
- 6. For every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$, $A' \cap F$ and $B' \cap F$ are filled in F.
- 7. For every cone $F \in \mathcal{G}_1 \cup \mathcal{G}_2$, if $F \subset C_i$, we have that e_n^F is perpendicular to the face of S contained in C_i .

5.3 Propositions

Again, before proving Theorem 5.3, we collect a list of results that will be used in the proof.

Proposition 5.4. Let $v_0, \ldots, v_n \subset \mathbb{R}^n$ be vectors not contained in a halfspace and let $A, B \subset \mathbb{R}^n$ be measurable sets with equal volume. Then there exists a vector $v \in \mathbb{R}^n$ such that for every cone $C \in \mathfrak{C}^{v_0, \ldots, v_n}$ we have

$$|A \cap C| = |(B+v) \cap C|.$$

Moreover, for every $\eta, \lambda > 0$, there is a computable constant $\eta'^{5.4} > 0$ such that the following holds. If $\{v_0, \ldots, v_n\} = \{e_0, \ldots, e_n\}$ (as in Definition 2.4) and if $A, B \subset \mathbb{R}^n$ is a λ -bounded $\eta'^{5.4}$ -sandwich, then A, B+v is a 2λ -bounded η -sandwich.

Proposition 5.5. Assume that $C \subset \mathbb{R}^n$ is a cone and assume that $A, B \subset \mathbb{R}^n$ are measurable sets with the property that

$$|A \cap C| = |B \cap C|$$
.

Then given a codimension-two subspace L (through the origin), there exists a hyperplane H through L which essentially partitions the cone

$$C = C_1 \cup C_2$$

with the property that

$$|A \cap C_1| = |B \cap C_1|$$
 and $|A \cap C_2| = |B \cap C_2|$

Proposition 5.6. There exists an $\ell_n^{5.6}$ so that for every $\varepsilon, \lambda, \eta > 0$ the following holds. Given $A, B \subset \mathbb{R}^n$ a simple λ_n -bounded η_n -sandwich with the property that for all cones $C_i \in \mathfrak{C}$, we have $|A \cap C_i| = |B \cap C_i|$. There exists a family of cones \mathcal{F} essentially partitioning \mathbb{R}^n refining \mathfrak{C} and a partition $\mathcal{F} = \mathcal{F}_0 \sqcup \mathcal{F}_1 \sqcup \mathcal{F}_2$ such that

- 1. Every cone $F \in \mathcal{F}$ has $|A \cap F| = |B \cap F|$.
- 2. For every cone $F \in \mathcal{F}$, the pair $A \cap F, B \cap F$ is a (λ, F) -bounded η -sandwich.
- 3. $\sum_{F \in \mathcal{F}_0} \mu_n(F) \leq \varepsilon$.
- 4. Every cone $F \in \mathcal{F}_1$ is $(1, \ell_n^{5.6}, \varepsilon)$ -good.
- 5. For every cone $F \in \mathcal{F}_2$ there exists a sub-cone F' of F with $\mu_n(F') \geq (1-\varepsilon)\mu_n(F)$ such that F' is $(2, \ell_n^{5.6}, \varepsilon)$ -good.
- 6. For every cone $F \in \mathcal{F}_1$ (or F' for $F \in \mathcal{F}_2$), if $F \subset C_i$, we have that e_n^F (or $e_n^{F'}$) is perpendicular to the face of S contained in C_i .

Proposition 5.7. For any $\ell, \xi, \lambda, \eta, \gamma > 0$, and $A, B \subset \mathbb{R}^n$ simple sets with equal volume, there exist $\varepsilon^{5.7} > 0$ such that the following holds. Let \mathcal{F} be a family of cones essentially partitioning \mathbb{R}^n refining \mathfrak{C} and let $\mathcal{F} = \mathcal{F}_0 \sqcup \mathcal{F}_1 \sqcup \mathcal{F}_2$ be a partition such that

- 1. Every cone $F \in \mathcal{F}$ has $|A \cap F| = |B \cap F|$.
- 2. For every cone $F \in \mathcal{F}$, the pair $A \cap F$, $B \cap F$ is a (λ, F) -bounded η -sandwich.
- 3. $\sum_{F \in \mathcal{F}_0} \mu_n(F) \leq \varepsilon^{5.7}$.
- 4. Every cone $F \in \mathcal{F}_1$ is $(1, \ell, \varepsilon^{5.7})$ -good.
- 5. For every cone $F \in \mathcal{F}_2$ there exists a sub-cone F' of F with $\mu(F') \geq (1 \varepsilon^{5.7})\mu_n(F)$ such that F' is $(2, \ell, \varepsilon^{5.7})$ -good.
- 6. For every cone $F \in \mathcal{F}_1$ (or F' for $F \in \mathcal{F}_2$), if $F \subset C_i$, we have that e_n^F (or $e_n^{F'}$) is perpendicular to the face of S contained in C_i .

Then there exist measurable subsets $A' \subset A$ and $B' \subset B$ and there exists a family of cones \mathcal{G} essentially partitioning \mathbb{R}^n refining \mathcal{F} and a partition $\mathcal{G} = \mathcal{G}_0 \sqcup \mathcal{G}_1 \sqcup \mathcal{G}_2$ such that

- 1. $|A'| = |B'| \ge (1 \xi)|A|$
- 2. Every cone $F \in \mathcal{G}_0$ has $|A' \cap F| = |B' \cap F| = 0$.
- 3. Every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$ has $|A' \cap F| = |B' \cap F|$.
- 4. Every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$ has $A' \cap F$ and $B' \cap F$ are $(2\lambda, F)$ -bounded (η, γ) -approximate sandwiches inside F.
- 5. Every cone $F \in \mathcal{G}_1$ is $(1, \ell, \infty)$ -good and every cone $F \in \mathcal{G}_2$ is $(2, \ell, \infty)$ -good.
- 6. For every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$, $A' \cap F$ and $B' \cap F$ are filled in F.

5.4 Proof of Theorem

5.4.1 Proof of Theorem 5.3

Proof of Theorem 5.3. Let $\ell_n^{5.3} := \ell_n^{5.6}$. Choose ϵ sufficiently small to be able to apply Proposition 5.7. Apply Proposition 5.6 with this ϵ and then apply Proposition 5.7.

5.5 Proofs of Propositions

5.5.1 Proof of Proposition 5.4

Proof of Proposition 5.4. We begin by proving the first conclusion. Applying an affine transformation if necessary, we may assume $S^{v_0,\dots,v_n}=S$ and $\mathfrak{C}^{v_0,\dots,v_n}=\mathfrak{C}=\{C_0,\dots,C_n\}$. Recall that S is the simplex containing the origin in its interior with vertices $V(S)=\{e_0,\dots,e_n\}$ (as in Definition 2.4). Denote by $F_i=\operatorname{co}(\{e_j\colon i\neq j\})$ the face of S opposite e_i .

First, assume for a contradiction that for all $v \in \mathbb{R}^n$, there exists $i \in [0, n]$ such that $|(B - v) \cap C_i| \neq |A \cap C_i|$. Note that for any $v \in \mathbb{R}^n$, we have

$$\sum_{i \in [0,n]} |(B-v) \cap C_i| = |B| = |A| = \sum_{i \in [0,n]} |A \cap C_i|.$$

Hence, if we define the closed sets

$$X_i := \{ v \in \mathbb{R}^n : |(B - v) \cap C_{f(v)}| \ge |A \cap C_{f(v)}| \}$$

for $i=0,\ldots,n$, then $\bigcup_i X_i=\mathbb{R}^n$. Consider rS, a large blow-up of S, so that $A,B\subset rS$. Note that rS is a simplex containing the origin with vertices $V(rS)=\{rv_0,\ldots,rv_n\}$. Moreover, the face of rS opposite to vertex rv_i is rF_i . It is easy to check that if $v\in rF_i$, then $|(B-v)\cap C_i|=0$. By the definition of the X_i 's, this implies that $v\in\bigcup_{j\neq i}X_i$. By the Knaster-Kuratowski-Mazurkiewicz lemma [KKM29], we find a point $v\in\mathbb{R}^n$, such that for all $i\in[0,n], |(B+v)\cap C_i|=|A\cap C_i|$.

We now prove the second conclusion. Fix $v \in \mathbb{R}^n$ as above and assume, without loss of generality, $v \in C_i$ for some $i \in \{0, \dots, n\}$, $P \subset A$, $B \subset (1+\eta')P$, and $S \subset A$, $B \subset \lambda S$. It is easy to check that $(X \cap C_i) + v \subset (X+v) \cap C_i$. Also, there exist universal constants $\alpha_n, \beta_n > 0$ such that $|(S+v) \cap C_i| \ge (1+\min(\alpha_n, \beta_n||v||_2))|S \cap C_i|$, where we used that $\{v_0, \dots, v_n\} = \{e_0, \dots, e_n\}$.

From the second and third inclusions and the above inequality, we get that

$$|(B+v) \cap C_{i}| \geq |((B \setminus S) + v) \cap C_{i}| + |(S+v) \cap C_{i}|$$

$$\geq |(B \setminus S) \cap C_{i}| + (1 + \min(\alpha_{n}, \beta_{n}||v||_{2}))|S \cap C_{i}|$$

$$= |(B \setminus S) \cap C_{i}| + |S \cap C_{i}| + \min(\alpha_{n}, \beta_{n}||v||_{2})|S \cap C_{i}|$$

$$\geq |B \cap C_{i}| + (n+1)^{-1} \min(\alpha_{n}, \beta_{n}||v||_{2})|S|$$

$$\geq |B \cap C_{i}| + \lambda^{-n}(n+1)^{-1} \min(\alpha_{n}, \beta_{n}||v||_{2})|B|.$$

From the first inclusion and the choice of v, we get that

$$|(1+\eta'P)\setminus P| \geq |A\setminus B| \geq |(A\cap C_i)\setminus (B\cap C_i)| \geq |A\cap C_i| - |B\cap C_i| = |(B+v)\cap C_i| - |B\cap C_i|.$$

Finally, from the first inclusion, we also get that $((1+\eta')^n-1)|B| \ge |(1+\eta')P \setminus P|$. Combining the last three inequalities, we get that $(1+\eta')^n-1 \ge \lambda^{-n}(n+1)^{-1}\min(\alpha_n,\beta_n||v||_2)$. By choosing $\eta'>0$ such that $((1+\eta')^n-1)\lambda^n(n+1)\alpha_n^{-1}<1$, we further get that $||v||_2 \le ((1+\eta')^n-1)\lambda^n(n+1)\beta_n^{-1}$.

Let Z is the unit volume ball centered at the origin, let τ_n be its radius and set $\zeta = ((1 + \eta')^n - 1)\lambda^n(n + 1)\beta_n^{-1}\tau_n^{-1}$. The above inequality implies that $v \in \zeta Z$. As S is a unit volume regular simplex and Z is a unit volume ball, both centered at the origin, we find $v \in \zeta Z \subset n\zeta S$. Moreover, from the first two inclusions, it easily follows that $S \subset 2P$ which implies $v \in 2n\zeta P$.

From the first two inclusions and the last two inclusions, we get

$$(1 - n\zeta)S \subset B + v \subset (\lambda + n\zeta)S$$
 and $(1 - 2n\zeta)P \subset B + v \subset (1 + \eta' + 2n\zeta)P$.

Provided $\zeta \leq 10^{-1} n^{-1} \min(1, \eta)$, and thus η' small, the conclusion follows. Thus, we shall choose η' such that $((1 + \eta')^n - 1)\lambda^n(n+1)^1\alpha_n^{-1} < 1$ and $((1 + \eta')^n - 1)\lambda^n(n+1)^1\beta_n^{-1}\tau_n^{-1} < 10^{-1}n^{-1}\min(1, \eta)$.

Proof of Proposition 5.5 5.5.2

Proof of Proposition 5.5. For notational convenience, assume that $A, B \subset C$, so that |A| = |B|. Consider any hyperplane H_0 containing L. For $\theta \in [0, 2\pi]$, let $\rho_{\theta} : \mathbb{R}^n \to \mathbb{R}^n$ be the rotation of the space fixing L by angle θ . Let $H_{\theta} := \rho_{\theta}(H_0)$ and let H_{θ}^+ and H_{θ}^- be the halfspaces generated by H_{θ} . Now consider the function $f : [0, \pi] \to \mathbb{R}$ defined by $f(\theta) := |H_{\theta}^+ \cap A| - |H_{\theta}^+ \cap B|$. Note that as A and B are bounded sets, f is continuous. Note moreover that as $H_{\pi}^{+} = H_{0}^{-}$, we find:

$$f(\pi) = |H_{\pi}^{+} \cap A| - |H_{\pi}^{+} \cap B| = |H_{0}^{-} \cap A| - |H_{0}^{-} \cap B| = (|A| - |H_{0}^{+} \cap A|) - (|B| - |H_{0}^{+} \cap B|) = -f(0).$$

By continuity, this implies the existence of a $\theta_0 \in [0,\pi]$ so that $f(\theta_0) = 0$. Hence, taking $H = H_{\theta_0}$, and $C_1 := H^+ \cap C$ and $C_2 := H^- \cap C$, we find

$$|A \cap C_1| = |B \cap C_1|$$
 and $|A \cap C_2| = |B \cap C_2|$.

This concludes the proof of the lemma.

5.5.3 Proof of Proposition 5.6

Let $\ell_n := \ell_n^{4.11}(n)$, where $\ell_n^{4.11}(n)$ is the constant from Theorem 4.11. For this proposition, we apply Theorem 4.11 to the context of A and B.

Proof of Proposition 5.6. We will construct a respectful function $f: \mathcal{C}^n \times \mathcal{T}_2^n \to \mathcal{T}_1^n$ as follows. Given a cone $C \in \mathcal{C}_n$ and a codimension-two subspace $L \in \mathcal{T}_2^n$, distinguish two case; either $|C \cap A| = |C \cap B|$ or not. In the latter case, let f(C, L) be any hyperplane containing L; this will not be a case that is going to affect us. If $|C \cap A| = |C \cap B|$, let f(C, L) be the hyperplane given by Proposition 5.5.

Apply Theorem 4.11 to each of the $C_i \in \mathfrak{C}$ with parameter $\varepsilon/(n+1)$, to find a partitions $\mathcal{F}^{C_i} = \mathcal{F}_0^{C_i} \sqcup \mathcal{F}_1^{C_i} \sqcup \mathcal{F}_2^{C_i}$. Let $\mathcal{F}_j = \bigcup_{i=0}^n \mathcal{F}_j^{C_i}$ for j=0,1,2, which then satisfies the conclusions 3, 4, 5, and 6, with the note that each of the C_i 's is defined by n lines. It remains to check the first two conclusions.

For conclusion 1, note that by Proposition 5.5, a valid partition of C_i contains only cones F so that $|A \cap F| =$ $|B \cap F|$, so the same holds in particular for all $F \in \mathcal{F}$.

For conclusion 2, note that being a λ -bounded η sandwich is inherited by taking subcones, so in particular for all $F \in \mathcal{F}$, we have that $A \cap F, B \cap F$ is a (λ, F) -bounded η_n sandwich.

5.5.4 Proof of Proposition 5.7

Proof of Proposition 5.7. We first construct $\mathcal{G}' = \mathcal{G}'_0 \sqcup \mathcal{G}'_1 \sqcup \mathcal{G}'_2$. First set $\mathcal{G}'_1 := \mathcal{F}_1$. For all $F \in \mathcal{F}_2$, let $F' \subset F$ be the subcone with $\mu_n(F') \geq (1 - \varepsilon)\mu_n(F)$, which is $(2, \ell, \varepsilon)$ -good. Then let $\mathcal{G}'_2 := \{F' : F \in \mathcal{F}_2\}$. For every $F \in \mathcal{F}_2$, we can partition $F \setminus F'$ into a number of convex cones. Let $\mathcal{F}(F)$ be that collection of convex cones. Let $\mathcal{G}'_0 := \mathcal{F}_0 \cup \bigcup_{F \in \mathcal{F}_2} \mathcal{F}(F)$.

Clearly, \mathcal{G}' is a refinement of \mathcal{F} . Note that $\sum_{G \in \mathcal{G}'_0} \mu_n(G) = \sum_{F \in \mathcal{F}_0} \mu_n(F) + \sum_{F \in \mathcal{F}_2} \mu_n(F \setminus F') \leq 2\varepsilon$. We construct $A \supset A_1 \supset A'$ and $B \supset B_1 \supset B'$ with increasingly more structure as follows.

Let $A_1 \subset A$ and $B_1 \subset B$ maximal so that $A_1 \cap F$ and $B_1 \cap F$ are filled for all $F \in \mathcal{G}'_1 \cup \mathcal{G}'_2$. To this end, for a given $F \in \mathcal{G}'_i$ with i = 1, 2, consider the cone $C \in \mathcal{C}^i$ and linear function $f : C \to \mathcal{K}^{n-i}$ so that $F = \bigcup_{x \in C} \{x\} \times f(x)$. Let $g, h : C \to \mathcal{P}(\mathbb{R}^{n-i})$ so that $A \cap F = \bigcup_{x \in C} \{x\} \times g(x)$ and $B \cap F = \bigcup_{x \in C} \{x\} \times h(x)$. Define $g', h': C \to \mathcal{K}^{n-i}$ by

$$g'(x) := \begin{cases} f(x) & \text{if } g(x) = f(x) \\ \emptyset & \text{otherwise} \end{cases}$$

and h' analogously. Let

$$A_1 \cap F := \bigcup_{x \in C} \{x\} \times g'(x)$$
 and $B_1 \cap F := \bigcup_{x \in C} \{x\} \times h'(x)$.

For $F \in \mathcal{G}'_0$, remove everything in the interior of F from A and B, so that $A_1 \cap F^{\circ} := \emptyset$ and $B_1 \cap F^{\circ} := \emptyset$. By λ -boundedness, we get

$$\sum_{F \in \mathcal{G}_0} |F \cap (A \setminus A_1)| \le \sum_{F \in \mathcal{G}_0'} O_n(|A|) \mu_n(F) \le O_n(\varepsilon |A|),$$

and analogously for B. To control $|A \setminus A_1|$ in $\mathcal{G}'_1 \cup \mathcal{G}'_2$, we use the fact that the cones $F \in \mathcal{G}'_1 \cup \mathcal{G}'_2$ are (i, ℓ, ε) -good and that A and B are simple. Find $Q_A \subset \mathbb{Z}^n$ so that $A = \bigcup_{x \in Q_A} x + [0,1]^n$. Note that if $x \in F \cap (A \setminus A_1)$ for

some cone $F \in \mathcal{G}_1' \cup \mathcal{G}_2'$, then consider the basis e_1^F, \ldots, e_n^F and write $x = x' \times x''$ so that $x \in x' \times g(x')$, with $g: C \to \mathcal{K}^{n-i}$ as before. As $x \notin A_1$, we find that $g(x') \neq \emptyset$ and $g(x') \neq f(x')$, so that g(x') contains an element of ∂A . As F was (i, ℓ, ε) -good, we find that the radius of f(x') is at most $\varepsilon||x'||_2 \leq O_n(\varepsilon|A|^{1/n})$. This implies that all points x in $A \setminus A_1$ must be close to the boundary of A, i.e., $F \cap (A \setminus A_1) \subset \partial A + B(o, O_n(\varepsilon|A|^{1/n}))$. Hence,

$$\sum_{F \in \mathcal{G}_1 \cup \mathcal{G}_2} |F \cap (A \setminus A_1)| \le |\partial A + B(o, O_n(\varepsilon |A|^{1/n}))| = O_n(\varepsilon |A|^{1/n} \cdot |Q_A|) = O_n(\varepsilon |A|^{(n+1)/n}),$$

so that we find

$$|A \setminus A_1| = \sum_{F \in \mathcal{G}_0} |F \cap (A \setminus A_1)| + \sum_{F \in \mathcal{G}_1 \cup \mathcal{G}_2} |F \cap (A \setminus A_1)| = O_n(\varepsilon |A|^{(n+1)/n}) \le \varepsilon',$$

for any ε' by choosing ε sufficiently small in terms of ε' , |A| and n. Let $B_1 \subset B$ be constructed analogously. Note that the sets A_1 and B_1 satisfy conclusions 1, 2, 5, and 6 in the lemma.

We continue to construct $A_2 \subset A_1$ and $B_2 \subset B_1$. We aim to remove part of $A_1 \cap F'$ or $B_1 \cap F'$ (whichever is larger), outside of $\frac{1}{2}rS$, where r is such that $rS \subset A, B \subset \lambda rS$, while maintaining the property that the sets remain filled in the cones. Indeed note that as

$$A \setminus A_1 \cap F' \subset \partial A + B(o, O_n(\varepsilon |A|^{1/n}))$$
 and $rS \subset A$,

we find that choosing ε sufficiently small in terms of |A|, r, and n, we have $\frac{2}{3}r(S \cap F') \subset A_1, B_1$. Hence, (assuming that $A_1 \cap F'$ is larger than $B_1 \cap F'$), there exists a filled subset $A_2 \cap F' \subset A_1 \cap F'$ so that if we let $B_2 \cap F' = B_1 \cap F'$, then $|A_2 \cap F'| = |B_2 \cap F'|$ and $\frac{1}{2}r(S \cap F') \subset A_2 \cap F', B_2 \cap F'$. Let A_2 and B_2 be defined thus in all $F' \in \mathcal{G}'_1 \cup \mathcal{G}'_2$.

To estimate $|\hat{A_2} \setminus \hat{A_1}| + |B_1 \setminus B_2|$, for every $F' \in \mathcal{G}_1 \cup \mathcal{G}_2$, compare $|A_1 \cap F'|$ and $|B_1 \cap F'|$. Find the $F \in \mathcal{F}_1 \cup \mathcal{F}_2$ so that $F' \subset F$. Note that $|A_1 \cap F'|, |B_1 \cap F'| \leq |F \cap B| = |F \cap A|$, so that we find

$$\sum_{F' \in \mathcal{G}_1' \cup \mathcal{G}_2'} \left| |A_1 \cap F'| - |B_1 \cap F'| \right| \le \sum_{F' \in \mathcal{G}_1' \cup \mathcal{G}_2'} |F \cap A| - |F' \cap A_1| + |F \cap B| - |F' \cap B_1|.$$

We consider the contribution from A and B independently to find

$$\sum_{F' \in \mathcal{G}_1' \cup \mathcal{G}_2'} |F \cap A| - |F' \cap A_1| \le \sum_{F' \in \mathcal{G}_1' \cup \mathcal{G}_2'} |(F \setminus F') \cap A| + |F' \cap (A \setminus A_1)| \le |A \setminus A_1| \le \varepsilon'.$$

Hence, we find

$$|A_2 \setminus A_1| + |B_1 \setminus B_2| \le \sum_{F' \in \mathcal{G}'_1 \cup \mathcal{G}'_2} \left| |A_1 \cap F'| - |B_1 \cap F'| \right| = 2\varepsilon'.$$

Note that A_2 and B_2 satisfy conclusions 1, 2, 3, 5, and 6 from the lemma. To get conclusion 4 as well, we construct $A' \subset A_2$ and $B' \subset B_2$ as follows. Let $\mathcal{H} \subset \mathcal{G}'_1 \cup \mathcal{G}'_2$ be the collection of those F' so that $A_2 \cap F'$ and $B_2 \cap F'$ is not a (η, γ) -approximate sandwich in F'. If we let $F \in \mathcal{F}$ so that $F' \subset F$, then we find that $A \cap F$ and $B \cap F$ is a (η, F) sandwich, so there exists a P so that $P \subset A \cap F$, $B \cap F \subset (1 + \eta)P$. This immediately implies that $A_2 \cap F'$, $B_2 \cap F' \subset (1 + \eta)P \cap F'$, so as $A_2 \cap F'$, $B_2 \cap F'$ is not (η, γ) -approximate sandwich, we must find $|(P \cap F') \setminus A_2| + |(P \cap F') \setminus B_2| \ge \gamma |P \cap F'|$. Notice that as $P \subset A$, B, we find that

$$|(A \cap F') \setminus A_2| + |(B \cap F') \setminus B_2| \ge \gamma |P \cap F'| \ge \frac{1}{2} (1 - \eta)^n \gamma (|\operatorname{co}(F' \cap A_2)| + |\operatorname{co}(F' \cap B_2)|)$$

$$\ge \Omega_n(|F' \cap A_2| + |F' \cap B_2|),$$

so that, summing over all F', we find

$$\sum_{F' \in \mathcal{H}} |F' \cap A_2| + |F' \cap B_2| = O_n \left(\sum_{F' \in \mathcal{H}} |(A \cap F') \setminus A_2| + |(B \cap F') \setminus B_2| \right) \le O_n(|A \setminus A_2| + |B \setminus B_2|) \le O_n(\varepsilon').$$

Hence, the result follows by choosing $\mathcal{G}_0 = \mathcal{G}_0' \cup \mathcal{H}$, $\mathcal{G}_1 = \mathcal{G}_1' \setminus \mathcal{H}$, $\mathcal{G}_2 = \mathcal{G}_2' \setminus \mathcal{H}$, and $A' \subset A_2$ and $B' \subset B_2$ are obtained by removing everything in the interior of the cones in \mathcal{H} , i.e., $A' = A_2 \setminus \bigcup_{F \in \mathcal{H}} F^{\circ}$ and $B' = B_2 \setminus \bigcup_{F \in \mathcal{H}} F^{\circ}$.

6 Intermediate results for the Quadratic Theorem (Theorem 1.5): Part III

6.1 Setup

Definition 6.1. Let e_1, \ldots, e_n be an orthogonal basis. Let $\pi_{i_1, \ldots, i_j} : \mathbb{R}^n \to \mathbb{R}^j$ be the projection onto the j-dimensional subspace spanned by e_1, \ldots, e_{i_j} . For a set Y and $x_{i_1}, \ldots, x_{i_j} \in \mathbb{R}$, let $Y_{x_{i_1}, \ldots, x_{i_j}}$ be the fibre of Y above $(x_{i_1}, \ldots, x_{i_j})$.

Definition 6.2. For a set Y let $Y' = Y \cap \pi_1^{-1}[1/4, 1/2]$.

Definition 6.3. Given a (n-1)-dimensional simplex $X \subset \mathbb{R}^n$, a cylinder over X is a set of the form $X + \mathbb{R}^+ v$ for some direction $v \in S^{n-1}$.

Definition 6.4. Given $\rho > 0$ and a subcone $C \subset \mathbb{R}^n$ of a cone in \mathfrak{C} , we say a cylinder U over a simplex $T \subset C \cap \frac{1}{2}\partial S$ is ρ -central if the n+1 defining hyperplanes H_1, \ldots, H_{n+1} (i.e., so that $U = \bigcap_{i=1}^{n+1} H_i^+$) of the cylinder have the property that for all choices $*_i \in \{+, -\}$, we have $|C \cap S \cap \bigcap_{i=1}^{n+1} H_i^{*_i}| \geq \rho |C \cap S|$.

Definition 6.5. Say a set $X \subset \mathbb{R}^n$ is α -almost convex if $|\operatorname{co}(X) \setminus X| \leq \alpha |X|$.

6.2 Propositions

Proposition 6.6. For any w > 1, there exists $c_{n,w} > 0$ such that the following holds. Let $T = \{o, e_1, \dots, e_{n-1}\}$ be an orthogonal simplex. Let $U = T \times \mathbb{R}$. Let $T \times [0,1] \subset A, B \subset T \times [0,w]$. Assume that A and B are filled in the sense that $A_{x_1,x_n} = U_{x_1,x_n}$ or $A_{x_1,x_n} = \emptyset$ and similarly $B_{x_1,x_n} = U_{x_1,x_n}$ or $B_{x_1,x_n} = \emptyset$. If $|tA + (1-t)B| \le (1+\delta)|A|$ with $0 < \delta \ll_{n,t} 1$, then $|A' \triangle B'| \le c_{n,w} t^{-1/2} \delta^{1/2} |A|$. Here A' and B' are as in Definition 6.2.

Proposition 6.7. Let $C \subset \mathbb{R}^n$ be a subcone of a cone in \mathfrak{C} , let U be a ρ -central cylinder over a simplex $T \subset C \cap \frac{1}{2}\partial S$, and let α -almost convex sets $X,Y \subset C$ be so that

- |X| = |Y|,
- $S \cap C \subset X, Y \subset \lambda S$,
- $|tX + (1-t)Y| \le (1+\delta)|X|$.

Then there exists a cylinder $V = (1 + \beta)U + x$ so that

- $|\beta| \le O_{n,\rho} \left(\delta^{1/2} t^{-1/2} + \alpha \right),$
- $|(U\triangle V)\cap S| = O_{n,\rho}\left(\delta^{1/2}t^{-1/2} + \alpha\right)|C\cap S|,$
- $|X \cap U| = |Y \cap V|$,
- $|t(X \cap U) + (1-t)(Y \cap V)| |X \cap U| \le |tX + (1-t)Y| |X|$.

Proposition 6.8. For all $n \in \mathbb{N}$, $\lambda \geq 1$, and $\epsilon > 0$, there exists $\rho = \rho_{n,\lambda,\epsilon}^{6.8} > 0$ so that the following holds. Let $C \subset \mathbb{R}^n$ be a subcone of a cone in \mathfrak{C} . Let f_1, \ldots, f_n be an orthonormal basis of \mathbb{R}^n so that f_n is perpendicular to the face of S intersecting C.

Then there exists a (n-1)-simplex $T \subset C \cap \frac{1}{2}\partial S$ of radius less than ϵ and a collection \mathcal{U} of ρ -central cylinders over T so that

- $|\mathcal{U}| = O_{n,\lambda,\epsilon}(1)$,
- $\lambda(C \cap S) \setminus (C \cap S) \subset \bigcup_{U \in \mathcal{U}} U$,
- one facet of T is parallel to the subspace spanned by f_1, \ldots, f_{n-2} .

6.3 Proof of Propositions

6.3.1 Proof of Quadratic Theorem: Proposition 6.6

Proof of Proposition 6.6. Let C, D be the sets A, B, respectively, Steiner symmetrized around e_1 , i.e., C_{x_1} , D_{x_1} are discs with the same size a A_{x_1} , B_{x_1} , respectively.

Claim 6.9.
$$|A' \triangle B'| \leq |C' \triangle D'| + O_{n,t}(\delta)|A|$$

Proof. By Theorem 1.4, we know that $|\operatorname{co}(A)\setminus A|, |\operatorname{co}(B)\setminus B| \leq O_{n,t}(\delta)|A|$. Therefore, it is enough to show that

$$|A' \triangle B'| \le |C' \triangle D'| + |\operatorname{co}(A) \setminus A| + |\operatorname{co}(B) \setminus B|.$$

For a fixed fiber, we have the elementary inequality

$$|A_{x_1,\dots,x_{n-1}} \triangle B_{x_1,\dots,x_{n-1}}| \le ||A_{x_1,\dots,x_{n-1}}| - |B_{x_1,\dots,x_{n-1}}|| + |\operatorname{co}(A)_{x_1,\dots,x_{n-1}}| - |A_{x_1,\dots,x_{n-1}}| + |\operatorname{co}(B)_{x_1,\dots,x_{n-1}}| - |B_{x_1,\dots,x_{n-1}}|.$$

By the hypothesis that A and B are filled, for fixed x_1 and varying x_2, \ldots, x_{n-1} , we have $|A_{x_1,\ldots,x_{n-1}}|$ is constant and $|B_{x_1,\ldots,x_{n-1}}|$ is constant. Hence, for fixed x_1 , we get that

$$\int \left| |A_{x_1,\dots,x_{n-1}}| - |B_{x_1,\dots,x_{n-1}}| \right| dx_2 \dots dx_{n-1} = \left| \int |A_{x_1,\dots,x_{n-1}}| - |B_{x_1,\dots,x_{n-1}}| dx_2 \dots dx_{n-1} \right| = \left| |A_{x_1}| - |B_{x_1}| \right|,$$

therefore,

$$\int |A_{x_1,\dots,x_{n-1}} \triangle B_{x_1,\dots,x_{n-1}}| dx_2 \dots dx_{n-1} \le \int ||A_{x_1,\dots,x_{n-1}}| - |B_{x_1,\dots,x_{n-1}}|| dx_2 \dots dx_{n-1}
+ \int |\operatorname{co}(A)_{x_1,\dots,x_{n-1}}| - |A_{x_1,\dots,x_{n-1}}| dx_2 \dots dx_{n-1}
+ \int |\operatorname{co}(B)_{x_1,\dots,x_{n-1}}| - |B_{x_1,\dots,x_{n-1}}| dx_2 \dots dx_{n-1}
= ||A_{x_1}| - |B_{x_1}|| + |\operatorname{co}(A)_{x_1}| - |A_{x_1}| + |\operatorname{co}(B)_{x_1}| - |B_{x_1}|.$$

Thus, we deduce

$$|A'\triangle B'| = \int_{T'} |A_{x_1,\dots,x_{n-1}} \triangle B_{x_1,\dots,x_{n-1}} | dx_1 dx_2 \dots dx_{n-1}$$

$$= \int_{1/4}^{1/2} \left(\left| |A_{x_1}| - |B_{x_1}| \right| + \left| \cos(A)_{x_1} \right| - |A_{x_1}| + \left| \cos(B)_{x_1} \right| - |B_{x_1}| \right) dx_1$$

$$= \int_{1/4}^{1/2} \left(\left| |C_{x_1}| - |D_{x_1}| \right| + \left| \cos(A)_{x_1} \right| - |A_{x_1}| + \left| \cos(B)_{x_1} \right| - |B_{x_1}| \right) dx_1$$

$$\leq |C'\triangle D'| + |\cos(A) \setminus A| + |\cos(B) \setminus B|.$$

This concludes the proof of the claim.

Recall that the sets C and D are Steiner symmetrized around e_1 . In particular, as $|tA+(1-t)B| \le (1+\delta)|A|$, we deduce $|tC+(1-t)D| \le (1+\delta)|C|$.

Finally, because $T \times [0,1] \subset A$, $B \subset T \times [0,w]$, there exists a parameter p such that $p(1-x_1)^n \leq |C_{x_1}|, |D_{x_1}| \leq wp(1-x_1)^n$ for $x_1 \in [0,1]$.

For a point $x \in [0,1]$ there exists a unique point $y_x \in [0,1]$ such that if we denote by I = [0,x] and J = [0,y], then $|C_I| = |D_J|$.

Claim 6.10. The function $y_x: [0,2/3] \to [0,1]$ is continuous and Lipschitz with parameter $\Theta_{n,w}(1)$.

Proof. This follows immediately from the fact that, for an interval $I \subset [0, 2/3]$, we have $|C_I|, |D_I| = \Theta_{n,w}(p)|I|$.

Claim 6.11. For $x \in [1/4, 1/2]$, $y = y_x$ satisfies $|y - x| = O_{d,w}(t^{-1/2}\delta^{1/2})$.

Proof. Let I = [0, x] and J = [0, y]. As $tC_I + (1-t)D_J$ and $tC_{I^c} + (1-t)D_{J^c}$ are disjoint, we get $|tC_I + (1-t)D_J| \le |C_I| + \delta |C|$. Moreover, as $|A| = O_w(|C_I|)$, we further get $|tC_I + (1-t)D_J| \le (1 + O_{n,w}(\delta))|C_I|$.

First note the qualitative bound that $y \leq 2/3$ (provided δ is small). Recall that for every $0 \leq z \leq 2/3$ we have $|C_z|, |D_z| = \Theta_{n,w}(p)$, and that the sets C_I and D_J are symmetrized. It is easy to see that, by doing d+1 parallel hyperplane cuts (containing the direction e_1), we can construct convex cylinders C_I' and D_J' inside C_I and D_J , respectively, such that

$$|C_I'| = |D_J'| = \Theta_{n,w}(1)|C_I|$$
 and $|tC_I' + (1-t)D_J'| - |C_I'| \le |tC_I + (1-t)D_J| - |C_I|$.

In particular, we get that the convex cylinders C_I' and D_J' have heights |I| and |J| and $|tC_I' + (1-t)D_J'| \le (1+\Theta_{n,w}(\delta))|C_I'|$. By the sharp stability of the Brunn-Minkowski for convex sets, we then deduce $|I| = (1+O(t^{-1/2}\delta^{1/2}))|J|$. As $x \in (1/4,1/2)$ We conclude $|y-x| = O_{n,w}(t^{-1/2}\delta^{1/2})$.

Claim 6.12. For points $x \in (1/4, 1/2)$, if $|y-x| = O_{n,w}(t^{-1/2}\delta^{1/2})$, then $|\operatorname{co}(C)_x| - |\operatorname{co}(C)_y| \le O_{n,w}(t^{-1/2}\delta^{1/2})$ and $|\operatorname{co}(D)_x| - |\operatorname{co}(D)_y| \le O_{n,w}(t^{-1/2}\delta^{1/2})p$.

Proof. It is enough to notice that for $x \in (1/4, 2/3)$, we have $|\operatorname{co}(C)_x|$ is Lipschitz with parameter $\Theta_{n,w}(p)$, which follows from the fact that $|\operatorname{co}(C)_x|^{1/n}$, $|\operatorname{co}(D)_x|^{1/n}$ are convex and $p(1-x_1)^n \leq |C_x|$, $|D_x| \leq wp(1-x_1)^n$ for $x_1 \in [0,1]$.

Consider the partitions into very small consecutive intervals $[0,1] = I_0 \cup \cdots \cup I_\ell$ and $[0,1] = J_0 \cup \cdots \cup J_\ell$ such that C_{I_k} and D_{J_k} are all cylinders with the same volume. Moreover, set $|tC_{I_k} + (1-t)D_{J_k}| = (1+\delta_k)|C_{I_k}|$. It is trivial to check that $\sum_k \delta_k |C_{I_k}| \leq \delta |C|$. In particular, $\sum_{k:I_k \subset [1/4,1/2]} \delta_k \leq O_{n,w}(\delta)$.

Claim 6.13. Fix k such that $I_k, J_k \subset [0, 2/3]$. For $x \in I_k$ and $y \in J_k$ we have $|C_x| = (1 + O_{n,w}(t^{-1/2}\delta_k^{1/2}))|D_y|$.

Proof. Because C_{I_k} and D_{J_k} are convex cylinders, the result follows immediately from the sharp stability of Brunn-Minkowski inequality for convex sets [FMP09].

It is easy to see that the function $x \to y_x$ is picewise linear and maps the interval I_k linearly into the interval J_k . Fix $x \in I_k \subset [1/4, 1/2]$. We have the elementary bound

$$||C_x| - |D_x|| \le ||C_x| - |D_{y_x}|| + |\operatorname{co}(D)_x \setminus D_x| + |\operatorname{co}(D)_{y_x} \setminus D_{y_x}| + ||\operatorname{co}(D)_x| - |\operatorname{co}(D)_{y_x}||.$$

From Claim 6.13, we deduce

$$||C_x| - |D_{y_x}|| = O_{n,w}(t^{-1/2}\delta_k^{1/2})p.$$

From Claim 6.11 and Claim 6.12, we get

$$\left| |\operatorname{co}(D)_x| - |\operatorname{co}(D)_{y_x}| \right| \le O_{n,w}(t^{-1/2}\delta^{1/2})p.$$

Therefore

$$\begin{aligned} |C'\Delta D'| &= \int_{1/4}^{1/2} \big| |C_x| - |D_x| \big| dx = \sum_{k:I_k \subset [1/4,1/2]} O_{n,w}(t^{-1/2}\delta_k^{1/2}) p \times |I_k| + O_{n,w}(t^{-1/2}\delta^{1/2}) p \times 1/4 \\ &+ \int_{1/4}^{1/2} \Big(|\operatorname{co}(D)_x \setminus D_x| + |\operatorname{co}(D)_{y_x} \setminus D_{y_x}| \Big) dx \\ &= \sum_{k:I_k \subset [1/4,1/2]} O_{n,w}(t^{-1/2}\delta_k^{1/2}) |C_{I_k}| + O_{n,w}(t^{-1/2}\delta^{1/2}) |C_{[1/4,1/2]}| \\ &+ \int_{1/4}^{1/2} \Big(|\operatorname{co}(D)_x \setminus D_x| + |\operatorname{co}(D)_{y_x} \setminus D_{y_x}| \Big) dx \\ &\leq O_{n,w}(t^{-1/2}\delta^{1/2}) |C| + \int_{1/4}^{1/2} \Big(|\operatorname{co}(D)_x \setminus D_x| + |\operatorname{co}(D)_{y_x} \setminus D_{y_x}| \Big) dx. \end{aligned}$$

By the linear stability result we have $|\operatorname{co}(D) \setminus D| \leq O_{n,w}(\delta)|C|$, and by Claim 6.10 we deduce

$$\int_{1/4}^{1/2} \left(|\operatorname{co}(D)_x \setminus D_x| + |\operatorname{co}(D)_{y_x} \setminus D_{y_x}| \right) dx \le O_{n,w}(\delta)|C|.$$

Therefore, we get $|C'\triangle D'| \leq O_{n,w}(t^{-1/2}\delta^{1/2})|C|$, that together with Claim 6.9 allows us to conclude that $|A'\triangle B'| \leq O_{n,w}(t^{-1/2}\delta^{1/2})|A|$.

6.3.2Proof of Proposition 6.7

Proof of Proposition 6.7. First affinely transform so that the simplex formed by the vertices of T together with the origin form a regular simplex.

Write H_0 for the hyperplane containing the face of $\frac{1}{2}S$ intersecting C. Writing H_0^+ for the halfspace defined by H_0 not containing the origin, we find that by $S \cap C \subset X, Y \subset \lambda S$, we have $|H_0^+ \cap X| = |H_0^+ \cap Y|$. Let $X' := H_0^+ \cap X$ and $Y' := H_0^+ \cap Y$, so that |X'| = |Y'|. By the Brunn-Minkowski inequality, we have

$$|t(H_0^- \cap X) + (1-t)(H_0^- \cap Y)| \ge |H_0^- \cap X|.$$

Since $t(H_0^- \cap X) + (1-t)(H_0^- \cap Y)$ is a subset of tX + (1-t)Y disjoint from tX' + (1-t)Y', we find that $|tX' + (1-t)Y'| - |X'| \le |tX + (1-t)Y| - |X|$.

We repeatedly apply this last argument for different hyperplane cuts.

Let H_1, \ldots, H_n be the hyperplane cuts defining U. Find parallel hyperplanes G_1, \ldots, G_n so that |X'| $\bigcap_{i=1}^{j} H_i^+|=|Y'\cap\bigcap_{i=1}^{j} G_i^+|$ for all $j=0,\ldots,n$, which exist by continuity. We will show that the distance between H_i and G_i is at most $O_n\left(\delta^{1/2}t^{-1/2}\right)$. We show this for i=1, the other cases follow analogously by

Consider the sets $X^{\pm} := X' \cap H_1^{\pm}$ and $Y^{\pm} := Y' \cap G_1^{\pm}$. By the fact that U is central, we know that $|X^- \cap S| \ge \rho |C \cap S|$. Write H' for the hyperplane $2H_0$ containing the face of S intersecting C. Let

$$X'' := X^- \cap (H')^- = S \cap H_0^+ \cap H_1^- \cap (H')^-$$

and find a plane G' parallel to H' so that $Y'' := Y^- \cap (G')^-$ satisfies |X''| = |Y''|. By the argument before, we find that

$$|tX'' + (1-t)Y''| - |X''| \le |tX' + (1-t)Y'| - |X'| \le |tX + (1-t)Y| - |X| \le \delta |X| \le \rho^{-1}\delta |X''|.$$

Note that X'' is convex, so by the stability result for the Brunn-Minkowski inequality in [FMP09] we find that

Note that X'' is convex, so by the stability result for the Brunn-Minkowski inequality in [FMF 63] we find that $|X'' \triangle (Y'' + x)| \le O_n \left(\delta^{1/2} t^{-1/2}\right) |X''|$ for some translate $x \in \mathbb{R}^n$.

Consider the component x^{\perp} of x perpendicular to H_0 . Note that the band between H_0 and $H_0 + x^{\perp}$ of $S \cap C \cap H_1^- \cap G_1^{-1}$ is completely contained in $X'' \triangle Y''$, so as U is central, this implies $||x^{\perp}||_2 = O_n \left(\delta^{1/2} t^{-1/2}\right)$. As X'' and Y'' are $\rho^{-1}\alpha$ -almost convex, we find that translating them by x^{\perp} changes the symmetric difference only little, i.e.,

$$|X''\triangle(Y'' + (x - x^{\perp}))| \le |X''\triangle(Y'' + x)| + O_n\left(\delta^{1/2}t^{-1/2} + \alpha\right)|X''| \le O_n\left(\delta^{1/2}t^{-1/2} + \alpha\right)|X''|.$$

It is now easy to see that the translate z with no component perpendicular to H_0 minimizing $|X''\triangle(Y''+z)|$ is z=0. Indeed, for z=0, in every plane P parallel to H_0 we have that $P \cap X''$ and $P \cap Y''$ are nested (which one is contained in which depends on the relative positions of H_1 and H'). Hence, we find

$$|X''\Delta Y''| \le O_n \left(\delta^{1/2} t^{-1/2} + \alpha\right) |X''|.$$

Finally, akin to the previous argument, we now find that $(G_1 \triangle H_1) \cap S \cap C \cap H_0^+ \cap H'^- \cap G'^- \subset X'' \triangle Y''$, so that $|(G_1 \triangle H_1) \cap S| \leq O_n \left(\delta^{1/2} t^{-1/2} + \alpha\right) |C \cap S|$. This concludes the induction.

Letting $V := \bigcap_{i=0}^n H_i^+$, we find that $U \triangle V \subset \bigcup_{i=1}^n G_i^+ \triangle H_i^+$, therefore

$$|(U\triangle V)\cap S| \le O_n \left(\delta^{1/2}t^{-1/2} + \alpha\right)|C\cap S|,$$

and consequently $|\beta| = O_n \left(\delta^{1/2} t^{-1/2} + \alpha\right)$.

6.3.3Proof of Proposition 6.8

Proof of Proposition 6.8. We first show that we may assume $C \cap \frac{1}{2}\partial S$ is a simplex.

Consider the set $P:=C\cap \frac{1}{2}\partial S$ which lies in a subspace spanned by f_1,\ldots,f_{n-1} . Let L be the subspace spanned by f_1, \ldots, f_{n-2} . Let $x, x' \in \partial P$ so that x + L and x' + L are the two translates of L tangent to P. Write $o' := \frac{x+x'}{2}$. Let P' be the projection of P onto o' + L along the direction xx'. Note that $P \cap (o' + L)$ contains $\frac{o'+P'}{2}$. Let T' be the largest simplex contained in that translate of $\frac{o'+P'}{2}$, so that $|T'| = \Omega_n(|P'|)$ and P' is contained in some translate of 2nT'. Let $T'' = \cos(T' \cup \{x\})$.

This construction gives a set T'' so that there exists a translate of 4nT'' which contains P. Indeed, for every point y on the line segment between o' and $\frac{o'+x}{2}$ we have that $L+y\cap T''$ is a homothetic copy of T' larger than $\frac{1}{2}T'$. Hence, for all points y on the line segment between 4no' and 2n(o'+x) we have that $L+y\cap 4nT''$ is a homothetic copy of T' larger than 2nT' (which contains a translate of P'). Hence, we can translate 4nT'' so that $P \subset z + 4nT''$ and $|T''| \ge \Omega_n(|P|)$.

Let T be the translate of $\min\{\epsilon, \frac{1}{4n}\}T''$ centred at the barycenter of T'', so that we still have $|T| \geq \Omega_{n,\epsilon}(|P|)$ (for notational convenience assume $\min\{\epsilon, \frac{1}{4n}\} = \frac{1}{4n}$). Note that one of the facets of T is parallel to T', i.e., to the subspace spanned by f_1, \ldots, f_{n-2} . Since the T is contained in the proper interior of $T'' \subset z + 4nT''$ (at a distance lower bounded in terms of n from $\partial(z + 4nT'')$), we find that there exists some σ_n so that z + 4nT'' is contained in the translate of $\sigma_n T$ centred at the barycenter of T. Write p for the barycenter of T, so that $H_{p,4n}(T) = T'' \subset P \subset H_{p,\sigma_n}(T)$, where $H_{q,\xi}$ is the homothety with ratio ξ centred at q.

Write S' for the halfspace containing S defined by the hyperplane which contains the face of S intersecting C. Note that $C \cap S = C \cap S'$. Write $C' := \bigcup_{s>0} sH_{p,\sigma_n}(T)$ for the cone generated by the set $H_{p,\sigma_n}(T) \supset P$ and write $C'' := \bigcup_{s>0} sH_{p,4n}(T)$ for the cone generated by the set $H_{p,4n}(T) = T'' \subset P$, so that $C'' \subset C \subset C'$. We have that

$$\lambda(C \cap S) \setminus (C \cap S) \subset \lambda(C' \cap S') \setminus (C' \cap S').$$

Up to an affine transformation, we may assume $\operatorname{co}(T \cup \{o\})$ is a regular simplex. Note that the problem of covering $\lambda(C' \cap S') \setminus (C' \cap S')$ with cylinders over T which are central in the cone C'' depends only on n, λ , and ϵ and no longer on the particular C, and f_1, \ldots, f_n that we started with. Hence, we can simply choose any collection \mathcal{U} of cylinders (which are ξ -central for some $\xi > 0$) so that $\lambda(C' \cap S') \setminus (C' \cap S') \subset \bigcup_{U \in \mathcal{U}} U$ and choose ρ' appropriately so that the cylinders are ρ' -central in C''. We automatically have $|\mathcal{U}| = O_{n,\lambda,\epsilon}(1)$ and if a cylinder is ρ' -central in C'' it is also ρ central in C for some ρ depending only on λ, n, ϵ , and ρ' .

7 Proof of the Quadratic Symmetric Difference result (Theorem 1.5)

Proof of Theorem 1.5. Choose parameters according to the following hierarchies. First choose

$$n \gg \lambda^{-1} \gg \rho \gg \epsilon \gg \lambda'^{-1} \gg w^{-1}$$

so that $\epsilon, \rho, \lambda, \lambda', w = \theta_n(1)$. Then choose

$$t, n, \epsilon, \rho, \lambda, \lambda', w \gg \Delta, \eta'' \gg \eta \gg \gamma \gg \eta',$$

where every variable is understood to be chosen sufficiently small in terms of all of the preceding variables.

By Proposition 2.9, we may assume that A, B form a simple $\lambda/2$ -bounded η' -sandwich. By Proposition 5.4 we may assume (after translating) that, for $C_i \in \mathfrak{C}$, $|C_i \cap A| = |C_i \cap B|$ at the cost of A, B form a simple λ -bounded η -sandwich.

Now apply Theorem 5.3 with parameters $\xi = \delta, \lambda, \eta, \gamma > 0$, to find $A' \subset A$, $B' \subset B$, and a family of cones \mathcal{G} essentially partitioning \mathbb{R}^n refining \mathfrak{C} and a partition $\mathcal{G} = \mathcal{G}_0 \sqcup \mathcal{G}_1 \sqcup \mathcal{G}_2$ such that:

- 1. $|A'| = |B'| \ge (1 \xi)|A|$;
- 2. For every cone $F \in \mathcal{G}_0$ it holds $|A' \cap F| = |B' \cap F| = 0$;
- 3. For every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$ it holds $|A' \cap F| = |B' \cap F|$;
- 4. For every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$, $A' \cap F$ and $B' \cap F$ are $(2\lambda, F)$ -bounded (η, γ) -approximate sandwiches inside F;
- 5. Every cone $F \in \mathcal{G}_1$ is $(1, \ell_n^{5.3}, \infty)$ -good, and every cone $F \in \mathcal{G}_2$ is $(2, \ell_n^{5.3}, \infty)$ -good;
- 6. For every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$, $A' \cap F$ and $B' \cap F$ are filled in F;
- 7. For every cone $F \in \mathcal{G}_1 \cup \mathcal{G}_2$, if $F \subset C_i$ then e_n^F is perpendicular to the face of S contained in C_i .

By our choice of ξ , it suffices to show $|A' \triangle B'| \le O_n(t^{-1/2}\delta^{1/2})|A|$, as $|A \triangle B| \le |A' \triangle B'| + |A \setminus A'| + |B \setminus B'| \le |A' \triangle B'| + 2\delta |A|$.

For every cone $F \in \mathcal{G}_1 \sqcup \mathcal{G}_2$, write $\delta_F := \frac{|t(A' \cap F) + (1-t)(B' \cap F)| - |A' \cap F|}{|A' \cap F|}$. For notational convenience, write $\delta_F = 0$ for $F \in \mathcal{G}_0$. In terms of this parameter, we find

$$\sum_{F \in \mathcal{G}} \delta_F |A' \cap F| = \sum_{F \in \mathcal{G}} |t(A' \cap F) + (1-t)(B' \cap F)| - |A' \cap F| \leq |tA' + (1-t)B'| - |A'| \leq |tA + (1-t)B| - |A'| \leq 2\delta |A|,$$

where we used that $t(A'\cap F)+(1-t)(B'\cap F)$ are essentially disjoint subsets of tA'+(1-t)B'. We also have the trivial observation that $\sum_{F\in\mathcal{G}}|(A'\cap F)\triangle(B'\cap F)|=\sum_{F\in\mathcal{G}}|(A'\triangle B')\cap F|=|A'\triangle B'|$. By concavity of the square root function, it thus suffices to show $|(A'\cap F)\triangle(B'\cap F)|\leq O_n(t^{-1/2}\delta^{1/2})|A'\cap F|$ for every cone $F\in\mathcal{G}$. Henceforth, fix a cone $C\in\mathcal{G}_2$ for which we will show this bound on the symmetric difference (the case that $C\in\mathcal{G}_0\sqcup\mathcal{G}_1$ follows analogously, though more easily). For notational convenience write $X:=A'\cap C$ and $Y:=B'\cap C$.

As X, Y form an (η, γ) -approximate sandwich inside C, we find that for sufficiently small η and γ , we have that δ_C is smaller than Δ for all $C \in \mathcal{G}$. Hence, by Theorem 1.4, we know that X and Y are $O_{n,t}(\delta)$ -almost convex.

Now apply Proposition 6.8 to the cone C and the basis e_1^C, \ldots, e_n^C to find a (n-1)-simplex $T \subset C \cap \frac{1}{2}\partial S$ and a collection \mathcal{U} of ρ -central cylinders over T of radius less than ϵ , so that

- $|\mathcal{U}| = O_{n,\lambda}(1) = O_n(1),$
- $\lambda(C \cap S) \setminus (C \cap S) \subset \bigcup_{U \in \mathcal{U}} U$,
- one facet of T is parallel to the subspace spanned by e_1^C, \ldots, e_{n-2}^C .

Since $A' \cap C$ and $B' \cap C$ are λ -bounded inside the cone C, we have

$$X\triangle Y\subset \lambda(C\cap S)\setminus (C\cap S)\subset \bigcup_{U\in \mathcal{U}}U,$$

so that

$$|X\triangle Y| \leq \sum_{U\in\mathcal{U}} |U\cap (X\triangle Y)| \leq |\mathcal{U}| \max_{U\in\mathcal{U}} |U\cap (X\triangle Y)| \leq O_n \left(\max_{U\in\mathcal{U}} |U\cap (X\triangle Y)| \right).$$

Hence, it suffices to show $|U \cap (X \triangle Y)| \leq O_n(t^{-1/2}\delta^{1/2})|X|$. Because one facet of T is parallel to the subspace spanned by e_1^C, \ldots, e_{n-2}^C , we find that $X \cap U$ filled.

Now apply Proposition 6.7 to C, U, and the $O_{n,t}(\delta)$ -almost convex pair X, Y, to find a cylinder $V = (1 + \beta)U + x$ so that:

- $|\beta| \le O_{n,\rho} \left(\delta^{1/2} t^{-1/2} + O_{n,t}(\delta) \right) = O_n \left(\delta^{1/2} t^{-1/2} \right),$
- $|(U\triangle V)\cap S| = O_{n,\rho}\left(\delta^{1/2}t^{-1/2} + O_{n,t}(\delta)\right)|C\cap S| = O_n\left(\delta^{1/2}t^{-1/2}\right)|C\cap S|,$
- $|X \cap U| = |Y \cap V|$,
- $|t(X \cap U) + (1-t)(Y \cap V)| |X \cap U| \le |tX + (1-t)Y| |X| = \delta_C|X|$.

Note that as V is homothetic to U, we also have that $V \cap Y$ is filled.

As T lies well within S and has radius less than ϵ , we can apply Lemma 10.10 to find $t' \in (t/2, 2t)$ and X', Y' that form a tubular λ' -bounded η'' -sandwich, where $\lambda' = \theta_n(\lambda) = \theta_n(1)$ and η'' tend to zero with η , so that

$$|t'X' + (1 - t')Y'| - t'|X'| - (1 - t')|Y'| \le |t(X \cap U) + (1 - t)(Y \cap V)| \le \delta_C|X|,$$

and $|(X \cap U)\triangle(Y \cap V)| \leq |X'\triangle Y'| + O_n\left(\delta_C^{1/2}t^{-1/2}\right)|X \cap U|$. Moreover, as we have only taken affine transformations, we find that X', Y' are both filled in their common tube, say W. After affine transformation, we may assume $W = S' \times \mathbb{R}$ for an orthogonal simplex S', so that $S' \times [0,1] \subset X', Y' \subset S' \times [0,w/2]$ for some $w = O_{n,\lambda'}(1) = O_n(1)$.

Note that by construction $||X'| - |Y'|| \le O_n \left(\delta_C^{1/2} t^{-1/2}\right) |X'|$. Extend (i.e., append a set of the form $S' \times [-\zeta, 0)$ for an appropriate $\zeta \le O_n \left(\delta_C^{1/2} t^{-1/2}\right)$ and then translating that set up by ζ) the smaller of the two sets to find $X'' \supseteq X'$ and $Y'' \supseteq Y'$ with |X''| = |Y''|, so that by Lemma 10.11, we find

$$|t'X'' + (1 - t')Y''| - |X''| = |t'X' + (1 - t')Y'| - t'|X'| - (1 - t')|Y'| \le \delta_C|X|.$$

Moreover, we have $|X'\triangle Y'|\leq |X''\triangle Y''|+O_n\left(\delta_C^{1/2}t^{-1/2}\right)|X''|$ and

$$S' \times [0,1] \subset X'', Y'' \subset S' \times [0, w/2 + \zeta] \subset S' \times [0, w].$$

Clearly, X'' and Y'' are still filled.

Recall that as U was ρ -central, we find that $|X''| \geq 2^{-n}|X'| \geq \Omega_{n,\rho}(|X|) = \Omega_n(|X|)$. Hence, as $\delta_C \leq \Delta$, we find that $\delta_C \frac{|X|}{|X''|}$ is sufficiently small in terms of n and t so that we can apply Proposition 6.6 to find that $|X''\Delta Y''| \leq O_{n,w}(t^{-1/2}\delta^{1/2})|X''| = O_n(t^{-1/2}\delta^{1/2})|X''|$.

8 Intermediate results for Linear Theorem with few vertices (Theorem 1.6)

For notational convenience, consider the following definitions. Given $A, B \subset \mathbb{R}^n$ with |A| = |B| and $t \in (0, 1)$, let

$$D_t(A, B) := tA + (1 - t)B$$
, and $\delta_t(A, B) := \frac{|D_t(A, B)|}{|A|} - 1$.

Throughout this section, we will always consider $t \in (0,1)$. We will show the following theorem.

Theorem 8.1. There exists a constants $c^{8.1}$ such that the following holds. Given $A, B \subset \mathbb{R}^n$ of equal volume and $t \in [\tau, 1 - \tau]$ such that co(A) is a simplex, A is the intersection between a simple set and a simplex and $\delta_t(A, B) \leq \min\{t, (1 - t)\}^n$, then

$$|\operatorname{co}(A) \setminus A| \leq \min\{t, (1-t)\}^{-c}^{8.1} n^{8} \delta_{t}(A,B) |A|.$$

Theorem 1.6 follows easily from Theorem 8.1 which is the same result with the stronger assumption that co(A) is a simplex.

8.1 Outline of the proof of Theorem 8.1

The proof of Theorem 8.1 follows the following steps.

1. Writing $S = co(x_0, \ldots, x_n)$, find a point $v \in A$ so that the density in each of the subsimplices

$$co(x_0,\ldots,x_{i-1},v,x_{i+1},\ldots,x_n)$$

does not decrease too much (cf Proposition 8.4).

2. The doubling of A in each of the subsimplices is subadditive (cf Proposition 8.2), i.e., there exists a matching partition $B = \bigcup_{S'} B_{S'}$ with $|B_{S'}| = |A \cap S'|$ so that

$$|tA + (1-t)B| \ge \sum |t(A \cap S') + (1-t)B_{S'}|.$$

- 3. Iterating this process, we end up with two types of simplices: those in which A has low density and those with small radius.
- 4. For a simplex in which A has a sufficiently low density (but not too low), i.e., $|S' \cap A|/|S'|$ is small, a recent result by van Hintum and Keevash [vHK23b] shows that the doubling is $\Omega_{n,t}(1)$ (cf Proposition 8.3). Since we can guarantee the density of A inside S' is $O_{n,t}(1)$, we find that |S'| is controlled by the doubling of $A \cap S'$.
- 5. Assuming that without loss of generality that A is a finite union of boxes intersected with a simplex (cf Proposition 2.9), the combined volume of simplices with small radius that are not completely filled by A goes to zero with the number of iterations.
- 6. Therefore, we conclude

$$\begin{aligned} |\operatorname{co}(A) \setminus A| &= \sum_{S'} |S' \setminus A| \leq \sum_{\text{low density } S'} O_{n,t}(|t(A \cap S') + (1-t)B_{S'}| - |A \cap S'|) + \sum_{\text{small radius } S'} |S'| \\ &= O_{n,t}(|tA + (1-t)B| - |A|). \end{aligned}$$

8.2 Auxiliary propositions

A crucial ingredient is the following proposition.

Proposition 8.2. If co(A) is a simplex S' with vertex set $\{x_0, \ldots x_n\}$ and $x \in A$ is some point in the interior of co(A), then we partition S' into simplices

$$S_i := co(x_0, \dots, x_{i-1}, x, x_{i+1}, \dots, x_n).$$

For any $B \subset \mathbb{R}^n$ with |B| = |A| there exist sets $B_i \subset \mathbb{R}^n$ with $|B_i| = |A \cap S_i|$, so that

$$\sum_{i} \frac{|A \cap S_i|}{|A|} \ \delta_t(A \cap S_i, B_i) \le \delta_t(A, B)$$

We recall here the following result of van Hintum and Keevash [vHK23b].

Proposition 8.3 ([vHK23b]). There exists a constant $c^{8.3}$ so that if $t \in (0, \frac{1}{2}]$ and $A, B \subset \mathbb{R}^n$ of equal volume satisfy $|\operatorname{co}(A)| + |\operatorname{co}(B)| \ge t^{-c^{8.3}n^2}|A|$, then $\delta_t(A, B) \ge t^n$.

The last crucial ingredient is the following proposition.

Proposition 8.4. For all $n \in \mathbb{N}$ and $\alpha > 0$ there exists a constant $\eta_{\alpha,n}^{8.4} > 0$ (we can take $\eta_{\alpha,n}^{8.4} = c_n^{8.4} \alpha^{c_n^{8.4} - 6}$ for some constants $c_n^{8.4}, c_n^{8.4} > 0$) so that the following holds. Let $X \subset \mathbb{R}^n$ be a set and $S' = \operatorname{co}\{x_0, \ldots, x_n\} \subset \mathbb{R}^n$ be a simplex with $X \subset S'$. If $|X| \geq \alpha |S'|$, then there exists a point $x \in X$ such that for all $0 \leq i \leq n$

$$|\cos\{x_0,\ldots,x_{i-1},x,x_{i+1},\ldots,x_n\}\cap X| \ge \eta_{\alpha,n}^{8.4}|S'|.$$

Moreover, there exists a constant $\rho_{\alpha,n}^{8.4} > 0$ (we can take $\rho_{\alpha,n}^{8.4} = \eta_{\alpha,n}^{8.4}$) such that

$$\max_{i} d(x, x_i) \le (1 - \rho_{\alpha, n}^{8.4}) \max_{i, j} d(x_i, x_j),$$

A weaker version of Proposition 8.4 in the dense domain follows later in Lemma 10.6, which has a simpler proof.

8.3 Auxiliary lemmas

Lemma 8.5. For all $n \in \mathbb{N}$ and $\alpha > 0$ there exist constants $\eta_{1,\alpha,n}^{8.5}, \eta_{2,\alpha,n}^{8.5} > 0$ (we can take $\eta_{1,\alpha,n}^{8.5} = \eta_{2,\alpha,n}^{8.5} = c_n^{8.5} \alpha^{c^{8.5} n^3}$, for some constants $c_n^{8.5}, c^{8.5} > 0$) so that the following holds. Let $X \subset \mathbb{R}^n$ be a set and $S' = co\{x_0, \ldots, x_n\} \subset \mathbb{R}^n$ be a simplex with $X \subset S'$. If $|X| \ge \alpha |S'|$, then there exists a subset $Y \subset X$ with $|Y| \ge \eta_{1,\alpha,n}^{8.5} |S'|$ such that for all $x \in Y$ and $1 \le i \le n$

$$|\cos\{x_0,\ldots,x_{i-1},x,x_{i+1},\ldots,x_n\}\cap X| = \eta_{2,\alpha,n}^{8.5}|S'|.$$

Lemma 8.6. For all $n \in \mathbb{N}$ and $\alpha > 0$ there exist constants $\eta_{1,\alpha,n}^{8.6}, \eta_{2,\alpha,n}^{8.6} > 0$ (we can take them $\eta_{1,\alpha,n}^{8.6} = \eta_{2,\alpha,n}^{8.6} = c_n^{8.6} \alpha^c^{8.6n^2}$, for some constants $c_n^{8.6n^2}, c^{8.6} > 0$) so that the following holds. Let $X \subset Q = [-1,1]^n$. If $|X| \ge \alpha |Q|$, then there exists a subset $Y \subset X$ with $|Y| \ge \eta_{1,\alpha,n}^{8.6} |Q|$ such that for all $x \in Y$ and all faces F of the box Q

$$|\operatorname{co}(\{x\} \cup F) \cap X| = \eta_{2,\alpha,n}^{8.6}|Q|.$$

Before the next lemma we need a definition.

Definition 8.7. For $\alpha > 0$ and $1 \le j \le n$ construct the (n-1)-dimensional box

$$Q_{j,\alpha} = \operatorname{co}\left(\left\{\sum_{i=1}^{n} \varepsilon_{i} e_{i} \colon \varepsilon_{i} \in \left\{-\alpha, \alpha\right\} \text{ for all } i \in [n] \setminus \left\{j\right\} \text{ and } \varepsilon_{j} = 1\right\}\right)$$

and construct the cone

$$C_{i,\alpha} = \bigcup_{t>0} tQ_{i,\alpha}$$

Moreover, for $-n \leq j \leq -1$ let $Q_{j,\alpha} = -Q_{-j,\alpha}$ and $C_{j,\alpha} = -C_{-j,\alpha}$.

Lemma 8.8. For all $n \in \mathbb{N}$ and $\alpha > 0$ there exist constants $\eta_{1,\alpha,n}^{8.8}, \eta_{2,\alpha,n}^{8.8} > 0$ (we can take $\eta_{1,\alpha,n}^{8.8} = \eta_{2,\alpha,n}^{8.8} = c_n^{8.8} \alpha^{c^{8.8}n^2}$, for some constants $c_n^{8.8}, c^{8.8} > 0$) so that the following holds. Let $X \subset Q = [-1,1]^n$. If $|X| \ge \alpha |Q|$, then there exists a subset $Y \subset X$ with $|Y| \ge \eta_{1,\alpha,n}^{8.8} |Q|$ such that for all $x \in Y$ and all $j \in [-n,n] \setminus \{0\}$

$$|(x+C_{j,\alpha})\cap X| \ge \eta_{2,\alpha,n}^{8.8}|Q|.$$

Lemma 8.9. For all $n \in \mathbb{N}$ and $\alpha > 0$ the following holds. Let $P = [0, a_1] \times \cdots \times [0, a_n] + x$ be a box with $x \in \mathbb{R}^n$ and $a_n \alpha^2 / 8 \le a_1, \ldots, a_{n-1} \le a_n \alpha^2 / 4$. Let X be a subset of the box with $|X| \ge \alpha |P|$. Then there exists a box

$$Q = [0, b_1] \times \cdots \times [0, b_n] + y$$
 with $y \in \mathbb{R}^n$ and $b_1(\alpha/4)^2/8 \le b_2, \dots, b_n \le b_1(\alpha/4)^2/4$

such that, if we set $Y = Q \cap X$, then the following holds:

$$|Q| \ge \alpha^{2n} |P|/2^{7n}$$
, $|Y| \ge \alpha |Q|/4$, and $|(y + C_{n,\alpha}) \cap X| \ge \alpha |P|/4$ for all $y \in Y$.

We shall need one more lemma, for the proof Theorem 1.6.

Lemma 8.10. Let $t \in (0,1)$ and $A,B,C \subset \mathbb{R}^n$ so that |A| = |B| and C is convex with a finite number of vertices, then there exists a subset $B' \subset B$ so that $|B'| = |A \cap C|$ and

$$|t(A \cap C) + (1-t)B'| - |A \cap C| \le |tA + (1-t)B| - |A|.$$

Proof of Theorem 8.1

Proof of Theorem 8.1. we will consider the case $t \leq \frac{1}{2}$, the other case follows analogously. Let $\varepsilon = t^{c}^{8.3}n^{2}$. Let $c^{8.3}$ be the constant from Proposition 8.3. Let $c^{8.4}$, $c^{8.4}_n$ and $\rho = \rho^{8.4}_{\varepsilon,n}$ the constants from Proposition 8.4. Consider the following iterative process. First set $\mathcal{T}_0 = \{\cos(A)\}$ and $\mathcal{S}_0 = \emptyset$, and note that $|\cos(A)| \leq \varepsilon^{-1}|A|$ by Proposition 8.3. At a given stand $|\cos(A)| \leq \varepsilon^{-1}|A|$

by Proposition 8.3. At a given stage i with $\mathcal{T}_i, \mathcal{S}_i$, look at every element $S' \in \mathcal{T}_i$ and distinguish two cases: either $|S' \cap A| \le \varepsilon |S'|$ or $|S' \cap A| > \varepsilon |S'|$.

For each simplex $S' = \operatorname{co}\{x_0, \dots, x_n\} \in \mathcal{T}_i$ with $|S' \cap A| \geq \varepsilon |S'|$ we construct the n+1 simplices $f_0(S'), \ldots, f_n(S')$ as follows. We apply Proposition 8.4 to find a central point $x \in S' \cap A$ and we construct the simplex $f_j(S') = co\{x_0, \dots, x_{j-1}, x, x_{j+1}, \dots, x_n\}.$

$$\mathcal{T}_{i+1} := \bigcup_{S' \in \mathcal{T}_i: \ |S' \cap A| \ge \varepsilon |S'|} \left\{ f_0(S'), \dots, f_n(S') \right\} \quad \text{and} \quad \mathcal{S}_{i+1} := \mathcal{S}_i \cup \left\{ S' \in \mathcal{T}_i: |S' \cap A| < \varepsilon |S'| \right\}.$$

Using the fact that A is closed, it follows by induction that for $i \in \mathbb{N}$ and $S' \in \mathcal{T}_i \sqcup \mathcal{S}_i$ we have $\operatorname{co}(A \cap S') = S'$. Moreover, $\mathcal{T}_i \sqcup \mathcal{S}_i$ forms an essential partition of co(A).

Claim 8.11. For all $S' \in \mathcal{T}_i$ and $j \in [0, n]$, we have $|f_j(S')| \ge c_n^{8.4} \varepsilon^{e^{8.4} n^6} |S'|$.

 $\textit{Proof.} \ \, \text{By Proposition 8.4 (and our choice of} \, \, x \in S' \cap A), \, \text{we have} \, |f_j(S')| \geq |f_j(S') \cap A| \geq c_n^{8.4} \varepsilon^{8.4} n^6 |S'|.$

Claim 8.12. For all $S' \in \mathcal{S}_i \cup \mathcal{T}_i$, we have $|S' \cap A| \geq c_n^{8.4} \varepsilon^{c^{8.4} n^6} |S'|$

Proof. Every simplex $S' \in \mathcal{S}_i \cup \mathcal{T}_i$ is $f_j(S'')$ for some S'' with $|S'' \cap A| \geq \varepsilon |S''|$. By Proposition 8.4 (and our choice of $x \in S' \cap A$), we have $|S' \cap A| \ge c_n^{8.4} \varepsilon^{c^{8.4} n^6} |S''|$.

Claim 8.13. At stage i there exists a function $g_i: \mathcal{T}_i \cup \mathcal{S}_i \to \mathcal{P}(\mathbb{R}^n)$, so that for all $S' \in \mathcal{T}_i \cup \mathcal{S}_i$, $g_i(S')$ is a measurable subset of B such that $|g_i(S')| = |S' \cap A|$ and

$$\sum_{S' \in \mathcal{T}_i \cup \mathcal{S}_i} |A \cap S'| \cdot \delta_t(A \cap S', g_i(S')) \le |A| \cdot \delta_t(A, B).$$

Proof. This follows by induction from Proposition 8.2.

Claim 8.14. For every i, we have that

$$\sum_{S' \in \mathcal{S}_i} |S'| \le \frac{1}{c_n^{8.4}} t^{-c^{8.4}} c^{8.3} n^{8-n} |A| \cdot \delta_t(A, B).$$

Proof of Claim. For $S' \in \mathcal{S}_i$, we know by construction and by the second claim that $c_n^{8.4} \varepsilon^{8.4} n^6 |S'| \leq |S' \cap A| < 1$ $\varepsilon |S'|$. Moreover, $\operatorname{co}(A \cap S') = S'$. Hence, by Proposition 8.3, $\delta_t(A \cap S', g_i(S')) \geq t^n$, where g_i is the function from the previous claim. Hence, combining the results from the previous two claims, we find

$$\sum_{S' \in \mathcal{S}_i} |S'| \le \frac{1}{c_n^{8.4}} \varepsilon^{-c^{8.4} n^6} \sum_{S' \in \mathcal{S}_i} |S' \cap A| \le \frac{1}{c_n^{8.4}} \varepsilon^{-c^{8.4} n^6} \sum_{S' \in \mathcal{S}_i} |S' \cap A| \cdot \frac{\delta_t(A \cap S', g_i(S'))}{t^n}$$

$$\le \frac{1}{c_n^{8.4}} t^{-n} \varepsilon^{-c^{8.4} n^6} |A| \cdot \delta_t(A, B) = \frac{1}{c_n^{8.4}} t^{-c^{8.4} c^{8.3} n^8 - n} |A| \cdot \delta_t(A, B),$$

which concludes the claim.

Before we conclude we need one more claim. Given a simplex S', let radius rad(S') be the maximal length among its edges.

Claim 8.15. For all r > 0 the following holds:

$$\sum_{S' \in \mathcal{T}_i: rad(T) > r} |S'| \to 0 \quad \text{as } i \to \infty.$$

Proof of Claim. We make the convention $\bigcup \mathcal{T}_i := \bigcup_{S' \in \mathcal{T}_i} S'$ Let $k = \lceil \log_{(1-\rho)}(r) \rceil$. We can distinguish two types of elements in \mathcal{T}_i . Either $S' \subset \bigcup \mathcal{T}_{i+k(n+1)}$ or not. Collect the former in \mathcal{T}_i' and the latter in \mathcal{T}_i'' . For $S' \in \mathcal{T}_i''$, at least some simplex originating from S' is in $\mathcal{S}_{i+k(n+1)}$. By the first claim, that simplex will

have size at least $(c_n^{8.4} \varepsilon^{c_n^{8.4} n^6})^{k(n+1)} |S'|$. Hence,

$$\sum_{S' \in \mathcal{T}_{i+k(n+1)}: \ S' \subset \bigcup \mathcal{T}_{i''}'} |S'| \le \left(1 - (c_n^{8.4} \varepsilon^{c^{8.4} n^6})^{k(n+1)}\right) \left| \bigcup \mathcal{T}_{i''}' \right|.$$

For $S' \in \mathcal{T}_i'$, we will find an element $S'' \in \mathcal{T}_{i+k(n+1)}$ with $S'' \subset S'$ and rad(S'') < r. Let $S' = S^0$, and consider $S^{j+1} := f_n(f_{n-1}(\dots f_0(S^j)\dots)$. Crucially, $rad(S^{j+1}) \leq (1-\rho)rad(S^j)$. Indeed, none of the edges of S^j remain and all of the edges added have length at most $(1-\rho)rad(S^j)$ by Proposition 8.4. Hence, $rad(S^k) \leq r \cdot rad(S^0) \leq r$. Note that again by the first claim $|S^k| \geq (c_n^{8.4} \varepsilon^{8.4} n^6)^{k(n+1)} |S'|$.

Combining these two cases gives

$$\sum_{S' \in \mathcal{T}_{i+k(n+1)}: \ rad(S') > r} |S'| \leq \left(1 - (c_n^{8.4} \varepsilon^{c^{8.4} n^6})^{k(n+1)}\right) \sum_{S' \in \mathcal{T}_i: \ rad(S') > r} |S'|.$$

The conclusion follows.

To bound the contribution from the simplices in \mathcal{T}_i , we use the assumption that A is the intersection between a finite union of cubes and a simplex. In particular, this implies that ∂A has finite (n-1)-dimensional Hausdorff measure. Hence, $|\partial A + B^n(o,r)| \to 0$ as $r \to 0$. Fix r to be such that $|\partial A + B^n(o,r)| \le \delta_t(A,B)^2|A|$. Using Claim 8.15, find an i, so that $\sum_{S' \in \mathcal{T}_i: rad(S') > r} |S'| \leq \delta_t(A, B)^2 |A|$. Note that for $S' \in \mathcal{T}_i$, we have $S' \cap A \neq \emptyset$, so if $S' \not\subset A$, we find that $S' \cap \partial A \neq \emptyset$. We conclude that

$$\sum_{S' \in \mathcal{T}_i} |S' \setminus A| \leq \sum_{S' \in \mathcal{T}_i: \ S' \subset A} |S' \setminus A| + \sum_{S' \in \mathcal{T}_i: \ rad(S') > r} |S'| + \sum_{S' \in \mathcal{T}_i: \ S' \cap \partial A \neq \emptyset, \ rad(S') \leq r} |S'|$$
$$\leq 0 + \delta_t(A, B)^2 |A| + |\partial A + B^n(o, r)| \leq 2\delta_t(A, B)^2 |A|.$$

Combining this with the bound on S_i , we find

$$|\operatorname{co}(A) \setminus A| = \sum_{S' \in S_t \cup T_t} |S' \setminus A| \le c_n^{8.4} t^{-c^{8.4} c^{8.3} n^8 - n} \delta_t(A, B) |A| + 2\delta_t(A, B)^2 |A| \le t^{-c^{8.1} n^8} \delta_t(A, B) |A|,$$

and the result follows.

Proof of Propositions 8.5

Sublinearity of doubling in subsimplices; Proposition 8.2

Proposition 8.2 hinges on a geometric lemma.

Let S_i as in the statement of Proposition 8.2 and translate so that x is the origin. Consider the cones generated by the S_i , i.e., $\mathfrak{C}^{x_0,\dots,x_n}$.

We will use Proposition 5.4 repeatedly and Proposition 8.2 will follow quickly.

Proof of Proposition 8.2. Apply Proposition 5.4, find a v so that $|A \cap S_i| = |A \cap C_i| = |(B - v) \cap C_i|$ for all $C_i \in \mathfrak{C}^{x_0,\dots,x_n}$. We will show that the union $\bigcup_{i=0}^n D_t(A \cap S_i, B \cap (v+C_i)) \subset D_t(A,B)$ is an essentially disjoint union. Indeed note that, as C_i is a convex set,

$$D_t(A \cap S_i, B \cap (v + C_i)) \subset D_t(C_i, v + C_i) = (1 - t)v + C_i$$

and $\bigcup_i (1-t)v + C_i$ is an essentially disjoint union. We can conclude by setting $B_i = B \cap (v + C_i)$ and expanding

$$\sum_{i} |A \cap S_{i}| \cdot \delta_{t}(A \cap S_{i}, B_{i}) = \sum_{i} |D_{t}(A \cap S_{i}, B \cap (v + C_{i}))| - |A \cap S_{i}| \le |D_{t}(A, B)| - |A| \le |A| \cdot \delta_{t}(A, B).$$

The proposition follows.

8.5.2 Proof of Proposition 8.4

Proof of Proposition 8.4. Set $\eta_{\alpha,n}^{8.4} = \min(\eta_{2,\alpha,n}^{8.5}, \eta_{2,\eta_{1,\alpha,n}^{8.5},n}^{8.5})$. We apply Lemma 8.5 to the set X together with the special vertex x_n . Thus we construct a subset $Y \subset X$ with $|Y| \geq \eta_{1,\alpha,n}^{8.5}|S'|$ such that for all $x \in Y$ and $0 \leq i \leq n-1$, $|\cos\{x_0,\ldots,x_{i-1},x,x_{i+1},\ldots,x_n\} \cap X| \geq \eta_{2,\alpha,n}^{8.5}|S'|$. Now we apply again Lemma 8.5 to the set Y together with the special vertex x_{n-1} . Thus we construct a subset $Z \subset Y$ with $|Z| \geq \eta_{1,\eta_{1,\alpha,n}^{8.5},n}^{8.5}|S'|$ such that for all $x \in Z$ and $0 \leq i \leq n-2$ or i=n,

$$|\operatorname{co}\{x_0,\ldots,x_{i-1},x,x_{i+1},\ldots,x_n\}\cap Y| \ge \eta_{2,\eta_{i}}^{8.5} \cdot 5_{1,n} |S'|.$$

Fix $z \in Z \subset Y \subset X$. By the above inequalities, we conclude that for $0 \le i \le n$,

$$|\cos\{x_0,\ldots,x_{i-1},x,x_{i+1},\ldots,x_n\}\cap X| \ge \min\left(\eta_{2,\alpha,n}^{8.5},\eta_{2,\eta_{1,\alpha,n}^{8.5},n}^{8.5}\right)|S'| \ge \eta_{\alpha,n}^{8.4}|S'|.$$

For the last part, fix $0 \le k \le n$. Let q_k be the intersection of the ray $x_k x$ with the face opposite x_k . On the one hand, a simple computation gives $\frac{|xq_k|}{|x_kq_k|} = \frac{|\cos\{x_0,\dots,x_{i-1},x,x_{i+1},\dots,x_n\}|}{|\cos\{x_0,\dots,x_n\}|}$. From the first part, we know that $\frac{|\cos\{x_0,\dots,x_{i-1},x,x_{i+1},\dots,x_n\}|}{|\cos\{x_0,\dots,x_n\}|} \ge \eta_{\alpha,n}^{8.4}$. Combining the last two inequalities, we get $\frac{|xq_k|}{|x_kq_k|} \ge \eta_{\alpha,n}^{8.4}$ i.e., $\frac{|xx_k|}{|x_kq_k|} \le 1 - \eta_{\alpha,n}^{8.4}$. On the other hand, because the diameter of a simplex is realized between two vertices, we have $|x_kq_k| \le \max_{i,j}(x_ix_j)$. Combining the last two inequalities, we find $d(x,x_i) \le (1-\rho_{\alpha,n}^{8.4})\max_{i,j}d(x_i,x_j)$, which concludes the proposition.

8.6 Proof of Auxiliary Lemmas

8.6.1 Proof of Lemma 8.5

Proof of Lemma 8.5. Set $\eta_{1,\alpha,n}^{8.5} = \eta_{1,\frac{\alpha^{n+1}}{(2n)^{n+1}},n}^{8.6} \frac{\alpha^{n+1}}{(2n)^{n+1}}$ and $\eta_{2,\alpha,n}^{8.5} = \eta_{2,\frac{\alpha^{n+1}}{(2n)^{n+1}},n}^{8.6} \frac{\alpha^{n+1}}{(2n)^{n+1}}$. After an affine transformation, without loss of generality, we can assume that $S' = \operatorname{co}(\{o,e_1,\ldots,e_n\})$. For $(\lambda_1,\ldots,\lambda_n) \in S'$ construct the box $Q_{\lambda_1,\ldots,\lambda_n} = \{(p_1,\ldots,p_n) \colon \lambda_1 \geq p_1 \geq 0,\ldots,\lambda_n \geq p_n \geq 0\}$.

Claim 8.16. There exist $\lambda_1 \geq 0, \ldots, \lambda_n \geq 0$ with $\lambda_1 + \cdots + \lambda_n = 1$ such that $Q := Q_{\lambda_1, \ldots, \lambda_n}$ satisfies $|X \cap Q| \geq \frac{\alpha^{n+1}}{(2n)^{n+1}} |S'|$.

Proof of Claim 8.16. It is enough to prove that for $\lambda = (\lambda_1, \dots, \lambda_n) \in S'$ chosen uniformly at random, we have

$$\mathbb{E}_{\lambda}|X \cap Q| \ge \frac{\alpha^{n+1}}{(2n)^{n+1}}|S'|.$$

Note that for $1 \ge s \ge 0$ we have $(1-s)S' \subset S'$ and $|(1-s)S'| = (1-s)^n |S'| \ge (1-ns)|S'|$. Therefore

$$\left|X \cap \left(1 - \frac{\alpha}{2n}\right)S'\right| \ge |X \cap S'| - \left|S' \setminus \left(1 - \frac{\alpha}{2n}\right)S'\right| \ge \alpha|S'| - |S'| + \left|\left(1 - \frac{\alpha}{2n}\right)S'\right|$$

$$\ge \alpha|S'| - |S'| + \left(1 - \frac{\alpha}{2}\right)|S'| = \frac{\alpha}{2}|S'|.$$

Fix $x = (x_1, \ldots, x_n) \in S'$ and choose a random $\lambda = (\lambda_1, \ldots, \lambda_n) \in F$. We have

$$\begin{split} \mathbb{P}_{\lambda}(x \in Q_{\lambda}) &= \frac{|\{(\lambda_{1}, \dots \lambda_{n}) \colon \lambda_{i} \geq x_{i} \text{ for all } i \text{ and } \lambda_{1} + \dots + \lambda_{n} \leq 1\}|}{|\{(\lambda_{1}, \dots \lambda_{n}) \colon \lambda_{i} \geq 0 \text{ for all } i \text{ and } \lambda_{1} + \dots + \lambda_{n} \leq 1\}|} \\ &= \frac{|\{(\lambda_{1}, \dots \lambda_{n}) \colon \lambda_{i} \geq 0 \text{ for all } i \text{ and } \lambda_{1} + \dots + \lambda_{n} \leq 1 - (x_{1} + \dots + x_{n})\}|}{|\{(\lambda_{1}, \dots \lambda_{n}) \colon \lambda_{i} \geq 0 \text{ for all } i \text{ and } \lambda_{1} + \dots + \lambda_{n} \leq 1\}|} \\ &= (1 - (x_{1} + \dots + x_{n}))^{n}. \end{split}$$

In particular, if $x \in (1 - \frac{\alpha}{2n})S'$, we have $x_1 + \dots + x_n \leq 1 - \frac{\alpha}{2n}$ and thus $\mathbb{P}_{\lambda}(x \in Q_{\lambda}) \geq \frac{\alpha^n}{(2n)^n}$. Putting all together, we conclude

$$\mathbb{E}_{\lambda}|X \cap Q| = \int_{x \in X} \mathbb{P}_{\lambda}(x \in Q_{\lambda}) \ge \int_{x \in X \cap (1 - \frac{\alpha}{2n})S'} \mathbb{P}_{\lambda}(x \in Q_{\lambda}) \ge \int_{x \in X \cap (1 - \frac{\alpha}{2n})S'} \frac{\alpha^{n}}{(2n)^{n}}$$
$$= \left| X \cap \left(1 - \frac{\alpha}{2n} \right) S' \right| \frac{\alpha^{n}}{(2n)^{n}} \ge \frac{\alpha}{2} \frac{\alpha^{n}}{(2n)^{n}} |S'| \ge \frac{\alpha^{n+1}}{(2n)^{n+1}}.$$

By Claim 8.16, we find $\lambda = (\lambda_1, \dots, \lambda_n) \in S'$ and $Q = Q_{\lambda}$ such that $|X \cap Q| \ge \frac{\alpha^{n+1}}{(2n)^{n+1}} |S'|$. Note that $Q \subset S'$. Hence $|X \cap Q| \ge \frac{\alpha^{n+1}}{(2n)^{n+1}} |Q|$. Set $X' = X \cap Q$.

We apply Lemma 8.6 to the set X'. Thus, we construct a subset $Y \subset X' \subset X$ with $|Y| \geq \eta_{1,\frac{\alpha^{n+1}}{(2n)^{n+1}},n}^{8.6}|Q|$ such that for all $x \in Y$ and all faces F of the box Q, we have $|\cos(\{x\} \cup F) \cap X'| = \eta_{2,\frac{\alpha^{n+1}}{(2n)^{n+1}},n}^{8.6}|Q|$. As $|Q| \geq |X \cap Q| \geq \frac{\alpha^{n+1}}{(2n)^{n+1}}|S'|$, this implies that

$$|Y| \ge \eta_{1,\frac{\alpha^{n+1}}{(2n)^{n+1}},n}^{8\cdot6} \frac{\alpha^{n+1}}{(2n)^{n+1}} |S'| \quad \text{ and } \quad |\operatorname{co}(\{x\} \cup F) \cap X| = \eta_{2,\frac{\alpha^{n+1}}{(2n)^{n+1}},n}^{8\cdot6} \frac{\alpha^{n+1}}{(2n)^{n+1}} |S'|.$$

Moreover, note that each face of S', except the face opposite to o, contains a face of Q inside it. Fix $1 \le i \le n$ and let F be the face of Q such that $F \subset co(0, e_1, \ldots, e_{i-1}, e_{i+1}, \ldots, e_n)$. We conclude that for $y \in Y$, we have

$$|\operatorname{co}(x_0,\ldots,x_{i-1},y,x_{i+1},\ldots,x_n)\cap X| \ge |\operatorname{co}(\{y\}\cup F)\cap X| \ge \eta_{2,\frac{\alpha^{n+1}}{(2n)^{n+1}},n}^{8.6} \frac{\alpha^{n+1}}{(2n)^{n+1}}|T|.$$

8.6.2 Proof of Lemma 8.6

Proof of Lemma 8.6. Set $\eta_{1,\alpha,n}^{8.6} = 2^{-1}\eta_{1,\alpha/16n,n}^{8.8}$ and $\eta_{2,\alpha,n}^{8.6} = 2^{-1}\eta_{2,\alpha/16n,n}^{8.8}$. By taking an affine transformation, without loss of generality, we can assume that $Q = [-(1+\frac{\alpha}{4n}),(1+\frac{\alpha}{4n})]^n$.

Let $Q' = [-1, 1]^n$. For $1 \le i \le n$ let F_i be the face of the box Q in direction e_i and for $-n \le i \le -1$ let F_i be the face of the box Q in direction $-e_{-i}$. So $F_{-n}, \ldots, F_{-1}, F_1, \ldots F_n$ are the faces of Q.

Let $X' = X \cap Q'$. Note that

$$|X'| \ge |X| + |Q'| - |Q| \ge \alpha |Q| + |Q'| - |Q| = \left(1 - (1 - \alpha)\left(1 + \frac{\alpha}{4n}\right)^n\right)|Q'| \ge \frac{\alpha}{2}|Q'|.$$

We apply Lemma 8.8 with parameter $\alpha/16n$ to the set X'. Thus, we construct a subset $Y \subset X'$ with $|Y| \ge \eta_{1,\alpha/16n,n}^{8.8}|Q'|$ such that for all $x \in Y$ and all $j \in [-n,n] \setminus \{0\}$, we have $|(x+C_{j,\alpha/16n}) \cap X'| \ge \eta_{2,\alpha/16n,n}^{8.8}|Q'|$. As $|Q'| = (1+\frac{\alpha}{4n})^{-n}|Q| \ge \frac{1}{2}|Q|$ this implies that

$$|Y| \ge 2^{-1} \eta_{1,\alpha/16n,n}^{8.8} |Q|$$
 and $|(x + C_{j,\alpha/16n}) \cap X| \ge 2^{-1} \eta_{2,\alpha/16n,n}^{8.8} |Q|$.

For $x \in Q'$ and $j \in [-n,n] \setminus \{0\}$ it is easy to check that $(x + C_{j,\alpha/16n}) \cap Q \subset \operatorname{co}(\{x\} \cup F_j)$ and as $X \subset Q$, this implies that $(x + C_{j,\alpha/16n}) \cap X \subset \operatorname{co}(\{x\} \cup F_j) \cap X$. We conclude that for $x \in Y$ and $j \in [-n,n] \setminus \{0\}$, we have

$$|\operatorname{co}(\{x\} \cup F_j) \cap X| \ge 2^{-1} \eta_{2,\alpha/16n,n}^{8.8} |\operatorname{co}(X)|.$$

8.6.3 Proof of Lemma 8.8

Proof of Lemma 8.8. Set $\eta_{1,\alpha,n}^{8.8} = \eta_{2,\alpha,n}^{8.8} = \alpha^{4n^2+2n-1}/2^{16n^3+14n^2+11n-5}$. Given a number i, let $\overline{i} \in [n]$ be the unique number such that $\overline{i} = i \mod n$. Set $\widehat{i} = \overline{i}$ if i < n and $\widehat{i} = -\overline{i}$ if $i \ge n$. Our strategy is to apply repeatedly analogues of Lemma 8.9 where instead of focusing on the pair of coordinates (n,1), we focus on the pair of coordinates $(\overline{i},\overline{i+1})$.

By Claim 8.18, there exists a box $Q' \subset Q$,

$$Q' = [0, b_1] \times \ldots \times [0, b_n] + y$$
 where $y \in \mathbb{R}^n$, such that $b_n = 2$ and $b_n \frac{(\alpha/4)^2}{8} \le b_1, \ldots, b_{n-1} \le b_n \frac{(\alpha/4)^2}{4}$,

and such that $X \cap Q'$ has density $|X \cap Q'| \ge \alpha |Q'|/4$. In particular $|Q'| \ge |Q|\alpha^{2n-2}/2^{7n-7}$

By applying Lemma 8.9, we construct inductively the sequence of pairs $(X^0, Q^0), \ldots, (X^{2n}, Q^{2n})$ with the following properties. Define $Q^0 = Q'$ and $X^0 = X \cap Q'$. Assume that at step i-1 we have constructed a box

$$Q^{i-1} = [0, a_1^{i-1}] \times \dots \times [0, a_n^{i-1}] + x^{i-1} \quad \text{ with } a_{i-1}^{i-1} \frac{(\alpha/4^i)^2}{8} \le a_j^{i-1} \le a_{i-1}^{i-1} \frac{(\alpha/4^i)^2}{4} \text{ for } j \in [n] \setminus \overline{i-1},$$

and a subset $X^{i-1} \subset Q^{i-1}$ with density $|X_{i-1}| \ge \alpha |Q^{i-1}|/4^i$. We apply Lemma 8.9 to the pair (X^{i-1}, Q^{i-1}) focusing on the pair of coordinates $(\overline{i-1}, \overline{i})$ to produce the pair (X^i, Q^i) . More precisely, we obtain a box $Q^i \subset Q^{i-1}$ with

$$Q^i = [0, a_1^i] \times \dots \times [0, a_n^i] + x^i \quad \text{ where } a_{\overline{i}}^i \frac{(\alpha/4^{i+1})^2}{8} \le a_j^i \le a_{\overline{i}}^i \frac{(\alpha/4^{i+1})^2}{4} \text{ for } j \in [n] \setminus \overline{i}.$$

We also obtain the set $X^i = X^{i-1} \cap Q^i$. These have the properties $|Q^i| \ge (\alpha/4^i)^{2n} |Q^{i-1}|/2^{7n}, |X^i| \ge \alpha |Q^i|/4^{i+1}$, and

$$|(x + C_{\widehat{i-1}\alpha}) \cap X^{i-1}| \ge |(x + C_{\widehat{i-1}\alpha/4^i}) \cap X^{i-1}| \ge \alpha |Q^{i-1}|/4^i$$
 for all $x \in X^i$.

Because $X^{2n} \subset \cdots \subset X^0 \subset X$, $Q^{2n} \subset \cdots \subset Q_0$ and $\{\widehat{0}, \widehat{1}, \ldots, \widehat{2n-1}\} = [-n, n] \setminus \{0\}$, it follows that for $x \in X^{2n}$ and $j \in [-n, n] \setminus \{0\}$ we have $|(x + C_{j,\alpha}) \cap X| \ge \alpha |Q^{2n}|/4^{2n}$. Moreover, $|X^{2n}| \ge \alpha |Q^{2n}|/4^{2n+1}$. It remains to evaluate $|Q^{2n}|$. Recall that $|Q^0| \ge |Q|\alpha^{2n-2}/2^{7n-7}$. Using the recurrence relation

$$|Q^i| \ge (\alpha/4^i)^{2n} |Q^{i-1}|/2^{7n} \ge \alpha^{2n} |Q^{i-1}|/2^{8n^2+7n},$$

we get $|Q^{2n}| \ge |Q^0|\alpha^{4n^2}/2^{16n^3+14n^2}$. These imply $|Q^{2n}| \ge \alpha^{4n^2+2n-2}|Q|/2^{16n^3+14n^2+7n-7}$. Thus, we conclude that the subset $X^{2n} \subset X$ satisfies

$$|X^{2n}| \ge \alpha |Q^{2n}|/4^{2n+1} \ge \alpha^{4n^2+2n-1}|Q|/2^{16n^3+14n^2+11n-5},$$

and for all $x \in X^{2n}$ and $j \in [-n, n] \setminus \{0\}$ we have

$$|(x+C_{j,\alpha})\cap X| \ge \alpha |Q^{2n}|/4^{2n} \ge \alpha^{4n^2+2n-1}|Q|/2^{16n^3+14n^2+11n-7}.$$

8.6.4 Proof of Lemma 8.9

Proof of Lemma 8.9.

Claim 8.17. Given a box $R = [0, c_1] \times \cdots \times [0, c_n] + z$ where $z \in \mathbb{R}^n$ and given a parameter $s \leq \min(c_1, \ldots, c_n)$, we can construct a partition into boxes with disjoint interiors $R = \bigsqcup_{i=1}^k R^i$ where $R^i = [0, c_1^i] \times \cdots \times [0, c_n^i] + z^i$ such that $z^i \in \mathbb{R}^n$ and $c_1^i = c_1$ and $s/2 \leq c_2^i, \ldots, c_n^i \leq s$.

Proof of Claim 8.17. For $2 \leq j \leq n$ partition the interval $[0, c_j]$ into consecutive intervals $[0, c_j] = I_1^j \sqcup \cdots \sqcup I_{\lfloor 2c_j/s \rfloor}^j$ where

$$I_k^j = [(k-1)s/2, ks/2] \quad \text{ for } 1 \leq k \leq \lfloor 2c_j/s \rfloor - 1, \qquad I_{\lfloor 2c_j/s \rfloor}^j = [(\lfloor 2c_j/s \rfloor - 1)s/2, c_j].$$

Let $[0, c_1] = I_1^1$. Finally, for $k_1 = 1, 1 \le k_2 \le \lfloor 2c_2/s \rfloor, \dots, 1 \le k_n \le \lfloor 2c_n/s \rfloor$ construct the box $R^{k_1, k_2, \dots, k_n} = I_{k_1}^1 \times \dots \times I_k^n + z$.

Note that $R = \bigsqcup_{k_1,\ldots,k_n} R^{k_1,k_2,\ldots,k_n}$ is a partition into boxes with disjoint interiors. Moreover, the box $R^{k_1,k_2,\ldots,k_n} = I^1_{k_1} \times \cdots \times I^n_{k_n} + z$ satisfies $|I^1_{k_1}| = c_1$, and for $2 \le j \le n$ we have $|I^j_{k_j}| = s/2$ if $1 \le k_j < \lfloor 2c_j/s \rfloor$, and $|I^j_{\lfloor 2c_j/s \rfloor}| = c_j - (\lfloor 2c_j/s \rfloor - 1)s/2 \in [s/2,s]$. We conclude that this partition into boxes has the desired properties.

Claim 8.18. Given a box $R = [0, c_1] \times \cdots \times [0, c_n] + z$ where $z \in \mathbb{R}^n$ and $c_1 \leq 2 \min(c_2, \ldots, c_n)$ and given a subset $Z \subset R$ with density $|Z| \geq \alpha |R|/4$, there exist a sub-box of R, $Q = [0, b_1] \times \cdots \times [0, b_n] + y$ where $y \in \mathbb{R}^n$ such that $b_1 = c_1$ and $b_1(\alpha/4)^2/8 \leq b_2, \ldots, b_n \leq b_1(\alpha/4)^2/4$, and such that $Y := Z \cap Q$ has density $|Y| \geq \alpha |Q|/4$.

Proof of Claim 8.18. Set $s = c_1(\alpha/4)^2/4$ and note that, because $c_1 \leq 2 \min(c_2, \ldots, c_n)$, we have $s \leq \min(c_1, \ldots, c_n)$. By Claim 8.17, we can construct a partition into boxes with disjoint interiors $R = \bigsqcup_{i=1}^k R^i$ where $R^i = [0, c_1^i] \times \cdots \times [0, c_n^i] + z^i$ such that $z^i \in \mathbb{R}^n$ and $c_1^i = c_1$ and $c_1(\alpha/4)^2/8 \leq c_2^i, \ldots, c_n^i \leq c_1(\alpha/4)^2/4$. A simple averaging argument shows that there exists an index i such that $\frac{|Z \cap R^i|}{|R_i|} \geq \frac{|Z \cap R|}{|R|} = \alpha/4$.

Set $Q = R^i$, $Y = Z \cap Q$, $b_1 = c_1^i$,..., $b_n = c_n^i$ and $y = z^i$. It is easy to check that $Q = [0, b_1] \times ... \times [0, b_n] + y$ where $b_1 = c_1$ and $b_1(\alpha/4)^2/8 \le b_2, ..., b_n \le b_1(\alpha/4)^2/4$ and $|Y| \ge \alpha |Q|/4$.

Fix $0 \le r \le a_n(1-\alpha/4)$ with the following property. Consider the partition of P into three box with disjoint interiors $P = R_1 \sqcup R_2 \sqcup R_3$ where $R_1 = [0, a_1] \times \cdots \times [0, r] + x$, $R_2 = [0, a_1] \times \cdots \times [r, r + a_n\alpha/4] + x$ and $R_3 = [0, a_1] \times \cdots \times [r + a_n\alpha/4, a_n] + x$. Let $Z_1 = X \cap R_1$, $Z_2 = X \cap R_2$ and $Z_3 = X \cap R_3$. We choose r such that $|Z_1| = |Z_3|$.

Claim 8.19. $a_1 \le 2 \min(a_2, \dots, a_{n-1}, r)$ and $|Z_1| = |Z_3| \ge \alpha |P|/4 \ge \alpha |R_1|/4$.

Proof of Claim 8.19. By construction $|Z_2| \le |R_2| \le \alpha |P|/4$. By hypothesis, $|Z_1| + |Z_2| + |Z_2| = |X| \ge \alpha |P|$. Therefore, $|Z_1| + |Z_3| \ge 3\alpha |P|/4$. As $|Z_1| = |Z_3|$, we conclude $|Z_1| = |Z_3| \ge \alpha |P|/4 \ge \alpha |R_1|/4$.

From hypothesis $a_n\alpha^2/8 \le a_1, \ldots, a_{n-1} \le a_n\alpha^2/4$, it immediately follows that $a_1 \le 2\min(a_2, \ldots, a_{n-1})$. For $a_1 \le 2r$, note that

$$r = a_n \frac{|R_1|}{|P|} \ge a_n \frac{|Z_1|}{|P|} \ge a_n \frac{\alpha}{4} > a_n \frac{\alpha^2}{8}.$$

Observation 8.20. It follows immediately from the definition of $C_{n,\alpha}$ and the hypothesis $a_1, \ldots, a_{n-1} \leq a_n \alpha^2/4$ that for $y \in R_1$ we have $y+C_{n,\alpha} \supset R_3$. In particular, by Claim 8.19, for $y \in Z_1$ we have $|(y+C_{n,\alpha}) \cap X| \geq \alpha |P|/4$.

To finish the proof, let recall $R_1 = [0, a_1] \times \cdots \times [0, r] + x$ and $Z_1 = R_1 \cap X$. By Claim 8.19, we have $a_1 \leq 2 \min(a_2, \ldots, a_{n_1}, r)$ and $|Z_1| \geq \alpha |R_1|/4$. By Claim 8.18 there exist a sub-box of R_1 ,

$$Q = [0, b_1] \times ... \times [0, b_n] + y$$
 where $y \in \mathbb{R}^n$, such that $b_1 = a_1$ and $b_1(\alpha/4)^2/8 \le b_2, ..., b_n \le b_1(\alpha/4)^2/4$,

and such that $Y := Z_1 \cap Q = X \cap Q$ has density $|Y| \ge \alpha |Q|/4$. As $Y \subset Z_1$, by Observation 8.20 for all $y \in Y |(y + C_{n,\alpha}) \cap X| > \alpha |P|/4$.

It remains to check that $|Q| \ge \alpha^{2n}/4^{3n-2}2^{n-1}|P|$. This easily follows from the aforementioned fact that $b_1 = a_1, a_1(\alpha/4)^2/4 \le b_2, \ldots, b_n$ and the fact that $a_1 \ge 2^{-1} \max(a_2, \ldots, a_{n-2})$ and $a_1 \ge a_n \alpha^2/8$ (the last fact follows from the hypothesis $a_n \alpha^2/8 \le a_1, \ldots, a_{n-1} \le a_n \alpha^2/4$).

8.6.5 Proof of Lemma 8.10

Proof of Lemma 8.10. Let the set $H_1, \ldots, H_m \subset \mathbb{R}^n$ be a finite collection of halfspaces so that $C = \bigcap H_i$. Consider the sequence $A_i := A \cap \bigcap_{j=1}^i H_i$. Construct another set of halfspaces H'_i and subsets $B_i \subset B$ defined by finding a halfspace H'_i parallel to H_i so that $|H'_i \cap B_{i-1}| = |A_i|$. Note that because H_i is parallel to H'_i we find that $tA_i + (1-t)B_i$ is disjoint from $t(A_{i-1} \setminus A_i) + (1-t)(B_{i-1} \setminus B_i)$. By the Brunn-Minkowksi inequality we find

$$|tA_{i-1} + (1-t)B_{i-1}| - |A_{i-1}| \ge |tA_i + (1-t)B_i| - |A_i| + |t(A_{i-1} \setminus A_i) + (1-t)(B_{i-1} \setminus B_i)| - |A_{i-1} \setminus A_i|$$

$$\ge |tA_i + (1-t)B_i| - |A_i|.$$

The lemma follows by induction.

9 Proof of the Linear Theorem for few vertices (Theorem 1.6)

Proof of Theorem 1.6. Let $c^{1.6} = c^{8.1}$ and $k_n^{1.6} = \binom{v}{n}$. Let λ_n be anything, and choose η_n so that $(1 + \eta_n)^{-n} = \frac{1}{2}$. By Proposition 2.9, we may assume that A, B form a simple λ_n -bounded η_n - sandwich.

Consider the origin $o \in A \cap B$ and a triangulation \mathcal{T}' of the boundary of co(A). Note that $|\mathcal{T}'| \leq {v \choose n}$. Now consider the triangulation \mathcal{T} of $\partial co(A)$ obtained by adding o to each of the simplices of \mathcal{T}' , i.e.

$$\mathcal{T} := \{\operatorname{co}(S' \cup \{o\}) : S' \in \mathcal{T}'\}.$$

We will consider $|\operatorname{co}(A)\setminus A|$ inside each of the simplices in \mathcal{T} . Write $A_{S'}$ for $A\cap S'$ and note that $\operatorname{co}(A_{S'})=S'$. Using Lemma 8.10, find $B_{S'}\subset B$ so that $|A_{S'}|=|B_{S'}|$ and $|tA_{S'}+(1-t)B_{S'}|\leq |tA+(1-t)B|\leq \delta|A|$. Distinguish two cases; either $\delta|A|<\min\{t,1-t\}^n|A_{S'}|$ or $\delta|A|\geq \min\{t,1-t\}^n|A_{S'}|$. In the former case, we can apply Theorem 8.1 to find $|\operatorname{co}(A_{S'})\setminus A_{S'}|\leq \min\{t,1-t\}^{-c}^{8.1}n^8\delta|A|$. In the latter case we use that A is a η_n sandwich so that $\frac{1}{1+\eta_n}\operatorname{co}(A_{S'})\subset A_{S'}$, which implies

$$|\operatorname{co}(A_{S'}) \setminus A_{S'}| \le \left| \operatorname{co}(A_{S'}) \setminus \frac{1}{1+\eta_n} \operatorname{co}(A_{S'}) \right| \le \left(1 - (1+\eta_n)^{-n}\right) |\operatorname{co}(A_{S'})|$$

$$\le \frac{1}{2} |\operatorname{co}(A_{S'})| \le |A_{S'}| \le \min\{t, 1-t\}^{-n} \delta |A|.$$

Hence, in both cases we find $|\operatorname{co}(A_{S'})\setminus A_{S'}| \leq \min\{t, 1-t\}^{-c^{8.1}n^8}\delta|A|$. We conclude by adding up the contributions over all $S' \in \mathcal{T}$,

$$|\operatorname{co}(A) \setminus A| = \sum_{S' \in \mathcal{T}} |\operatorname{co}(A_{S'}) \setminus A_{S'}| \le |\mathcal{T}| \min\{t, 1 - t\}^{-c^{8.1} n^8} \delta |A| \le {v \choose n} \min\{t, 1 - t\}^{-c^{8.1} n^8} \delta |A|.$$

This concludes the proof of the theorem.

10 Intermediate results for the Linear Theorem: from few vertices (Theorem 1.6) to the general case (Theorem 1.4)

10.1 Outline of the proof of Theorem 1.4

The proof of Theorem 1.4 follows the following steps.

- 1. By Proposition 2.9, we may assume $B(o, 0.01) \subset K \subset A, B \subset 1.01K \subset B(0, 100)$ for some convex set K.
- 2. Find a collection of disjoint convex regions X_i with small diameter ξ so that $\sum_i |\operatorname{co}(A \cap X_i) \setminus (A \cap X_i)| \ge \Omega_n(|\operatorname{co}(A) \setminus A|)$ (Proposition 10.3) along the following steps.
 - Cone off at the origin to find simplices T_i so that $\bigcup T_i = \operatorname{co}(A)$ and $\operatorname{co}(T_i \cap A) = T_i$. Note that $|T_i \cap A| \ge 0.99|T_i|$.
 - In each of the T_i run the following process: find a point in A centrally in T_i and cone off to the vertices of T_i to find smaller simplices with the same properties as T_i and not much lower density. (Lemma 10.6)
 - Iterate until either the diameter is small (in which case we have found our region X_i) or the density has dropped (say between 0.9 and 0.95). (Lemma 10.7)
 - For simplices with moderate density, it is easy to find a set of small diameter and a positive proportion of the missing region. (Lemma 10.8)
- 3. Consider the tube $U = S \times \mathbb{R}^+$ where $S \subset \mathbb{R}^{n-1}$ is a regular simplex with side length ϵ centered at the origin. We choose $\epsilon \ll 0.01$, so that S falls well within B(o, 0.01). On the other hand, we choose $\epsilon \gg \xi$, so that a random rotation of U contains a region X_i completely with probability $\Omega_n(1)$.
- 4. Note that $|\cos(A \cap U) \setminus (A \cap U)| \ge \sum_{X_i \subset U} |\cos(A \cap X_i) \setminus (A \cap X_i)|$, so that it remains to show $|\cos(A \cap U) \setminus (A \cap U)| \le O_{n,t}(|tA + (1-t)B| |A|)$. (Proposition 10.4)
- 5. To this ends find a homothetic tube $V = x + (1 + \beta)U$ so that $|A \cap U| = |B \cap V|$ and $|t(A \cap U) + (1 t)(B \cap V)| |A \cap U| \le |tA + (1 t)B| |A|$, which moreover is very similar to U, so that x and β are very small. (Lemma 10.9)
- 6. Slightly rescaling $A \cap U$ and $B \cap V$ (i.e., taking homotheties with factors very close to 1), we find A', B' in a tube over the same simplex W so that $2|\cos(A') \setminus A'| \ge |\cos(A \cap U) \setminus (A \cap U)|$ and a $t' \in (\frac{1}{2}t, 2t)$ so that $|t'A' + (1-t')B'| t'|A'| (1-t')|B'| \le |tA + (1-t)B| |A|$. (Lemma 10.10)
- 7. Partition the tube W into parallel simplicial tubes U_i according to the convex hull of A', so that $\bigcup_i \operatorname{co}(A' \cap U_i) = \operatorname{co}(A')$ and each of the $\operatorname{co}(A' \cap U_i)$ has exactly 2n vertices. Note that we have

$$\sum_{i} |t'(A' \cap U_i) + (1 - t')(B' \cap U_i)| - t'|A' \cap U_i| - (1 - t')|B' \cap U_i| \le |t'A' + (1 - t')B'| - t'|A'| - (1 - t')|B'|.$$

- 8. Consider the fibres in the direction of the tube and note that A' and B' start with a long interval in each of these fibres by the virtue of S lying well within B(o, 0.01). We might not have $|A' \cap U_i| = |B' \cap U_i|$, but because of the long interval at the beginning of each fibre, we may extend A' and B' without affecting $|t'(A' \cap U_i) + (1 t')(B' \cap U_i)| t'|A' \cap U_i| (1 t')|B' \cap U_i|$. (Lemma 10.11)
- 9. Because $K \subset A, B \subset 1.01K$, we never have to extend much to get $|A' \cap U_i| = |B' \cap U_i|$. Hence, we can apply Theorem 1.6 to find that

$$|\operatorname{co}(A' \cap U_i) \setminus (A' \cap U_i)| \le O_{n,t}(|t'(A' \cap U_i) + (1 - t')(B' \cap U_i)| - t'|A' \cap U_i| - (1 - t')|B' \cap U_i|),$$

so that summing over U_i gives the result.

10.2 Setup

Definition 10.1. A vertical tube with diameter ℓ is a set U of the form $U = S \times \mathbb{R}_+$ where $S \subset \mathbb{R}^{n-1}$ is a regular simplex with side length ℓ centered at the origin. A tube with diameter ℓ is a rotated vertical tube with diameter ℓ .

Definition 10.2. A pair $X, Y \subset \mathbb{R}^n$ is a tubular λ -bounded η -sandwich if it is the intersection between the same tube and a λ -bounded η -sandwich.

10.3 Propositions

Proposition 10.3. For $n \in \mathbb{N}$ and $\xi > 0$, there are constants $\theta > 0$ and $\eta > 0$ such that the following hold. Assume $A \subset \mathbb{R}^n$ is a measurable η -sandwich, then there exist disjoint measurable subsets A_1, \ldots, A_k of A with the following properties:

- 1. $A_i = A \cap \operatorname{co}(A_i)$,
- 2. $co(A_1), \ldots, co(A_k)$ are disjoint,
- 3. $\sum_{i} |\operatorname{co}(A_i) \setminus A_i| \ge \theta |\operatorname{co}(A) \setminus A|$
- 4. for every i, $diam(co(A_i)) < \xi diam(co(A))$.

Proposition 10.4. For $n \in \mathbb{N}$, $\lambda > 1$, $\varepsilon > 0$ sufficiently small in terms of n and λ , and $0 < t \le \frac{1}{2}$, there exists $\eta > 0$ such that the following hold. Assume $U \subset \mathbb{R}^n$ is a tube with diamater ε and assume $A, B \subset \mathbb{R}^n$ form a λ -bounded, η -sandwich of measurable sets of volume 1, then

$$|\operatorname{co}(A \cap U) \setminus (A \cap U)| \le k_n^{10.4} t^{-c^{10.4} n^8} (|tA + (1-t)B| - |A|).$$

Proposition 10.5. For $n \in \mathbb{N}$, $\lambda > 1$, $\varepsilon > 0$ sufficiently small in terms of n and λ , and $t \in (0,1)$, there exists $\eta_{n,\lambda,\varepsilon,t}^{10.5}, k_n^{10.5}, c^{10.5} > 0$ such that the following holds. Assume $W \subset \mathbb{R}^n$ is a tube or diameter ϵ and assume the subsets $A', B' \subset W$ form a tubular λ -bounded η -sandwich. Then

$$|\operatorname{co}(A') \setminus A'| \le k_n^{10.5} (t')^{-c^{10.5} n^8} (|t'A' + (1 - t')B'| - t'|A'| - (1 - t')|B'|).$$

10.4 Lemmas

Lemma 10.6. For all $n \in \mathbb{N}$ there exists $\alpha > 0$ depending n such that the following holds. If A is a subset of \mathbb{R}^n with co(A) a simplex with vertices $V(co(A)) = \{x_0, \ldots, x_n\}$ and if $|co(A)| \le (1+\alpha)|A|$, then there exists a point $x \in A$ such that for all $0 \le i \le n$

$$\frac{|\cos\{x_0,\ldots,x_{i-1},x,x_{i+1},\ldots,x_n\}|}{|\cos\{x_0,\ldots,x_{i-1},x_i,x_{i+1},\ldots,x_n\}|} \ge \frac{1}{n+2} \quad and \quad d(x,x_i) \le \frac{n+1}{n+2} \max_{j,k} d(x_j,x_k).$$

Lemma 10.7. For all $n \in \mathbb{N}$ and $\alpha > 0$ sufficiently small in terms of n, there exists an $\eta > 0$ so that the following holds. Given an η -sandwich A there exists an essential partition into convex sets $co(A) = P_1 \sqcup \cdots \sqcup P_k$ so that denoting $A_i = P_i \cap A$ we have:

- 1. $P_i = co(A_i),$
- 2. $|\operatorname{co}(A_i) \cap \operatorname{co}(A_j)| = 0 \text{ if } i \neq j,$
- 3. $co(A_i)$ is a simplex,
- 4. Each A_i fall into two categories, i.e., $[k] = I \sqcup J$ so that:
 - for all $i \in I$, we have $\frac{|\operatorname{co}(A_i)|}{|A_i|} \in (1 + \alpha, 1 + n!\alpha)$,
 - $\sum_{I \in I} |\operatorname{co}(A_i) \setminus A_i| \leq \alpha |\operatorname{co}(A) \setminus A|$.

The same conclusion holds if we replace the hypothesis that A is an η -sandwich with the hypothesis that co(A) is a simplex with $|co(A)| \le (1+\alpha)|A|$.

Lemma 10.8. For all $n \in \mathbb{N}$, $\alpha > 0$ sufficiently small in terms of n, there exist $\zeta, \beta > 0$ so that the following holds. Given $A \subset \mathbb{R}^n$ so that co(A) is a simplex and $|co(A)| = (1 + \alpha)|A|$, there exists a subset $A' \subset A$ and $\psi \geq \zeta$, so that

- 1. $diam(A') \leq (1 \psi)diam(A)$,
- 2. $|\operatorname{co}(A')| \ge (1 \psi)^{3n^2} |\operatorname{co}(A)|,$
- 3. co(A') is a simplex,
- 4. $|\operatorname{co}(A') \setminus A'| \ge \beta |A'|$.

Lemma 10.9. For $n \in \mathbb{N}$, $\lambda > 1$, for all $\varepsilon > 0$ sufficiently small in terms of λ and n, so that for all $0 < t \le \frac{1}{2}$, $\alpha > 0$ there exists $\eta > 0$ such that the following hold. Assume $U \subset \mathbb{R}^n$ is a tube with diameter 2ε and assume $A, B \subset \mathbb{R}^n$ form a λ -bounded, η -sandwich measurable sets of volume 1, then there exists a homothetic tube $V = x + (1 + \beta)U$ with $||x||_2 \le \alpha$ and $|\beta| \le \alpha$ such that $A^{\circ} = A \cap U$ and $B^{\circ} = B \cap V$ satisfy $|A^{\circ}| = |B^{\circ}|$ and

$$|tA^{\circ} + (1-t)B^{\circ}| - |A^{\circ}| \le |tA + (1-t)B| - |A|.$$

Lemma 10.10. For all $\eta' > 0$, $n \in \mathbb{N}$, $t \in (0,1/2]$, $\lambda > 1$ and sufficiently small ϵ in terms of n and λ , there exist $\eta, \alpha > 0$, so that the following holds. Let $t' = \frac{t}{1+\beta-\beta t}$. Assume $U \subset \mathbb{R}^n$ is a tube of diameter ϵ , V is a homothetic tube $V = x + (1+\beta)U$ with $|\beta|, ||x||_2 \le \alpha$ and assume A, B is a λ -bounded η -sandwich with |A| = |B| = 1 and $A^{\circ} := A \cap U$ and $B^{\circ} := V \cap B$ have equal size. Set $W = \frac{t}{t'}U$, $W' = \frac{1-t}{1-t'}V$, $A' = \frac{t}{t'}A^{\circ}$ and $B' = \frac{1-t}{1-t'}B^{\circ}$. Then the tube $W' = W + \frac{1-t}{1-t'}x$ has diameter less than 2ϵ , $A' \subset W$ and $B' \subset W'$. Moreover, we have

$$|t'A' + (1 - t')B'| - t'|A'| - (1 - t')|B'| \le |tA^{\circ} + (1 - t)B^{\circ}| - |A^{\circ}|.$$

Furthermore, $A', B' - \frac{1-t}{1-t'}x$ form a tubular 2λ -bounded η' -sandwich. Finally, if A°, B° are α -almost convex, then $|A^{\circ} \triangle B^{\circ}| \leq |A' \triangle (B' - \frac{1-t}{1-t'}x)| + O_{n,\epsilon}(\alpha |A^{\circ}|)$

Lemma 10.11. For $n \in \mathbb{N}$, $\lambda > 1$, $\varepsilon > 0$ sufficiently small in terms of n and λ , and $t \in (0,1)$, there exists $\eta > 0$ such that the following holds. Given a convex $K \subset \mathbb{R}^n$ which is λ -bounded and a tube $U = S \times \mathbb{R}_+$ of radius ε . Then for all $x \in S \times \mathbb{R}_-$, we have $(1-t)x + t(U \cap ((1+\eta)K \setminus K)) \subset K \cup (S \times \mathbb{R}_-)$.

10.5 Proof of Propositions

10.5.1 Proof of Proposition 10.3

Proof of Proposition 10.3. Choose parameters according to the following hierarchy

$$n, \xi \gg \alpha \gg \zeta \gg \beta \gg \theta \gg \eta$$
.

We create a sequence of families \mathcal{F}_i of sets which all have properties 1 and 2. They have property 3 with ever decreasing θ while the diameter of the sets decreases until property 4 is satisfied. All sets in the families will have a simplex as their convex hull. Given a family \mathcal{F} of subsets of A satisfying properties 1 and 2, write $D(\mathcal{F}) := \max_{X \in \mathcal{F}} \frac{diam(X)}{diam(A)}$ and $P(\mathcal{F}) := \frac{\sum_{X \in \mathcal{F}} |\cos(X) \setminus X|}{|\cos(A) \setminus A|}$, so that we want to find a family with $D(\mathcal{F}) < \xi$ and $P(\mathcal{F}) \ge \theta$.

First apply Lemma 10.7 with parameter α to find a partition and let \mathcal{F}_0 be the set of parts so that $\frac{|\cos(A_i)|}{|A_i|} \in (1 + \alpha, 1 + n!\alpha)$. Note that Lemma 10.7 says that $P(\mathcal{F}_0) \geq 1 - \alpha$.

Consider the following process of obtaining a new family \mathcal{F}_{2i+1} from \mathcal{F}_{2i} and \mathcal{F}_{2i+2} from \mathcal{F}_{2i+1} . All elements X in \mathcal{F}_i with $diam(X) \leq \xi diam(A)$ remain fixed and stay in all consequent families, we will call them *finished*. The idea will be to apply Lemma 10.8 to all other elements in \mathcal{F}_{2i} to reduce the diameter. Then we apply Lemma 10.7 to some of the elements in \mathcal{F}_{2i+1} so that the density of the sets in \mathcal{F}_i in their convex hull always stays below $1/(1+\alpha)$.

All non-finished elements $X \subset A$ in \mathcal{F}_{2i} have $\frac{|\operatorname{co}(X)\backslash X|}{|\operatorname{co}(X)|} \geq \alpha$. If we consider the last subset $Y \in \bigcup_{j<2i} \mathcal{F}_j$, so that $X \subset Y$ and $\frac{|\operatorname{co}(Y)\backslash Y|}{|\operatorname{co}(Y)|} \leq n!\alpha$, then X derives from Y by a sequence of applications of Lemma 10.8. Hence, as $\operatorname{diam}(X) > \xi \operatorname{diam}(Y)$, we know that

$$\frac{|\operatorname{co}(X)\setminus X|}{|\operatorname{co}(X)|} \le \frac{|\operatorname{co}(Y)\setminus Y|}{|\operatorname{co}(Y)|} \cdot \frac{|\operatorname{co}(Y)|}{|\operatorname{co}(X)|} < n!\alpha\xi^{-3n^2}.$$

Choosing α sufficiently small in terms of ξ and n, we can apply Lemma 10.8 to all non-finished elements of \mathcal{F}_{2i} and construct \mathcal{F}_{2i+1} with $D(\mathcal{F}_{2i+1}) \leq (1-\zeta)D(\mathcal{F}_{2i})$ and $P(\mathcal{F}_{2i+1}) \geq \beta P(\mathcal{F}_{2i})$.

To construct \mathcal{F}_{2i+2} from \mathcal{F}_{2i+1} , apply Lemma 10.7 to all non-finished elements with $|\operatorname{co}(A)\setminus A|<\alpha|A|$.

Note that $D(\mathcal{F}_{2i+2}) \leq D(\mathcal{F}_{2i+1})$ and $P(\mathcal{F}_{2i+2}) \geq (1-\alpha)P(\mathcal{F}_{2i+1})$. This construction implies that there exists a $i \leq \frac{\log(\xi)}{\log(1-\zeta)} + 1$, so that $D(\mathcal{F}_{2i+1}) \leq \xi$. For this i, we thus find $P(\mathcal{F}_{2i+1}) \geq \beta^{i+1}(1-\alpha)^i \geq \theta$, where the last inequality follows from choosing θ sufficiently small in terms of n, ξ, α, ζ , and β . Hence, this family satisfies all the conclusions and thus confirms the proposition.

Proof of Proposition 10.4 10.5.2

Proof of Proposition 10.4. Let ε sufficiently small in terms of n and λ to apply Lemma 10.9, Lemma 10.10, and Proposition 10.5. Let $\eta' := \eta_{n,\lambda,\varepsilon,t}^{10.5}$ so that we can apply Proposition 10.5. Choose α sufficiently small to apply Lemma 10.10. Let η sufficiently small so that we can apply Lemma 10.9, and Lemma 10.10. Let $A^{\circ} := U \cap A$. First use Lemma 10.9 to find $V = x + (1 + \beta)U$ with $||x||_2 \le \alpha$ and $|\beta| \le \alpha$. Let $B^{\circ} = B \cap V$, so that $|tA^{\circ} + (1-t)B^{\circ}| - |A^{\circ}| \le |tA + (1-t)B| - |A|.$

As in Lemma 10.10, let $t' = \frac{t'}{1+\beta-\beta t}$. Set $W = \frac{t}{t'}U$, $W' = \frac{1-t}{1-t'}V$, $A' = \frac{t}{t'}A^{\circ}$ and $B' = \frac{1-t}{1-t'}B^{\circ}$. By Lemma 10.10, the tube $W' = W + \frac{1-t}{1-t'}x$ has diameter less than 2ϵ , $A' \subset W$ and $B' \subset W'$, and we have

$$|t'A' + (1-t')B'| - t'|A'| - (1-t')|B'| \le |tA^{\circ} + (1-t)B^{\circ}| - |A^{\circ}| \le |tA + (1-t)B| - |A|.$$

Furthermore, $A', B' - \frac{1-t}{1-t'}x$ form a tubular 2λ -bounded η' -sandwich

By Proposition 10.5 (applied to A' and $B' - \frac{1-t}{1-t'}x$), we find

$$\begin{split} |\operatorname{co}(A') \setminus A'| &\leq k_n^{10.5}(t')^{-c^{10.5}n^8} (|t'A' + (1-t')B'| - t'|A'| - (1-t')|B'|) \\ &\leq k_n^{10.5}(t/2)^{-c^{10.5}n^8} (|tA + (1-t)B| - |A|). \end{split}$$

By definition of A', we have:

$$|\operatorname{co}(A\cap U)\setminus (A\cap U)|=|\operatorname{co}(A^\circ)\setminus A^\circ|=\left(\frac{t'}{t}\right)^n|\operatorname{co}(A')\setminus A'|\leq 2^n|\operatorname{co}(A')\setminus A'|.$$

Hence, we find some constant $k_n^{10.4} := k_n^{10.5} 2^n 2^{c^{10.5}} n^s$, so that

$$|\operatorname{co}(A \cap U) \setminus (A \cap U)| = k_n^{10.4} t^{-c^{10.5} n^8} (|tA + (1-t)B| - |A|).$$

Proof of Proposition 10.5

Proof of Proposition 10.5. Let $k_n^{10.5} = k_n^{1.6}(2n)$ and $c^{10.5} = c^{1.6}$. Let $\eta = \eta_{n,\lambda,\varepsilon,t}^{10.5}$ sufficiently small to apply Lemma 10.11.

As the problem is rotation invariant, we can assume wlog that $W = W' = S \times \mathbb{R}_+$ is a vertical tube.

We can assume wlog that co(A') has 2n vertices (co(A')) is the intersection of T with a halfspace). Indeed, partition S into simplices $S = S_1 \cup \cdots \cup S_m$ by projecting a triangulation of the upper boundary of co(A'). Consider the corresponding partition $T = T_1 \cup \cdots \cup T_m$, where $T_i = S_i \times \mathbb{R}_+$ and set $A_i = A' \cap T_i$ and $B_i = B' \cap T_i$. Note that $A' = \bigcup_i A_i$, $B' = \bigcup_i B_i$, and $\operatorname{co}(A') = \bigcup_i \operatorname{co}(A_i)$ are partitions (but $\operatorname{co}(B') \supset \bigcup_i \operatorname{co}(B_i)$ is essentially a disjoint union). Unlike many other partitions in this paper, we generally do not have $|A_i| = |B_i|$. Moreover, as $t'A_i + (1-t')B_i$ are essentially disjoint, we find

$$\sum_{i} |t'A_i + (1-t')B_i| - t'|A_i| - (1-t')|B_i| \le |t'A' + (1-t')B'| - t'|A'| - (1-t')|B'|.$$

Hence, it is enough to show that

$$|\operatorname{co}(A_i) \setminus A_i| \le |t'A_i + (1-t')B_i| - t'|A_i| - (1-t')|B_i|.$$

As A', B' form a tubular λ -bounded η -sandwich, we deduce that A_i , B_i form a 2λ -bounded η -sandwich. Moreover, by construction $co(A_i)$ has 2n vertices $(co(A_i)$ is the intersection of $T_i = S_i \times \mathbb{R}_+$ with a half-space). By taking an affine transformation, we can also assume $S_i \subset \mathbb{R}^{n-1}$ is a regular simplex centered at the origin. This concludes the reduction to the case when co(A') has 2n vertices.

Now assume wlog co(A') has 2n vertices and say $|A'| \leq |B'|$ (the other case is identical). Let $\alpha = (|B'| - |A'|)/|S|$ and note that $0 \leq \alpha = O_{n,\lambda}(\eta)$. Let $\beta > 0$ large. Construct the sets $A'' = A' \sqcup S \times [-\alpha - \beta, 0)$ and $B'' = B' \sqcup S \times [-\beta, 0)$ such that |A''| = |B''|.

By Lemma 10.11, we find that

$$t'A'' + (1 - t')B'' = S \times [-t'\alpha - \beta, 0) \sqcup (t'A' + (1 - t')B').$$

Recalling the definition of A'' and B'' and the fact that |A''| = |B''|, we deduce

$$|t'A'' + (1-t')B'| - |A''| = |t'A' + (1-t')B'| - t'|A'| - (1-t')|B'|$$

and $|\operatorname{co}(A'') \setminus A''| = |\operatorname{co}(A') \setminus A'|$. Therefore, it is enough to show that

$$|\operatorname{co}(A'') \setminus A''| \le k_n^{10.5} (t')^{-c^{10.5}} n^8 (|t'A'' + (1-t')B''| - |A''|).$$

Moreover, by making β arbitrarily large, we can make the ratio $\frac{|t'A''+(1-t')B''|-|A''|}{|A''|}$ arbitrarily close to 0 (as the numerator is constant but the denominator can be arbitrarily large). As co(A'') = co(A') has 2n vertices, this is a simple application of Theorem 1.6.

10.6 Proof of Lemmas

10.6.1 Proof of Lemma 10.6

Proof of Lemma 10.6. By taking an affine transformation, we can assume without loss of generality that $S = co(A) = co\{x_0, \ldots, x_n\}$ is a regular simplex with volume 1 centered at the origin o. Clearly, for all $0 \le i \le n$

$$\frac{|\cos\{x_0,\ldots,x_{i-1},o,x_{i+1},\ldots,x_n\}|}{|\cos\{x_0,\ldots,x_{i-1},x_i,x_{i+1},\ldots,x_n\}|} = \frac{1}{n+1}.$$

By continuity, there exists $\varepsilon > 0$ sufficiently small in terms of n such that for all $x \in \varepsilon S$ and $0 \le i \le n$

$$\frac{|\cos\{x_0,\ldots,x_{i-1},o,x_{i+1},\ldots,x_n\}|}{|\cos\{x_0,\ldots,x_{i-1},x_i,x_{i+1},\ldots,x_n\}|} \ge \frac{1}{n+2}.$$

As $|\varepsilon S| = \varepsilon^n |S|$, provided $\alpha \le \varepsilon^n / 10$ we deduce that there exists $x \in \varepsilon S \cap A$ and it follows that x has the desired properties.

For the last part, fix $0 \le k \le n$. Let q_k be the intersection of the ray $x_k x$ with the face opposite x_k . On the one hand, a simple computation gives $\frac{|xq_k|}{|x_kq_k|} = \frac{|\cos\{x_0,\dots,x_{i-1},x,x_{i+1},\dots,x_n\}|}{|\cos\{x_0,\dots,x_n\}|}$. From the first part, we know that $\frac{|\cos\{x_0,\dots,x_{i-1},x,x_{i+1},\dots,x_n\}|}{|\cos\{x_0,\dots,x_n\}|} \ge \frac{1}{n+2}$. Combining the last two inequalities, we get $\frac{|xq_k|}{|x_kq_k|} \ge \frac{1}{2(n+1)}$, i.e., $\frac{|xx_k|}{|x_kq_k|} \le \frac{n+1}{n+2}$. On the other hand, because the diameter of a simplex is realized between two vertices, we have $|x_kq_k| \le \max_{i,j}(x_ix_j)$. Combining the last two inequalities, we find $d(x,x_i) \le \frac{n+1}{n+2} \max_{i,j} d(x_i,x_j)$, which concludes the proposition.

10.6.2 Proof of Lemma 10.7

Proof of Lemma 10.7. First we argue that we can assume without loss of generality that P := co(A) is a simplex.

Consider the essential partition $P = S^1 \sqcup \cdots \sqcup S^r$ into simplices with a vertex at the origin o, obtained by partitioning the boundary ∂P into (n-1)-dimensional simplices and conning off at o. Then the sets $A^j = A \cap S^j$ are n-sandwiches.

Note that given A is a finite union of boxes, we have $V(S^j) \subset A_j$ i.e., $co(A^j) = S^j$. In particular, $|co(A) \setminus A| = \sum_j |co(A^j) \setminus A^j|$.

Now assume that for each j we can find an essential partition into convex sets $co(A^j) = P_1^j \sqcup \cdots \sqcup P_{k_j}^j$ and a corresponding partition of indices $[k_j] = I^j \sqcup J^j$ with the desired properties. Then it is easy to check that the essential partition into convex sets of $co(A) = \sqcup_{j \leq r, i \leq k_j} P_i^j$ and the corresponding partition of indices $I = \sqcup_j I^j$ and $J = \sqcup_j J^j$ have the desired properties.

Thus from now on we can assume co(A) to be a simplex. Moreover, from now on we only retain the weaker hypothesis that $\frac{|co(A)|}{|A|} \le (1+\eta)^n \le 1+\alpha$.

Given a simplex co(A), consider the following iterative process. First set $\mathcal{T}_0 = \{co(A)\}$ and $\mathcal{S}_0 = \emptyset$, and note that $|\operatorname{co}(A)| \leq (1+\alpha)|A|$ by hypothesis. At a given stage i with $\mathcal{T}_i, \mathcal{S}_i$, look at every element $S' \in \mathcal{T}_i$ and distinguish two cases: either $|S'| \leq (1+\alpha)|S' \cap A|$ or $|S'| \geq (1+\alpha)|S' \cap A|$.

For each simplex $S' = co\{x_0, \ldots, x_n\} \in \mathcal{T}_i$ with $|S'| \leq (1+\alpha)|S' \cap A|$ we construct the n+1 simplices $f_0(S'), \ldots, f_n(S')$ as follows. We apply Lemma 10.6 to find a central point $x \in S' \cap A$ and we construct the simplex $f_j(S') = co\{x_0, \dots, x_{j-1}, x, x_{j+1}, \dots, x_n\}.$

$$\mathcal{T}_{i+1} := \bigcup_{S' \in \mathcal{T}_i: \ |S'| \le (1+\alpha)|S' \cap A|} \{f_0(S'), \dots, f_n(S')\} \quad \text{and} \quad \mathcal{S}_{i+1} := \mathcal{S}_i \cup \{S' \in \mathcal{T}_i: |S'| \ge (1+\alpha)|S' \cap A|\}.$$

Using the fact that A is closed, it follows by induction that for $i \in \mathbb{N}$ and $S' \in \mathcal{T}_i \sqcup \mathcal{S}_i$ we have $\operatorname{co}(A \cap S') = S'$. Moreover, $\mathcal{T}_i \sqcup \mathcal{S}_i$ forms an essential partition of co(A).

Claim 10.12. For all $S' \in \mathcal{T}_i$ and $j \in [0, n]$, we have $|f_i(S')| \ge |S'|/n + 2$.

Proof of Claim. By Lemma 10.6 (and our choice of $x \in S' \cap A$), we have $|f_j(S')| \ge |f_j(S') \cap A| \ge |S'|/n + 2$. \square

Claim 10.13. For all $S' \in \mathcal{T}_i$, we have $|S'| \leq (1 + (n+3)\alpha)|S' \cap A|$ and for all $S' \in \mathcal{S}_i$ we have $(1+\alpha)|S' \cap A| \leq (1+\alpha)|S' \cap A|$ $|S'| \le (1 + (n+3)\alpha)|S' \cap A|$

Proof of Claim. Every simplex $S' \in \mathcal{S}_i \cup \mathcal{T}_i$ is $f_i(S'')$ for some S'' with $|S'| \leq (1+\alpha)|S' \cap A|$. By Lemma 10.6 and the fact that $\alpha > 0$ is small, we have $|S''| \leq (1 + (n+3)\alpha)|S'' \cap A|$. In addition, by construction of S_i , for all $S' \in \mathcal{S}_i$ we have $(1 + \alpha)|S' \cap A| \leq |S'|$.

Before we conclude we need two more claims. Given a simplex S', let radius rad(S') be the maximal length among its edges.

Claim 10.14. For all r > 0 the following holds

$$\sum_{S' \in \mathcal{T}_i: rad(T) > r} |S'| \to 0 \text{ as } i \to \infty.$$

Proof of Claim. We make the convention $\bigcup \mathcal{T}_i := \bigcup_{S' \in \mathcal{T}_i} S'$ Let $k = \lceil \log_{(1-\rho)}(r) \rceil$. We can distinguish two types of elements in \mathcal{T}_i . Either $S' \subset \bigcup \mathcal{T}_{i+k(n+1)}$ or not. Collect the former in \mathcal{T}_i' and the latter in \mathcal{T}_i'' . For $S' \in \mathcal{T}_i''$, at least some simplex originating from S' is in $\mathcal{S}_{i+k(n+1)}$. By the first claim, that simplex will

have size at least $\frac{1}{(n+2)^k(n+1)}|S'|$. Hence,

$$\sum_{S' \in \mathcal{T}_{i+k(n+1)}: \ S' \subset \bigcup \mathcal{T}''_i} |S'| \le \left(1 - \frac{1}{(n+2)^{k(n+1)}}\right) \left| \bigcup \mathcal{T}''_i \right|.$$

For $S' \in \mathcal{T}_i'$, we will find an element $S'' \in \mathcal{T}_{i+k(n+1)}$ with $S'' \subset S'$ and rad(S'') < r. Let $S' = S^0$, and consider $S^{j+1} := f_n(f_{n-1}(\dots f_0(S^j)\dots)$. Crucially, $rad(S^{j+1}) \leq ((n+1)/(n+2))rad(S^j)$. Indeed, none of the edges of S^j remain and all of the edges added have length at most $(1-\rho)rad(S^j)$ by Lemma 10.6. Hence, $rad(S^k) \leq r \cdot rad(S^0) \leq r$. Note that again by the first claim $|S^k| \geq \frac{1}{(n+2)^{k(n+1)}} |S'|$.

Combining these two cases gives

$$\sum_{S' \in \mathcal{T}_{i+k(n+1)}: \ rad(S') > r} |S'| \le \left(1 - \frac{1}{(n+2)^{k(n+1)}}\right) \sum_{S' \in \mathcal{T}_i: \ rad(S') > r} |S'|.$$

The conclusion follows.

Claim 10.15. Assuming A is a finite union of boxes, for every $\varepsilon > 0$ there exists r > 0 depending on A and ε such that for all $i \in \mathbb{N}$ the following holds

$$\sum_{\substack{S' \in \mathcal{T}_i: \\ rad(T) \leq r \text{ and } S' \not\subset A}} |S'| \leq \varepsilon.$$

Proof of Claim. Note that $\bigcup_{\substack{S' \in \mathcal{T}_i: \\ rad(T) \leq r \text{ and } S' \not\subset A}} S' \subset B(o,r) + \partial A$ and hence

$$\sum_{\substack{S' \in \mathcal{T}_i: \\ rad(T) \leq r \text{ and } S' \not\subset A}} |S'| \leq |B(o,r) + \partial A|.$$

Moreover, we have $\lim_{r\to 0} \frac{|B(o,r)+\partial A|}{r} = |\partial A|$. Therefore, for r>0 sufficiently small in terms of A and ε

$$\sum_{\substack{S' \in \mathcal{T}_i: \\ rad(T) \leq r \text{ and } S' \not\subset A}} |S'| \leq |B(o,r) + \partial A| \leq r |\partial A| \leq \varepsilon.$$

Returning to the proof of the Lemma, we can combine Claim 10.14 and Claim 10.15 to obtain that for some r sufficiently small and some i sufficiently large (both depending on A, α), we have

$$\sum_{\substack{S' \in \mathcal{T}_i: \\ S' \not\subset A}} |S'| \le \sum_{\substack{S' \in \mathcal{T}_i: \\ rad(T) \le r \text{ and } S' \not\subset A}} |S'| + \sum_{\substack{S' \in \mathcal{T}_i: \\ rad(T) \ge r}} |S'| \le \alpha |\operatorname{co}(A) \setminus A|.$$

We assumed here that A is not convex as otherwise we are immediately done. We further get that

$$\sum_{S' \in \mathcal{T}_i} |S' \setminus A| \le \alpha |\operatorname{co}(A) \setminus A|.$$

On the other hand, by Claim 10.13, for all $S' \in \mathcal{S}_i$, we have $\frac{|S'|}{|S' \cap A|} \in (1 + \alpha, 1 + n!\alpha)$.

By construction, $\mathcal{T}_i \cup \mathcal{S}_i$ gives an essential partition into convex sets $co(A) = P_1 \sqcup \cdots \sqcup P_k$ that satisfy properties 1, 2 and 3. Moreover, by the last two centered equations, the partition of the indices $[k] = I \sqcup J$ where $I = \{j \in [k]: P_i \in \mathcal{S}_i\}$ and $J = \{j \in [k]: P_i \in \mathcal{S}_i\}$ satisfies properties 4 and 5. The conclusion follows. \square

10.6.3 Proof of Lemma 10.8

Proof of Lemma 10.8. Choose parameters according to the following hierarchy:

$$n \gg \gamma \gg \alpha_0 \ge \alpha \gg \zeta \gg \eta \gg \beta$$
.

Here $\alpha_0 \ge \alpha$ reflects that α is assumed to be sufficiently small in terms of n. We proceed by contradiction, so that every subset $X \subset A$ which satisfies conclusions 1, 2, and 3 has $|\cos(X) \setminus X| \le \beta |X|$.

Let v_0, \ldots, v_n denote the vertices of $T = \operatorname{co}(A)$ and assume T is centred, i.e., that its barycenter is the origin. Consider the slightly shrunk simplex $(1-\gamma)T$ with vertices $(1-\gamma)v_0, \ldots, (1-\gamma)v_n$. Our first step will be to find points in $A \cap (1-\gamma)T$ close to each of these vertices. Indeed, consider the simplex $S_i := \gamma T + (1-2\gamma)v_i$ which is a translate of γT whose i-th vertex coincides with $(1-\gamma)v_i$. Note that since $|\operatorname{co}(A) \setminus A| = \alpha |A|$ and $|\gamma T| = \gamma^n |T| \ge \gamma^n |A|$, there must be a point $u_i \in S_i \cap A$ for all i.

We iterate a similar construction to show that in fact we can find points considerably closer to the vertices v_i . Indeed consider the simplex $T_i := \eta T + (1 - \zeta - \eta)v_i$ which is a translate of ηT whose *i*-th vertex lies on $(1 - \zeta)v_i$. Note that for η sufficiently small in terms of ζ , we find that $T_i \subset \text{co}(u_0, \ldots, u_{i-1}, v_i, u_{i+1}, \ldots, u_n)$ for any points $u_i \in S_i$. Hence, if we consider

$$A_i := A \cap \operatorname{co}(u_0, \dots, u_{i-1}, v_i, u_{i+1}, \dots, u_n),$$

we find that $co(A_i) = co(u_0, \ldots, u_{i-1}, v_i, u_{i+1}, \ldots, u_n)$ is a simplex, $diam(A_i) \leq (1 - \gamma/n)diam(A)$, and $|co(A_i)| \geq (1 - \gamma)^{3n} |co(A)| \geq (1 - \gamma/n)^{3n^2} |co(A)|$, so by our contradiction assumption, we have that $|co(A_i)| \leq \beta |A_i|$. Since

$$T_i \subset \operatorname{co}(u_0, \dots, u_{i-1}, v_i, u_{i+1}, \dots, u_n)$$
 and $|T_i| = \eta^n |T| > \beta |A_i|$

(assuming that β is sufficiently small in terms of η), there exists $p_i \in T_i \cap A_i \subset A$. Let $A' := A \cap \operatorname{co}(p_0, \dots, p_n)$ so that $A' \subset (1-\zeta)T$ and $\operatorname{diam}(A') \leq (1-\zeta)\operatorname{diam}(A)$. Clearly, $\operatorname{co}(A') = \operatorname{co}(p_0, \dots, p_n)$ is a simplex. In the limit $\zeta, \eta \to 0$, we have $|\operatorname{co}(A')| = (1-o(1))|\operatorname{co}(A)|$. Hence, for ζ, η , and β sufficiently small in terms of α and n we find:

$$|\operatorname{co}(A) \setminus A| < |\operatorname{co}(A) \setminus \operatorname{co}(A')| + |\operatorname{co}(A') \setminus A'| < \alpha |A|,$$

a contradiction. This proves the result.

10.6.4 Proof of Lemma 10.9

Proof of Lemma 10.9. As A, B form a λ -bounded η -sandwich, consider the convex $K \subset \mathbb{R}^n$, so that

$$B(o, (3\lambda n)^{-1}) \subset K \subset A, B \subset (1+\eta)K \subset B(o, 3\lambda n),$$

where the balls follow from Observation 2.6 combined with $1 + \eta < 3/2$.

We construct V as follows. Let H_1, \ldots, H_n be the defining hyperplanes of U, and let H_i^+ be the corresponding halfspaces so that $U = \bigcap_i H_i^+$. We construct parallel hyperplanes G_i as follows: Given G_1, \ldots, G_{i-1} , find G_i so that $|A \cap \bigcap_{j=1}^i H_j^+| = |B \cap \bigcap_{j=1}^i G_j^+|$. Note that this is possible since, by construction $|A \cap \bigcap_{j=1}^{i-1} H_j^+| = |B \cap \bigcap_{j=1}^{i-1} G_j^+|$ and $|B \cap \bigcap_{j=1}^i G_j^+|$ changes continuously with G_j .

We will show by induction that the distance between G_i and H_i tends to zero as η tends to zero. By induction $|(H_i^+ \triangle G_i^+) \cap B| \to 0$ as $\eta \to 0$ for j < i. This implies that also

$$\left|\left|\left(B\cap H_i^+\right)\cap\bigcap_{i=1}^{i-1}H_j^+\right|-\left|\left(B\cap H_i^+\right)\cap\bigcap_{i=1}^{i-1}G_j^+\right|\right|\leq \left|\left(B\cap H_i^+\right)\cap\bigcup_{i=1}^{i-1}(H_j^+\triangle G_j^+)\right|\to 0\qquad\text{as }\eta\to0.$$

In the same vein, we have that because A, B is a λ -bounded η sandwich

$$\left| |A \cap \bigcap_{j=1}^{i} H_j^+| - |B \cap \bigcap_{j=1}^{i} H_j^+| \right| \le \left| (A \triangle B) \cap \bigcap_{j=1}^{i} H_j^+ \right| \le \left| (1+\eta)K \setminus K \right| \to 0 \quad \text{as } \eta \to 0.$$

Hence, using that $\left| (B \cap G_i^+) \cap \bigcap_{j=1}^{i-1} G_j^+ \right| = |A \cap \bigcap_{j=1}^i H_j^+|$, we find

$$\left|\left|(B\cap G_i^+)\cap\bigcap_{j=1}^{i-1}G_j^+\right|-\left|(B\cap H_i^+)\cap\bigcap_{j=1}^{i-1}G_j^+\right|\right|=\left|\left|A\cap\bigcap_{j=1}^iH_j^+\right|-\left|(B\cap H_i^+)\cap\bigcap_{j=1}^{i-1}G_j^+\right|\right|\to 0\quad\text{ as }\eta\to 0.$$

On the other hand, we find

$$\left| \left| (B \cap G_i^+) \cap \bigcap_{j=1}^{i-1} G_j^+ \right| - \left| (B \cap H_i^+) \cap \bigcap_{j=1}^{i-1} G_j^+ \right| \right| = \left| B \cap (G_i^+ \triangle H_i^+) \cap \bigcap_{j=1}^{i-1} G_j^+ \right| \ge \left| (G_i^+ \triangle H_i^+) \cap B(o, (3\lambda n)^{-1}) \cap \bigcap_{j=1}^{i-1} G_j^+ \right|.$$

By induction, we have $\left| (G_i^+ \triangle H_i^+) \cap B(o, (3\lambda n)^{-1}) \cap \bigcap_{j=1}^{i-1} G_j^+ \right| = \left| (G_i^+ \triangle H_i^+) \cap B(o, (3\lambda n)^{-1}) \cap \bigcap_{j=1}^{i-1} H_j^+ \right| + o(1)$, where $o(1) \to 0$ as $\eta \to 0$. As ϵ is sufficiently small in terms of λ and n, we find that

$$\left| (G_i^+ \triangle H_i^+) \cap B(o, (3\lambda n)^{-1}) \cap \bigcap_{j=1}^{i-1} H_j^+ \right| = \Omega_n(\left| (G_i^+ \triangle H_i^+) \cap B(o, (3\lambda n)^{-1}) \right|).$$

We conclude that $|(G_i^+ \triangle H_i^+) \cap B(o, (3\lambda n)^{-1})| \to 0$ as $\eta \to 0$, so that indeed the distance between G_i and H_i tends to zero as $\eta \to 0$.

Let $V := \bigcap_{i=1}^n G_i^+$. Since the G_i 's are parallel to the H_i 's, we find that V is homothetic to U, i.e., there are $x \in \mathbb{R}^n$ and $\beta \in \mathbb{R}$ so that $V = x + (1+\beta)U$. As the defining hyperplanes tend to each other as $\eta \to 0$, we find that for sufficiently small η , $||x||_2 \le \alpha$ and $|\beta| \le \alpha$.

For the last part of the theorem, we proceed by induction to show that $A_i := A \cap \bigcap_{j=1}^i H_j^+$ and $B_i := B \cap \bigcap_{j=1}^i H_j^+$ satisfy

$$|tA_i + (1-t)B_i| - |A_i| < |tA + (1-t)B| - |A|.$$

For i=0 this is vacuously true. Note that $A_i=A_{i-1}\cap H_i^+$ and $B_i=B_{i-1}\cap G_i^+$. By construction $|A_i|=|B_i|$ and $|A_{i-1}|=|B_{i-1}|$, so that we also have $A_i':=A_{i-1}\setminus A_i=A_{i-1}\cap H_i^-$ and the analogously defined B_i' have the same volume. We find that

$$tA_i + (1-t)B_i \subset tH_i^+ + (1-t)G_i^+$$
 and $tA_i' + (1-t)B_i' \subset tH_i^- + (1-t)G_i^-$,

which are two halfspaces separated by the hyperplane $tH_i + (1-t)G_i$, so that $tA_i + (1-t)B_i$ and $tA'_i + (1-t)B'_i$ are disjoint subsets of $tA_{i-1} + (1-t)B_{i-1}$ and thus

$$|tA_i + (1-t)B_i| + |tA_i' + (1-t)B_i'| \le |tA_{i-1} + (1-t)B_{i-1}|.$$

By the Brunn-Minkowski inequality we know that $|tA'_i + (1-t)B'_i| \ge |A'_i|$, so we can conclude

$$|tA_i + (1-t)B_i| - |A_i| \le |tA_i + (1-t)B_i| - |A_i| + |tA_i' + (1-t)B_i'| - |A_i'| \le |tA_{i-1} + (1-t)B_{i-1}| - |A_{i-1}| < |tA + (1-t)B| - |A|,$$

where the last inequality follows from induction. Finally, $A_n = A^{\circ}$ and $B_n = B^{\circ}$, so the theorem follows.

10.6.5 Proof of Lemma 10.10

Proof of Lemma 10.10. The first two conclusions are straightforward to checks so we turn our attention to the third conclusion. Note that by construction $t'A' + (1-t')B' = tA^{\circ} + (1-t)B^{\circ}$, so it is enough to show that

$$t'|A'| + (1 - t')|B'| \ge |A^{\circ}|.$$

As $|A'| = (t/t')^n |A^{\circ}|$ and $|B'| = ((1-t)/(1-t'))^n |B^{\circ}|$ and $|A^{\circ}| = |B^{\circ}|$, this is equivalent to

$$t'(t/t')^n + (1-t')((1-t)/(1-t'))^n \ge 1,$$

which is true by the convexity of the function $\tau \mapsto \tau^n$. We turn our attention to the last conclusion.

Recall that $K \subset A, B \subset (1+\eta)K$ for some convex set K. Let $K' = \min\left\{\frac{t}{t'}, \frac{1-t}{1-t'}\right\}K$, so that $K' \cap W \subset A'$ and $K' \cap W' \subset B'$. Similarly, let $K'' = \max\left\{\frac{t}{t'}, \frac{1-t}{1-t'}\right\}(1+\eta)K$, so that $K'' \cap W \supset A'$ and $K'' \cap W' \supset B'$. Note that for α sufficiently small in terms of η , we have t and t' sufficiently similar that $K'' \subset (1+2\eta)K'$. To deal with the translation, note that K' contains a smaller homothetic copy centred at $\frac{1-t}{1-t'}x$. Indeed, let p

be the intersection of the ray \mathbb{R}^+x with $\partial K'$. Since K is λ -bounded, K' is 1.5λ -bounded, and so $||p||_2 = \Omega_n(1/\lambda)$. Consider the homothety H centred at p with ratio $\frac{||p||_2 - \frac{1-t}{1-t'}||x||_2}{||p||_2} = 1 + O_n(\alpha\lambda^{-1})$. Note that $H(o) = \frac{1-t}{1-t'}x$. Let K''' := H(K'). As $p' \in K'$, we find $K''' \subset K'$. Hence, $K''' - \frac{1-t}{1-t'}x \subset A'$, $B' - \frac{1-t}{1-t'}x$. Similarly, let q be the point in the intersection between \mathbb{R}^-x and $\partial(1+2\eta)K'$, so that $||q||_2 = \Omega_n(1/\lambda)$. Consider the homothety H' centred at q with ratio $\frac{||q||_2 + \frac{1-t}{1-t'}||x||_2}{||q||_2} = 1 + O_n(\alpha\lambda^{-1})$, so that $H'(o) = \frac{1-t}{1-t'}x$ and

 $H((1+2\eta)K')\supset (1+2\eta)K'$ is centred at $\frac{1-t}{1-t'}x$. Hence, we find $A',B'-\frac{1-t}{1-t'}x\subset \frac{||q||_2+\frac{1-t}{1-t'}||x||_2}{||q||_2}(1+2\eta)K'$. Combining these two, and taking η and α sufficiently small in terms of λ , n, and η' , we find that $A',B'-\frac{1-t}{1-t'}x$ form a tubular 2λ -bounded η' -sandwich.

For the final conclusion on the symmetric difference, first note that for translating does not change the symmetric difference too much:

$$|A^{\circ}\triangle(B^{\circ}+x)| \leq |\operatorname{co}(A^{\circ}) \setminus A^{\circ}| + |\operatorname{co}(B^{\circ}+x) \setminus (B^{\circ}+x)| + |\operatorname{co}(A^{\circ})\triangle \operatorname{co}(B^{\circ}+x)|$$

$$\leq 2\alpha |A^{\circ}| + |\operatorname{co}(A^{\circ})\triangle(\operatorname{co}(B^{\circ})+x)|$$

$$\leq 2\alpha |A^{\circ}| + O_{n}(||x||_{2}) + |\operatorname{co}(A^{\circ})\triangle \operatorname{co}(B^{\circ})|$$

$$\leq (2\alpha + O_{n,\epsilon}(||x||_{2}))|A^{\circ}| + |\operatorname{co}(A^{\circ})\triangle \operatorname{co}(B^{\circ})|$$

$$\leq O_{n,\epsilon}(\alpha |A^{\circ}|) + (|\operatorname{co}(A^{\circ}) \setminus A^{\circ}| + |\operatorname{co}(B^{\circ}) \setminus B^{\circ}| + |A^{\circ}\triangle B^{\circ}|)$$

$$\leq O_{n,\epsilon}(\alpha |A^{\circ}|) + |A^{\circ}\triangle B^{\circ}|.$$

Hence, we may assume x=0. Similarly we find that the homotheties do not affect the symmetric difference too much. For simplicity assume $\beta > 0$ (the case $\beta \leq 0$ follows analogously), so that $|A' \triangle B'| \leq |(1+\beta)B^{\circ} \triangle A^{\circ}|$. As above, it suffices to show that $|(1+\beta)\cos(B^{\circ})\triangle\cos(A^{\circ})|$ is small. Because the origin is in both $\cos(A^{\circ})$ and $co(B^{\circ})$ we find that $|co(A^{\circ}) \setminus (1+\beta) co(B^{\circ})| \leq |co(A^{\circ}) \setminus co(B^{\circ})|$. For the other term, note that

$$|(1+\beta)\cos(B^{\circ})\setminus\cos(A^{\circ})| \leq |(1+\beta)\cos(B^{\circ})\setminus\cos(B^{\circ})| + |\cos(B^{\circ})\setminus\cos(A^{\circ})| \leq O_n(\beta)|A^{\circ}| + |\cos(B^{\circ})\setminus\cos(A^{\circ})|.$$

Combining these two bounds, we find

$$|(1+\beta)\cos(B^{\circ})\triangle\cos(A^{\circ})| = |\cos(A^{\circ})\setminus(1+\beta)\cos(B^{\circ})| + |(1+\beta)\cos(B^{\circ})\setminus\cos(A^{\circ})| \le O_n(\alpha)|A^{\circ}| + |\cos(B^{\circ})\triangle\cos(A^{\circ})|.$$

We conclude as above that $|(1+\beta)B^{\circ}\triangle A^{\circ}| \leq O_n(\alpha)|A^{\circ}| + |\cos(B^{\circ})\triangle\cos(A^{\circ})|$, which finishes the proof.

10.6.6 Proof of Lemma 10.11

Proof of Lemma 10.11. Note that by monotonicity it suffices to show that for points $p \in (\partial(1+\eta)K) \cap U$ and $x \in S \times \{0\}$, (where S is the regular simplex from Definition 10.1), we have $(1-t)x + tp \in K \cup (S \times \mathbb{R}_{-})$.

Consider the angle between px and op. Since $|ox| \le \epsilon$ and $|op| \ge (2\lambda n)^{-1}$ by Observation 2.6, we find that for sufficiently small ϵ , the angle between px and op is less than any $\alpha_{n,\lambda}$.

Let q be the intersection between the line xp and ∂K that lies between x and p. Similarly, let $p':=\frac{1}{1+n}p\in$ $op \cap \partial K$. By the above, we have $\angle qpp' \leq \alpha_{n,\lambda}$.

Now we will show that $\angle op'q$ is bounded away from zero in terms of n and λ . Let p'_n and q_n denote the n-th coordinates of p' and q respectively. Since both points lie in $U \setminus B(0, (2\lambda n)^{-1})$, we find that $p'_n, q_n \ge 1$

 $(2\lambda n)^{-1} - \epsilon > (3\lambda n)^{-1}$, so that the line extending p'q does not intersect $B(o, (3\lambda n)^{-1})$ inside U. Since, that line intersects K exactly in the segment p'q, it also does not intersect $B(o, (3\lambda n)^{-1}) \subset K$ outside of K. Now note that for every line ℓ through any point $r \in B(o, 2\lambda n) \setminus B(o, (3\lambda n)^{-1})$ that does not intersect $B(o, (3\lambda n)^{-1})$, we find that $\angle \ell$, $or \ge \arcsin((3\lambda n)^{-2}) \ge (3\lambda n)^{-2}$. Hence, $\angle op'q \ge (3\lambda n)^{-2}$, and thus

$$\angle p'qp = \angle op'q - \angle qpp' \ge (3\lambda n)^{-2} - \alpha_{n,\lambda} \ge (4\lambda n)^{-2}.$$

This implies that $|pq| = O_{n,\lambda}(|p'p|) = O_{n,\lambda}(\eta|op'|)$. Note that as $p' \in \partial K$ and $K \subset B(o,2\lambda n)$, we have $|op'| \leq 2\lambda n$, so that $|pq| = O_{n,\lambda}(|p'p|) = O_{n,\lambda}(\eta)$. On the other hand, we have by the triangle inequality that $|xp| \geq |op| - |ox| \geq 2\lambda n - \epsilon \geq \lambda n$. Hence, $\frac{|pq|}{|qx|} = O_{n,\lambda}(\eta)$, so that choosing η sufficiently small in terms of n, λ , and t, we find that $(1-t)x + tp \in xp \subset K$. The lemma follows.

11 Proof of general Linear Theorem (Theorem 1.4)

Proof of Theorem 1.4. Let $\epsilon = \epsilon_n$ sufficiently small that we can apply Proposition 10.4 with $\lambda = 2n^3$. Choose ξ sufficiently small in terms of n and ϵ . Finally, choose $\theta = \theta_n$ and $\eta = \eta_{n,t}$ sufficiently small so that we can apply Proposition 10.4 and Proposition 10.3. Note that the choice for θ does not depend on t.

By Proposition 2.9, we may assume that A, B form a $2n^3$ -bounded η -sandwich, by picking $d_{n,t}$ sufficiently small. After re-normalization so that |A| = |B| = 1, this implies by Observation 2.6, that $B(o, (4n^4)^{-1}) \subset A, B \subset B(o, 4n^4)$.

we will show the bound for $|\operatorname{co}(A)\setminus A|$. The bound on $|\operatorname{co}(B)\setminus B|$ follows analogously.

Apply Proposition 10.3 with parameter ξ to A to find A_1, \ldots, A_k . Note that as $A \subset B(o, 4n^4)$, we have $diam(\operatorname{co}(A)) \leq 8n^4$, so that $diam(\operatorname{co}(A_i)) \leq 8n^4\xi$.

Consider a uniformly random direction $v \in S^{n-1}$ and consider the tube $U \subset \mathbb{R}^n$ of diameter ϵ in that direction. Note that a simplex of diameter ϵ contains a ball of diameter ϵ' , where ϵ' depends only on n as ϵ depends only on n. Hence, U contains the circular $B(o,\epsilon') + \mathbb{R}v$. Fix one point a_i in each of the $\operatorname{co}(A_i)$ and note that if $a_i \in B(o,\epsilon'-8n^4\xi) + \mathbb{R}v$ then $A_i \subset U$. Choosing $\xi \leq \epsilon'/(16n^4)$, we find that the probability $\mathbb{P}(A_i \subset U) \geq \mathbb{P}(a_i \in B(o,\epsilon'-8n^4\xi) + \mathbb{R}v)$ is lower bounded in terms of n for any $a_i \in A \subset B(o,4n^4)$. Say $\rho = \rho_n > 0$ is so that $\mathbb{P}(A_i \subset U) \geq \rho$.

Note that if $A_i \subset U$, then $co(A \cap U) \setminus (A \cap U) \supset co(A_i) \setminus A_i$. Hence, as the $co(A_i)$ are disjoint, we find

$$\mathbb{E}\Big[|\operatorname{co}(U\cap A)\setminus (U\cap A)|\Big] \geq \sum_i |\operatorname{co}(A_i)\setminus A_i| \cdot \mathbb{P}(A_i\subset U) \geq \rho \sum_i |\operatorname{co}(A_i)\setminus A_i| \geq \rho \theta |\operatorname{co}(A)\setminus A|.$$

Combining this with the bound we have by Proposition 10.4, we find

$$|\operatorname{co}(A) \setminus A| \le \frac{1}{\rho \theta} \mathbb{E} \Big[|\operatorname{co}(U \cap A) \setminus (U \cap A)| \Big] \le \frac{k_n^{10.4}}{\rho \theta} t^{-c^{10.4} n^8} \delta |A|.$$

Letting $k_n^{1.4} = \frac{k_n^{10.4}}{\rho\theta}$ gives the theorem.

12 Proof of Symmetric difference vs Common convex hull result (Theorem 1.7)

12.1 Propositions

The proof of these results separates into two parts, viz if X, Y very similar, i.e., $|X \triangle Y|$ small, or if X, Y very dissimilar, i.e., $|X \cap Y|$ small. The former is more difficult.

Proposition 12.1. There exists an absolute constants $c_n, d_n > 0$ such that the following holds. Assume $\delta \in [0, d_n]$ and let $A, B \subset \mathbb{R}^n$ be convex sets such that $|A \cap B| \ge (1 - \delta) \max(|A|, |B|)$. Then

$$|\operatorname{co}(A \cup B)| \le (1 + c_n \delta) \min(|A|, |B|).$$

Proposition 12.2. There exist constants c_n so that, given convex sets $X, Y \subset \mathbb{R}^n$, we have

$$|\operatorname{co}(X \cup Y)| \le c_n \frac{|X| \cdot |Y|}{|X \cap Y|}.$$

12.2 Proofs of Propositions

12.2.1 Proof of Proposition 12.1

For this proof we use a particular case of the main result from [vHST22], which asserts the following.

Theorem 12.3. For all $n \in \mathbb{N}$, there are computable constants $c_n^{12.3}, d_n^{12.3} > 0$, such that the following holds. Let $X \subset \mathbb{R}^n$ so that $\left| \frac{X+X}{2} \right| \leq (1+d_n^{12.3})|X|$, then

$$|\operatorname{co}(X) \setminus X| \le c_n^{12.3} \left| \frac{X+X}{2} \setminus X \right|.$$

With this theorem in hand, the proposition follows quickly.

Proof of Proposition 12.1. Let $c_n^{12.3}$ and $d_n^{12.3}(t)$ be the output of Theorem 12.3. Choose $c_n = 10c_n^{12.3}$. Choose $d_n \leq 10^{-1}d_n^{12.3}(\frac{1}{2})$

By standard approximation arguments we can assume that co(A) and co(B) have a finite number of vertices. Construct the finite set $F = V(co(A)) \cup V(co(B))$ and construct the measurable sets $X = (A \cap B) \cup F$. By construction $co(A \cup B) = co(X)$ and $|X| \le min(|A|, |B|)$. Therefore, it is enough to prove

$$|\operatorname{co}(X) \setminus X| \le c_n \delta |X| = 10c_n^{12.3} \delta |X|.$$

Given that $10\delta \leq d_n^{12.3}(\frac{1}{2})$, by Theorem 12.3 applied to the set X, it suffices to prove that $\left|\frac{1}{2}(X+X)\right| \leq (1+10\delta)|X|$. Recall that by construction

$$\begin{split} \frac{1}{2}(X+X) &= \frac{1}{2}\bigg([(A\cap B)\cup V(\operatorname{co}(A))\cup V(\operatorname{co}(B))] + [(A\cap B)\cup V(\operatorname{co}(A))\cup V(\operatorname{co}(B))]\bigg) \\ &\subset \frac{1}{2}\bigg([(A\cap B)\cup V(\operatorname{co}(A))] + [A\cap B]\bigg)\bigcup\frac{1}{2}\bigg([(A\cap B)\cup V(\operatorname{co}(B))] + [A\cap B]\bigg) \\ &\qquad \qquad \bigcup\frac{1}{2}\bigg([V(\operatorname{co}(A))\cup V(\operatorname{co}(B))] + [V(\operatorname{co}(A))\cup V(\operatorname{co}(B))]\bigg) \\ &\subset \frac{1}{2}\bigg(A+A\bigg)\bigcup\frac{1}{2}\bigg(B+B\bigg)\bigcup\frac{1}{2}\bigg([V(\operatorname{co}(A))\cup V(\operatorname{co}(B))] + [V(\operatorname{co}(A))\cup V(\operatorname{co}(B))]\bigg) \\ &\subset A\cup B\cup\frac{1}{2}\big(V(\operatorname{co}(A))+V(\operatorname{co}(B))\big). \end{split}$$

Therefore, by hypothesis and as $\left|\frac{1}{2}(V(co(A)) + V(co(B)))\right| = 0$, we get

$$\left| \frac{1}{2}(X+X) \right| \le |A \cup B| \le 2\max(|A|,|B|) - |A \cap B| \le (1+\delta)\max(|A|,|B|).$$

Also by hypothesis, we get $|X| = |A \cap B| \ge (1 - \delta) \max(|A|, |B|)$. Combining the last two inequalities, we conclude

$$\left| \frac{1}{2}(X+X) \right| \le (1+\delta) \max(|A|,|B|) \le (1+\delta)(1-\delta)^{-1}|X| \le (1+10\delta)|X|.$$

The last inequality follows by noting $\delta \leq \frac{1}{2}$ and $(1+x)(1-x)^{-1} \leq 1+10x$ for $x \leq 1/2$.

12.2.2 Proof of Proposition 12.2

We first prove the result for axis aligned boxes and then reduce the general case to that case.

Lemma 12.4. For axis aligned $R, T \subset \mathbb{R}^n$, we have

$$|\operatorname{co}(R \cup T)| \le 2^n \frac{|R| \cdot |T|}{|R \cap T|}$$

Proof. Linearly transforming and translating if needed we may assume that R is a translated unit cube and $T = [0, t_1] \times \cdots \times [0, t_n]$. Given this setup we find that $|R \cap T| \leq \prod_{i=1}^n \min\{t_i, 1\}$. On the other hand,

$$|\operatorname{co}(R \cup T)| \le |R + T| = \prod_{i=1}^{n} (t_i + 1) \le 2^n \prod_{i=1}^{n} \max\{t_i, 1\}.$$

Combining these bounds with |R| = 1 and $|T| = \prod_{i=1}^{n} t_i$, we get

$$|\operatorname{co}(R \cup T)| \le 2^n \prod_{i=1}^n \max\{t_i, 1\} = 2^n \prod_{i=1}^n \frac{t_i}{\min\{t_i, 1\}} \le 2^n \frac{|R| \cdot |T|}{|R \cap T|},$$

which concludes the lemma.

With this lemma in hand, the proof of Proposition 12.2 is just a quick reduction.

Proof of Proposition 12.2. Let E and F the John ellipsoids of X and Y, respectively, so that

$$X \subset E$$
, $Y \subset F$, $|E| \le O_n(|X|)$, and $|F| \le O_n(|Y|)$.

Note that $|E \cap F| \ge |X \cap Y|$. Affinely transforming if needed we may assume E is a ball. Rotating if necessary, we may assume the axes of symmetry of F are the basis vectors e_1, \ldots, e_n . Let R and T be the smallest axis aligned boxes containing E and F respectively, so that $|R| = O_n(|E|) = O_n(|X|)$ and $|T| = O_n(|F|) = O_n(|Y|)$. Now use $co(X \cup Y) \subset co(R \cup T)$ and $X \cap Y \subset R \cap T$, together with Lemma 12.4, to find

$$|\operatorname{co}(X \cup Y)| \le |\operatorname{co}(R \cup T)| \le 2^n \frac{|R| \cdot |T|}{|R \cap T|} \le O_n \left(\frac{|X| \cdot |Y|}{|X \cap Y|}\right).$$

12.3 Proof of Theorem Theorem 1.7

Proof of Theorem 1.7. Let $d_n^{12.1}, c_n^{12.1}, c_n^{12.2}$, be the constants from Proposition 12.1 and Proposition 12.2. Let $c_n := \max\{c_n^{12.1}, c_n^{12.2}/d_n^{12.1}\}$. Let $\delta = 1 - \frac{|X \cap Y|}{\max(|X|, |Y|)}$. Consider two cases; either $\delta \leq d_n^{12.1}$ or $\delta > d_n^{12.1}$. In the former case, by Proposition 12.1 we have

$$\frac{|\operatorname{co}(X \cup Y)|}{\min(|X|,|Y|)} - 1 \le c_n^{12 \cdot 1} \delta = c_n \left(1 - \frac{|X \cap Y|}{\max(|X|,|Y|)} \right) \le c_n \frac{|X \triangle Y|}{|X \cap Y|}.$$

In the latter case, we use

$$d_n^{12.1} < \delta \le 1 - \frac{|X \cap Y|}{\max(|X|, |Y|)} \le \frac{|X \triangle Y|}{\max\{|X|, |Y|\}},$$

to find that $\max\{|X|, |Y|\} < |X \triangle Y|/d_n^{12.1}$. Hence, by Proposition 12.2 we find

$$\frac{|\operatorname{co}(X \cup Y)|}{\min\{|X|,|Y|\}} \le c_n^{12.2} \frac{\max\{|X|,|Y|\}}{|X \cap Y|} \le c_n \frac{|X \triangle Y|}{|X \cap Y|},$$

which concludes the proof.

13 Putting it all together: Proof of Theorem 1.3

Proof of Theorem 1.3. Choose $d_{n,t}^{1.3}$ sufficiently small in terms of n and t to make various statements throughout the proof true, e.g. to allow the applications of Theorem 1.5, Theorem 1.4, and Proposition 12.1.

By Theorem 1.5, we find that (after a translation)

$$|A \triangle B| \leq c_n^{1.5} t^{-1/2} \delta^{\frac{1}{2}} |A|, \quad \text{ so that } \quad |A \cap B| \geq (1 - c_n^{1.5} t^{-1/2} \delta^{\frac{1}{2}}) |A|.$$

By Theorem 1.4, we find that

$$|\operatorname{co}(A) \setminus A| + |\operatorname{co}(B) \setminus B| \le t^{-c^{1.4} n^8} \delta |A|.$$

Combining these two bounds and using that $d_{n,t}^{1.3}$ is small, we find $|\cos(A)\triangle\cos(B)| \le 2c_n^{1.5}t^{-1/2}\delta^{\frac{1}{2}}|A|$. Hence, using Proposition 12.1 and the fact that $|A| \le \min\{|\cos(A)|, |\cos(B)|\}$, we have

$$\begin{aligned} |\operatorname{co}(A \cup B)| &= |\operatorname{co}(\operatorname{co}(A) \cup \operatorname{co}(B))| \le \left(1 + c_n^{12.1} 2 c_n^{1.5} t^{-1/2} \delta^{\frac{1}{2}}\right) \max\{|\operatorname{co}(A)|, |\operatorname{co}(B)|\} \\ &\le \left(1 + c_n^{12.1} 2 c_n^{1.5} t^{-1/2} \delta^{\frac{1}{2}}\right) \left(1 + t^{-1.4n^8} \delta\right) |A| \le \left(1 + c_n^{1.3} t^{-1/2} \delta^{1/2}\right) |A|. \end{aligned}$$

This concludes the proof of the theorem.

14 Open Problems

It is natural to ask for the t-dependence in Theorem 1.4. To this end we make the following conjecture.

Conjecture 14.1. For all $n \in \mathbb{N}$ and $t \in (0, 1/2]$, there are computable constants $c_n^{14.1}, d_{n,t}^{14.1} > 0$ such that the following holds. Assume $\delta \in [0, d_{n,t}^{14.1}]$, and assume $A, B \subset \mathbb{R}^n$ are measurable sets of equal volume, so that

$$|tA + (1-t)B| = (1+\delta)|A|.$$

Then

$$|\operatorname{co}(A) \setminus A| \le c_n^{14.1} t^{-1} \delta |A|$$
 and $|\operatorname{co}(B) \setminus B| \le c_n^{14.1} t^{-n+1} \delta |A|$

If true, in Conjecture 14.1, the exponents of δ and t, prioritised in this order, are optimal. Indeed, to bound $|\operatorname{co}(A)\setminus A|$, take $B=[-1,1]^n$ and $A=[-1,1]^n\cup\{p\}$, where $p=(1+h,0,\ldots,0)$. To bound $|\operatorname{co}(B)\setminus B|$, take $A=[-1,1]^n$ and $B=[-1,1]^n\cup\{p\}$, where $p=(1+h,0,\ldots,0)$. In both cases $h\ll_{n,t}1$.

It is worth noting that Conjecture 14.1 was established in dimension two [vHSTarb].

For doubling further from minimal, we recall the following conjecture from [vHK23a].

Conjecture 14.2. There is an absolute constant $\Delta > 0$ so that if $\delta < \Delta$ and $A \subset \mathbb{R}^n$ with $\left| \frac{A+A}{2} \right| \leq (1+\delta)|A|$ then there is some convex $K \subset \mathbb{R}^n$ with $|K \triangle A| \leq O_{\delta}(1)|A|$.

Finally, we recall the following conjecture suggested by [BFRar].

Conjecture 14.3. Let $t \in (0,1)$ $f,g,h: \mathbb{R}^n \to \mathbb{R}_{\geq 0}$ be measurable functions with the property that $h(tx+(1-t)y) \geq f(x)^t g(y)^{1-t}$ for all $x,y \in \mathbb{R}^n$ and $\int f = \int g = 1$. If $\int h \leq 1 + \delta$ with δ sufficiently small in terms of t and n, then there exists a log-concave function $\ell: \mathbb{R}^n \to \mathbb{R}_{\geq 0}$ so that $\int |h-\ell| + |f-\ell| + |g-\ell| \leq O_{t,n}(\delta^{1/2})$.

References

- [BFRar] Károly J Böröczky, Alessio Figalli, and João PG Ramos. A quantitative stability result for the Prékopa-Leindler inequality for arbitrary measurable functions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, to appear.
- [BJ17] Marco Barchiesi and Vesa Julin. Robustness of the Gaussian concentration inequality and the Brunn-Minkowski inequality. Calc. Var. Partial Differential Equations, 56, 05 2017.
- [Chr12a] Michael Christ. Near equality in the Brunn–Minkowski inequality. arXiv preprint arXiv:1207.5062, 2012.
- [Chr12b] Michael Christ. Near equality in the two-dimensional Brunn–Minkowski inequality. $arXiv\ preprint\ arXiv:1206.1965,\ 2012.$
- [CM17] Eric Carlen and Francesco Maggi. Stability for the Brunn-Minkowski and Riesz rearrangement inequalities, with applications to Gaussian concentration and finite range non-local isoperimetry. *Canad. J. Math.*, 69(5):1036–1063, 2017.
- [EK14] Ronen Eldan and Bo'az Klartag. Dimensionality and the stability of the Brunn-Minkowski inequality. Annali della Scuola Normale Superiore di Pisa. Classe di scienze, 13(4):975-1007, 2014.
- [Fig14] Alessio Figalli. Quantitative stability results for the Brunn-Minkowski inequality. In *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. III*, pages 237–256. Kyung Moon Sa, Seoul, 2014.

- [Fig15] Alessio Figalli. Stability results for the Brunn–Minkowski inequality. In *Colloquium De Giorgi 2013* and 2014, pages 119–127. Springer, 2015.
- [FJ15] Alessio Figalli and David Jerison. Quantitative stability for sumsets in \mathbb{R}^n . J. Eur. Math. Soc. (JEMS), 17(5):1079–1106, 2015.
- [FJ17] Alessio Figalli and David Jerison. Quantitative stability for the Brunn–Minkowski inequality. Adv. Math., 314:1–47, 2017.
- [FJ21] Alessio Figalli and David Jerison. A sharp Freiman type estimate for semisums in two and three dimensional Euclidean spaces. Ann. Sci. Éc. Norm. Supér., 54(4):235–257, 2021.
- [FMM18] Alessio Figalli, Francesco Maggi, and Connor Mooney. The sharp quantitative Euclidean concentration inequality. *Camb. J. Math.*, 6:59–87, 3 2018.
- [FMP09] Alessio Figalli, Francesco Maggi, and Aldo Pratelli. A refined Brunn-Minkowski inequality for convex sets. Ann. Inst. H. Poincaré Anal. Non Linéaire,, 26:2511–2519, 11 2009.
- [FMP10a] Alessio Figalli, Francesco Maggi, and Aldo Pratelli. A mass transportation approach to quantitative isoperimetric inequalities. *Invent. Math.*, 182(1):167–211, 2010.
- [FMP10b] Alessio Figalli, Francesco Maggi, and Aldo Pratelli. A mass transportation approach to quantitative isoperimetric inequalities. *Invent. Math*, pages 167–211, 2010.
- [Fre59] Gregory A Freiman. The addition of finite sets. I. *Izv. Vysš. Učebn. Zaved. Matematika*, (6):202–213, 1959.
- [FvHT] Alessio Figalli, Peter van Hintum, and Marius Tiba. Sharp stability of the Brunn-Minkowski inequality via optimal mass transportation. *preprint*.
- [vHK23a] Peter van Hintum and Peter Keevash. Locality in sumsets. arXiv preprint arXiv:2304.01189, 2023.
- [vHK23b] Peter van Hintum and Peter Keevash. The sharp doubling threshold for approximate convexity. arXiv preprint arXiv:2304.01176, 2023.
- [vHST22] Peter van Hintum, Hunter Spink, and Marius Tiba. Sharp stability of Brunn–Minkowski for homothetic regions. J. Eur. Math. Soc. (JEMS), 24(12):4207–4223, 2022.
- [vHSTara] Peter van Hintum, Hunter Spink, and Marius Tiba. Sharp L^1 Inequalities for Sup-Convolution. Discrete Anal., to appear.
- [vHSTarb] Peter van Hintum, Hunter Spink, and Marius Tiba. Sharp quantitative stability of the planar Brunn–Minkowski inequality. J. Eur. Math. Soc. (JEMS), to appear.
- [KKM29] Bronisław Knaster, Kazimierz Kuratowski, and Stefan Mazurkiewicz. Ein beweis des fixpunktsatzes für n-dimensionale simplexe (in German). Fund. Math., 14(1):132–137, 1929.
- [LS95] Vsevolod F Lev and Pavel Y Smeliansky. On addition of two distinct sets of integers. *Acta Arith.*, 70(1):85–91, 1995.
- [Sta96] Yonutz Stanchescu. On addition of two distinct sets of integers. Acta Arith., 75(2):191–194, 1996.