

BMJ Best Practice

Abdominal aortic aneurysm

The right clinical information, right where it's needed



Table of Contents

Summary	3
Basics	4
Definition	4
Epidemiology	4
Aetiology	4
Pathophysiology	4
Classification	5
Prevention	6
Screening	6
Diagnosis	7
Case history	7
Step-by-step diagnostic approach	7
Risk factors	8
History & examination factors	10
Diagnostic tests	11
Differential diagnosis	13
Treatment	16
Step-by-step treatment approach	16
Treatment details overview	24
Treatment options	26
Emerging	33
Follow up	34
Recommendations	34
Complications	34
Prognosis	36
Guidelines	37
Diagnostic guidelines	37
Treatment guidelines	37
Evidence scores	40
References	41
Images	57
Disclaimer	64

Summary

- ◇ Patients are usually asymptomatic and their abdominal aortic aneurysm is detected incidentally. In the minority of patients who experience symptoms, abdominal, back, and groin pain are typical.
- ◇ Ultrasound remains the definitive test for initial diagnosis and screening. Imaging with computed tomography angiography or magnetic resonance angiography is used for anatomical mapping to assist with operative planning.
- ◇ For AAA detected as an incidental finding, surveillance is preferred to repair until the theoretical risk of rupture exceeds the estimated risk of operative mortality. Repair is indicated in patients with large asymptomatic AAA.
- ◇ Mortality during elective surgical repair is higher for women than men for both open repair (7.0% vs. 5.2%) and endovascular aneurysm repair (EVAR) (2.1% vs. 1.3%).
- ◇ Complications of treatment include acute kidney injury, limb ischaemia, spinal cord ischaemia, anastomotic pseudoaneurysm, graft infection, graft limb occlusion, and distal embolisation. Endoleak is a complication of EVAR.

Definition

Abdominal aortic aneurysm (AAA) is a permanent pathological dilation of the aorta with a diameter >1.5 times the expected anteroposterior (AP) diameter of that segment, given the patient's sex and body size.[1]
[2] The most commonly adopted threshold is a diameter of 3 cm or more.[3] More than 90% of aneurysms originate below the renal arteries.[4]
[Fig-1]

Epidemiology

Epidemiology varies by region, age, and sex, with large screening studies offering the closest approximation. A screening study of US veterans aged 50 to 79 years found the highest prevalence of AAA (AAA >3 cm) among white male smokers (5.9%).[10] Prevalence among men is 4 to 6 times higher than in women.[1]
[11] Current prevalence in the UK National Health Service AAA screening programme is 1.34% (AAA >2.9 cm) among men invited for screening in their 65th year.[12]

A systematic review of studies reporting screen-detected AAA in women found that the pooled prevalence of AAA increased with age ($>1\%$ for women >70 years of age) and with smoking status ($>1\%$ for female ever-smokers; $>2\%$ for current smokers).[13] The prevalence of aneurysms among men increases by about 6% per decade of life.[9] In 2014 there were 9863 deaths attributed to AAA in the US.[14]

In the UK, deaths from AAA declined sharply from 1997 to 2009, with mortality in men aged >65 years falling from 65.9 to 44.6 per 100,000 population.[15] Admissions for ruptured AAA fell from 18.6 to 13.5 per 100,000 in all age groups. The reduction in mortality and admissions has been attributed to the fall in smoking prevalence and the rise in elective AAA repair in older age groups.[15]

Aetiology

Traditionally, arterial aneurysms were thought to arise from atherosclerotic disease, and certainly intimal atherosclerosis reliably accompanies AAA.[16] Subsequent data suggest that altered tissue metalloproteinases may diminish the integrity of the arterial wall.[6]

Smoking remains the most important risk factor.[1] [9] [11] [17] [18]

A systematic review and meta-analysis found that diabetes protects against the growth and enlargement of AAA.[19] However, the protective mechanism is yet to be determined. Operative and long-term survival was lower among AAA repair patients with diabetes than those without, suggesting an increased cardiovascular burden.[19]

Pathophysiology

Histologically there is obliteration of collagen and elastin in the media and adventitia, smooth muscle cell loss with resulting tapering of the medial wall, infiltration of lymphocytes and macrophages, and neovascularisation.[16] There are 4 mechanisms relevant to AAA development:[20]

- Proteolytic degradation of aortic wall connective tissue: matrix metalloproteinases (MMPs) and other proteases are derived from macrophages and aortic smooth muscle cells and secreted into

the extracellular matrix. Disproportionate proteolytic enzyme activity in the aortic wall may promote deterioration of structural matrix proteins (e.g., elastin and collagen).[4] Increased expression of collagenases MMP-1 and -13 and elastases MMP-2, -9, and -12 has been demonstrated in human AAAs.[21] [22] [23] [24]

- Inflammation and immune responses: an extensive transmural infiltration by macrophages and lymphocytes is present on aneurysm histology and these cells may release a cascade of cytokines that subsequently activate many proteases.[16] Additionally, deposition of immunoglobulin G into the aortic wall supports the hypothesis that AAA formation may be an autoimmune response. There is currently interest in the role of reactive oxygen species and antioxidants in AAA formation.[21] [25] [26] [27] [28]
- Biomechanical wall stress: elastin levels and the elastin-collagen ratio decrease progressively distal down the aorta. Diminished elastin is associated with aortic dilation, and collagen degradation predisposes to rupture.[11] Additionally, data support increased MMP-9 expression and activity, disordered flow and an increase in wall tension, and relative tissue hypoxia in the distal aorta (i.e., infrarenal).[21] [29] [30]
- Molecular genetics: AAA exhibits significant heritability (the proportion of phenotype due to genotype). A Swedish Twin Registry study found that the twin of a monozygotic twin with AAA had a risk of AAA approximately 70 times greater than that of the twin of a monozygotic twin without AAA.[31] While there is no single genetic defect or polymorphism responsible, familial clustering suggests a polygenic aetiology. Genome-wide approaches have identified 10 risk loci for AAA, 4 of which are not shared with other cardiovascular diseases.[32]

Classification

Types of AAA

Specific types of AAA are:[5] [6]

- Congenital: while medial degeneration occurs naturally with age, it is accelerated in patients with bicuspid aortic valves and Marfan syndrome.
- Infectious: infection of the aortic wall (mycotic aneurysm) is a rare aetiology. *Staphylococcus* and *Salmonella* are the most common pathogens. *Chlamydia pneumoniae* has been postulated as an infectious aetiology for conventional aneurysms. Tertiary syphilis may manifest as aortic aneurysm, but this is an exceptionally rare presentation.[7]
- Inflammatory: the aetiology of inflammatory AAAs remains controversial. This variant is characterised by an abnormal accumulation of macrophages and cytokines in diseased tissue. Pathologically there is perianeurysmal fibrosis, thickened walls, and dense adhesions.

Screening

Screening 65-year-old men for AAA significantly reduces AAA-specific mortality.^[71]

Current US Preventive Services Task Force (USPSTF) recommendations include:^[3]

- One-time screening for AAA with ultrasonography in men aged 65 to 75 years who have ever smoked
- Selective screening (based on an assessment of risk factors, including older age and a first-degree relative with an AAA) for AAA in men aged 65 to 75 years who have never smoked.

The USPSTF concludes that there is currently insufficient evidence to assess the balance of benefits and harms of screening for AAA in women aged 65 to 75 years who have ever smoked.^[3] In women who have never smoked, the USPSTF recommends against routine screening for AAA.

Routine screening for AAA among all men aged 65 and over has been available in the UK since 2013, and in England since 2009.^[72] Aortic diameter at the screening appointment determines the subsequent management pathway:^[73]

Aortic diameter (anterior-posterior, inner to inner)	Re-scanning interval
<3 cm	Discharged from programme
3-4.4 cm	Annual surveillance programme
4.5-5.4 cm	3-monthly surveillance programme
>5.5 cm	Referral to vascular surgeon*

*Re-scanning intervals associated with aortic diameters
NHS population screening: care pathways, Public Health England*

*European guidelines recommend review by a vascular surgeon within 2 weeks of an AAA >5.5 cm being detected, to prevent interval rupture.^[74]

Meta-analysis of AAA surveillance studies suggest that longer surveillance intervals than those currently employed in the UK AAA screening programme could be safely implemented, thereby reducing cost.^[75]

Case history

Case history #1

A 65-year-old man presents to his local aneurysm surveillance team for a screening ultrasound scan. He has been feeling well and in his usual state of good health. His medical history is notable for mild hypertension and he has a 100-pack-year tobacco history. On ultrasound an infrarenal AAA is identified.

Case history #2

A 55-year-old man with a history of hypertension (well controlled with medication) and tobacco use presents to his primary care physician with a 2-day history of constant and gnawing hypogastric pain. The pain has been steadily worsening in intensity. He says the pain radiates to his lower back and both groins at times. While he cannot identify any aggravating factors (such as movement), he feels the pain improves with his knees flexed. There is a palpable pulsatile mass just left of midline below the umbilicus. He is immediately referred for definitive management, but during transfer becomes hypotensive and unresponsive.

Other presentations

The triad of abdominal pain, weight loss, and elevated erythrocyte sedimentation rate suggests inflammatory AAA.[6] A tender, palpable pulsatile mass on examination and elevated C-reactive protein may also be present. Abdominal or back pain with fever is suggestive of mycotic or infectious AAA. Typically there is history of arterial trauma, intravenous drug abuse, local or concurrent infection, bacterial endocarditis, or impaired immunity. Osteomyelitis of the thoracic or lumbar spine may develop. Anaemia, leukocytosis, and positive blood cultures are common.[8] Diagnosis may be aided by complications of unruptured aneurysms, including distal embolisation, acute thrombosis, or symptoms caused by ureterohydronephrosis.[9]

Step-by-step diagnostic approach

Patients most commonly lack any symptoms and their aneurysm is noted on physical examination or imaging studies performed for other reasons.

History

In the minority of patients who experience symptoms, abdominal, back, and groin pain are typical. Medical history is directed towards risk factors:

- Development (i.e., hyperlipidaemia, connective tissue disorder, COPD, and hypertension)[1] [4] [9] [11] [17] [49] [50] [52]
- Expansion (i.e., previous cardiac or renal transplant, previous stroke, advanced age [>70 years], and severe cardiac disease)[53] [54]
- Rupture (i.e., female sex, previous cardiac or renal transplant, hypertension).[9] [45] [46] [51] [53] [55]

A history of cigarette smoking increases a patient's risk of AAA development, expansion, and rupture.[9] [17] [18] [33] [53] In men who have never smoked, important risk factors for AAA include older age and a first-degree relative with an AAA.[3]

A history of previous abdominal surgery or previous endovascular aortic aneurysm repair can be elicited as well as family history of AAA.

Physical examination

The abdomen can be palpated for a pulsatile abdominal mass and abdominal tenderness. Physical examination should include an assessment for peripheral artery aneurysm (femoral and popliteal).[56]

Aneurysm palpation on clinical examination has only been shown to be sensitive in thin patients and those with AAA >5 cm, with an overall sensitivity and specificity of 68% and 75%, respectively.[1] [57]

Ruptured aneurysm presents with the triad of abdominal and/or back pain, pulsatile abdominal mass, and hypotension.

The presence of fever may increase suspicion for infectious AAA in the appropriate clinical setting.

Key tests

Ultrasonography is the initial method of choice for AAA detection (sensitivity and specificity of 92% to 99% and nearly 100%, respectively).[1] [2] [57] [58] Once the diagnosis is made, further imaging with computed tomography angiography (CTA) or magnetic resonance angiography (MRA) is used for anatomical mapping to assist with operative planning (open or endovascular).[59]

Elevated erythrocyte sedimentation rate and C-reactive protein support a diagnosis of possible inflammatory AAA. Leukocytosis and a relative anaemia on full blood count with positive blood cultures are indicative of infectious AAA.

Predictors of rupture risk include AAA expansion rate, increase in intraluminal thrombus thickness, wall stiffness, wall tension, and peak AAA wall stress.[56] [60]

Risk factors

Strong

cigarette smoking

- This is the risk factor most strongly associated.[1] [9] [11] [17] [18]
- Active cigarette smoking is independently associated with histological high-grade tissue inflammation.[33]
- The duration of smoking is significantly associated with an increased risk in a linear dose-response relationship. Each year of smoking increases the relative risk by 4%.[18]

hereditary/family history

- Studies support a familial aggregation of and genetic predisposition to AAA.[1] [9] [34] [35] [36] [37] [38] [39] [40] [41]
- In a large population-based study, a positive first-degree family history for AAA was more common among cases than among controls (8.4% vs. 4.6%, $P = 0.0001$).[42] The risk of AAA associated with

family history was approximately doubled compared with no family history (odds ratio [OR] 1.9, 95% CI 1.6 to 2.2).

increased age

- Prevalence increases with age.[1] [11]
- Most frequently diagnosed in men >55 years of age, and rupture rarely occurs before 65 years of age.
- AAA is discovered approximately 10 years later in women.[9] [43] [44]

male sex (prevalence)

- AAAs are 4 to 6 times more prevalent in men than women.[1] [9] [11]

female sex (rupture)

- Risk of rupture is greater in women than in men.[45] [46]

congenital/connective tissue disorders

- Aortic degeneration is accelerated in patients with bicuspid aortic valves, Marfan syndrome, and during pregnancy.[47] [48] [49]
- Marfan syndrome specifically is associated with cystic medial necrosis of the aorta secondary to an autosomal dominant anomaly in fibrillin type 1, a structural protein that directs and orients elastin in the developing aorta.[47] [48] As a result, the mature aorta demonstrates abnormal elastic properties, progressive stiffening, and dilation.[49]

Weak

hyperlipidaemia

- Lipoproteins are elevated in patients with AAA independent of cardiovascular risk factors and extent of atherosclerosis.[4] [9]
- AAA patients have significantly lower levels of apolipoprotein AI and HDL cholesterol than matched controls with aorto-iliac occlusive disease.[1] [4]
- High serum total cholesterol is a relatively weak risk factor for AAA, whereas high HDL cholesterol was strongly associated with a low risk of AAA.[11] [44]

COPD

- This is attributed to tobacco-induced elastin degradation.[4] [9]
- Studies suggest that the association between reduced respiratory function and AAA may be due to the activation of inflammation and haemostasis in response to injury.[50]

atherosclerosis (i.e., coronary artery disease [CAD], peripheral arterial occlusive disease)

- CAD is an independent associated risk factor.[1] [17]

hypertension

- Hypertension is a weak independent risk factor.[1] [9] [11]
- There is a relation between systolic BP and AAA in women and an association with ever-use antihypertensive medication and AAA risk for both sexes.[11] [44]

increased height

- Increased height is an independent associated risk factor, although after adjustment for age and sex the association was no longer significant.[\[1\]](#) [\[17\]](#) [\[51\]](#)

central obesity

- One study of more than 12,000 men demonstrated an independent association between central obesity and AAA.[\[44\]](#)

non-diabetic

- A systematic review and meta-analysis found that diabetes protects against the growth and enlargement of AAA.[\[19\]](#) However, the protective mechanism is yet to be determined. Operative and long-term survival was lower among AAA repair patients with diabetes than those without, suggesting an increased cardiovascular burden.[\[19\]](#)

History & examination factors

Key diagnostic factors

presence of risk factors (common)

- Key risk factors include cigarette smoking, family history, increased age, male sex for prevalence and female sex for rupture, and congenital/connective tissue disorders.

palpable pulsatile abdominal mass (uncommon)

- Aneurysm palpation on clinical examination has been shown to be sensitive only in thin patients and those with AAA >5 cm (sensitivity and specificity of 68% and 75%, respectively).[\[1\]](#) [\[57\]](#)

Other diagnostic factors

abdominal, back, or groin pain (uncommon)

- Patients are usually asymptomatic and their aneurysm is detected incidentally.

hypotension (uncommon)

- Patients with ruptured aneurysm present with the triad of abdominal and/or back pain, pulsatile abdominal mass, and hypotension.

Diagnostic tests

1st test to order

Test	Result
abdominal ultrasound <ul style="list-style-type: none"> Definitive test (sensitivity and specificity of 92% to 99% and nearly 100%, respectively).[1] [2] [57] [58] The ultrasound is performed perpendicular to the aortic axis as oblique views may overestimate the true aortic diameter.[2] Intra-observer correlation may be better near the aortic bifurcation than in the proximal infrarenal aorta.[61] Unfortunately, ultrasound offers little utility in imaging aneurysms close to the origins of, or proximal to, the renal arteries.[62] [63] 	abdominal aortic dilation of >1.5 times the expected anterior-posterior diameter of that segment, given the patient's sex and body size; the most commonly adopted threshold is a diameter of 3 cm or more 24957320 LeFevre ML; US Preventive Services Task Force. Screening for abdominal aortic aneurysm: U.S. Preventive Services Task Force recommendation statement. Ann Intern Med. 2014 Aug 19;161(4):281-90. http://annals.org/aim/fullarticle/1883339/screening-abdominal-aortic-aneurysm-u-s-preventive-services-task-force

Other tests to consider

Test	Result
ESR/CRP <ul style="list-style-type: none"> Suggests inflammatory AAA. 	elevated
FBC <ul style="list-style-type: none"> Leukocytosis and a relative anaemia on FBC with positive blood cultures are indicative of infectious AAA. 	leukocytosis, anaemia
blood cultures <ul style="list-style-type: none"> Leukocytosis and a relative anaemia on FBC with positive blood cultures are indicative of infectious AAA. 	positive

Test	Result
computed tomography angiography (CTA)/CT <ul style="list-style-type: none"> May demonstrate retroperitoneal haematoma, discontinuity of the aortic wall, or extravasation of contrast into the peritoneal cavity, which are all signs of rupture.[9] [64] Also useful in diagnosing aortic aneurysms close to the origins of, or proximal to, the renal arteries.[63] [64] The first choice for intraoperative planning; 0.5 mm slices are desirable and 3D reconstruction is essential for accurate planning. 	abdominal aortic dilation of >1.5 times the expected anterior-posterior diameter of that segment, given the patient's sex and body size; the most commonly adopted threshold is a diameter of 3 cm or more 24957320 LeFevre ML; US Preventive Services Task Force. Screening for abdominal aortic aneurysm: U.S. Preventive Services Task Force recommendation statement. Ann Intern Med. 2014 Aug 19;161(4):281-90. http://annals.org/aim/fullarticle/1883339/screening-abdominal-aortic-aneurysm-u-s-preventive-services-task-force
magnetic resonance angiography (MRA)/MRI <ul style="list-style-type: none"> Preoperative study of choice for operative strategy if a patient has an iodinated contrast allergy. 	abdominal aortic dilation of >1.5 times the expected anterior-posterior diameter of that segment, given the patient's sex and body size; the most commonly adopted threshold is a diameter of 3 cm or more 24957320 LeFevre ML; US Preventive Services Task Force. Screening for abdominal aortic aneurysm: U.S. Preventive Services Task Force recommendation statement. Ann Intern Med. 2014 Aug 19;161(4):281-90. http://annals.org/aim/fullarticle/1883339/screening-abdominal-aortic-aneurysm-u-s-preventive-services-task-force

Differential diagnosis

Condition	Differentiating signs / symptoms	Differentiating tests
Diverticulitis	<ul style="list-style-type: none"> • Obstipation; abdominal pain is more common and typically localises to the left lower quadrant. • No pulsatile abdominal mass on clinical examination. Instead, abdominal or perirectal 'fullness' may be appreciated. Fever is possible.[65] 	<ul style="list-style-type: none"> • Stool guaiac testing may be trace positive. • Leukocytosis may be present. • CT scan will demonstrate a normal-calibre aorta and possibly diverticula, inflammation of the pericolic fat or other tissues, bowel-wall thickness >4 mm, or a peridiverticular abscess.[65]
Ureteric colic	<ul style="list-style-type: none"> • Severe abdominal pain that starts in the flank and radiates anteriorly to the groin. • Associated with nausea, emesis, haematuria, dysuria, and urinary frequency or urgency.[66] • Men >55 years of age presenting with ureteric colic should be considered to have a leaking/ruptured AAA until proven otherwise. 	<ul style="list-style-type: none"> • Urinalysis positive for blood and may demonstrate crystals and/or evidence of infection. • Ultrasound and CT scan will demonstrate a normal-calibre aorta and possibly ureteral or renal stones.[66]
Irritable bowel syndrome (IBS)	<ul style="list-style-type: none"> • Intermittent abdominal discomfort with flares lasting 2 to 4 days. • Associated symptoms may include bloating, stool frequency, and abnormal defecation. • Women aged 20 to 40 years are affected more often than men. • General examination is usually normal, although some patients may appear anxious. There may be poorly localised abdominal tenderness to palpation.[67] 	<ul style="list-style-type: none"> • Imaging modalities are often inconclusive, but will demonstrate a normal-calibre aorta.

Condition	Differentiating signs / symptoms	Differentiating tests
Inflammatory bowel disease	<ul style="list-style-type: none"> Abdominal pain is often 'crampy' and left-sided. Patients typically suffer from diarrhoea (bloody and non-bloody), urgency of defecation, and tenesmus. Extra-intestinal manifestations are common in Crohn's disease. Abdominal examination may demonstrate abnormal bowel sounds, detection of an abdominal mass, and pain on palpation. Mucocutaneous lesions may be visible. Perianal fistulae, fissures, or abscesses may be present on rectal examination.[68] 	<ul style="list-style-type: none"> Anaemia is common. Ultrasound or CT scan will demonstrate a normal-calibre aorta. Endoscopic evaluation with biopsy shows typical lesions of ulcerative colitis or Crohn's disease.[68]
Appendicitis	<ul style="list-style-type: none"> Pain is typically periumbilical with localisation to the right lower quadrant. Associated nausea, emesis, and anorexia are common. Patients are classically febrile with tenderness in the right lower quadrant or rebound tenderness on abdominal examination. 	<ul style="list-style-type: none"> Leukocytosis and sterile pyuria on urinalysis is common. Ultrasound or CT scan will demonstrate a normal-calibre aorta with an inflamed appendix or evidence of perforation.
Ovarian torsion	<ul style="list-style-type: none"> Women suffer sudden, continuous, non-specific pain in the lower abdomen; nausea and emesis are common. Patients may demonstrate fever on clinical examination, and an adnexal mass may be palpable.[69] 	<ul style="list-style-type: none"> Leukocytosis may be present. Ultrasound will demonstrate a normal-calibre aorta and possibly reduced or absent adnexal vascular flow.[69]
Gastrointestinal (GI) haemorrhage	<ul style="list-style-type: none"> Patients presenting with haemorrhagic shock may mimic aortic rupture. A history of previous GI bleed, haematemesis, melaena, or bright red blood per rectum is common. Historical risk factors for GI malignancy or peptic ulcer disease may be elicited. On rectal examination gross blood may be visible, or coffee ground haematemesis may be returned with nasogastric tube placement. 	<ul style="list-style-type: none"> Stool is likely to be guaiac positive. Endoscopic evaluation may demonstrate the luminal bleeding source along with mucosal ulcerations, polyps, or tumour. Ultrasound or CT scan will demonstrate a normal-calibre aorta.

Condition	Differentiating signs / Differentiating tests symptoms	
Splanchnic artery aneurysms/acute occlusion	<ul style="list-style-type: none"> • Acute embolic or thrombotic occlusion of the splanchnic vessels results in a marked disparity between acute excruciating mid-abdominal pain and a paucity of early physical findings. • Patients typically suffer unremitting, intense mid-abdominal pain with nausea and vomiting that might be accompanied by explosive diarrhoea. • Most splanchnic artery aneurysms are asymptomatic until rupture.^[70] 	<ul style="list-style-type: none"> • Leukocytosis, haemoconcentration, and systemic acidosis are common with acute splanchnic vessel occlusion. Elevated levels of serum amylase, inorganic phosphorus, creatinine phosphokinase, and alkaline phosphatase may accompany frank bowel infarction. • Angiography is diagnostic and potentially therapeutic in the case of vascular occlusion. • Ultrasound and CT scan will demonstrate a normal-calibre aorta and will diagnose any splanchnic artery aneurysms.^[70]

Step-by-step treatment approach

Patients presenting with a ruptured aneurysm require urgent repair.

For patients with symptomatic aortic aneurysms, repair is indicated regardless of diameter.^[76]

For asymptomatic AAA detected as an incidental finding, surveillance is preferred to repair until the theoretical risk of rupture exceeds the estimated risk of operative mortality. Generally, repair is indicated in patients with large asymptomatic AAA (e.g., with a diameter exceeding 5.5 cm in men or 5.0 cm in women in the US, although treatment decisions based on greater size may differ in other countries, e.g., UK).^{[1] [77] [78] [79] [80] [81]}

Repair of asymptomatic, symptomatic, and ruptured aneurysms can be accomplished using either an endovascular or open surgical technique; the selection of surgical technique should take account of patient preference, patient age, sex, perioperative risk factors, and anatomical factors.

Ruptured AAA

Patients with the triad of abdominal and/or back pain, pulsatile abdominal mass, and hypotension warrant immediate resuscitation and surgical evaluation as repair offers the only potential cure.^{[56] [82]} However, most patients with rupture will not survive to reach theatre.

^[Fig-3]

In patients with confirmed ruptured AAA, 3-year mortality was lower among those randomised to endovascular aneurysm repair (EVAR) than to an open repair strategy (48% vs. 56%; hazard ratio [HR] 0.57, 95% CI 0.36 to 0.90).^[83] The difference between treatment groups was no longer evident after 7 years of follow-up (HR 0.92, 95% CI 0.75 to 1.13). Re-intervention rates were not significantly different between the randomised groups at 3 years (HR 1.02, 95% CI 0.79 to 1.32).^[83] There is some evidence to suggest that an endovascular strategy for repair of ruptured AAA may reduce mortality more effectively in women than in men.^{[83] [84] [85]}

Supportive treatment of ruptured AAA

Standard resuscitation measures are initiated immediately. These include:

- Airway management (supplemental oxygen or endotracheal intubation and assisted ventilation if the patient is unconscious).

[VIDEO: Tracheal intubation animated demonstration]

[VIDEO: Bag-valve-mask ventilation animated demonstration]

- Intravenous access (central venous catheter).
- Arterial catheter; urinary catheter.
- Hypotensive resuscitation: aggressive fluid replacement may cause dilutional and hypothermic coagulopathy and secondary clot disruption from increased blood flow, increased perfusion pressure, and decreased blood viscosity, thereby exacerbating bleeding.^{[89] [90] [91] [92]} Infusing more than 3.5 litres of fluid preoperatively may increase the relative risk of death.^[89] A target systolic BP of 50 to 70 mmHg and withholding fluids is advocated preoperatively.^{[90] [91] [92]}
- Blood product (packed red cells, platelets, and fresh frozen plasma) availability and transfusion for resuscitation, severe anaemia, and coagulopathy.

- Notifying anaesthetic, ICU, and operating teams.

Data from one multi-centre randomised clinical trial support the use of CT scans in cases of ruptured AAA.[93] It did not result in significant delays, and allowed appropriate preoperative planning.

Symptomatic but not ruptured AAA

In patients with symptomatic aortic aneurysm, repair is indicated regardless of diameter.[76] [94]

The development of new or worsening pain may herald aneurysm expansion and impending rupture.

Symptomatic, non-ruptured aneurysm is, therefore, best treated urgently.[56] Under some circumstances, intervention may be delayed for several hours to optimise conditions to ensure successful repair; these patients should be closely monitored in the ICU.[56]

EVAR is increasingly used in the management of patients with symptomatic AAA.[95] [96] In observational studies, short-term all-cause mortality rates did not differ between endovascular and open repair of symptomatic AAA.[95] [96] [97]

Data from the 2011-2013 American College of Surgeons National Surgical Quality Improvement Program suggest that 30-day mortality among patients with symptomatic AAA was approximately double that of patients with asymptomatic AAA following EVAR (symptomatic 3.8% vs. asymptomatic 1.4%, $P = 0.001$) or open repair (symptomatic 7.7% vs. asymptomatic 4.3%, $P = 0.08$).[98] Smaller patient numbers probably contribute to the non-statistically significant finding reported for open repair.

Incidental finding of small asymptomatic AAA

For AAA detected as an incidental finding, surveillance is preferred to repair until the theoretical risk of rupture exceeds the estimated risk of operative mortality.[3] [94] Early surgery for the treatment of smaller AAAs does not reduce all-cause or AAA-specific mortality.[3] [99] One systematic review (4 trials, 3314 participants) found high-quality evidence to demonstrate that immediate repair of small AAA (4 cm to 5.5 cm) did not improve long-term survival compared with surveillance (adjusted HR 0.88, 95% CI 0.75 to 1.02, mean follow-up 10 years).[99] The lack of benefit attributable to immediate surgery was consistent regardless of patient age, diameter of small aneurysm, and whether repair was endovascular or open.[99]

Surgical referral of smaller AAA is usually reserved for rapid growth, or once the threshold diameter for aneurysm repair is reached on repeated ultrasonography.[3]

Medical goals for asymptomatic small aneurysms include:

1. Surveillance:

- American College of Cardiology Foundation/American Heart Association guidelines recommend that infra-/juxtarenal AAAs measuring 4.0 to 5.4 cm in diameter by ultrasound/CT should be monitored every 6 to 12 months.[76] AAAs <4.0 cm require ultrasonography every 2 to 3 years.[76]
- The UK National Health Service recommends that annual screening intervals are employed for 3.0 to 4.4 cm AAAs and 3-month intervals for 4.5 to 5.4 cm AAAs.[100]
- One systematic review and meta-analysis of individual patient data concluded that surveillance intervals of 2 years for 3.0 to 4.4 cm AAA, and 6 months for 4.5 to 5.4 cm AAA, are safe and cost-effective.[75]
- Analysis of AAA growth and rupture rates indicated that in order to maintain an AAA rupture risk <1%, an 8.5-year surveillance interval is required for men with baseline AAA diameter of 3.0 cm.[75] The corresponding estimated surveillance interval for men with an initial aneurysm

diameter of 5.0 cm was 17 months. Despite having similar small aneurysm growth rates, rupture rates were 4 times higher in women than in men.^[75] Surveillance programmes and criteria for considering surgery need to be tailored for women with opportunistically detected AAA.

2. Control of modifiable risk factors for expansion and rupture:

- Smoking cessation - nicotine-replacement therapy, nortriptyline, and bupropion, or counselling.^{[1] [9] [11] [17] [18] [101] [102] [103]}
- Short-term treatment with beta-blockers does not appear to reduce the rate of AAA expansion.^[3] Trials in which patients with small AAAs were randomised to propranolol with the intention of reducing the rate of aneurysm expansion failed to demonstrate significant protective effects.^[104] Propranolol was poorly tolerated in these studies.^[104]

3. Aggressive management of other cardiovascular disease:

- Modifiable cardiovascular risk factors such as hypertension and hyperlipidaemia should be treated. Statins should be started at least 1 month before surgery to reduce cardiovascular morbidity, and continued indefinitely.^[74]

Incidental finding of large asymptomatic AAA

Generally, repair is indicated in patients with large asymptomatic AAA (e.g., with a diameter exceeding 5.5 cm in men or 5.0 cm in women in the US, although treatment decisions based on greater size may differ in other countries, e.g., UK). Repair of aneurysms ≥ 5.5 cm offers a survival advantage.^{[1] [78] [79] [80] [81]}

Decisions regarding repair should be individualised, taking account of patient preference, patient age, sex, perioperative risk factors, and anatomical risk factors. Care should be taken to evaluate patient quality of life, and careful counselling undertaken regarding the risks of surgery (e.g., informing patients of their Vascular Quality Initiative perioperative mortality risk score) and subsequent quality of life. EVAR should be considered in patients who are unfit for open surgery.^{[56] [77] [78] [79] [80] [99] 1[B]Evidence}

Data suggest that in patients with large AAAs (ranging from 5 to 5.5 cm) undergoing elective repair, EVAR is equivalent to open repair in terms of overall survival, although the rate of secondary interventions is higher for EVAR.^[108] EVAR reduces AAA-related mortality (but not longer-term overall survival) in patients with large AAA (≥ 5.5 cm) who are unsuitable for open repair.^{[109] 2[A]Evidence} Post-repair, larger AAAs appear to be associated with worse late survival than smaller aneurysms (pooled HR 1.14 per 1-cm increase in AAA diameter, 95% CI 1.09 to 1.18; 12- to 91.2-month follow-up).^[111] The association is more pronounced with EVAR than with open repair.

Elective repair in asymptomatic patients allows for preoperative assessment, cardiac risk stratification, and medical optimisation of other comorbidities. Coronary artery disease remains the leading cause of early and late mortality after AAA repair.

Endovascular aneurysm repair (EVAR)

EVAR involves the transfemoral endoluminal delivery of a covered stent graft into the aorta, thus sealing off the aneurysm wall from systemic pressures, preventing rupture, and allowing for sac shrinkage. The endograft can be deployed percutaneously through low-profile devices, or after exposing the femoral arteries surgically. A Cochrane review found no difference between the techniques after short follow-up (6

months), except that the percutaneous approach may be faster.[112] Long-term data for the durability of low-profile devices are lacking.

[Fig-4]

[Fig-5]

Assessment of suitability for EVAR should be through the use of 0.5 mm-slice CT angiography. It is essential that the operator is familiar with the specific instructions for use of the endograft to be used.

Lifelong, annual surveillance with ultrasonography or CT is recommended following EVAR of AAAs.[56] [74]

Open repair

Open repair may be transperitoneal or retroperitoneal. With proximal and distal aortic control obtained, the aneurysm is opened, back-bleeding branch arteries are ligated, and a prosthetic graft is sutured from normal proximal aorta to normal distal aorta (or iliac segments). Once flow is restored to the bilateral iliac arteries, the aneurysm sac is closed over the graft.[113] A retroperitoneal approach should be considered for patients in which aneurysmal disease extends to the juxtarenal and/or visceral aortic segment, or in the presence of an inflammatory aneurysm, horseshoe kidney, or hostile abdomen.[56] [114] [115]

Straight tube grafts are recommended for repair in the absence of significant disease of the iliac arteries.[56] The proximal aortic anastomosis should be performed as close to the renal arteries as possible.[56] It is recommended that all portions of an aortic graft should be excluded from direct contact with the intestinal contents of the peritoneal cavity.[56] Re-implantation of a patent inferior mesenteric artery (IMA) should be considered under circumstances that suggest an increased risk of colonic ischaemia (i.e., associated coeliac or superior mesenteric artery occlusive disease, an enlarged meandering mesenteric artery, a history of prior colon resection, inability to preserve hypogastric perfusion, substantial blood loss or intraoperative hypotension, poor IMA back-bleeding when graft open, poor Doppler flow in colonic vessels, or should the colon appear ischaemic).[56] [116]

Choice of elective repair

EVAR accounts for more than 70% of all AAA repairs in the US.[117] In the UK, 69% of elective infrarenal AAAs and 89% of complex AAAs were treated with EVAR during 2014-15.[118] However, not all patients are suitable candidates for EVAR. Guidelines therefore recommend an individualised approach to surgical choice.[56] [74] [76] [119] [120] Factors that will influence the decision include: anatomical determinants (e.g., aneurysm diameter, neck length, neck diameter); life expectancy, sex, comorbidities; and perioperative risk.

EVAR may be preferred in patients who:

- Have high perioperative risk, and
- Have anatomy that is congruent with the relevant stent-graft manufacturer's eligibility criteria as determined in the instructions for use, and
- Are able to satisfy the mandatory surveillance regimen following surgery.

Patients with lower perioperative risk and favourable anatomy may also be candidates for EVAR.[121] [122] Consideration should be given to safety and durability of repair (need for re-intervention).

Open repair may be preferred in patients who:

- Have lower perioperative risk, and
- Are relatively younger.

American College of Cardiology/American Heart Association Practice Guidelines (compilation of 2005 and 2011 guideline recommendations) recommend open aneurysm repair for patients who cannot comply with the long-term surveillance required after EVAR.[76] Open repair is also indicated in patients who may otherwise be considered for EVAR, but who have unfavourable anatomy.[123]

Elective repair outcomes

Data regarding the comparative safety and efficacy of EVAR and open repair differ depending on the outcome of interest. Evidence to date suggests that:

- Short-term postoperative mortality (≤ 30 days) is lower for endovascular than open repair
- Operative, perioperative, and postoperative mortality (≤ 30 days) is greater among women than men for both open repair and EVAR
- Aneurysm-related mortality (≥ 3 years post repair) is higher after EVAR than open repair
- Overall rates of re-intervention are more common following EVAR than open repair.

Six-month postoperative mortality appears to be lower among patients with AAA who undergo EVAR than those who have open surgery, but this is primarily attributable to lower 30-day operative mortality.[124] Pooled analysis of data from 4 high-quality randomised trials (that enrolled patients with AAA diameter >5 cm) found that short-term mortality (30-day or in-hospital mortality) was significantly lower among those randomised to EVAR than among those who underwent open repair (1.4% vs. 4.2%, odds ratio [OR] 0.33, 95% CI 0.20 to 0.55; $P < 0.0001$).[125] However, the early benefit of EVAR is diminished with follow-up.[124] [125] Longer-term mortality (>4 years) did not differ between patients randomised to EVAR or to open surgery (37.3% vs. 37.8%, OR 0.98, 95% CI 0.83 to 1.15; $P = 0.78$). Open repair was significantly associated with lower all-cause mortality than EVAR, after more than 8 years of follow-up, in the UK endovascular aneurysm repair trial 1 (46% vs. 53%, $P = 0.048$).[126]

Perioperative and short-term mortality is higher among women than among men. During elective AAA repair, operative mortality among women exceeds that of men for both open (7.0% vs. 5.2%) and endovascular approaches (2.1% vs. 1.3%).[127] In the UK, women undergoing elective AAA repair were found to have increased short-term mortality compared with men for open repair (30-day mortality: OR 1.39; 95% CI 1.25 to 1.56) and EVAR (30-day mortality: OR 1.57; 95% CI 1.23 to 2.00), despite having fewer preoperative cardiovascular risk factors.[128] Female sex was an independent risk factor for all-cause mortality among women who had an open repair at 1 year (crude cumulative all-cause mortality 15.9% vs. 12.1%, $P < 0.001$) and at 5 years (22.2% vs. 19.6%, $P < 0.001$).[128] Long-term all-cause survival did not differ significantly between women and men in the EVAR group ($P = 0.356$).

One meta-analysis of individual-patient data from 4 trials found that, after 3 years, aneurysm-related mortality was significantly higher in the EVAR group than in those with open repair (pooled hazard ratio [HR] 5.16, 95% CI 1.49 to 17.89; $P = 0.010$).[124] Data from the UK endovascular aneurysm repair trial 1 indicate that, after >8 years of follow-up, aneurysm-related mortality is greater among EVAR patients than open repair patients (5% vs. 1%, $P = 0.0064$).[126] Aneurysm rupture was more common in patients after EVAR than open repair (5.4% vs. 1.4%, $P < 0.001$) in a large cohort study with 8 years of follow-up.[129] Meta-analysis found no significant difference between EVAR and open surgery in the incidence of myocardial death (OR 1.14, 95% CI 0.86 to 1.52; $P = 0.36$), fatal stroke (OR 0.81, 95% CI 0.42 to 1.55; $P = 0.52$), or non-fatal stroke (OR 0.81, 95% CI 0.50 to 1.31; $P = 0.39$).[125] Patients with moderate renal dysfunction or cardiovascular disease do not appear to derive an early survival benefit (to 6 months) from

EVAR, while those with peripheral arterial disease may benefit from open repair.[124] In another meta-analysis, long-term survival following elective AAA repair (EVAR or open) was worst among patients with end-stage renal disease (HR 3.15, 95% CI 2.45 to 4.04) and COPD requiring supplementary oxygen (HR 3.05, 95% CI 1.93 to 4.80).[130]

Overall rates of re-intervention were higher with EVAR than with open surgery; however, rates have been reported heterogeneously in clinical trials.[124] [125] In pooled analysis of individual-patient data, re-intervention was reported in 65.8% of EVAR patients with type I endoleak (79 of 120) and 22.8% of EVAR patients with type II endoleak (99 of 435) over 5 years of follow-up.[124] Observational data suggest that interventions related to the management of aneurysm or its complications are more common following EVAR than open surgery (18.8% vs. 3.7%, $P < 0.001$) over 8 years of follow-up.[129]

Low-quality evidence from 4 small RCTs suggests that elective open repair performed retroperitoneally can reduce blood loss and hospital stay compared with a transperitoneal approach.[131] However, there was no difference in mortality between retroperitoneal and transperitoneal elective open AAA repair (very low-quality evidence). Moreover, the retroperitoneal approach may increase the risk of haematoma, chronic wound pain, and abdominal wall hernia compared with transperitoneal.[131]

Adverse effects

Complications of EVAR may include endoleak, graft occlusion, and graft migration with aortic neck expansion.[119] A systematic review reported aortic neck dilation in 24.6% of EVARs (9439 men included), which led to higher rates of type I endoleak, graft migration, and re-intervention.[132]

As an adjunct to EVAR, bilateral hypogastric artery occlusion may be acceptable in certain anatomical situations for patients at high risk for open surgical repair. Buttock claudication and erectile dysfunction may occur in up to 40% of patients after unilateral embolisation; these symptoms may persist in 11% to 13% of patients following bilateral occlusion.[56] [133] [134] Internal iliac artery revascularisation techniques, involving specialised iliac branch devices, have high technical success rates and are associated with low morbidity (e.g., buttock claudication rate of 4.1%).[135]

In patients with synchronous intra-abdominal malignancy, EVAR reduces mortality and delay between the treatment of the 2 pathologies despite a significant risk for thrombotic events.[136] [137]

Complications of open repair include cardiac and pulmonary events, mesenteric ischaemia, renal failure, bleeding, wound and graft infection, spinal cord ischaemia/paraplegia, embolisation/limb ischaemia, and late graft complications (i.e., aorto-enteric fistula and aortic pseudoaneurysm).[1] [138]

Perioperative management

Treatment of co-existing cardiac disease:

- Cardiovascular risk prevention should be prioritised in patients in AAA screening programmes.[139]
- Non-invasive stress testing should be considered for patients with a history of ≥ 3 clinical risk factors (i.e., coronary artery disease, congestive heart failure, stroke, diabetes mellitus, chronic renal insufficiency) and an unknown or poor functional capacity ($MET < 4$) if it will change management.
- Coronary revascularisation is indicated for those patients who present with acute ST elevation myocardial infarction, unstable angina, or stable angina with left main coronary artery or 3-vessel disease, as well as those patients with 2-vessel disease that includes the proximal left anterior descending artery and either ischaemia on non-invasive testing or an ejection fraction < 0.5 . [56]

Regarding blood transfusion:[56]

- Preoperative autologous blood donation may be beneficial for patients undergoing open aneurysm repair.
- Cell salvage or an ultrafiltration device is recommended if large blood loss is anticipated or the risk of disease transmission from banked blood is considered high.
- Blood transfusion is recommended if the intraoperative haemoglobin level is <10 g/dL in the presence of ongoing blood loss. Consider use of fresh frozen plasma and platelets in a ratio with packed blood cells of 1:1:1.

Pulmonary artery catheters should not be used routinely in aortic surgery, unless there is a high risk for a major haemodynamic disturbance.[56] Central venous access is recommended for all patients undergoing open aneurysm repair.[56] DVT prophylaxis consisting of intermittent pneumatic compression and early ambulation are recommended for all patients undergoing open repair or EVAR.[56] [140]

Avoiding hypothermia during open repair and EVAR can reduce hospital length of stay, length of stay in the ITU, and rates of organ dysfunction.[141]

Preoperative cardiovascular risk reduction:

- Addressing modifiable cardiovascular risk factors preoperatively improves long-term survival after AAA repair.[142]
- Preoperative exercise training reduced post-surgical cardiac complications in a small randomised controlled trial of patients undergoing open or endovascular AAA repair.[143]
- Perioperative statin use reduces cardiovascular events during non-cardiac surgery.[144] Statins should be started at least 1 month before surgery to reduce cardiovascular mortality, and continued indefinitely.[74]
- In the absence of contraindications, patients should receive low-dose aspirin and this should be continued during the perioperative period.[74]
- Hypertension should be controlled to reduce cardiovascular morbidity.[74]
- Preoperative beta-blockade may be reasonable in patients at high risk of myocardial ischaemia (ischaemic heart disease or myocardial ischaemia present on stress testing), if the therapy can be commenced more than 1 month before surgery.[74] [145] Large trials where beta-blockade was started a few days before surgery have indicated no benefit, or even harm, from perioperative beta-blockade.[146] [147] [148]
- A large, multi-centre study of patients undergoing non-cardiac surgery found that clonidine did not reduce the rate of death or non-fatal myocardial infarction.[149] Alpha-2 agonists are not, therefore, recommended for non-cardiac surgery patients.[144]

Antibiotic cover:

- Antibiotic therapy is indicated for patients undergoing elective and urgent repair of ruptured AAA to cover gram-positive and gram-negative organisms (i.e., *Staphylococcus aureus*, *Staphylococcus epidermidis*, and enteric gram-negative bacilli) and prevent graft infection.
- Broad-spectrum antibiotic coverage is tailored to patient clinical presentation and cultures, and in accordance with local protocols.

Endovascular aneurysm repair (EVAR) leak

Endoleak is persistent blood flow outside the graft and within the aneurysm sac.[150] [151] It is not a complication following open repair.

Postoperative surveillance can detect major endoleaks and aneurysm sac expansion.

Risk of endoleak following EVAR is 24%.[150] There are 5 types of endoleak.

Type I:

- Leak at the attachment site (proximal/distal end of the endograft or iliac occluder); usually immediate, but delayed leaks may occur.
[Fig-6]
- Every effort should be made to repair type I endoleak before completing the procedure (e.g., balloon moulding of the proximal seal zone, placement of a proximal cuff, endostaples). Persistent type IA endoleak may necessitate conversion to open repair.[56]
[Fig-7]
[Fig-8]

Type II:

- Patent branch leak.
[Fig-9]
- Spontaneous resolution may occur, although persistence may result in sac growth.[152]
- If a type II endoleak or other abnormality of concern is observed on contrast-enhanced CT imaging at 1 month after EVAR, postoperative imaging at 6 months is recommended.[56]
- Treatment remains controversial and is advocated either if persistent at 6 to 12 months or when aneurysm sac size increases such that proximal and/or distal sealing zones may be compromised.[153] [154] [155] [156]
- Treatment of choice is transarterial coil embolisation, although laparoscopic ligation of collateral branches, direct percutaneous translumbar puncture of the sac, translumbar embolisation, and transcatheter transcaval embolisation have been reported.[151] [153] [154] [155] [157] [158] [159] [160]

Type III:

- Graft defect with leak through fabric tears, graft disconnection, or disintegration of the fabric.[150] [151]
- Repair is indicated upon discovery (endovascular stent graft extension).[56] [154]

Type IV:

- Leak from graft wall porosity.[150] [151]
- These leaks are uncommon with newer stent grafts and are self-limiting.[56] [154]

Type V (endotension):

- Endotension is increased intrasac pressure after EVAR without visualised endoleak on delayed contrast CT scans.

- Endotension is less common with the newer-generation grafts.[56]
- There is no standardised method to measure endotension or consensus on indicated therapy in the absence of aneurysm enlargement; however, treatment of endotension to prevent aneurysm rupture is suggested in selected patients with continued aneurysm expansion.[56] [151]

[\[VIDEO: Bag-valve-mask ventilation animated demonstration \]](#)

[\[VIDEO: Central venous catheter insertion animated demonstration \]](#)

[\[VIDEO: Female urethral catheterisation animated demonstration \]](#)

[\[VIDEO: Male urethral catheterisation animated demonstration \]](#)

Treatment details overview

Consult your local pharmaceutical database for comprehensive drug information including contraindications, drug interactions, and alternative dosing. (see [Disclaimer](#))

Acute (summary)		
Patient group	Tx line	Treatment
ruptured AAA	1st	standard resuscitation measures
	plus	urgent surgical repair
	plus	perioperative antibiotic therapy
symptomatic, but not ruptured AAA	1st	urgent surgical repair
	plus	preoperative cardiovascular risk reduction
	plus	perioperative antibiotic therapy

Ongoing (summary)		
Patient group	Tx line	Treatment
incidental finding: small asymptomatic AAA	1st	surveillance
	plus	aggressive cardiovascular risk management
incidental finding: large asymptomatic AAA	1st	elective surgical repair
	plus	preoperative cardiovascular risk reduction

Ongoing (summary)		
	plus	perioperative antibiotic therapy
endovascular repair leak requiring treatment	1st	corrective procedure
	plus	preoperative cardiovascular risk reduction
	plus	perioperative antibiotic therapy

Treatment options

Acute

Patient group	Tx line	Treatment
ruptured AAA	1st	<p>standard resuscitation measures</p> <ul style="list-style-type: none"> » The airway is managed with supplemental oxygen or endotracheal intubation and assisted ventilation if the patient is unconscious. <p>[VIDEO: Tracheal intubation animated demonstration]</p> <p>[VIDEO: Bag-valve-mask ventilation animated demonstration]</p> <ul style="list-style-type: none"> » A central venous catheter is inserted. » Monitoring requires insertion of an arterial catheter and urinary catheter. » A target systolic BP of 50 to 70 mmHg and withholding fluids is advocated preoperatively.^{[90] [91] [92]} » Aggressive fluid replacement may cause dilutional and hypothermic coagulopathy and secondary clot disruption from increased blood flow, increased perfusion pressure, and decreased blood viscosity, thereby exacerbating bleeding.^{[89] [90] [91] [92]} Infusing more than 3.5 L of fluid preoperatively may increase the relative risk of death.^[89] <p>plus</p> <p>urgent surgical repair</p> <ul style="list-style-type: none"> » Aorto-iliac anatomy permitting, endovascular aneurysm repair (EVAR) is preferred.^[83] » In patients with confirmed ruptured AAA, 3-year mortality was lower among those randomised to EVAR than to an open repair strategy (48% vs. 56%; hazard ratio [HR] 0.57, 95% CI 0.36 to 0.90).^[83] The difference between treatment groups was no longer evident after 7 years of follow-up (HR 0.92, 95% CI 0.75 to 1.13). Re-intervention rates were not significantly different between the randomised groups at 3 years (HR 1.02, 95% CI 0.79 to 1.32).^[83] » There is some evidence to suggest that an endovascular strategy for repair of ruptured AAA

Acute

Patient group

Tx line

Treatment

may reduce mortality more effectively in women than in men.[83] [84] [85]

plus

perioperative antibiotic therapy

» Antibiotic therapy is indicated for patients undergoing emergency repair of ruptured AAA to cover gram-positive and gram-negative organisms and prevent graft infection.

» Broad-spectrum antibiotic coverage is tailored to patient clinical presentation and cultures, and in accordance with local protocols.

symptomatic, but not ruptured AAA

1st

urgent surgical repair

» In patients with symptomatic aortic aneurysm, repair is indicated regardless of diameter.[76] [94] The development of new or worsening pain may herald aneurysm expansion and impending rupture. Symptomatic, non-ruptured aneurysm is, therefore, best treated urgently.[56] Under some circumstances, intervention may be delayed for several hours to optimise conditions to ensure successful repair; these patients should be closely monitored in the ICU.[56]

» Endovascular aneurysm repair (EVAR) is increasingly used in the management of patients with symptomatic AAA.[95] [96] In observational studies, short-term all-cause mortality rates did not differ between endovascular and open repair of symptomatic AAA.[95] [96] [97]

plus

preoperative cardiovascular risk reduction

» In the absence of contraindications, patients should receive low-dose aspirin and this should be continued during the perioperative period.[74]

» Hypertension should be controlled to reduce cardiovascular mortality.[74]

plus

perioperative antibiotic therapy

» Perioperative antibiotic therapy is given. Broad-spectrum antibiotic coverage is necessary, in accordance with local protocols.

Ongoing

Patient group	Tx line	Treatment
incidental finding: small asymptomatic AAA	1st	<p>surveillance</p> <ul style="list-style-type: none"> » For AAA detected as an incidental finding, surveillance is preferred to repair until the theoretical risk of rupture exceeds the estimated risk of operative mortality.[3] [94] » Early open surgery for the treatment of smaller AAAs does not reduce all-cause or AAA-specific mortality.[3] [99] One systematic review (4 trials, 3314 participants) found high-quality evidence to demonstrate that immediate repair of small AAA (4 cm to 5.5 cm) did not improve long-term survival compared with surveillance (adjusted HR 0.88, 95% CI 0.75 to 1.02, mean follow-up 10 years).[99] The lack of benefit attributable to immediate surgery was consistent regardless of patient age, diameter of small aneurysm, and whether repair was endovascular or open.[99] » Surgical referral of smaller AAA is usually reserved for rapid growth, or once the threshold diameter for aneurysm repair is reached on repeated ultrasonography.[3] <p>plus</p> <p>aggressive cardiovascular risk management</p> <ul style="list-style-type: none"> » Patients should be encouraged to stop smoking and offered drug therapy to assist with this if needed. » Cardiovascular risk prevention should be prioritised in patients in AAA screening programmes.[139] » Addressing modifiable risk factors preoperatively improves long-term survival after AAA repair.[142] » Preoperative exercise training reduced post-surgical cardiac complications in a small randomised controlled trial of patients undergoing open or endovascular AAA repair.[143] » In the absence of contraindications, patients should receive low-dose aspirin and this should be continued during the perioperative period.[74] » Hypertension should be controlled to reduce cardiovascular mortality.[74]

Ongoing

Patient group

Tx line

Treatment

» Perioperative statin use reduces cardiovascular events during non-cardiac surgery.^[144] Statins should be started at least 1 month before surgery to reduce cardiovascular mortality, and continued indefinitely.^[74]

» Perioperative beta-blockade may be reasonable in patients at high risk of myocardial ischaemia (ischaemic heart disease or myocardial ischaemia present on stress testing), if the therapy can be commenced more than 1 month before surgery.^{[74] [145]}

incidental finding: large asymptomatic AAA

1st

elective surgical repair

» Generally, repair is indicated in patients with large asymptomatic AAA (e.g., with a diameter exceeding 5.5 cm in men or 5.0 cm in women in the US, although treatment decisions based on greater size may differ in other countries, e.g., UK). Repair of aneurysms ≥ 5.5 cm offers a survival advantage.^{[1] [78] [79] [80] [81]}

» Decisions regarding repair should be individualised, taking account of patient preference, patient age, sex, perioperative risk factors, and anatomical risk factors. Care should be taken to evaluate patient quality of life, and careful counselling undertaken regarding the risks of surgery and subsequent quality of life. Endovascular aneurysm repair (EVAR) should be considered in patients who are unfit for open surgery.^{[56] [77] [78] [79] [80] [99] 1[B]Evidence}

» Data suggest that in patients with large AAAs (≥ 5.5 cm) undergoing elective repair, EVAR is equivalent to open repair in terms of overall survival, although the rate of secondary interventions is higher for EVAR.^[108] EVAR also reduces AAA-related mortality (but not longer-term overall survival) in patients with large AAA (≥ 5.5 cm) who are unsuitable for open repair.^{[109] 2[A]Evidence}

» Post repair, larger AAAs appear to be associated with worse late survival than smaller aneurysms (pooled HR 1.14 per 1-cm increase in AAA diameter, 95% CI 1.09 to 1.18; 12- to 91.2-month follow-up).^[111] The association is more pronounced with EVAR than with open repair.

plus

preoperative cardiovascular risk reduction

Ongoing

Patient group

Tx line

Treatment

- » Addressing modifiable risk factors preoperatively improves long-term survival after AAA repair.[142]
- » Preoperative exercise training reduced post-surgical cardiac complications in a small randomised controlled trial of patients undergoing open or endovascular AAA repair.[143]
- » In the absence of contraindications, patients should receive low-dose aspirin and this should be continued during the perioperative period.[74]
- » Hypertension should be controlled to reduce cardiovascular mortality.[74]
- » Perioperative statin use reduces cardiovascular events during non-cardiac surgery.[144] Statins should be started at least 1 month before surgery to reduce cardiovascular mortality, and continued indefinitely.[74]
- » Perioperative beta-blockade may be reasonable in patients at high risk of myocardial ischaemia (ischaemic heart disease or myocardial ischaemia present on stress testing), if the therapy can be commenced more than 1 month before surgery.[74] [145]

plus

perioperative antibiotic therapy

- » Perioperative antibiotic therapy is given. Broad-spectrum antibiotic coverage is necessary, in accordance with local protocols.

endovascular repair leak requiring treatment

1st

corrective procedure

- » Endoleak is persistent blood flow outside the graft and within the aneurysm sac.[150] [151] It is not a complication following open repair.
- » Postoperative surveillance can detect major endoleaks and aneurysm sac expansion. Risk of endoleak following endovascular aneurysm repair (EVAR) is 24%.[150]
- » Type I: clinically significant; every effort should be made to repair type I endoleak before completing the procedure (e.g., balloon moulding of the proximal seal zone, placement of a proximal cuff, endostaples). Persistent type IA endoleak may necessitate conversion to open repair.[56]

Ongoing

Patient group

Tx line

Treatment

[Fig-7]

[Fig-8]

» Type II: treatment remains controversial and is advocated either if persistent at 6 to 12 months or when aneurysm sac size increases such that proximal and/or distal sealing zones may be compromised.[153] [154] [155] [156] Treatment of choice is transarterial coil embolisation, although laparoscopic ligation of collateral branches, direct percutaneous translumbar puncture of the sac, translumbar embolisation, and transcatheter transcaval embolisation have been reported.[151] [153] [154] [155] [157] [158] [159] [160]

» Type III: repair is indicated upon discovery (endovascular stent graft extension).[56] [154]

» Type IV: these leaks are uncommon with newer stent grafts and are self-limiting, requiring no treatment.[56] [154]

» Type V (endotension): less common with the newer-generation grafts;[56] there is no standardised method to measure endotension or consensus on indicated therapy in the absence of aneurysm enlargement; however, treatment of endotension to prevent aneurysm rupture is suggested in selected patients with continued aneurysm expansion.[56] [151]

plus

preoperative cardiovascular risk reduction

» In the absence of contraindications, patients should receive low-dose aspirin and this should be continued during the perioperative period.[74]

» Hypertension should be controlled to reduce cardiovascular mortality.[74]

» Perioperative statin use reduces cardiovascular events during non-cardiac surgery.[144] Statins should be started at least 1 month before surgery to reduce cardiovascular mortality, and continued indefinitely.[74]

» Perioperative beta-blockade may be reasonable in patients at high risk of myocardial ischaemia (ischaemic heart disease or myocardial ischaemia present on stress testing), if the therapy can be commenced more than 1 month before surgery.[74] [145]

Ongoing

Patient group

Tx line

Treatment

plus

perioperative antibiotic therapy

» Perioperative antibiotic therapy is given. Broad-spectrum antibiotic coverage is necessary, in accordance with local protocols.

Emerging

Doxycycline

Doxycycline is a non-specific inhibitor of matrix metalloproteinases (MMPs). MMPs promote degradation of collagen and elastin and are integral to aneurysm formation.[1] [21] One clinical trial found that prolonged administration of doxycycline for 6 months is safe and well tolerated by patients with small asymptomatic AAAs and is associated with a gradual reduction in plasma MMP-9 levels.[1] [170] Another small randomised study reported no growth of aortic aneurysms in doxycycline-treated patients at 6 and 12 months.[171] [172] N-TA³CT is a phase II randomised, double-blind, placebo-controlled study to determine whether doxycycline can reduce the rate of increase of small (3.5 to 5.0 cm among men and 3.5 to 4.5 cm among women) AAAs in patients ≥55 years of age.[173] The primary outcome is AAA maximum transverse diameter determined by CT scans at 2-year follow-up (with allowance for baseline diameter). The expected completion date of the N-TA³CT study is September 2019. Currently, there is insufficient evidence to recommend use of doxycycline.[174]

Ticagrelor

A phase II randomised placebo-controlled trial is investigating whether treatment with ticagrelor (a P2Y₁₂ receptor antagonist) inhibits growth of small AAA in aspirin-naïve individuals (aged 50 to 85 years) with confirmed AAA between 35 and 49 mm.[175] The primary outcome of this study is mean reduction in AAA volume growth rate (%) measured with magnetic resonance imaging (MRI) at 12 months. Estimated study completion date is July 2017.

Endovascular aneurysm sealing

Endovascular aneurysm sealing (EVAS) has been developed to address the issues of endoleak and stent graft failure-related complications associated with endovascular aneurysm repair (EVAR). It involves inserting expandable stents, each with an attached bag (endobag), into the aneurysm via both femoral arteries simultaneously. A polymer is injected into the endobag to fill the aneurysm sac, thereby excluding it from the circulation. EVAS is intended to stabilise the stent graft position and reduce the rate of endoleaks and repeat interventions.[176] There is a paucity of good-quality evidence regarding the use of EVAS; however, an industry-sponsored trial and registry report study demonstrated high technical success rates, low perioperative mortality, and high rates of freedom from adverse events at 1 month.[177] [178] Endoleak rates were low, but they are more difficult to detect after EVAS and the treatment options are more complex than current EVAR endoleak treatments.[177] [178] Device migration was reported in a retrospective case series of EVAS patients (17% grafts had migrated caudally).[179] This was without clinical sequelae, but the long-term stability and reliability of the EVAS grafts requires further study and close observation.

Fenestrated EVAR (FEVAR)

FEVAR is a viable alternative to open repair for juxta- and suprarenal AAA, or for those with AAA where a short or diseased neck precludes conventional repair. However, there is no level 1 evidence for FEVAR.[180] The largest published single-centre series demonstrated mortality of 0.8% and renal occlusion rate of 4.3%.[181]

Recommendations

Monitoring

In the US, guidelines recommend that infra-/juxtarenal AAAs measuring 4.0 to 5.4 cm in diameter with ultrasonography/CT should be monitored every 6 to 12 months.[76] AAAs <4.0 cm require ultrasonography every 2 to 3 years.[76]

In the UK, annual intervals are employed for small AAAs (3.0 cm-4.4 cm), and 3-month intervals for 4.5- to 5.4-cm AAAs.[100]

A systematic review and meta-analysis of individual patient data concluded that surveillance intervals of 2 years for 3.0- to 4.4-cm AAAs, and 6 months for 4.5- to 5.4-cm AAAs, are safe and cost-effective.[75]

Analysis of AAA growth and rupture rates indicated that, in order to maintain an AAA rupture risk <1%, an 8.5-year surveillance interval is required for men with baseline AAA diameter of 3.0 cm.[75] The corresponding estimated surveillance interval for men with an initial aneurysm diameter of 5.0 cm was 17 months. Despite having similar growth rates of small aneurysms, rupture rates were 4 times greater in women than in men.[75] Surveillance programmes and criteria for considering surgery need to be tailored for women with opportunistically detected AAA.

Post repair

The US Society for Vascular Surgery recommends follow-up non-contrast CT imaging at 5-year intervals after open repair or endovascular aneurysm repair (EVAR).[56]

Post EVAR, baseline surveillance is recommended at 1 month with contrast-enhanced CT and colour duplex ultrasound imaging.[56] If neither endoleak nor AAA enlargement is documented, imaging should be repeated at 12 months using either contrast-enhanced CT or colour duplex ultrasound imaging.[56]

The European Society for Vascular Surgery recommends that all patients should have computed tomography angiography (CTA) and plain radiographs 30 days after the initial repair.[74] If there is endoleak, CTA at 6 months and 12 months with plain radiographs is warranted. Patients with no early endoleak and good component overlap may omit the 6-month CTA, but should have CTA and plain radiographs at 12 months. If there is no endoleak and a stable/shrinking AAA, annual duplex ultrasonography is recommended with plain radiographs.[74]

Antibiotic prophylaxis of graft infection is required prior to bronchoscopy, gastrointestinal or genitourinary endoscopy, and any dental procedure that may lead to bleeding.[56]

Generalised sepsis, groin drainage, pseudoaneurysm formation, or ill-defined pain after open repair or EVAR should prompt evaluation of graft infection.[56] Gastrointestinal bleeding after open repair or EVAR should prompt evaluation of an aorto-enteric fistula.[56]

Patient instructions

Patients should be educated on the importance of smoking cessation (including counselling and pharmacotherapy as needed), and of blood pressure and cholesterol control.

Complications

Complications	Timeframe	Likelihood
abdominal compartment syndrome	short term	high

Complications	Timeframe	Likelihood
Rates of abdominal compartment syndrome (ACS) may approach 55% after open repair of ruptured AAA.[74] A small retrospective review reported ACS in 34% of patients following open repair of ruptured AAA and 21% after endovascular aneurysm repair (EVAR).[183] A nationwide population-based study (Swedish Vascular Registry) found that after ruptured AAA repair, ACS developed in 6.8% following open repair compared with 6.9% after EVAR.[184]		
ileus, intestinal obstruction, and ischaemic colitis	short term	high
Ileus has been reported in 11% of patients, with intestinal obstruction and colitis each occurring in 1% of patients undergoing open repair.[182] [185] Ischaemic colitis requiring colectomy is rare.		
acute kidney injury	short term	high
Following EVAR and open repair of AAA, there is a significant incidence of acute kidney injury (AKI).[186] [187] [188] In open repair, this seems to be transient. However, following EVAR the causes are multifactorial, and decline in renal function is significantly greater (especially with suprarenal fixation) than in open surgery.[189] AKI following EVAR is associated with medium-term increased morbidity and mortality.[186]		
post-implantation syndrome	short term	medium
May last for up to 10 days following EVAR.[74] Fever, malaise, and back pain, which may be due to cytokine release, are typical.		
amputation due to limb ischaemia	short term	low
Rates of amputation due to limb ischaemia were very low in a large series of patients who underwent open repair.[182]		
spinal cord ischaemia	short term	low
Spinal cord ischaemia is rare after EVAR, with an incidence in the EUROSTAR collaborators registry of 0.21%.[190] In a retrospective analysis of emergency endovascular treatment for ruptured AAA, 4 of 35 patients (11.5%) developed spinal cord ischaemia postoperatively.[191] Delayed spinal cord ischaemia (developing 2 days after EVAR) has been reported.[192]		
impaired sexual function	long term	high
Damage to the autonomic nerves present at the aorto-iliac bifurcation during dissection, as well as reduction in pelvic blood supply, can result in impotence and retrograde ejaculation. In one trial, 10% of men developed new impotence in the first year after open repair.[193]		
anastomotic pseudoaneurysm	long term	high
One case series reported para-anastomotic aneurysms in 10% of patients after aortic bypass grafting.[194] The rate of femoral anastomotic pseudoaneurysm may be as high as 20% at 10 years after aortobifemoral reconstruction for AAA.[195]		
aortic neck dilation	long term	high
Aortic neck dilation occurred in 24.6% of EVAR patients during 15 months to 9 years of follow-up.[132] A composite clinical event of endoleak, migration, and re-intervention was significantly more common in this group than in those patients without aortic neck dilation.		

Complications	Timeframe	Likelihood
graft infection	long term	medium
Can result from infection during implantation, or haematogenous seeding following dental procedures or endoscopic procedures with biopsy. Incidence is low: a retrospective cohort study found that the 2-year rate of graft infection was 0.19% following open repair versus 0.16% with EVAR.[196]		
ureteric obstruction	long term	low
Ureteric obstruction is related to encasement of the ureters in an inflammatory perianeurysmal fibrosis of unresolved aetiology rather than secondary to aneurysm compression.[197] Most often, ureteral compression is associated with inflammatory aortic aneurysm. Extensive retroperitoneal adhesions may result in ureteral obstruction in 18% of patients. The inferior vena cava may become involved as well.[198]		
functional gastric outlet obstruction	long term	low
Duodenal obstruction is a consequence of compression of the duodenum in its fixed retroperitoneal course between the aneurysmal aorta and the superior mesenteric artery.[197]		
graft limb occlusion	long term	low
The incidence of graft limb occlusion up to 10 years after open AAA repair has been reported as being between 2.6% and 3.0%.[199] [200] The risk of graft occlusion is greater with EVAR, with a reported incidence of up to 7.2% in follow-up studies.[201] Kinking is a risk factor for graft limb occlusion following EVAR.[202]		
endoleak	variable	high
Risk of endoleak following EVAR is 24%.[150] Type II endoleaks are the most common. Repair may be indicated upon discovery of postoperative type I endoleak. Endoleak is not a complication following open repair.		
distal embolisation	variable	low
Incidence is 3% to 29%, most commonly affecting the digits (blue toe syndrome). There is a 5% incidence of distal embolisation resulting in limb-threatening ischaemia, digital ischaemia, and calf myonecrosis.[203]		

Prognosis

The natural course involves slow and steady growth with ultimate progression to rupture. Most patients with rupture will not survive to reach the operating theatre. Given the morbidity and mortality associated with surgical intervention, repair is typically deferred until the theoretical risk of rupture exceeds the estimated risk of operative mortality. The majority of patients undergoing open repair remain without significant graft-related complications during the remainder of their lives (0.4% to 2.3% incidence of late graft-related complications).[1] [182] Five-year survival rates after intact aneurysm repair average 60% to 75%. Those undergoing endovascular aneurysm repair (EVAR) are more likely to have a delayed complication and require re-intervention.

Diagnostic guidelines

Europe

Abdominal aortic aneurysm screening: how it works

Published by: Public Health England

Last published: 2015

Summary: Includes information about the NHS AAA screening programme, and the tests and processes involved.

2014 ESC guidelines on the diagnosis and treatment of aortic diseases

Published by: European Society of Cardiology

Last published: 2014

Summary: Includes a critical evaluation of diagnostic imaging and screening for AAA.

North America

The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm

Published by: Society for Vascular Surgery

Last published: 2018

Summary: Addresses patient evaluation and risk of rupture in AAA.

ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery

Published by: American College of Cardiology; American Heart Association

Last published: 2014

Summary: Recommendations regarding preoperative risk assessment and cardiac testing in the adult patient undergoing non-cardiac surgery.

Screening for abdominal aortic aneurysm: US Preventive Services Task Force recommendation statement

Published by: US Preventive Services Task Force

Last published: 2014

Summary: Describes risk factors and assessment, and includes recommendations regarding screening.

Treatment guidelines

Europe

2014 ESC guidelines on the diagnosis and treatment of aortic diseases

Published by: European Society of Cardiology

Last published: 2014

Summary: Includes recommendations for the management of asymptomatic and symptomatic AAA.

Europe

Management of abdominal aortic aneurysms: clinical practice guidelines of the European Society for Vascular Surgery**Published by:** European Society for Vascular Surgery**Last published:** 2011**Summary:** Evidence-based recommendations regarding the clinical care of preoperative, perioperative, and postoperative AAA patients.**Endovascular stent-grafts for the treatment of abdominal aortic aneurysms****Published by:** National Institute for Health and Care Excellence**Last published:** 2009**Summary:** Appraises the use of grafts in the treatment of AAA.**Laparoscopic repair of abdominal aortic aneurysm****Published by:** National Institute for Health and Care Excellence**Last published:** 2007**Summary:** Recommendations regarding the laparoscopic repair of AAA, including who should carry out the procedure, and the information given to patients and their Trust.

North America

The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm**Published by:** Society for Vascular Surgery**Last published:** 2018**Summary:** Reviews peri- and intra-operative strategies, follow-up, and treatment of complications.**ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery****Published by:** American College of Cardiology; American Heart Association**Last published:** 2014**Summary:** Guideline addressing pharmacotherapeutic and anaesthetic considerations for non-cardiac surgery patients.**Screening for abdominal aortic aneurysm: US Preventive Services Task Force recommendation statement****Published by:** US Preventive Services Task Force**Last published:** 2014**Summary:** Identifies patients who are candidates for open surgical repair or endovascular aneurysm repair (EVAR), and those who should be managed conservatively via surveillance.**Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA guideline recommendations)****Published by:** American College of Cardiology Foundation; American Heart Association**Last published:** 2013**Summary:** Recommendations for the screening and management of patients with aneurysms of the abdominal aorta.

North America

Guidelines for the treatment of abdominal aortic aneurysms

Published by: American Association for Vascular Surgery; Society for Vascular Surgery **Last published:** 2003

Summary: Decisions relating to elective repair of AAA require careful assessment of factors that influence rupture risk, operative mortality, and life expectancy.

Evidence scores

1. Rupture: there is medium-quality evidence that the risk of rupture was 20% for aneurysms measuring larger than 5.0 cm to 6.0 cm in diameter, and 2 prospective randomised trials comparing early intervention versus expectant observation for infrarenal AAAs measuring 4.0 cm to 5.4 cm in diameter were conducted in the UK and the US Department of Veterans Affairs during the past decade.[\[105\]](#) [\[106\]](#) [\[107\]](#)

Evidence level B: Randomized controlled trials (RCTs) of <200 participants, methodologically flawed RCTs of >200 participants, methodologically flawed systematic reviews (SRs) or good quality observational (cohort) studies.

2. Reduction in AAA-related mortality in patients unsuitable for open repair: there is good-quality evidence from a large randomised controlled trial that, in patients with large abdominal aortic aneurysms 5.5 cm or greater physically ineligible for open repair, endovascular repair (EVAR) was associated with a significantly lower AAA-related mortality compared with no repair, although no reduction in overall mortality was seen after 8 years. However, the rate of graft-related complications and secondary interventions following EVAR was high.[\[110\]](#)

Evidence level A: Systematic reviews (SRs) or randomized controlled trials (RCTs) of >200 participants.

Key articles

- Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg*. 2018 Jan;67(1):2-77.e2. [Full text](#) [Abstract](#)
- Moll FL, Powell JT, Fraedrich G, et al; European Society for Vascular Surgery. Management of abdominal aortic aneurysms: clinical practice guidelines of the European Society for Vascular Surgery. *Eur J Vasc Endovasc Surg*. 2011 Jan;41 Suppl 1:S1-58. [Full text](#) [Abstract](#)
- Anderson JL, Halperin JL, Albert NM, et al. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA guideline recommendations): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2013 Apr 2;127(13):1425-43. [Full text](#) [Abstract](#)
- Brewster DC, Cronenwett JL, Hallett JW Jr, et al. Guidelines for the treatment of abdominal aortic aneurysms: report of a subcommittee of the Joint Council of the American Association for Vascular Surgery and Society for Vascular Surgery. *J Vasc Surg*. 2003 May;37(5):1106-17. [Full text](#) [Abstract](#)
- IMPROVE Trial Investigators. Comparative clinical effectiveness and cost effectiveness of endovascular strategy v open repair for ruptured abdominal aortic aneurysm: three year results of the IMPROVE randomised trial. *BMJ*. 2017 Nov 14;359:j4859. [Full text](#) [Abstract](#)
- Greenhalgh RM, Brown LC, Powell JT, et al; United Kingdom EVAR Trial Investigators. Endovascular versus open repair of abdominal aortic aneurysm. *N Engl J Med*. 2010 May 20;362(20):1863-71. [Abstract](#)
- Greenhalgh RM, Brown LC, Powell JT, et al; United Kingdom EVAR Trial Investigators. Endovascular repair of aortic aneurysm in patients physically ineligible for open repair. *N Engl J Med*. 2010 May 20;362(20):1872-80. [Abstract](#)
- United Kingdom EVAR Trial Investigators, Greenhalgh RM, Brown LC, et al. Endovascular repair of aortic aneurysm in patients physically ineligible for open repair. *N Engl J Med*. 2010;362:1872-1880. [Abstract](#)
- Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014 Dec 9;130(24):e278-333. [Full text](#) [Abstract](#)

References

1. Dehlin JM, Upchurch GR. Management of abdominal aortic aneurysms. *Curr Treat Options Cardiovasc Med*. 2005 Jun;7(2):119-30. [Abstract](#)

2. Johnston KW, Rutherford RB, Tilson MD, et al. Suggested standards for reporting on arterial aneurysms. *J Vasc Surg.* 1991 Mar;13(3):452-8. [Abstract](#)
3. LeFevre ML; US Preventive Services Task Force. Screening for abdominal aortic aneurysm: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2014 Aug 19;161(4):281-90. [Full text](#) [Abstract](#)
4. McConathy WJ, Alaupovic P, Woolcock N, et al. Lipids and apolipoprotein profiles in men with aneurysmal and stenosing aorto-iliac atherosclerosis. *Eur J Vasc Surg.* 1989 Dec;3(6):511-4. [Abstract](#)
5. Saratzis A, Bown MJ. The genetic basis for aortic aneurysmal disease. *Heart.* 2014 Jun;100(12):916-22. [Abstract](#)
6. Tang T, Boyle JR, Dixon AK, et al. Inflammatory abdominal aortic aneurysms. *Eur J Vasc Endovasc. Surg.* 2005 Apr;29(4):353-62. [Abstract](#)
7. Stone JR, Bruneval P, Angelini A, et al. Consensus statement on surgical pathology of the aorta from the Society for Cardiovascular Pathology and the Association for European Cardiovascular Pathology: I. Inflammatory diseases. *Cardiovasc Pathol.* 2015 Sep-Oct;24(5):267-78. [Full text](#) [Abstract](#)
8. Gomes MN, Choyke PL, Wallace RB. Infected aortic aneurysms: a changing entity. *Ann Surg.* 1992 May;215(5):435-42. [Full text](#) [Abstract](#)
9. Zankl AR, Schumacher H, Krumsdorf U, et al. Pathology, natural history and treatment of abdominal aortic aneurysms. *Clin Res Cardiol.* 2007 Mar;96(3):140-51. [Abstract](#)
10. Silverstein MD, Pitts SR, Chaikof EL, et al. Abdominal aortic aneurysm (AAA): cost-effectiveness of screening, surveillance of intermediate-sized AAA, and management of symptomatic AAA. *Proc (Bayl Univ Med Cent).* 2005 Oct;18(4):345-67. [Full text](#) [Abstract](#)
11. Singh K, Bønaa H, Jacobsen BK, et al. Prevalence of and risk factors for abdominal aortic aneurysms in a population-based study: The Tromsø Study. *Am J Epidemiol.* 2001 Aug 1;154(3):236-44. [Full text](#) [Abstract](#)
12. Jacomelli J, Summers L, Stevenson A, et al. Impact of the first 5 years of a national abdominal aortic aneurysm screening programme. *Br J Surg.* 2016 Aug;103(9):1125-31. [Abstract](#)
13. Ulug P, Powell JT, Sweeting MJ, et al; SWAN Collaborative Group. Meta-analysis of the current prevalence of screen-detected abdominal aortic aneurysm in women. *Br J Surg.* 2016 Aug;103(9):1097-104. [Full text](#) [Abstract](#)
14. Centers for Disease Control and Prevention, Division for Heart Disease and Stroke Prevention. Aortic aneurysm fact sheet. Jun 2016 [internet publication]. [Full text](#)
15. Anjum A, von Allmen R, Greenhalgh R, et al. Explaining the decrease in mortality from abdominal aortic aneurysm rupture. *Br J Surg.* 2012 May;99(5):637-45. [Abstract](#)

16. López-Candales A, Holmes DR, Liao S, et al. Decreased vascular smooth muscle cell density in medial degeneration of human abdominal aortic aneurysms. *Am J Pathol.* 1997 Mar;150(3):993-1007. [Full text](#) [Abstract](#)
17. Lederle FA, Johnson GR, Wilson SE, et al; Aneurysm Detection and Management (ADAM) Veterans Affairs Cooperative Study Group. Prevalence and associations of abdominal aortic aneurysm detected through screening. *Ann Intern Med.* 1997 Mar 15;126(6):441-9. [Abstract](#)
18. Wilmink TB, Quick CR, Day NE. The association between cigarette smoking and abdominal aortic aneurysms. *J Vasc Surg.* 1999 Dec;30(6):1099-105. [Abstract](#)
19. De Rango P, Farchioni L, Fiorucci B, et al. Diabetes and abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg.* 2014 Mar;47(3):243-61. [Full text](#) [Abstract](#)
20. Wassef M, Baxter BT, Chisholm RL, et al. Pathogenesis of abdominal aortic aneurysms: a multidisciplinary research program supported by the National Heart, Lung, and Blood Institute. *J Vasc Surg.* 2001 Oct;34(4):730-8. [Abstract](#)
21. Ailawadi G, Eliason JL, Upchurch GR. Current concepts in the pathogenesis of abdominal aortic aneurysm. *J Vasc Surg.* 2003 Sep;38(3):584-8. [Abstract](#)
22. Davies MJ. Aortic aneurysm formation: lessons from human studies and experimental models. *Circulation.* 1998 Jul 21;98(3):193-5. [Full text](#) [Abstract](#)
23. Grange JJ, Davis V, Baxter BT. Pathogenesis of abdominal aortic aneurysm: an update and look toward the future. *Cardiovasc Surg.* 1997 Jun;5(3):256-65. [Abstract](#)
24. Takagi H, Manabe H, Kawai N, et al. Circulating matrix metalloproteinase-9 concentrations and abdominal aortic aneurysm presence: a meta-analysis. *Interact Cardiovasc Thorac Surg.* 2009 Sep;9(3):437-40. [Full text](#) [Abstract](#)
25. Stocker R, Keaney JF Jr. Role of oxidative modifications in atherosclerosis. *Physiol Rev.* 2004 Oct;84(4):1381-478. [Full text](#) [Abstract](#)
26. Griending KK, FitzGerald GA. Oxidative stress and cardiovascular injury: Part I: basic mechanisms and in vivo monitoring of ROS. *Circulation.* 2003 Oct 21;108(16):1912-6. [Full text](#) [Abstract](#)
27. Griending KK, FitzGerald GA. Oxidative stress and cardiovascular injury: Part II: animal and human studies. *Circulation.* 2003 Oct 28;108(17):2034-40. [Full text](#) [Abstract](#)
28. McCormick ML, Gavrila D, Weintraub NL. Role of oxidative stress in the pathogenesis of abdominal aortic aneurysms. *Arterioscler Thromb Vasc Biol.* 2007 Mar;27(3):461-9. [Abstract](#)
29. Ailawadi G, Knipp BS, Lu G, et al. A nonintrinsic regional basis for increased infrarenal aortic MMP-9 expression and activity. *J Vasc Surg.* 2003 May;37(5):1059-66. [Abstract](#)
30. Moore JE, Ku DN, Zarins CK, et al. Pulsatile flow visualization in the abdominal aorta under differing conditions: implications for increased susceptibility to atherosclerosis. *J Biomech Eng.* 1992 Aug;114(3):391-7. [Abstract](#)

31. Wahlgren CM, Larsson E, Magnusson PK, et al. Genetic and environmental contributions to abdominal aortic aneurysm development in a twin population. *J Vasc Surg.* 2010 Jan;51(1):3-7. [Full text](#) [Abstract](#)
32. Jones GT, Tromp G, Kuivaniemi H, et al. Meta-analysis of genome-wide association studies for abdominal aortic aneurysm identifies four new disease-specific risk loci. *Circ Res.* 2017 Jan 20;120(2):341-53. [Full text](#) [Abstract](#)
33. Rasmussen TE, Hallett JW Jr, Tazelaar HD, et al. Human leukocyte antigen class II immune response genes, female gender, and cigarette smoking as risk and modulating factors in abdominal aortic aneurysms. *J Vasc Surg.* 2002 May;35(5):988-93. [Abstract](#)
34. Shibamura H, Olson JM, van Vlijmen-Van Keulen C, et al. Genome scan for familial abdominal aortic aneurysm using sex and family history as covariates suggests genetic heterogeneity and identifies linkage to chromosome 19q13. *Circulation.* 2004 May 4;109(17):2103-8. [Full text](#) [Abstract](#)
35. Kuivaniemi H, Shibamura H, Arthur C, et al. Familial abdominal aortic aneurysms: collection of 233 multiplex families. *J Vasc Surg.* 2003 Feb;37(2):340-5. [Abstract](#)
36. Majumder PP, St Jean PL, Ferrell RE, et al. On the inheritance of abdominal aortic aneurysm. *Am J Hum Genet.* 1991 Jan;48(1):164-70. [Full text](#) [Abstract](#)
37. Verloes A, Sakalihasan N, Koulischer L, et al. Aneurysms of the abdominal aorta: familial and genetic aspects in three hundred thirteen pedigrees. *J Vasc Surg.* 1995 Apr;21(4):646-55. [Full text](#) [Abstract](#)
38. Johansen K, Koepsell T. Familial tendency for abdominal aortic aneurysms. *JAMA.* 1986 Oct 10;256(14):1934-6. [Abstract](#)
39. Ogata T, MacKean GL, Cole CW, et al. The lifetime prevalence of abdominal aortic aneurysms among siblings of aneurysm patients is eightfold higher than among siblings of spouses: an analysis of 187 aneurysm families in Nova Scotia, Canada. *J Vasc Surg.* 2005 Nov;42(5):891-7. [Full text](#) [Abstract](#)
40. Darling RC 3rd, Brewster DC, Darling RC, et al. Are familial abdominal aortic aneurysms different? *J Vasc Surg.* 1989 Jul;10(1):39-43. [Abstract](#)
41. Webster MW, Ferrell RE, St Jean PL, et al. Ultrasound screening of first-degree relatives of patients with an abdominal aortic aneurysm. *J Vasc Surg.* 1991 Jan;13(1):9-13. [Abstract](#)
42. Larsson E, Granath F, Swedenborg J, et al. A population-based case-control study of the familial risk of abdominal aortic aneurysm. *J Vasc Surg.* 2009 Jan;49(1):47-50. [Full text](#) [Abstract](#)
43. Van der Vliet JA, Boll AP. Abdominal aortic aneurysm. *Lancet.* 1997 Mar 22;349(9055):863-6. [Abstract](#)
44. Golledge J, Clancy P, Jamrozik K, et al. Obesity, adipokines, and abdominal aortic aneurysm: Health in Men study. *Circulation.* 2007 Nov 13;116(20):2275-9. [Abstract](#)
45. Skibba AA, Evans JR, Hopkins SP, et al. Reconsidering gender relative to risk of rupture in the contemporary management of abdominal aortic aneurysms. *J Vasc Surg.* 2015 Dec;62(6):1429-36. [Full text](#) [Abstract](#)

46. Gokani VJ, Sidloff D, Bath MF, et al. A retrospective study: factors associated with the risk of abdominal aortic aneurysm rupture. *Vascul Pharmacol*. 2015 Feb-Mar;65-66:13-6. [Abstract](#)
47. Hollister DW, Godfrey M, Sakai LY, et al. Immunohistologic abnormalities of the microfibrillar-fiber system in the Marfan syndrome. *N Engl J Med*. 1990 Jul 19;323(3):152-9. [Abstract](#)
48. Dietz HC, Cutting GR, Pyeritz RE, et al. Marfan syndrome caused by a recurrent de novo missense mutation in the fibrillin gene. *Nature*. 1991 Jul 25;352(6333):337-9. [Abstract](#)
49. Jeremy RW, Huang H, Hwa J, et al. Relation between age, arterial distensibility, and aortic dilatation in the Marfan syndrome. *Am J Cardiol*. 1994 Aug 15;74(4):369-73. [Abstract](#)
50. Fowkes FG, Anandan CL, Lee AJ, et al. Reduced lung function in patients with abdominal aortic aneurysm is associated with activation of inflammation and hemostasis, not smoking or cardiovascular disease. *J Vasc Surg*. 2006 Mar;43(3):474-80. [Abstract](#)
51. Brown LC, Powell JT. Risk factors for aneurysm rupture in patients kept under ultrasound surveillance. *Ann Surg*. 1999 Sep;230(3):289-96. [Full text](#) [Abstract](#)
52. Takagi H, Umemoto T; ALICE (All-Literature Investigation of Cardiovascular Evidence) Group. A meta-analysis of the association of chronic obstructive pulmonary disease with abdominal aortic aneurysm presence. *Ann Vasc Surg*. 2016 Jul;34:84-94. [Abstract](#)
53. Upchurch GR, Schaub TA. Abdominal aortic aneurysm. *Am Fam Physician*. 2006 Apr 1;73(7):1198-204. [Full text](#) [Abstract](#)
54. Chang JB, Stein TA, Liu JP, et al. Risk factors associated with rapid growth of small abdominal aortic aneurysms. *Surgery*. 1997 Feb;121(2):117-22. [Abstract](#)
55. Englesbe MJ, Wu AH, Clowes AW, et al. The prevalence and natural history of aortic aneurysms in heart and abdominal organ transplant patients. *J Vasc Surg*. 2003 Jan;37(1):27-31. [Full text](#) [Abstract](#)
56. Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg*. 2018 Jan;67(1):2-77.e2. [Full text](#) [Abstract](#)
57. Fink HA, Lederle FA, Roth CS, et al. The accuracy of physical examination to detect abdominal aortic aneurysm. *Arch Intern Med*. 2000 Mar 27;160(6):833-6. [Full text](#) [Abstract](#)
58. Cosford PA, Leng GC. Screening for abdominal aortic aneurysm. *Cochrane Database Syst Rev*. 2007 Apr 18;(2):CD002945. [Full text](#) [Abstract](#)
59. Truijers M, Resch T, Van Den Berg JC, et al. Endovascular aneurysm repair: state-of-art imaging techniques for preoperative planning and surveillance. *J Cardiovasc Surg (Torino)*. 2009 Aug;50(4):423-38. [Abstract](#)
60. Fillinger MF, Marra SP, Raghavan ML, et al. Prediction of rupture risk in abdominal aortic aneurysm during observation: wall stress versus diameter. *J Vasc Surg*. 2003 Apr;37(4):724-32. [Abstract](#)

61. Lindholt JS, Vammen S, Juul S, et al. The validity of ultrasonographic scanning as screening method for abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg*. 1999 Jun;17(6):472-5. [Abstract](#)
62. Vowden P, Wilkinson D, Ausobsky JR, et al. A comparison of three imaging techniques in the assessment of an abdominal aortic aneurysm. *J Cardiovasc Surg (Torino)*. 1989 Nov-Dec;30(6):891-6. [Abstract](#)
63. Taylor SM, Mills JL, Fujitani RM. The juxtarenal abdominal aortic aneurysm: a more common problem than previously realized? *Arch Surg*. 1994 Jul;129(7):734-7. [Abstract](#)
64. Rakita D, Newatia A, Hines JJ, et al. Spectrum of CT findings in rupture and impending rupture of abdominal aortic aneurysms. *Radiographics*. 2007 Mar-Apr;27(2):497-507. [Abstract](#)
65. Jacobs DO. Clinical practice: diverticulitis. *N Engl J Med*. 2007 Nov 15;357(20):2057-66. [Abstract](#)
66. Scheinman SJ. Urinary calculi. *Medicine*. 2003 Jun 1;31(6):77-80. [Full text](#)
67. Spiller R. Clinical update: irritable bowel syndrome. *Lancet*. 2007 May 12;369(9573):1586-8. [Abstract](#)
68. Nikolaus S, Schreiber S. Diagnostics of inflammatory bowel disease. *Gastroenterology*. 2007 Nov;133(5):1670-89. [Abstract](#)
69. Oelsner G, Shashar D. Adnexal torsion. *Clin Obstet Gynecol*. 2006 Sep;49(3):459-63. [Abstract](#)
70. Stanley JC. Mesenteric arterial occlusive and aneurysmal disease. *Cardiol Clin*. 2002 Nov;20(4):611-22. [Abstract](#)
71. Wanhainen A, Hultgren R, Linné A, et al; Swedish Aneurysm Screening Study Group (SASS). Outcome of the Swedish Nationwide Abdominal Aortic Aneurysm Screening Program. *Circulation*. 2016 Oct 18;134(16):1141-8. [Abstract](#)
72. Public Health England. Population screening programmes: NHS abdominal aortic aneurysm (AAA) programme. Jul 2015 [internet publication]. [Full text](#)
73. Public Health England. Abdominal aortic aneurysm screening: care pathway. Nov 2015 [internet publication]. [Full text](#)
74. Moll FL, Powell JT, Fraedrich G, et al; European Society for Vascular Surgery. Management of abdominal aortic aneurysms: clinical practice guidelines of the European Society for Vascular Surgery. *Eur J Vasc Endovasc Surg*. 2011 Jan;41 Suppl 1:S1-58. [Full text](#) [Abstract](#)
75. Thompson S, Brown L, Sweeting M, et al; RESCAN Collaborators. Systematic review and meta-analysis of the growth and rupture rates of small abdominal aortic aneurysms: implications for surveillance intervals and their cost-effectiveness. *Health Technol Assess*. 2013 Sep;17(41):1-118. [Full text](#) [Abstract](#)
76. Anderson JL, Halperin JL, Albert NM, et al. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA guideline recommendations): a report of the American

- College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. Circulation. 2013 Apr 2;127(13):1425-43. [Full text](#) [Abstract](#)
77. Lederle FA, Wilson SE, Johnson GR, et al; Aneurysm Detection and Management Veterans Affairs Cooperative Study Group. Immediate repair compared with surveillance of small abdominal aortic aneurysms. N Engl J Med. 2002 May 9;346(19):1437-44. [Abstract](#)
 78. UK Small Aneurysm Trial Participants. Mortality results for randomized controlled trial of early elective surgery or ultrasonographic surveillance for small abdominal aortic aneurysms. Lancet. 1998 Nov 21;352(9141):1649-55. [Abstract](#)
 79. Powell JT, Brady AR, Brown LC, et al; United Kingdom Small Aneurysm Trial Participants. Long-term outcomes of immediate repair compared with surveillance of small abdominal aortic aneurysms. N Engl J Med. 2002 May 9;346(19):1445-52. [Full text](#) [Abstract](#)
 80. Powell JT, Brown LC, Forbes JF, et al. Final 12-year follow-up of surgery versus surveillance in the UK Small Aneurysm Trial. Br J Surg. 2007 Jun;94(6):702-8. [Abstract](#)
 81. Brewster DC, Cronenwett JL, Hallett JW Jr, et al. Guidelines for the treatment of abdominal aortic aneurysms: report of a subcommittee of the Joint Council of the American Association for Vascular Surgery and Society for Vascular Surgery. J Vasc Surg. 2003 May;37(5):1106-17. [Full text](#) [Abstract](#)
 82. Harkin DW, Dillon M, Blair PH, et al. Endovascular ruptured abdominal aortic aneurysm repair (EVRAR): a systematic review. Eur J Vasc Endovasc Surg. 2007 Dec;34(6):673-81. [Abstract](#)
 83. IMPROVE Trial Investigators. Comparative clinical effectiveness and cost effectiveness of endovascular strategy v open repair for ruptured abdominal aortic aneurysm: three year results of the IMPROVE randomised trial. BMJ. 2017 Nov 14;359:j4859. [Full text](#) [Abstract](#)
 84. Sweeting MJ, Balm R, Desgranges P, et al; Ruptured Aneurysm Trialists. Individual-patient meta-analysis of three randomized trials comparing endovascular versus open repair for ruptured abdominal aortic aneurysm. Br J Surg. 2015 Sep;102(10):1229-39. [Full text](#) [Abstract](#)
 85. Sweeting MJ, Ulug P, Powell JT, et al; Ruptured Aneurysm Trialists. Ruptured aneurysm trials: the importance of longer-term outcomes and meta-analysis for 1-year mortality. Eur J Vasc Endovasc Surg. 2015 Sep;50(3):297-302. [Full text](#) [Abstract](#)
 86. Soar J, Nolan JP, Böttiger BW, et al; Adult advanced life support section collaborators. European Resuscitation Council Guidelines for resuscitation 2015: section 3. Adult advanced life support. Resuscitation. 2015;95:100-147.
 87. Colquhoun MC, Handley AJ, Evans TR, eds. ABC of resuscitation. 5th ed. Wiley-Blackwell; 2003.
 88. Soar J, Nolan JP, Böttiger BW, et al. European Resuscitation Council guidelines for resuscitation 2015: Section 3. Adult advanced life support. Resuscitation. 2015;95:100-147.
 89. Crawford ES. Ruptured aortic aneurysm. J Vasc Surg. 1991 Feb;13(2):348-50. [Abstract](#)

90. Roberts K, Revell M, Youssef H, et al. Hypotensive resuscitation in patients with ruptured abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg*. 2006 Apr;31(4):339-44. [Abstract](#)
91. Hardman DT, Fisher CM, Patel MI, et al. Ruptured abdominal aortic aneurysms: who should be offered surgery. *J Vasc Surg*. 1996 Jan;23(1):123-9. [Abstract](#)
92. Ohki T, Veith FJ. Endovascular grafts and other image-guided catheter-based adjuncts to improve the treatment of ruptured aortoiliac aneurysms. *Ann Surg*. 2000 Oct;232(4):466-79. [Abstract](#)
93. Powell JT, Hinchcliffe RJ, Thompson MM, et al; IMPROVE Trial Investigators. Observations from the IMPROVE trial concerning the clinical care of patients with ruptured abdominal aortic aneurysm. *Br J Surg*. 2014 Feb;101(3):216-24. [Full text](#) [Abstract](#)
94. Erbel R, Aboyans V, Boileau C, et al; ESC Committee for Practice Guidelines. 2014 ESC guidelines on the diagnosis and treatment of aortic diseases: document covering acute and chronic aortic diseases of the thoracic and abdominal aorta of the adult. *Eur Heart J*. 2014 Nov 1;35(41):2873-926. [Full text](#) [Abstract](#)
95. De Martino RR, Nolan BW, Goodney PP, Chang CK, et al; Vascular Study Group of Northern New England. Outcomes of symptomatic abdominal aortic aneurysm repair. *J Vasc Surg*. 2010 Jul;52(1):5-12.e1. [Full text](#) [Abstract](#)
96. Chandra V, Trang K, Virgin-Downey W, et al. Management and outcomes of symptomatic abdominal aortic aneurysms during the past 20 years. *J Vasc Surg*. 2017 Dec;66(6):1679-85. [Abstract](#)
97. Ten Bosch JA, Willigendael EM, Kruidenier LM, et al. Early and mid-term results of a prospective observational study comparing emergency endovascular aneurysm repair with open surgery in both ruptured and unruptured acute abdominal aortic aneurysms. *Vascular*. 2012 Apr;20(2):72-80. [Abstract](#)
98. Soden PA, Zettervall SL, Ultee KH, et al. Outcomes for symptomatic abdominal aortic aneurysms in the American College of Surgeons National Surgical Quality Improvement Program. *J Vasc Surg*. 2016 Aug;64(2):297-305. [Full text](#) [Abstract](#)
99. Filardo G, Powell JT, Martinez MA, et al. Surgery for small asymptomatic abdominal aortic aneurysms. *Cochrane Database Syst Rev*. 2015 Feb 8;(2):CD001835. [Full text](#) [Abstract](#)
100. NHS England. NHS public health functions agreement 2016-17: service specification No. 23. NHS abdominal aortic aneurysm screening programme. Apr 2016 [internet publication]. [Full text](#)
101. Stead LF, Perera R, Bullen C, et al. Nicotine replacement therapy for smoking cessation. *Cochrane Database Syst Rev*. 2012 Nov 14;(11):CD000146. [Full text](#) [Abstract](#)
102. Rigotti NA, Clair C, Munafò MR, et al. Interventions for smoking cessation in hospitalised patients. *Cochrane Database Syst Rev*. 2012 May 16;(5):CD001837. [Full text](#) [Abstract](#)
103. Hughes JR, Stead LF, Hartmann-Boyce J, et al. Antidepressants for smoking cessation. *Cochrane Database Syst Rev*. 2014 Jan 8;(1):CD000031. [Full text](#) [Abstract](#)

104. Rughani G, Robertson L, Clarke M. Medical treatment for small abdominal aortic aneurysms. Cochrane Database Syst Rev. 2012 Sep 12;(9):CD009536. [Full text](#) [Abstract](#)
105. Lederle FA, Wilson SE, Johnson GR, et al. Immediate repair compared with surveillance of small abdominal aortic aneurysms. N Engl J Med. 2002;346:1437-1444. [Abstract](#)
106. The UK Small Aneurysm Trial Participants. Mortality results for randomized controlled trial of early elective surgery or ultrasonographic surveillance for small abdominal aortic aneurysms. Lancet. 1998;352:1649-1655. [Abstract](#)
107. United Kingdom Small Aneurysm Trial Participants. Long-term outcomes of immediate repair compared with surveillance of small abdominal aortic aneurysms. N Engl J Med. 2002;346:1445-1452. [Full text](#) [Abstract](#)
108. Greenhalgh RM, Brown LC, Powell JT, et al; United Kingdom EVAR Trial Investigators. Endovascular versus open repair of abdominal aortic aneurysm. N Engl J Med. 2010 May 20;362(20):1863-71. [Abstract](#)
109. Greenhalgh RM, Brown LC, Powell JT, et al; United Kingdom EVAR Trial Investigators. Endovascular repair of aortic aneurysm in patients physically ineligible for open repair. N Engl J Med. 2010 May 20;362(20):1872-80. [Abstract](#)
110. United Kingdom EVAR Trial Investigators, Greenhalgh RM, Brown LC, et al. Endovascular repair of aortic aneurysm in patients physically ineligible for open repair. N Engl J Med. 2010;362:1872-1880. [Abstract](#)
111. Khashram M, Hider PN, Williman JA, et al. Does the diameter of abdominal aortic aneurysm influence late survival following abdominal aortic aneurysm repair? A systematic review and meta-analysis. Vascular. 2016 Dec;24(6):658-67. [Abstract](#)
112. Gimzewska M, Jackson AI, Yeoh SE, et al. Totally percutaneous versus surgical cut-down femoral artery access for elective bifurcated abdominal endovascular aneurysm repair. Cochrane Database Syst Rev. 2017 Feb 21;(2):CD010185. [Full text](#) [Abstract](#)
113. Eliason JL, Upchurch GR. Endovascular abdominal aortic aneurysms repair. Circulation. 2008 Apr 1;117(13):1738-44. [Full text](#) [Abstract](#)
114. Cambria RP, Brewster DC, Abbott WM, et al. Transperitoneal versus retroperitoneal approach for aortic reconstruction: a randomized prospective study. J Vasc Surg. 1990 Feb;11(2):314-24. [Abstract](#)
115. Sicard GA, Reilly JM, Rubin BG, et al. Transabdominal versus retroperitoneal incision for abdominal aortic surgery: report of a prospective randomized trial. J Vasc Surg. 1995 Feb;21(2):174-81. [Abstract](#)
116. Senekowitsch C, Assadian A, Assadian O, et al. Replanting the inferior mesentery artery during infrarenal aortic aneurysm repair: influence on postoperative colon ischemia. J Vasc Surg. 2006 Apr;43(4):689-94. [Abstract](#)

117. Beck AW, Sedrakyan A, Mao J, et al; International Consortium of Vascular Registries. Variations in abdominal aortic aneurysm care: a report from the International Consortium of Vascular Registries. *Circulation*. 2016 Dec 13;134(24):1948-58. [Full text](#) [Abstract](#)
118. Royal College of Surgeons and Vascular Society of Great Britain and Ireland. National Vascular Registry: 2016 annual report. Nov 2016 [internet publication]. [Full text](#)
119. Hirsch AT, Haskal ZJ, Hertzner NR, et al. ACC/AHA 2005 practice guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic). *Circulation*. 2006 Mar 21;113(11):e463-654. [Full text](#) [Abstract](#)
120. Kristensen SD, Knuuti J, Saraste A, et al; Authors/Task Force Members. 2014 ESC/ESA guidelines on non-cardiac surgery: cardiovascular assessment and management. *Eur Heart J*. 2014 Sep 14;35(35):2383-431. [Full text](#) [Abstract](#)
121. Siracuse JJ, Gill HL, Graham AR, et al. Comparative safety of endovascular and open surgical repair of abdominal aortic aneurysms in low-risk male patients. *J Vasc Surg*. 2014 Nov;60(5):1154-8. [Full text](#) [Abstract](#)
122. Kontopodis N, Antoniou SA, Georgakarakos E, et al. Endovascular vs open aneurysm repair in the young: systematic review and meta-analysis. *J Endovasc Ther*. 2015 Dec;22(6):897-904. [Abstract](#)
123. Joels CS, Langan EM 3rd, Daley CA, et al. Changing indications and outcomes for open abdominal aortic aneurysm repair since the advent of endovascular repair. *Am Surg*. 2009 Aug;75(8):665-9. [Abstract](#)
124. Powell JT, Sweeting MJ, Ulug P, et al; EVAR, DREAM, OVER and ACE Trialists. Meta-analysis of individual-patient data from EVAR-1, DREAM, OVER and ACE trials comparing outcomes of endovascular or open repair for abdominal aortic aneurysm over 5 years. *Br J Surg*. 2017 Feb;104(3):166-78. [Full text](#) [Abstract](#)
125. Paravastu SC, Jayarajasingam R, Cottam R, et al. Endovascular repair of abdominal aortic aneurysm. *Cochrane Database Syst Rev*. 2014 Jan 23;(1):CD004178. [Full text](#) [Abstract](#)
126. Patel R, Sweeting MJ, Powell JT, et al; EVAR Trial Investigators. Endovascular versus open repair of abdominal aortic aneurysm in 15-years' follow-up of the UK endovascular aneurysm repair trial 1 (EVAR trial 1): a randomised controlled trial. *Lancet*. 2016 Nov 12;388(10058):2366-74. [Full text](#) [Abstract](#)
127. Schermerhorn ML, Bensley RP, Giles KA, et al. Changes in abdominal aortic aneurysm rupture and short-term mortality, 1995-2008: a retrospective observational study. *Ann Surg*. 2012 Oct;256(4):651-8. [Full text](#) [Abstract](#)
128. Desai M, Choke E, Sayers RD, et al. Sex-related trends in mortality after elective abdominal aortic aneurysm surgery between 2002 and 2013 at National Health Service hospitals in England: less benefit for women compared with men. *Eur Heart J*. 2016 Dec 7;37(46):3452-60. [Full text](#) [Abstract](#)
129. Schermerhorn ML, Buck DB, O'Malley AJ, et al. Long-term outcomes of abdominal aortic aneurysm in the Medicare population. *N Engl J Med*. 2015 Jul 23;373(4):328-38. [Full text](#) [Abstract](#)

130. Khashram M, Williman JA, Hider PN, et al. Systematic review and meta-analysis of factors influencing survival following abdominal aortic aneurysm repair. *Eur J Vasc Endovasc Surg.* 2016 Feb;51(2):203-15. [Full text](#) [Abstract](#)
131. Ma B, Wang YN, Chen KY, et al. Transperitoneal versus retroperitoneal approach for elective open abdominal aortic aneurysm repair. *Cochrane Database Syst Rev.* 2016 Feb 5;(2):CD010373. [Full text](#) [Abstract](#)
132. Kouvelos GN, Oikonomou K, Antoniou GA, et al. A systematic review of proximal neck dilatation after endovascular repair for abdominal aortic aneurysm. *J Endovasc Ther.* 2017 Feb;24(1):59-67. [Abstract](#)
133. Farahmand P, Becquemin JP, Desgranges P, et al. Is hypogastric artery embolization during endovascular aortoiliac aneurysm repair (EVAR) innocuous and useful? *Eur J Vasc Endovasc Surg.* 2008 Apr;35(4):429-35. [Abstract](#)
134. Mehta M, Veith FJ, Ohki T, et al. Unilateral and bilateral hypogastric artery interruption during aortoiliac aneurysm repair in 154 patients: a relatively innocuous procedure. *J Vasc Surg.* 2001 Feb;33(2 Suppl):S27-32. [Abstract](#)
135. Kouvelos GN, Katsargyris A, Antoniou GA, et al. Outcome after interruption or preservation of internal iliac artery flow during endovascular repair of abdominal aorto-iliac aneurysms. *Eur J Vasc Endovasc Surg.* 2016 Nov;52(5):621-34. [Abstract](#)
136. Kumar R, Dattani N, Asaad O, et al. Meta-analysis of outcomes following aneurysm repair in patients with synchronous intra-abdominal malignancy. *Eur J Vasc Endovasc Surg.* 2016 Dec;52(6):747-56. [Abstract](#)
137. Kouvelos GN, Patelis N, Antoniou GA, et al. Management of concomitant abdominal aortic aneurysm and colorectal cancer. *J Vasc Surg.* 2016 May;63(5):1384-93. [Abstract](#)
138. Johnston KW. Multicenter prospective study of nonruptured abdominal aortic aneurysm. Part II. Variables predicting morbidity and mortality. *J Vasc Surg.* 1989 Mar;9(3):437-47. [Abstract](#)
139. Saratzis A, Sidloff D, Bown MJ. Cardiovascular risk in patients with small abdominal aortic aneurysms. Spring meeting for clinician scientists in training 2017: poster abstract. *Lancet.* 2017 Feb 23;389(S89). [Full text](#)
140. Bani-Hani MG, AL Khaffaf H, Titi MA, et al. Interventions for preventing venous thromboembolism following abdominal aortic surgery. *Cochrane Database Syst Rev.* 2008 Jan 23;(1):CD005509. [Full text](#) [Abstract](#)
141. Samoilă G, Ford RT, Glasbey JC, et al. The significance of hypothermia in abdominal aortic aneurysm repair. *Ann Vasc Surg.* 2017 Jan;38:323-31. [Abstract](#)
142. Khashram M, Williman JA, Hider PN, et al. Management of modifiable vascular risk factors improves late survival following abdominal aortic aneurysm repair: a systematic review and meta-analysis. *Ann Vasc Surg.* 2017 Feb;39:301-11. [Abstract](#)

143. Barakat HM, Shahin Y, Khan JA, et al. Preoperative supervised exercise improves outcomes after elective abdominal aortic aneurysm repair: a randomized controlled trial. *Ann Surg*. 2016 Jul;264(1):47-53. [Abstract](#)
144. Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014 Dec 9;130(24):e278-333. [Full text](#) [Abstract](#)
145. Chen RJ, Chu H, Tsai LW. Impact of beta-blocker initiation timing on mortality risk in patients with diabetes mellitus undergoing noncardiac surgery: a nationwide population-based cohort study. *J Am Heart Assoc*. 2017 Jan 10;6(1):e004392. [Full text](#) [Abstract](#)
146. Brady AR, Gibbs JS, Greenhalgh RM, et al; POBBLE Trial Investigators. Perioperative beta-blockade (POBBLE) for patients undergoing infrarenal vascular surgery: results of a randomized double-blind controlled trial. *J Vasc Surg*. 2005 Apr;41(4):602-9. [Full text](#) [Abstract](#)
147. Devereaux PJ, Yang H, Yusuf S, et al; POISE Study Group. Effects of extended-release metoprolol succinate in patients undergoing non-cardiac surgery (POISE trial): a randomised controlled trial. *Lancet*. 2008 May 31;371(9627):1839-47. [Full text](#) [Abstract](#)
148. Yang H, Raymer K, Butler R, et al. The effects of perioperative beta-blockade: results of the Metoprolol after Vascular Surgery (MaVS) study, a randomized controlled trial. *Am Heart J*. 2006 Nov;152(5):983-90. [Abstract](#)
149. Devereaux PJ, Mrkobrada M, Sessler DI, et al. Aspirin in patients undergoing noncardiac surgery. *N Engl J Med*. 2014 Apr 17;370(16):1494-503. [Full text](#) [Abstract](#)
150. Schurink GW, Aarts NJ, vanBockel JH. Endoleak after stent-graft treatment of abdominal aortic aneurysm: a meta-analysis of clinical studies. *Br J Surg*. 1999 May;86(5):581-7. [Abstract](#)
151. Veith FJ, Baum RA, Ohki T, et al. Nature and significance of endoleaks and endotension: summary of opinions expressed at an international conference. *J Vasc Surg*. 2002 May;35(5):1029-35. [Abstract](#)
152. Higashiura W, Greenberg RK, Katz E, et al. Predictive factors, morphologic effects, and proposed treatment paradigm for type II endoleaks after repair of infrarenal abdominal aortic aneurysms. *J Vasc Interv Radiol*. 2007 Aug;18(8):975-81. [Abstract](#)
153. Mansueto G, Cenzi D, Scuro A, et al. Treatment of type II endoleak with a transcatheter transcaval approach: results at 1-year follow-up. *J Vasc Surg*. 2007 Jun;45(6):1120-7. [Abstract](#)
154. Baum RA, Stavropoulos SW, Fairman RM, et al. Endoleaks after endovascular repair of abdominal aortic aneurysms. *J Vasc Interv Radiol*. 2003 Sep;14(9 Pt 1):1111-7. [Abstract](#)
155. Van Marrewijk CJ, Fransen G, Laheij RJ, et al. Is a type II endoleak after EVAR a harbinger of risk? Causes and outcome of open conversion and aneurysm rupture during follow-up. *Eur J Vasc Endovasc Surg*. 2004 Feb;27(2):128-37. [Abstract](#)

156. Harris PL, Vallabhaneni SR, Desgranges P, et al. Incidence and risk factors of late rupture, conversion, and death after endovascular repair of infrarenal aortic aneurysms: the EUROSTAR experience. *J Vasc Surg.* 2000 Oct;32(4):739-49. [Abstract](#)
157. Steinmetz E, Rubin BG, Sanchez LA, et al. Type II endoleak after endovascular abdominal aortic aneurysm repair: a conservative approach with selective intervention is safe and cost-effective. *J Vasc Surg.* 2004 Feb;39(2):306-13. [Abstract](#)
158. Baum RA, Cope C, Fairman, et al. Translumbal embolization of type 2 endoleaks after endovascular repair of abdominal aortic aneurysms. *J Vasc Interv Radiol.* 2001 Jan;12(1):111-6. [Abstract](#)
159. Baum RA, Carpenter JP, Golden MA, et al. Treatment of type 2 endoleaks after endovascular repair of abdominal aortic aneurysms: comparison of transarterial and translumbal techniques. *J Vasc Surg.* 2002 Jan;35(1):23-9. [Abstract](#)
160. Schmid R, Gurke L, Aschwanden M, et al. CT-guided percutaneous embolization of a lumbar artery maintaining a type II endoleak. *J Endovasc Ther.* 2002 Apr;9(2):198-202. [Abstract](#)
161. Kusminsky RE. Complications of central venous catheterization. *J Am Coll Surg.* 2007;204:681-696.
162. McGee DC, Gould MK. Preventing complications of central venous catheterization. *N Engl J Med.* 2003;348:1123-1133. [Full text](#)
163. Smith RN, Nolan JP. Central venous catheters. *BMJ.* 2013;347:f6570.
164. Reich DL. Monitoring in anesthesia and perioperative care. Cambridge: Cambridge University Press; 2011.
165. Abbott Northwestern Hospital Internal Medicine Residency. Internal jugular central venous line. 2015. <http://www.anwresidency.com> (last accessed 27 October 2017). [Full text](#)
166. Bishop L, Dougherty L, Bodenham A, et al. Guidelines on the insertion and management of central venous access devices in adults. *Int J Lab Hematol.* 2007;29:261-278.
167. Fletcher SJ, Bodenham AR. Safe placement of central venous catheters: where should the tip of the catheter lie? *Br J Anaesth.* 2000;85:188-191.
168. Gibson F, Bodenham A. Misplaced central venous catheters: applied anatomy and practical management. *Br J Anaesth.* 2013;110:333-346. [Full text](#)
169. Schuster M, Nave H, Piepenbrock S, Pabst R, Panning B. The carina as a landmark in central venous catheter placement. *Br J Anaesth.* 2000;85:192-194.
170. Baxter BT, Pearce WH, Waltke EA, et al. Prolonged administration of doxycycline in patients with small asymptomatic abdominal aortic aneurysms: report of a prospective (Phase II) multicenter study. *J Vasc Surg.* 2002 Jul;36(1):1-12. [Abstract](#)
171. Rentschler M, Baxter BT. Pharmacological approaches to prevent abdominal aortic aneurysm enlargement and rupture. *Ann N Y Acad Sci.* 2006 Nov;1085:39-46. [Abstract](#)

172. Mosorin M, Juvonen J, Biancari F, et al. Use of doxycycline to decrease the growth rate of abdominal aortic aneurysms: a randomized, double-blind, placebo-controlled pilot study. *J Vasc Surg.* 2001 Oct;34(4):606-10. [Abstract](#)
173. ClinicalTrials.gov. Non-invasive treatment of abdominal aortic aneurysm clinical trial (N-TA³CT). NCT01756833. Dec 2017 [internet publication]. [Full text](#)
174. Dodd BR, Spence RA. Doxycycline inhibition of abdominal aortic aneurysm growth: a systematic review of the literature. *Curr Vasc Pharmacol.* 2011 Jul 1;9(4):471-8. [Abstract](#)
175. ClinicalTrials.gov. The efficacy of ticagrelor on abdominal aortic aneurysm (AAA) expansion (TicAAA). NCT02070653. Oct 2016 [internet publication]. [Full text](#)
176. National Institute for Health and Care Excellence. Endovascular aneurysm sealing for abdominal aortic aneurysm. Feb 2016 [internet publication]. [Full text](#)
177. Carpenter JP, Cuff R, Buckley C, et al. One-year pivotal trial outcomes of the Nellix system for endovascular aneurysm sealing. *J Vasc Surg.* 2017 Feb;65(2):330-6.e4. [Abstract](#)
178. Thompson MM, Heyligers JM, Hayes PD, et al; EVAS FORWARD Global Registry Investigators. Endovascular aneurysm sealing: early and midterm results from the EVAS FORWARD Global Registry. *J Endovasc Ther.* 2016 Oct;23(5):685-92. [Abstract](#)
179. England A, Torella F, Fisher RK, et al. Migration of the Nellix endoprosthesis. *J Vasc Surg.* 2016 Aug;64(2):306-12. [Abstract](#)
180. Cross J, Gurusamy K, Gadhvi V, et al. Fenestrated endovascular aneurysm repair. *Br J Surg.* 2012 Feb;99(2):152-9. [Abstract](#)
181. O'Neill S, Greenberg RK, Haddad F, et al. A prospective analysis of fenestrated endovascular grafting: intermediate-term outcomes. *Eur J Vasc Endovasc Surg.* 2006 Aug;32(2):115-23. [Abstract](#)
182. Hertzner NR, Mascha EJ, Karafa MT, et al. Open infrarenal abdominal aortic aneurysm repair: the Cleveland Clinic experience from 1989 to 1998. *J Vasc Surg.* 2002 Jun;35(6):1145-54. [Abstract](#)
183. Rubenstein C, Bietz G, Davenport DL, et al. Abdominal compartment syndrome associated with endovascular and open repair of ruptured abdominal aortic aneurysms. *J Vasc Surg.* 2015 Mar;61(3):648-54. [Full text](#) [Abstract](#)
184. Ersryd S, Djavani-Gidlund K, Wanhainen A, et al. Editor's choice - abdominal compartment syndrome after surgery for abdominal aortic aneurysm: a nationwide population based study. *Eur J Vasc Endovasc Surg.* 2016 Aug;52(2):158-65. [Full text](#) [Abstract](#)
185. Johnston KW, Scobie TK. Multicenter prospective study of nonruptured abdominal aortic aneurysms. I. Population and operative management. *J Vasc Surg.* 1988 Jan;7(1):69-81. [Abstract](#)
186. Saratzis A, Melas N, Mahmood A, et al. Incidence of acute kidney injury (AKI) after endovascular abdominal aortic aneurysm repair (EVAR) and impact on outcome. *Eur J Vasc Endovasc Surg.* 2015 May;49(5):534-40. [Full text](#) [Abstract](#)

187. Karthikesalingam A, Bahia SS, Patel SR, et al. A systematic review and meta-analysis indicates underreporting of renal dysfunction following endovascular aneurysm repair. *Kidney Int.* 2015 Feb;87(2):442-51. [Abstract](#)
188. Dariane C, Coscas R, Boulitrop C, et al. Acute kidney injury after open repair of intact abdominal aortic aneurysms. *Ann Vasc Surg.* 2017 Feb;39:294-300. [Abstract](#)
189. Saratzis A, Bath MF, Harrison S, et al. Long-term renal function after endovascular aneurysm repair. *Clin J Am Soc Nephrol.* 2015 Nov 6;10(11):1930-6. [Full text](#) [Abstract](#)
190. Berg P, Kaufmann D, van Marrewijk CJ, et al. Spinal cord ischaemia after stent-graft treatment for infra-renal abdominal aortic aneurysms. Analysis of the Eurostar database. *Eur J Vasc Endovasc Surg.* 2001 Oct;22(4):342-7. [Full text](#) [Abstract](#)
191. Peppelenbosch N, Cuypers PW, Vahl AC, et al. Emergency endovascular treatment for ruptured abdominal aortic aneurysm and the risk of spinal cord ischemia. *J Vasc Surg.* 2005 Oct;42(4):608-14. [Full text](#) [Abstract](#)
192. Kwok PC, Chung TK, Chong LC, et al. Neurologic injury after endovascular stent-graft and bilateral internal iliac artery embolization for infrarenal abdominal aortic aneurysm. *J Vasc Interv Radiol.* 2001 Jun;12(6):761-3. [Abstract](#)
193. Lederle FA, Johnson GR, Wilson SE, et al; Aneurysm Detection and Management Veterans Affairs Cooperative Study. Quality of life, impotence, and activity level in a randomized trial of immediate repair versus surveillance of small abdominal aortic aneurysm. *J Vasc Surg.* 2003 Oct;38(4):745-52. [Full text](#) [Abstract](#)
194. Edwards JM, Teefey SA, Zierler RE, et al. Intraabdominal paraanastomotic aneurysms after aortic bypass grafting. *J Vasc Surg.* 1992 Feb;15(2):344-50. [Abstract](#)
195. Ylönen K, Biancari F, Leo E, et al. Predictors of development of anastomotic femoral pseudoaneurysms after aortobifemoral reconstruction for abdominal aortic aneurysm. *Am J Surg.* 2004 Jan;187(1):83-7. [Abstract](#)
196. Vogel TR, Symons R, Flum DR. The incidence and factors associated with graft infection after aortic aneurysm repair. *J Vasc Surg.* 2008 Feb;47(2):264-9. [Full text](#) [Abstract](#)
197. Hodgson KJ, Webster DJ. Abdominal aortic aneurysm causing duodenal and ureteric obstruction. *J Vasc Surg.* 1986 Feb;3(2):364-8. [Abstract](#)
198. Kashyap VS, Fang R, Fitzpatrick CM, et al. Caval and ureteral obstruction secondary to an inflammatory abdominal aortic aneurysm. *J Vasc Surg.* 2003 Dec;38(6):1416-21. [Abstract](#)
199. Conrad MF, Crawford RS, Pedraza JD, et al. Long-term durability of open abdominal aortic aneurysm repair. *J Vasc Surg.* 2007 Oct;46(4):669-75. [Full text](#) [Abstract](#)
200. Hallett JW Jr, Marshall DM, Petterson TM, et al. Graft-related complications after abdominal aortic aneurysm repair: reassurance from a 36-year population-based experience. *J Vasc Surg.* 1997 Feb;25(2):277-84. [Abstract](#)

201. Cochenne F, Becquemin JP, Desgranges P, et al. Limb graft occlusion following EVAR: clinical pattern, outcomes and predictive factors of occurrence. *Eur J Vasc Endovasc Surg*. 2007 Jul;34(1):59-65. [Full text](#) [Abstract](#)

202. Faure EM, Becquemin JP, Cochenne F; ENGAGE Collaborators. Predictive factors for limb occlusions after endovascular aneurysm repair. *J Vasc Surg*. 2015 May;61(5):1138-45.e2. [Full text](#) [Abstract](#)

203. Baxter BT, McGee GS, Flinn WR, et al. Distal embolization as a presenting symptom of aortic aneurysms. *Am J Surg*. 1990 Aug;160(2):197-201. [Abstract](#)

Images




Figure 1: Ultrasound of a 3.8 cm x 4.2 cm AAA

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology

Aortic diameter (anterior-posterior, inner to inner)	Re-scanning interval
<3 cm	Discharged from programme
3-4.4 cm	Annual surveillance programme
4.5-5.4 cm	3-monthly surveillance programme
>5.5 cm	Referral to vascular surgeon*

Figure 2: Re-scanning intervals associated with aortic diameters

NHS population screening: care pathways, Public Health England




Figure 3: CT scan of a ruptured AAA

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology




Figure 4: Various endovascular stent grafts used for endovascular aneurysm repair (EVAR)

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology




Figure 5: Endovascular aneurysm repair (EVAR)

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology




Figure 6: Type I endoleak at the distal left iliac anastomosis (leak encircled)

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology



Figure 7: Extension stent graft deployed for the same type I endoleak (encircled)

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology



Figure 8: Resolution of the type I endoleak resolved after extension deployed

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology



Figure 9: Type II endoleak (encircled) discovered on follow-up CT

University of Michigan, specifically the cases of Dr Upchurch reflecting the Departments of Vascular Surgery and Radiology

Disclaimer

This content is meant for medical professionals situated outside of the United States and Canada. The BMJ Publishing Group Ltd ("BMJ Group") tries to ensure that the information provided is accurate and up-to-date, but we do not warrant that it is nor do our licensors who supply certain content linked to or otherwise accessible from our content. The BMJ Group does not advocate or endorse the use of any drug or therapy contained within nor does it diagnose patients. Medical professionals should use their own professional judgement in using this information and caring for their patients and the information herein should not be considered a substitute for that.

This information is not intended to cover all possible diagnosis methods, treatments, follow up, drugs and any contraindications or side effects. In addition such standards and practices in medicine change as new data become available, and you should consult a variety of sources. We strongly recommend that users independently verify specified diagnosis, treatments and follow up and ensure it is appropriate for your patient within your region. In addition, with respect to prescription medication, you are advised to check the product information sheet accompanying each drug to verify conditions of use and identify any changes in dosage schedule or contraindications, particularly if the agent to be administered is new, infrequently used, or has a narrow therapeutic range. You must always check that drugs referenced are licensed for the specified use and at the specified doses in your region. This information is provided on an "as is" basis and to the fullest extent permitted by law the BMJ Group and its licensors assume no responsibility for any aspect of healthcare administered with the aid of this information or any other use of this information.

View our full [Website Terms and Conditions](#).

BMJ Best Practice

Contributors:

// Authors:

Matt J. Bown, MB BCH, MD, FRCS, PGCert (Bioinformatics)

Professor of Vascular Surgery

Department of Cardiovascular Sciences, University of Leicester, Leicester Royal Infirmary, Leicester, UK

DISCLOSURES: MJB has grant funding from the British Heart Foundation and the National Institute for Health Research.

Andrew Duncan, MBBS, BSc, MRCS

Honorary Research Fellow

Department of Cardiovascular Sciences, University of Leicester, Leicester, UK

DISCLOSURES: AD declares that he has no competing interests.

// Acknowledgements:

Professor Matt Bown and Mr Andrew Duncan would like to gratefully acknowledge Dr Maureen K. Sheehan, Dr Dawn M. Barnes, and Dr Gilbert R. Upchurch, previous contributors to this monograph.

DISCLOSURES: MKS, DMB, and GRU declare that they have no competing interests.

// Peer Reviewers:

Ross Naylor, MBBS

Professor of Vascular Surgery

Vascular Surgery Group, Division of Cardiovascular Sciences, Leicester Royal Infirmary, UK

DISCLOSURES: RN declares that he has no competing interests.

William Pearce, MD

Chief of Division of Vascular Surgery

Department of Surgery, Northwestern Memorial Hospital, Chicago, IL

DISCLOSURES: WP declares that he has no competing interests.