



Genetics of vascular malformation and therapeutic implications

Miguel Zúñiga-Castillo, Christopher L. Teng, and Joyce M.C. Teng

Purpose of review

Vascular malformations (VaMs) are a consequence of disrupted morphogenesis that may involve arterial, capillary, venous, or lymphatic endothelium alone or in a combination. VaMs can have serious health impacts, leading to life-threatening conditions sometimes. Genetic mutations affecting proliferation, migration, adhesion, differentiation, and survival of endothelial cells, as well as integrity of extracellular matrix are believed to be the pathogenesis of these disorders. Here, we present an updated review of genetic mutations and potential therapeutic targets for VaMs.

Recent findings

Increased number of genetic mutations have been discovered in vascular anomalies via targeted deep sequencing. When a genetic defect is identified, it often presents in only a small percentage of cells within the malformation. In addition, mutations within the same gene may result in different clinical phenotypes. Management of VaMs can be challenging depending on the severity and functional impairment associated. There are no standard treatment algorithms available to date for VaMs, therefore the disorder has significant unmet clinical needs. Currently, the focus of therapeutic development is to target constitutively activated intracellular signaling pathways resulted from genetic mutations.

Summary

Knowledge about the genetic mutations and altered signaling pathways related to VaMs have improved our understanding about the pathogenesis of vascular anomalies and provided insights to the development of new targeted therapies.

Keywords

genetic mutations, signaling pathways, targeted therapy, vascular malformations

INTRODUCTION

Vascular malformations (VaMs) are a consequence of defective morphogenesis involving arterial, capillary, venous, or lymphatic endothelium alone or in combination [1]. These are lifelong conditions affecting approximately 1.5% of the general population [2]. VaMs often result in disfigurement, functional impairment, and recurrent infections. They can have a profound impact on quality of life and may increase lifetime risk of morbidity and mortality. VaMs can also arise with extracutaneous abnormalities as a part of genetic syndromes [1].

Genetic mutations affecting proliferation, migration, adhesion, differentiation, and survival of endothelial cells, as well as the integrity of extracellular matrix are believed to contribute to the pathogenesis of VaMs. These mutations are often identified via targeted deep sequencing in a percentage of cells within the malformation, and majority of VaMs are caused by monogenetic mutations. It is not uncommon that VaMs with different clinical

phenotypes result from mutations within the same gene [3,4[■]]. As a mosaic disorder, VaMs can be attributed to somatic mutation alone or loss of second allele in the presence of a preexisting germline mutation [4[■],5,6]. This explains why such patients can present with multifocal lesions evolving over time [7].

Here, we present an updated comprehensive review about the genetic pathogenesis (Table 1), current managements and potential targeted

Department of Dermatology, Division of Pediatric Dermatology, Stanford University School of Medicine, Lucile Packard Children's Hospital at Stanford, Stanford, California, USA

Correspondence to Joyce M.C. Teng, MD, PhD, Department of Dermatology, Division of Pediatric Dermatology, Stanford University School of Medicine, Lucile Packard Children's Hospital at Stanford, 700 Welch Road, Suite 301G, MC5896, Stanford, CA 94304, USA.

Tel: +1 650 724 9627; fax: +1 650 724 2491;

e-mail: jteng3@stanford.edu

Curr Opin Pediatr 2019, 31:498–508

DOI:10.1097/MOP.0000000000000794

KEY POINTS

- VaMs are a consequence of defective morphogenesis involving arterial, capillary, venous, or lymphatic endothelium alone or in a combination during embryonic development. Pathogenic mutations of VaM result in altered proliferation, migration, adhesion, differentiation of endothelial cells, as well as integrity of extracellular matrix.
- VaMs can be attributed to either somatic mutation alone or somatic mutation of the second allele in the presence of an inherited germline mutation of the other allele.
- Majority of VaMs are monogenic disorders, although VaMs with mutations in the same gene may have different clinical phenotypes.
- Recent discovery of specific pathogenic mutations associated with VaMs has revealed possible targets for therapeutic interventions. These therapies are designed against constitutively activated proangiogenic pathways in VaMs.

therapies for VaMs (Table 2). We have only included VaMs based on 2018 classification by the International Society for the Study of Vascular Anomalies (<https://www.issva.org/classification>), excluding the entities provisionally unclassified. Mutations with unknown functional effect or unknown clinical significance are included but not discussed.

GENETICS OF VASCULAR MALFORMATION

Mutations associated with VaMs often occur in genes encoding proteins that are part of the RAS/MEK/ERK (also known as MAPK/ERK) and/or PI3 kinase/mTOR signaling pathways (Fig. 1). Such genetic changes lead directly to altered endothelial cell proliferation, differentiation, and survival [3]. Un-controlled angiogenesis because of increased endothelial cell proliferation results in dysmorphogenesis of vascular network and vascular malformation. Even though many VaMs share commonly signaling pathways, their clinical phenotypes clearly vary. Although genotype–phenotype correlation remains an area of active investigation currently, studies have suggested that the origins of cells affected during embryogenesis may play a role in determining the final clinical phenotype of VaMs [3]. Recent discovery of many inhibitors targeting key players of these signaling pathways have offered unprecedented therapeutic opportunities for vascular anomalies.

G-protein subunit α Q, G-protein subunit α 11 in capillary malformation

Mutation of G-protein subunit α Q (*GNAQ*) presents in endothelial cells of nonsyndromic and syndromic capillary malformations. For example both non-complicated cutaneous capillary malformation and capillary malformation associated with Sturge-Weber syndrome were found to have same *GNAQ* mutations [4[¶],8,9–13]. The direct downstream effect of the *GNAQ* mutation is to induce activation of the MEK/ERK pathway, therefore promoting cell proliferation, migration, angiogenesis and malformation [9,11,13]. In addition, mutations within G-protein subunit α 11 have also been discovered among individuals with diffuse capillary malformations. These mutations are likely to activate MEK/ERK pathway as well, but more frequently associated with overgrowth syndrome [3,7,14].

PIK3CA, AKT1/2/3, PTEN in lymphatic malformation and Proteus syndrome

Recent studies have demonstrated that more than 70% of lymphatic malformation are associated with *PIK3CA* mutations. Gain of function mutations in *PIK3CA* lead to increased phosphorylation of AKT (AKT1, AKT2, or AKT3), therefore subsequent activation of the AKT/mTOR pathway (Fig. 1) as well as formation of disorganized vascular network because of uncontrolled cell proliferation [4[¶],7,15]. Overactivation of *PIK3CA* has also been shown to cause loss of fibronectin in extracellular matrix, which promote increased cell migration [7]. Aside from localized lymphatic malformation, *PIK3CA* mutation was also discovered in patients with CLOVES syndrome, CLAPO syndrome, capillary malformation with megalencephaly, Klippel-Trenaunay syndrome, or mucocutaneous venous malformations [3,7,16,17]. Similarly, mutations in *AKT1*, *AKT2*, and *AKT3* have also been discovered in Proteus syndrome, venous malformations, and capillary malformation with megalencephaly patients, respectively [17–19]. PTEN exerts an inhibitory effect on AKT via dephosphorylation (Fig. 1). Mutation of *PTEN* therefore results in loss of AKT inhibition and increased activation of AKT/mTOR pathway. Bannayan-Riley-Ruvalcaba syndrome, a genetic condition with macrocephaly and multiple hamartomas, is known to associate with *PTEN* mutations [15,18,20].

FLT-4 (VEGFR3), VEGFC, CCBE1, ADAMTS3 in lymphatic anomalies

VEGFR3, encoded by *FLT4*, and its ligand VEGFC play a major role in lymphangiogenesis during the embryonic development. Loss of function (LoF)

Table 1. Vascular malformations and associated genetic mutations

Vascular malformation	Gene	Mutation type	Inheritance	References
CM				
<i>Nevus simplex/salmon patch</i>	Unknown	Unknown	Unknown	
<i>Port-wine stain</i>				
Nonsyndromic CM	<i>GNAQ</i>	GoF	Somatic	[3,4 ²² ,7]
Sturge-Weber syndrome	<i>GNAQ</i>	GoF	Somatic	[3,4 ²² ,7,10]
CM with overgrowth	<i>GNA11</i>	GoF	Somatic	[3,7,14,66]
<i>Reticulate CM</i>				
CM of MIC	<i>STAMBP</i>	LoF	Germline. Unknown Mendelian pattern of inheritance	[67,68]
CM of MAC-polymicrogyria	1) <i>PIK3CA</i> 2) <i>AKT3</i> 3) <i>PIK3R2</i>	GoF	1)Somatic and somatic mosaicism 2)Germline and mosaic mutations. Unknown Mendelian pattern of inheritance 3)Germline. Unknown Mendelian pattern of inheritance	[3,19,55,59]
CMTC	<i>ARL6IP6</i> . Novel candidate gene	LoF	AD or mosaic ^a	[69–71]
Lymphatic Anomaly				
<i>Micro or Macrocytic LM</i>	<i>PIK3CA</i>	GoF	Somatic	[3,7,62]
<i>Generalized lymphatic anomaly</i>	<i>NRAS</i>	GoF	Somatic	[72]
Kaposiform lymphangiomatosis	1) <i>NRAS</i> 2) <i>BAD</i> 3) <i>TSC1</i>	1)GoF 2–3)LoF ^a	1)Somatic 2–3)Somatic	[73,74]
<i>LM in Gorham-Stout disease</i>	1) <i>TNFRSF11A</i> 2) <i>TREM2</i> 3) <i>PTEN</i>	Unknown	1–2)Somatic ^a 3)Germline. Unknown Mendelian pattern of inheritance	[75]
<i>Channel type LM</i>	<i>EPHB4</i>	LoF	AD	[76]
<i>Acquired progressive lymphatic anomaly</i>	Unknown	Unknown	Unknown	— — —
<i>Primary lymphedema</i>				
1)Nonne-Milroy syndrome or Milroy's disease and 2)Milroy-like lymphoedema	1) <i>FLT4(VEGFR3)</i> 2) <i>VEGFC</i>	LoF	1)AD, AR or <i>de novo</i> 2)AD	[22,77,78]
Primary hereditary lymphedema or Meige disease	1) <i>GJC2</i> 2) <i>MET</i> 3) <i>HGF</i>	1)GoF ^a 2–3)LoF	1)AD ^a 2–3)Unknown	[77,79]
Lymphedema-distichiasis	<i>FOXC2</i>	LoF	AD	[77,80]
Hypotrichosis-lymphedema-telangiectasia	<i>SOX18</i>	LoF	AD or AR or <i>de novo</i>	[24,81,82]
Primary lymphedema with myelodysplasia or Emberger syndrome	<i>GATA2</i>	LoF	AD	[77,79]
Primary generalized lymphatic anomaly or dysplasia (Hennekam lymphangiectasia-lymphedema syndrome)	1) <i>CCBE1</i> 2) <i>ADAMTS3</i> 3) <i>FAT4</i> 4) <i>PIEZO1</i> 5) <i>EPHB4</i>	1–2)LoF 3)LoF ^a 4–5)Unknown	1–4)AR 5)AD	[21,22,77,83]

Table 1 (Continued)

Vascular malformation	Gene	Mutation type	Inheritance	References
Microcephaly with or without chorioretinopathy, lymphedema, or mental retardation syndrome	<i>KIF11</i>	LoF ^a	AD	[77,84]
Lymphedema-choanal atresia	<i>PTPN14</i>	LoF	AR	[81,85]
VM				
- Common VM	1) <i>TEK</i> (<i>TIE2</i>) 2) <i>PIK3CA</i> 3) <i>GANQ</i> 4) <i>IRS2</i> 5) <i>MAP2K1/MEK1</i> 6) <i>AKT2</i> 7) <i>AKT3</i> 8) <i>NF1</i>	1–7) GoF 8) LoF	Somatic	[4 [■] ,12]
- Familial VMCM	<i>TEK</i> (<i>TIE2</i>)	GoF	AD	[3,4 [■]]
- BRBNS	<i>TEK</i> (<i>TIE2</i>)	GoF	Somatic	[4 [■]]
- GVM	<i>Glomulin</i>	LoF	AD and somatic (second hit)	[3,4 [■]]
- CCM	1) <i>CCM1/KRIT1</i> 2) <i>CCM2/malacavernin</i> 3) <i>CCM3/PDCD10</i>	LoF	AD or sporadic	[3,4 [■]]
- FVM	<i>ELMO2</i>	LoF	AR	[86]
- VVM	<i>MAP3K3/MEKK3</i>	GoF	Somatic	[3,7]
-MVM	<i>TEK</i> (<i>TIE2</i>)	GoF	Mosaic and sporadic somatic	[7]
AVM or AVF				
- Sporadic and non syndromic cerebrospinal AVM	1) <i>MAP2K1/MEK1</i> 2) <i>SMAD9</i> 3) <i>KRAS</i> 4) <i>BRAF</i> 5) <i>NARFL</i> 6) <i>PITPNM3</i> 7) <i>SARS</i> 8) <i>LEMD3</i> 9) <i>ALK1/ACVRL1</i> 10) <i>CLDN14</i>	1) GoF 2) LoF 6–9) LoF 10) GoF ^a	1) Somatic 2) Germline mutation Unknown Mendelian pattern of inheritance 6–9) De novo 10) Germline. Unknown Mendelian pattern of inheritance	[3,43,48,87–89]
- HHT	1) <i>ENG</i> 2) <i>ALK1/ACVRL1</i> 3) <i>BMP9/GDF2</i> 4) <i>SMAD4/MADH4</i> 5) <i>BMPR2</i>	1–5) LoF	1–2) AD and mosaicism 3–4) AD 5–6) Germline mutation. Unknown Mendelian pattern of inheritance	[3–7,88]
- CM of CM-AVM 1 and 2	1) <i>RASA1</i> 2) <i>EPHB4</i>	LoF	AD or somatic	[3,4 [■] ,7]
- Extracranial AVM	<i>MAP2K1/MEK1</i>	GoF	Somatic	[3,43,44]
- pulmonary AVM	<i>NARFL</i>	LoF	Germline. Unknown Mendelian pattern of inheritance	[46]
Other vascular malformation associated with other anomalies				
Klippel-Trenaunay syndrome	<i>PIK3CA</i>	GoF	Somatic and somatic mosaicism	[4 [■] ,62]
Parkes Weber syndrome	<i>RASA1</i>	LoF	AD and somatic	[4 [■] ,7]
Limb CM + congenital nonprogressive limb overgrowth	<i>GNA11</i>	GoF	Somatic	[14,66]
Maffucci syndrome	<i>Isocitrate dehydrogenase 1 and/or 2</i>	Unknown	Somatic and mosaic	[3,20]

Table 1 (Continued)

Vascular malformation	Gene	Mutation type	Inheritance	References
CLOVES syndrome	PIK3CA	GoF	Somatic and somatic mosaicism	[3,17,62]
Proteus syndrome	AKT1	GoF	Somatic and somatic mosaicism	[3,15,17,55]
Bannayan-Riley-Ruvalcaba syndrome	PTEN	LoF	AD and mosaic	[15,81,90]
CLAPO syndrome	PIK3CA	GoF	Somatic	[17]
MOVLDS	DDX24	Unknown	Germline. Unknown Mendelian pattern of inheritance	[2]

AD, autosomal dominant; AR, autosomal recessive; AVF, arteriovenous fistula; AVM, arteriovenous malformation; BRBNS, Blue rubber bleb nevus syndrome; CCM, cerebral cavernous malformation; CM, capillary malformation; CMTC, cutis marmorata telangiectatica congenita; FIVM, familial intraosseous vascular malformation; GoF, gain of function; GVM, glomuvenous malformation; HHT, hereditary hemorrhagic telangiectasia; LM, lymphatic malformation; LoF, loss of function; MAC, macrocephaly; MIC, microcephaly; MOVLDS, multiorgan venous and lymphatic defect syndrome; MVM, multifocal venous malformation; VM, venous malformation; VMCM, venous malformation cutaneo-mucosal; VVM, verrucous venous malformation.

*it needs more studies to be conducted in this field.

mutations of VEGFR3 and VEGFC have been reported in Milroy's disease and Milroy-like lymphedema, respectively. CCBE1 and ADAMTS3 are also known to be required for the activation of VEGFC, as autosomal recessive LoF mutations in *CCBE1* and *ADAMTS3* have been identified in Hennekam lymphangiectasia-lymphedema syndrome [21,22]. Mutations within any of the four genes (*FLT-4* (*VEGFR3*), *VEGFC*, *CCBE1*, *ADAMTS3*) may impair signaling via the MAPK/ERK and AKT/PI3K/mTOR pathways, which present potential targets for therapeutic interventions [23,24].

TEK (TIE2) in Venous Malformation

Blue rubber bleb nevus syndrome (BRBNS), as well as familial mucocutaneous venous malformation and more than 50% sporadic venous malformations can be attributed to activating mutations in *TEK*, which encodes the TIE2 receptor expressed primarily on venous endothelial cells [4^{***},16,25]. Individuals with venous malformation and BRBNS may carry activating mutation (s) in the same allele, though double mutations are more common in BRBNS [26]. Multifocal venous malformation and familial venous malformation cutaneomucosal are because of a somatic second-hit mutation among individuals who already carry a germline *TEK* mutation [7,16]. Mutation of *TEK* results in ligand-independent phosphorylation of TIE2 receptor and constitutive activation of TIE2-mediated signaling via pathways including AKT/PI3K/mTOR and MAPK/ERK [7,27]. Pathogenic *TEK* mutations also give rise to activities of c-ABL tyrosine kinase that results in subsequent activation of PI3K/AKT/mTOR and PLC γ /ERK1/2 pathways. The activation of these pathways directly

enhance angiogenesis and reduce endothelial cell apoptosis, therefore leads to abnormal vascular morphogenesis [25].

Glomulin and glumovenous malformation

Recent study has shown that germline glomulin (*GLMN*) LoF mutation and somatic mutation of second allele are often associated with glumuvenous malformation [3,7]. Germline mutations affecting both alleles are most likely incompatible with life. *GLMN* are expressed in endothelial and perivascular smooth muscle cells. Mutation in *GLMN* blocks TGF- β signaling, but enhance signaling via the PI3K/mTOR pathway that results in abnormal differentiation of the vascular smooth muscle cells and defective formation of vascular bed because of inappropriate remodeling [4^{***},7].

DDX24 in extacutaneous veno-lymphatic malformation

A missense mutation of *DDX24* was recently described in a large family with multiorgan venous and lymphatic defect syndrome. Veno-lymphatic malformation lesions were found in multiples organs of the affected individuals, in addition to portal and hepatic vein stenosis. The same mutation was reported in another group of patients with sporadic portal and hepatic vein stenosis and increased risk of inferior vena cava occlusion. *DDX24* regulates endothelial cell migration and tube formation, thus affecting generalized venous and lymphatic development especially in liver and heart [2].

Table 2. Targeted therapies for vascular anomalies

Signaling pathway	Vascular anomalies	Targeted therapy ^a	References
PI3K-AKT-mTOR	HHT, GVM, Proteus, CMMAC, BRRS, CLM, LM in Gorham-Stout disease CLOVESS, CLAPOS, and KTS.	<i>PI3K inhibitors:</i> Alpelisib, Dactolisib, Idelalisib, copanlisib/ BAY 80–6946, and taselisib/ GDC-0032 <i>PI3K p110α Inhibitor:</i> Pictilisib/ GDC-0941 <i>PIK3CA Inhibitors:</i> Buparlisib/ BKM-120, Wortmannin, LY294002, and Alpelisib <i>AKT inhibitor:</i> MK2206 <i>Pan-AKT inhibitor:</i> ARQ092 <i>mTOR inhibitors:</i> sirolimus, everolimus and temsirolimus <i>mTORC1/C2-PI3K inhibitor:</i> BEZ235	[7,16,54–56,59,62,63,65]
RAS-BRAF-MEK-ERK	NSCM, CM with OG, CM with bone and/or OG, DCM with or without OG, LCMCNPLOG, SWS, PWwM, CCM, Kaposiform lymphangiomatosis, VVM, EAVM, and SAVM	<i>BRAF inhibitor:</i> Vemurafenib <i>MEK inhibitors:</i> Trametinib, Cobimetinib, PD98059, U0126 <i>MEK5 inhibitor:</i> BIX02189 <i>ERK5 inhibitors:</i> XMD8–92, XMD17–105 <i>Multiple kinase inhibitors (VEGFR1, RAF-ERK, others):</i> Sorafenib <i>VEGF inhibitor:</i> Bevacizumab <i>ALK5 inhibitor:</i> SB-431542	[7,16,31,44,47,60,61]
PI3K-AKT-mTOR and RAS-BRAF-MEK-ERK	PWS, CM-AVM 1 and 2, Channel type LM, MD, HLLS, MLL, BAVM, CVM, MVM, FVMCM, and BRBNS.	<i>TIE2 inhibitor:</i> CAS 948557–43–5 <i>ABL inhibitor:</i> Ponatinib	[16,25]
Smad1/5/8	HHT	<i>Calcineurin inhibitor and activator of Smad1/5/8:</i> Tacrolimus	[54]

BAVM, brain arteriovenous malformation; BRBNS, blue rubber bleb nevus syndrome; BRRS, Bannayan-Riley-Ruvalcaba syndrome; CCM, cerebral cavernous malformation; CLAPOS, CLAPO syndrome; CLM, cystic lymphatic malformation; CLOVESS, CLOVES syndrome; CM, capillary malformation; CM-AVM, capillary malformation – arteriovenous malformation; CMMAC, capillary malformation of macrocephaly; CVM, common venous malformation; DCM, diffuse capillary malformation; EAVM, extracranial arteriovenous malformation; FVMCM, familial venous malformation cutaneous-mucosal; GVM, glomuvenous malformation; HHT, hereditary hemorrhagic telangiectasia; HLLS, Hennekam lymphangiectasia-lymphedema syndrome; KTS, Klippel-Trenaunay syndrome; LCMCNPLOG, limb capillary malformation + congenital nonprogressive limb overgrowth; LM, lymphatic malformation; MD, Milroy's disease; MLL, Milroy-like lymphedema; MVM, multifocal venous malformation; NSCM, non syndromic capillary malformation; OG, overgrowth; Proteus, Proteus syndrome; PWS, Parkes Weber syndrome; PWwM, port-wine stain with macrocheilia; SAVM, spinal arteriovenous malformation; SWS, Sturge-Weber syndrome; VVM, verrucous venous malformation.

^aTherapies that have shown some efficacy in human studies or current in clinical trials are highlighted in bold.

CCM1/KRIT1, CCM2/malcavernin, CCM3/PDCD10 in cerebral cavernous malformation

Most of the patients with cerebral cavernous malformation (CCM) have sporadic mutations in the CCM genes [3,28–30]. LoF mutations in any of the CCM genes result in similar clinical phenotypes. However, the most severe diseases are often because of mutations of CCM3 gene [31]. CCM genes encode cytosolic proteins that form a complex downregulating MAP3K3 functions, therefore, inhibits the activity of MEK/ERK pathway. The inhibitory effect maintains quiescent state of endothelial cells and mitigates VEGF proangiogenic signaling [4^{••},7,32]. Mutations of CCM gene result in loss of regulation on MEK/ERK pathway that leads to proangiogenic

endothelial cells migration, impaired endothelium integrity and disrupted vascular morphogenesis [7,32]. The intracranial vasculature is therefore fragile and prone to hemorrhage [4^{••},32,33].

RASA1, EPHB4 and capillary malformation-arteriovenous malformation

Germline autosomal dominant RASA1 mutations have been identified in 50% of capillary malformation-arteriovenous malformation (CM-AVM1) patients including those with Parkes-Weber syndrome [3,7,34]. Germline RASA1 mutations as 'first hit' produce haploinsufficiency, and a somatic mutation of the second allele as "second hit" results




FIGURE 1. Schematic diagram of a signal transduction in endothelial cells and the main genetic mutations associated to vascular malformation. Commonly activated signaling pathways associated with vascular anomalies are color coded in green and blue. Transmembrane receptors and intracellular proteins play a role in angiogenesis are shown in color. Proteins mutated in different vascular disorders are shown. Arrows indicate direct or indirect interactions and red lines indicate inhibition. PIK3CA is part of PI3K; RAS includes HRAS and KRAS; RASA1 is one of the RASGAPs. BAVM, brain arteriovenous malformation; BRBNS, blue rubber bleb nevus syndrome; BRRS, Bannayan-Riley-Ruvalcaba syndrome; CCM, cerebral cavernous malformation; CLAPOS, CLAPO syndrome; CLM, cystic lymphatic malformation; CLOVESS, CLOVES syndrome; CM-AVM, capillary malformation-arteriovenous malformation; CMMAC, capillary malformation of Macrocephaly; CVM, common venous malformation; DCM, diffuse capillary malformation; EAVM, extracranial arteriovenous malformation; FVMCM, familial venous malformation cutaneous-mucosal; GoFM, gain of function mutation; GVM, glomuvenous malformation; HHT, hereditary hemorrhagic telangiectasia; HLLS, Hennekam lymphangiectasia-lymphedema syndrome; KTS, Klippel-Trenaunay syndrome; LCMCNPLOG, limb capillary malformation + congenital nonprogressive limb overgrowth; LoFM, loss of function mutation; MD, Milroy's disease; MLL, Milroy-like lymphedema; NSCM, non syndromic capillary malformation; OG, overgrowth; ProteusS, Proteus syndrome; PWS, Parkes Weber syndrome; PWwM, port-wine stain with macrocheilia; SAVM, spinal arteriovenous malformation; SWS, Sturge-Weber syndrome; VVM, verrucous venous malformation.

in VaMs in this disorder [4²²,35,36,37]. *RASA1* encodes p120-RasGAP protein that inhibits activity of RAS protein, an upstream regulator of the MEK/ERK1/2 and AKT/PI3K/mTOR pathways [7,38–40]. LoF mutations of *RASA1* therefore may lead to activation of RAS and increased downstream signaling via MEK/ERK1/2 and AKT/mTOR pathways [4²²,7,41] that can be targeted potentially.

Additional studies demonstrated recently that autosomal dominant *EPHB4* mutations were found in CM-AVM2 [42]. This gene encodes a transmembrane receptor expressed primarily in venous endothelial cells during vascular development interacting with its ligand, EphrinB2, on arterial endothelial cells. The interaction between *EPHB4* and EphrinB2 is essential in venous and arterial

differentiation. EPHB4 also activates p120RasGAP, therefore exerts similar downstream effects to MEK/ERK pathway as RASA1. The phenotypic similarity between CM-AVM1 and CM-AVM2 suggests *RASA1* and *EPHB4* play an overlapping role in vascular development during embryogenesis [7,38].

MAP2K1 (MEK1), nuclear prelamin A recognition factor-like and extracranial arteriovenous malformation

MAP2K1 or MEK1 is a well-known, downstream effector of BRAF. Mutation of *MAP2K1* has been identified in more than 50% cases of extracranial AVMs, though it has also been described in some cases of intracranial AVM [3,43,44]. The discovery has presented the possibility of using targeted MEK inhibitors that are currently available to manage complex extracranial AVMs. This has been confirmed by a recent proof-of-concept case report by Lekwuttikarn *et al.* [45].

Nuclear prelamin A recognition factor-like protein is encoded by *CIAO3* gene. LoF mutation of *CIAO3* has been described in two siblings with diffuse pulmonary AVMs from a Chinese consanguineous family. In animal model, nuclear prelamin A recognition factor-like knockdown presents with upregulation of transferrin receptor 1 expression and intracellular iron storage, which leads to increased angiogenesis via activated VEGF signaling [46].

KRAS, BRAF and SMAD9 in AVM of central nervous system

Somatic mutations of KRAS and BRAF were found in 80–100% of patients with brain and spinal AVMs [44]. These mutations activate signaling of the MAPK/ERK pathway, which leads to unregulated endothelial cells proliferation, growth, migration, and differentiation [7,44,47].

SMAD9 mutation was recently described in a 14-year-old girl with recurrent brain AVM. Phosphorylated wild type SMAD9 interacts with SMAD4, before being translocated into the nucleus to exert its proangiogenic function [48].

ENG, ALK1/ACVRL1, BMP9/GDF2, and SMAD4/MADH4 in hereditary hemorrhagic telangiectasia

Autosomal dominant mutations of *ENG* and *ALK1* affected approximately 96% of hereditary hemorrhagic telangiectasia (HHT) patients. The remaining populations affected were attributed to mutations of *SMAD4* and *BMP9* [3,5].

Genotype-phenotype correlation is better defined in HHT patients. Individuals with *ENG* mutations have a higher incidence of pulmonary AVMs, therefore increased risk of complications. *ALK1* mutations, on the other hand, are associated more frequently with hepatic AVMs and gastrointestinal bleeds [49]. Patients with aortic dilation often carry *SMAD4* mutations [50].

ENG is an endothelial transmembrane protein that can bind different types of TGF- β and BMP, and interact with ALK1 upon activation [7,51,52]. The *ENG*-ALK1 complex phosphorylates SMADs, before they translocate into the nucleus to promote the transcription of proangiogenic genes [51,53]. Tacrolimus was specifically proposed as a therapy for HHT1, as it helps to mitigate aberrant AKT activation [54].

THERAPEUTIC IMPLICATIONS

Vascular anomalies with mutations affecting the AKT/PI3K/mTOR pathway (e.g. venous malformation, venous malformation cutaneo-mucosal, multifocal venous malformation, BRBNS, and lymphatic malformation) are known to respond to mTOR inhibitors (Fig. 1, Table 2) (sirolimus, everolimus, and temsirolimus) [55,56]. Recent phase II study has clearly demonstrated the efficacy of sirolimus for a variety of vascular anomalies in majority of participants [57,58]. Theoretically, PI3K inhibitors (alpelisib, dactolisib, copanlisib, idelasib, copanlisib/BAY 80-6946, and taselisib/GDC-0032) as well as AKT inhibitor (MK2206) [7,16,59] can block upstream signaling of mTOR pathway and exert better therapeutic effects. However, well-controlled perspective studies are needed to evaluate the safety as well as efficacy.

Malformations because of mutations affecting RAS/BRAF/MEK/ERK pathway (e.g. capillary malformation, CM-AVM, cerebrospinal AVM) could perhaps be targeted by BRAF inhibitor (e.g. vemurafenib) and/or MEK inhibitors (trametinib, cobimetinib, U0126 or PD98059) that are available [7,16,44,47]. The MEK inhibitor U0126 restores cadherin localization in the junction of endothelial cells in human umbilical vein endothelial cells; therefore has been proposed as a potential therapy for central nervous system AVMs [60,61]. Additional therapeutics, including sorafenib, multiple kinase inhibitors (VEGFR1, RAF-ERK), ALK5 inhibitor (SB-431542), ERK5 inhibitors (XMD8–92, XMD17–105), and MEK5 inhibitor (BIXO2189) are currently investigated for treatment of CCMs [31].

In addition, other targeted treatments for VaMs using different mechanism of actions are being studied. For example, sorafenib, PIK3CA inhibitors

(buparlisib/BKM-120, wortmannin, and Ly294002), selective PI3K p110 α inhibitor (pictilisib/GDC-0941) and MK2206 have been evaluated *in vitro* for the treatment of lymphatic malformation [62,63]. In-vitro and in-vivo studies with everolimus, MK2206 and mTOR-PI3K duo inhibitor (BEZ235) are being evaluated for the treatment of venous malformation because of PIK3CA mutations [56,63]. Alpelisib (BYL719), a highly specific PIK3CA inhibitor is found to be superior than sirolimus *in vitro* as well as in-vivo studies. The drug was also shown to be effective in treating cutaneous venous malformation in a mouse allotransplantation model when delivered as a topical cream. Case report of two patients with CLOVES appear to have benefited from treatment with alpelisib [16,64]. A subsequent large case series published recently [64] demonstrated marked improvement among patients with PIK3-related overgrowth syndrome. Phase 1 clinical trial is currently underway in the United States to further investigate its safety and benefit. The efficacy of new pan-AKT inhibitor, ARQ092, for the treatment of Proteus, CLOVES, and capillary malformation of macrocephaly is also being examined [59,65]. Ponatinib, a potent ABL inhibitor, was shown to be efficacious inhibiting angiogenesis in comparison to PI3K or MEK inhibitors *in vitro*. Combined therapy with ponatinib and sirolimus induces regression of the venous malformation synergistically [25]. It has been hypothesized that ponatinib blocks signaling via both TIE2 and c-ABL, therefore downregulates multiple pathways including PI3K/AKT/mTOR and PLC γ /ERK1/2. Furthermore, a new TIE2 kinase inhibitor is currently in development (CAS 948557-43-5) [16].

CONCLUSION

Knowledge about the genetics of VaMs has improved our understanding about the pathogenesis of these disorders, and provided a foundation for the development of new targeted therapy. As many of these malformations activate similar downstream angiogenic pathways, targeted therapies developed may be used for treatment of several vascular anomalies. Proof-of-concept studies are underway to investigate the safety and efficacy of these treatments. Furthermore combination therapies may be considered in the future to improve the management of complex vascular disorders.

Acknowledgements

None.

Financial support and sponsorship

None.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Carqueja IM, Sousa J, Mansilha A. Vascular malformations: classification, diagnosis and treatment. *Int Angiol* 2018; 37:127–142.
2. Pang P, Hu X, Zhou B, *et al*. DDX24 mutations associated with malformations of major vessels to the viscera. *Hepatology* 2019; 69:803–816.
3. Greene AK, Goss JA. Vascular anomalies: from a clinicohistologic to a genetic framework. *Plast Reconstr Surg* 2018; 141:709e–717e.
4. Wetzel-Strong SE, Detter MR, Marchuk DA. The pathobiology of vascular malformations: insights from human and model organism genetics. *J Pathol* 2017; 241:281–293.

This article emphasizes the role of the genetic factors in the pathogenesis of vascular malformations.

5. McDonald J, Wooderchak-Donahue WL, Henderson K, *et al*. Tissue-specific mosaicism in hereditary hemorrhagic telangiectasia: implications for genetic testing in families. *Am J Med Genet A* 2018; 176:1618–1621.
6. Tørring PM, Kjeldsen AD, Ousager LB, Brusgaard K. ENG mutational mosaicism in a family with hereditary hemorrhagic telangiectasia. *Mol Genet Genomic Med* 2018; 6:121–125.
7. Queisser A, Boon LM, Vikkula M. Etiology and genetics of congenital vascular lesions. *Otolaryngol Clin North Am* 2018; 51:41–53.
8. Huang L, Couto JA, Pinto A, *et al*. Somatic GNAQ mutation is enriched in brain endothelial cells in Sturge-Weber syndrome. *Pediatr Neurol* 2017; 67:59–63.
9. Sundaram SK, Michelhaugh SK, Klinger NV, *et al*. GNAQ mutation in the venous vascular malformation and underlying brain tissue in sturge-weber syndrome. *Neuropediatrics* 2017; 48:385–389.
10. Martins L, Giovani PA, Rebouças PD, *et al*. Computational analysis for GNAQ mutations: new insights on the molecular etiology of Sturge-Weber syndrome. *J Mol Graph Model* 2017; 76:429–440.
11. Higueros E, Roe E, Granell E, Baselga E. Sturge-Weber syndrome: a review. *Actas Dermosifiliogr* 2017; 108:407–417.
12. Castel P, Carmona FJ, Grego-Bessa J, *et al*. Somatic PIK3CA mutations as a driver of sporadic venous malformations. *Sci Transl Med* 2016; 8:332ra42.
13. Ma G, Yu Z, Liu F, *et al*. Somatic GNAQ mutation in different structures of port-wine macrocheilia. *Br J Dermatol* 2018; 179:1109–1114.
14. Couto JA, Ayturk UM, Konczyk DJ, *et al*. A somatic GNA11 mutation is associated with extremity capillary malformation and overgrowth. *Angiogenesis* 2017; 20:303–306.
15. Nathan N, Keppler-Noreuil KM, Biesecker LG, *et al*. Mosaic Disorders of the PI3K/PTEN/AKT/TSC/mTORC1 Signaling Pathway. *Dermatol Clin* 2017; 35:51–60.
16. Kangas J, Nätyнки M, Eklund L. Development of molecular therapies for venous malformations. *Basic Clin Pharmacol Toxicol* 2018; 123(Suppl 5):6–19.
17. Rodríguez-Laguna L, Ibañez K, Gordo G, *et al*. CLAPO syndrome: identification of somatic activating PIK3CA mutations and delineation of the natural history and phenotype. *Genet Med* 2018; 20:882–889.
18. Castillo SD, Vanhaesebroeck B, Sebire NJ. Phosphoinositide 3-kinase: a new kid on the block in vascular anomalies. *J Pathol* 2016; 240:387–396.
19. Negishi Y, Miya F, Hattori A, *et al*. A combination of genetic and biochemical analyses for the diagnosis of PI3K-AKT-mTOR pathway-associated megalencephaly. *BMC Med Genet* 2017; 18:4.
20. Kirkorian AY, Grossberg AL, Püttgen KB. Genetic basis for vascular anomalies. *Semin Cutan Med Surg* 2016; 35:128–136.
21. Brouillard P, Dupont L, Helaers R, *et al*. Loss of ADAMTS3 activity causes Hennekam lymphangiectasia-lymphedema syndrome 3. *Hum Mol Genet* 2017; 26:4095–4104.
22. Scheuerle AE, Sweed NT, Timmons CF, *et al*. An additional case of Hennekam lymphangiectasia-lymphedema syndrome caused by loss-of-function mutation in ADAMTS3. *Am J Med Genet A* 2018; 176:2858–2861.
23. Sevic-Muraca EM, King PD. Lymphatic vessel abnormalities arising from disorders of Ras signal transduction. *Trends Cardiovasc Med* 2014; 24:121–127.
24. Brouillard P, Boon L, Vikkula M. Genetics of lymphatic anomalies. *J Clin Invest* 2014; 124:898–904.
25. Li X, Cai Y, Goines J, *et al*. Ponatinib combined with rapamycin causes regression of murine venous malformation. *Arterioscler Thromb Vasc Biol* 2019; ATVB.AHA118312315.
26. Soblet J, Kangas J, Nätyнки M, *et al*. Blue rubber bleb nevus (BRBN) syndrome is caused by somatic TEK (TIE2) mutations. *J Invest Dermatol* 2017; 137:207–216.

27. Kennedy MA, Xu Z, Wu Y, Sohl CD. A Tie2 kinase mutation causing venous malformations increases phosphorylation rates and enhances cooperativity. *Biochem Biophys Res Commun* 2019; 509:898–902.
28. Spiegler S, Rath M, Hoffjan S, *et al.* First large genomic inversion in familial cerebral cavernous malformation identified by whole genome sequencing. *Neurogenetics* 2018; 19:55–59.
29. Rinaldi C, Bramanti P, Scimone C, *et al.* Relevance of CCM gene polymorphisms for clinical management of sporadic cerebral cavernous malformations. *J Neurol Sci* 2017; 380:31–37.
30. Wang H, Pan Y, Zhang Z, *et al.* A novel KRIT1/CCM1 gene insertion mutation associated with cerebral cavernous malformations in a Chinese family. *J Mol Neurosci* 2017; 61:221–226.
31. Lampugnani MG, Malinverno M, Dejana E, Rudini N. Endothelial cell disease: emerging knowledge from cerebral cavernous malformations. *Curr Opin Hematol* 2017; 24:256–264.
32. Russo A, Neu MA, Theruvath J, *et al.* Novel loss of function mutation in KRIT1/CCM1 is associated with distinctly progressive cerebral and spinal cavernous malformations after radiochemotherapy for intracranial malignant germ cell tumor. *Childs Nerv Syst* 2017; 33:1275–1283.
33. de Vos IJ, Vreeburg M, Koek GH, van Steensel MA. Review of familial cerebral cavernous malformations and report of seven additional families. *Am J Med Genet A* 2017; 173:338–351.
34. Gourier G, Audebert-Bellanger S, Vourc'h P, *et al.* Multiple capillary malformations of progressive onset: capillary malformation-arteriovenous malformation syndrome (CM-AVM). *Ann Dermatol Venereol* 2018; 145:486–491.
35. Lapinski PE, Doosti A, Salato V, *et al.* Somatic second hit mutation of RASA1 in vascular endothelial cells in capillary malformation-arteriovenous malformation. *Eur J Med Genet* 2018; 61:11–16.
36. Cai R, Liu F, Liu Y, *et al.* RASA-1 somatic 'second hit' mutation in capillary malformation-arteriovenous malformation. *J Dermatol* 2018; 45:1478–1480.
37. Cai R, Liu F, Hua C, *et al.* A novel RASA1 mutation causing capillary malformation-arteriovenous malformation (CM-AVM): the first genetic clinical report in East Asia. *Hereditas* 2018; 155:24.
38. Yu J, Streicher JL, Medne L, *et al.* EPHB4 Mutation Implicated in capillary malformation-arteriovenous malformation syndrome: a case report. *Pediatr Dermatol* 2017; 34:e227–e230.
39. Amyere M, Revencu N, Helaers R, *et al.* Germline loss-of-function mutations in EPHB4 cause a second form of capillary malformation-arteriovenous malformation (CM-AVM2) deregulating RAS-MAPK signaling. *Circulation* 2017; 136:1037–1048.
40. Cao H, Alrejaye N, Klein OD, *et al.* A review of craniofacial and dental findings of the RASopathies. *Orthod Craniofac Res* 2017; 20(Suppl 1):32–38.
41. Edwards LR, Blechman AB, Zlotoff BJ. RASA1 mutation in a family with capillary malformation-arteriovenous malformation syndrome: a discussion of the differential diagnosis. *Pediatr Dermatol* 2018; 35:e9–e12.
42. Saliou G, Eyries M, Iacobucci M, *et al.* Clinical and genetic findings in children with central nervous system arteriovenous fistulas. *Ann Neurol* 2017; 82:972–980.
43. Couto JA, Huang AY, Konczyk DJ, *et al.* Somatic MAP2K1 mutations are associated with extracranial arteriovenous malformation. *Am J Hum Genet* 2017; 100:546–554.
44. Hong T, Yan Y, Li J, *et al.* High prevalence of KRAS/BRAF somatic mutations in brain and spinal cord arteriovenous malformations. *Brain* 2019; 142:23–34.
45. Lekwuttikarn R, Lim YH, Admani S, *et al.* Genotype-guided medical treatment of an arteriovenous malformation in a child. *JAMA Dermatol* 2018; doi: 10.1001/jamadermatol.2018.4653. [Epub ahead of print]
46. Liu HZ, Du CX, Luo J, *et al.* A novel mutation in nuclear prelamin A recognition factor-like causes diffuse pulmonary arteriovenous malformations. *Oncotarget* 2017; 8:2708–2718.
47. Starke RM, McCarthy D, Komotar RJ, Connolly ES. Somatic KRAS mutation found in sporadic arteriovenous malformations. *Neurosurgery* 2018; 83:E14–E15.
48. Walcott BP, Winkler EA, Zhou S, *et al.* Identification of a rare. *Hum Genome Var* 2018; 5:18001.
49. Mu W, Cordner ZA, Yuqi Wang K, *et al.* Characterization of pulmonary arteriovenous malformations in ACVRL1 versus ENG mutation carriers in hereditary hemorrhagic telangiectasia. *Genet Med* 2018; 20:639–644.
50. Vorselaars VM, Diederik A, Prabhudesai V, *et al.* SMAD4 gene mutation increases the risk of aortic dilation in patients with hereditary haemorrhagic telangiectasia. *Int J Cardiol* 2017; 245:114–118.
51. Ollauri-Ibañez C, López-Novoa JM, Pericacho M. Endoglin-based biological therapy in the treatment of angiogenesis-dependent pathologies. *Expert Opin Biol Ther* 2017; 17:1053–1063.
52. Sugden WW, Siekmann AF. Endothelial cell biology of Endoglin in hereditary hemorrhagic telangiectasia. *Curr Opin Hematol* 2018; 25:237–244.
53. Albiñana V, Zafra MP, Colau J, *et al.* Mutation affecting the proximal promoter of Endoglin as the origin of hereditary hemorrhagic telangiectasia type 1. *BMC Med Genet* 2017; 18:20.
54. Ruiz S, Chandakkar P, Zhao H, *et al.* Tacrolimus rescues the signaling and gene expression signature of endothelial ALK1 loss-of-function and improves HHT vascular pathology. *Hum Mol Genet* 2017; 26:4786–4798.
55. Martínez-López A, Blasco-Morente G, Pérez-López I, *et al.* CLOVES syndrome: review of a PIK3CA-related overgrowth spectrum (PROS). *Clin Genet* 2017; 91:14–21.
56. di Blasio L, Puliafito A, Gagliardi PA, *et al.* PI3K/mTOR inhibition promotes the regression of experimental vascular malformations driven by PIK3CA-activating mutations. *Cell Death Dis* 2018; 9:45.
57. Wiegand S, Wichmann G, Dietz A. Treatment of lymphatic malformations with the mTOR inhibitor sirolimus: a systematic review. *Lymphat Res Biol* 2018; 16:330–339.
58. Nadal M, Giraudeau B, Tavernier E, *et al.* Efficacy and safety of mammalian target of rapamycin inhibitors in vascular anomalies: a systematic review. *Acta Derm Venereol* 2016; 96:448–452.
59. Akgumus G, Chang F, Li MM. Overgrowth syndromes caused by somatic variants in the phosphatidylinositol 3-kinase/akt/mammalian target of rapamycin pathway. *J Mol Diagn* 2017; 19:487–497.
60. Cheng F, Nussinov R. KRAS activating signaling triggers arteriovenous malformations. *Trends Biochem Sci* 2018; 43:481–483.
61. Nikolaev SI, Vetiska S, Bonilla X, *et al.* Somatic activating KRAS mutations in arteriovenous malformations of the brain. *N Engl J Med* 2018; 378:250–261.
62. Blesinger H, Kaulfuß S, Aung T, *et al.* PIK3CA mutations are specifically localized to lymphatic endothelial cells of lymphatic malformations. *PLoS One* 2018; 13:e0200343.
63. Suzuki Y, Enokido Y, Yamada K, *et al.* The effect of rapamycin, NVP-BEZ235, aspirin, and metformin on PI3K/AKT/mTOR signaling pathway of PIK3CA-related overgrowth spectrum (PROS). *Oncotarget* 2017; 8:45470–45483.
64. Venot Q, Blanc T, Rabia SH, *et al.* Targeted therapy in patients with PIK3CA-related overgrowth syndrome. *Nature* 2018; 558:540–546.
65. Ranieri C, Di Tommaso S, Loconte DC, *et al.* In vitro efficacy of ARQ 092, an allosteric AKT inhibitor, on primary fibroblast cells derived from patients with PIK3CA-related overgrowth spectrum (PROS). *Neurogenetics* 2018; 19:77–91.
66. Liu KX, Prajapati VH, Liang MG, *et al.* A cross-sectional survey of long-term outcomes for patients with diffuse capillary malformation with overgrowth. *J Am Acad Dermatol* 2018; 78:1023–1025.
67. Hori I, Miya F, Negishi Y, *et al.* A novel homozygous missense mutation in the SH3-binding motif of STAMBP causing microcephaly-capillary malformation syndrome. *J Hum Genet* 2018; 63:957–963.
68. Naseer MI, Sogaty S, Rasool M, *et al.* Microcephaly-capillary malformation syndrome: Brothers with a homozygous STAMBP mutation, uncovered by exome sequencing. *Am J Med Genet A* 2016; 170:3018–3022.
69. Jia D, Rajadurai VS, Chandran S. Cutis marmorata telangiectatica congenita with skin ulceration: a rare benign skin vascular malformation. *BMJ Case Rep* 2018; pii: bcr-2018-226763. doi: 10.1136/bcr-2018-226763.
70. Mon RA, Mozurkewich E, Treadwell MC, Berman DR. Cutis marmorata telangiectatica congenita presenting as a fetal hemothorax. *Fetal Diagn Ther* 2019; 45:281–284.
71. Amaral J, Peixoto S, Mimoso G, Pereira D. Cutis marmorata telangiectatica congenita and major lower limb asymmetry. *BMJ Case Rep* 2018; 2018:pii: bcr-2017-222269. doi: 10.1136/bcr-2017-222269.
72. Michio O. Generalized Lymphatic Anomaly and Gorham–Stout Disease: Overview and Recent Insights. In: Toshiyuki, F., editor. *Advances in Wound Care* 2019. DOI: 10.1089/wound.2018.0850.
73. Barclay SF, Inman KW, Luks VL, *et al.* A somatic activating NRAS variant associated with kaposiform lymphangiomatosis. *Genet Med* 2018; doi: 10.1038/s41436-018-0390-0. [Epub ahead of print]
74. Glaser K, Dickie P, Dickie BH. Proliferative cells from kaposiform lymphangiomatosis lesions resemble mesenchymal stem cell-like pericytes defective in vessel formation. *J Pediatr Hematol Oncol* 2018; 40:e495–e504.
75. Li MH, Zhang HQ, Lu YJ, *et al.* Successful management of gorham-stout disease in scapula and ribs: a case report and literature review. *Orthop Surg* 2018; 10:276–280.
76. Li D, Wenger TL, Seiler C, *et al.* Pathogenic variant in EPHB4 results in central conducting lymphatic anomaly. *Hum Mol Genet* 2018; 27:3233–3245.
77. Jones GE, Mansour S. An approach to familial lymphoedema. *Clin Med (Lond)* 2017; 17:552–557.
78. Nadarajah N, Schulte D, McConnell V, *et al.* A novel splice-site mutation in VEGFC is associated with congenital primary lymphoedema of gordon. *Int J Mol Sci* 2018; 19; pii: E2259. doi: 10.3390/ijms19082259.
79. Michelini S, Vettori A, Maltese PE, *et al.* Genetic screening in a large cohort of Italian patients affected by primary lymphedema using a next generation sequencing (NGS) approach. *Lymphology* 2016; 49:57–72.
80. De Niear MA, Breazzano MP, Mawn LA. Novel FOXC2 mutation and distichiasis in a patient with lymphedema-distichiasis syndrome. *Ophthalmic Plast Reconstr Surg* 2018; 34:e88–e90.
81. Perkins JA. New frontiers in our understanding of lymphatic malformations of the head and neck: natural history and basic research. *Otolaryngol Clin North Am* 2018; 51:147–158.
82. Valenzuela I, Fernández-Alvarez P, Plaja A, *et al.* Further delineation of the SOX18-related hypotrichosis, lymphedema, telangiectasia syndrome (HTLS). *Eur J Med Genet* 2018; 61:269–272.
83. Pujol F, Hodgson T, Martínez-Corral I, *et al.* Dachous1-Fat4 signaling controls endothelial cell polarization during lymphatic valve morphogenesis-brief report. *Arterioscler Thromb Vasc Biol* 2017; 37:1732–1735.
84. Güneş N, Taşdemir E, Jeffery H, *et al.* A Novel Mutation of KIF11 in a Child with 22q11.2 Deletion Syndrome Associated with MCLMR. *Mol Syndromol* 2019; 9:266–270.

85. Au AC, Hernandez PA, Lieber E, *et al*. Protein tyrosine phosphatase PTPN14 is a regulator of lymphatic function and choanal development in humans. *Am J Hum Genet* 2010; 87:436–444.
86. Cetinkaya A, Xiong JR, Vargel İ, *et al*. Loss-of-function mutations in ELMO2 cause intraosseous vascular malformation by impeding RAC1 signaling. *Am J Hum Genet* 2016; 99:299–317.
87. Yılmaz B, Toktaş ZO, Akakin A, *et al*. Familial occurrence of brain arteriovenous malformation: a novel ACVRL1 mutation detected by whole exome sequencing. *J Neurosurg* 2017; 126:1879–1883.
88. Wang K, Zhao S, Liu B, *et al*. Perturbations of BMP/TGF- β and VEGF/VEGFR signalling pathways in nonsyndromic sporadic brain arteriovenous malformations (BAVM). *J Med Genet* 2018; 55:675–684.
89. Duran D, Zeng X, Jin SC, *et al*. Mutations in chromatin modifier and ephrin signaling genes in vein of galen malformation. *Neuron* 2019; 101:429.e4–43.e4.
90. Ozsu E, Sen A, Ceylaner S. A case of Riley Ruvalcaba syndrome with a novel PTEN mutation accompanied by diffuse testicular microlithiasis and precocious puberty. *J Pediatr Endocrinol Metab* 2018; 31:95–99.