## CS6190: Probabilistic Modeling Homework 2 Bayesian Networks

Gopal Menon 09 March, 2018

## Written Part

1. The total probability of the network is given by

```
p(Beysian\ Network) = p(income)p(smoke|income)p(exercise|income)p(bmi|income, exercise) \\ \times p(blood\ pressure|exercise, income, smoking)p(cholesterol|exercise, income, smoking) \\ \times p(diabetes|bmi)p(stroke|bmi, bp, cholesterol)p(attack|bmi, bp, cholesterol) \\ \times p(angina|bmi, bp, cholesterol)
```

The total probability distribution requires the product of 10 random variables. If the probabilities are multiplied as shown above, the probability table will have 10 variables. Number of allowed states for the random variables are:

Table 1: CDC Survey Variable states

Survey Variable	Number of states
income	8
exercise	2
smoke	2
bmi	4
bp	4
cholesterol	2
angina	2
stroke	2
attack	2
diabetes	4

The maximum number of rows in the probability distribution table for the above variables will be the product of the number of allowed states -  $8 \times 2 \times 2 \times 4 \times 4 \times 2 \times 2 \times 2 \times 2 \times 4 = 32768$ . This is the number of probabilities that are needed to store the full joint distribution.

- 2. For each of the four health outcomes (diabetes, stroke, heart attack, angina)
  - (a) Probability of the outcome if I have bad habits (smoke and don't exercise)? How about if I have good habits (don't smoke and do exercise)?

Table 2: Health Outcome Probabilities based on Habits

Health Outcome	Bad Habits	Good Habits
diabetes	0.1593304	0.1350624
stroke	0.0501267	0.0368082


Health Outcome	Bad Habits	Good Habits
attack	0.0724154	0.0510272
angina	0.0777561	0.0523189

(b) What is the probability of the outcome if I have poor health (high blood pressure, high cholesterol, and overweight)? What if I have good health (low blood pressure, low cholesterol, and normal weight)?

Table 3: Health Outcome Probabilities based on Health

Health Outcome	Bad Health	Good Health
diabetes	0.1222794	0.0616345
stroke	0.0839749	0.0138704
attack	0.1343300	0.0158879
angina	0.1531853	0.0128387

3. Evaluate the effect a person's income has on their probability of having one of the four health outcomes (diabetes, stroke, heart attack, angina). For each of these four outcomes, plot their probability given income status (your horizontal axis should be i = 1, 2, ..., 8, and your vertical axis should be P(y = 1 | income = i), where y is the outcome). What can you conclude?



It looks like overall the probability of a bad health outcome reduces with increasing income. However, it looks like the probability first increases from the lowest to the next highest income and then reduces.

4. Notice there are no links in the graph between the habits (smoking and exercise) and the outcomes. What assumption is this making about the effects of smoking and exercise on health problems? Let's

test the validity of these assumptions. Create a second Bayesian network as above, but add edges from smoking to each of the four outcomes and edges from exercise to each of the four outcomes. Now redo the queries in Question 2. What was the effect, and do you think the assumptions of the first graph were valid or not?

I think the assumption made in the graph is that the habits (smoking and exercise) have an impact on the health (bmi, bp and cholesterol) of the person and thus indirectly impact the outcomes (diabetes, stroke, attack and angina).

Table 4: Health Outcome Probabilities based on Habits

Health Outcome	Bad Habits	Good Habits
diabetes	0.2267429	0.1025276
stroke	0.0790277	0.0253386
attack	0.1175417	0.0303845
angina	0.1142626	0.0359644

Table 5: Health Outcome Probabilities based on Health

Health Outcome	Bad Health	Good Health
diabetes	0.1310737	0.0576544
stroke	0.0857540	0.0133544
attack	0.1361617	0.0152652
angina	0.1548693	0.0124368

5. Also notice there are no edges between the four outcomes. What assumption is this making about the interactions between health problems? Make a third network, starting from the network in Question 4, but adding an edge from diabetes to stroke. For both networks, evaluate the following probabilities:

$$P(\text{stroke} = 1|\text{diabetes} = 1) \text{ and } P(\text{stroke} = 1|\text{diabetes} = 3)$$

Again, what was the effect, and was the assumption about the interaction between diabetes and stroke valid?

Table 6: Probability for stroke when diabetes is present

Probability	stroke
$ \begin{array}{c} 0.0764278 \\ 0.9235722 \end{array} $	Will have stroke Will not have stroke

Table 7: Probability for stroke when diabetes is not present

Probability	stroke
0.0358626	Will have stroke
0.9641374	Will not have stroke

6. Finally, make sure that your code runs correctly on all of the examples in BayesNetExamples.r. Your code will be graded for correctness on these also.

Table 8: Bishop 8.30 - p(G = 0)

probs	gauge
0.685	1
0.315	0

Table 9: Bishop 8.31 - p(G = 0|F = 0)

probs	gauge	fuel
0.19	1	0
0.81	0	0

Table 10: Bishop 8.32 - p(F = 0|G = 0)

probs	gauge	fuel
0.7428571	0	1
0.2571429	0	0

Table 11: Bishop 8.33 - p(F = 0|G = 0, B = 0)

probs	gauge	battery	fuel
0.8888889	0	0	1
0.1111111	0	0	0

Table 12: Kevin Murphy - Pr(S=1|W=1)

probs	wet	sprinkler
0.5702364	Т	F
0.4297636	${\rm T}$	Τ

Table 13: Kevin Murphy - Pr(R=1|W=1)

probs	wet	rain
0.2920723	Τ	F
0.7079277	${ m T}$	$\mathbf{T}$

Table 14: Kevin Murphy - Pr(W=1)

probs	we
0.3529	F
0.6471	Τ

Table 15: Kevin Murphy -  $\Pr(S{=}1|W{=}1{,}R{=}1)$ 

probs	wet	sprinkler	rain
0.805501	Т	F	Т
0.194499	$\mathbf{T}$	T	Τ