Written Assignment #1 CS 6530 Fall 2016

Gopal Menon

September 4, 2016

- 1. **Problem 1.** Relational Algebra [28pts] Consider the dvdrental schema from the dvdrental database on our database server: http://www.postgresqltutorial.com/postgresql-sample-database/. Answer the following queries in relational algebra:
 - a) Find the customer ids who have rented all films.

The customer ids who have rented all films can be found by dividing a relation containing customer id and film rentals for the customer, by a relation containing all film ids. Let R1, I1 and F1 be instances of the rental, inventory and film relations. The following will give the customer ids who have rented all films.

```
\pi_{customer\_id,film\_id}(R1 \bowtie_{R1.inventory\_id=I1.inventory\_id} I1)/\pi_{film\_id}(F1)
```

b) Print the store id and the phone number for each store that has only one staff member.

Let *STO*, *ADR* and *STA* be instances of the store, address and staff relations. In order to find stores with only one staff member, I will subtract the set of stores that have at least two staff members from the set of all stores. I am assuming that since each store has a manager staff id, each store will at least have one staff member who is a manager.

The StorePhone relation contains stores id and associated phone for all stores. $\rho(StorePhone, \pi_{store_id, phone}(STO \bowtie_{STO.address_id=ADR.address_id}ADR))$

The StoreStaff relation contains all staff members associated with a store

 $\rho(StoreStaff, \pi_{store_id,phone,staff_id}(StorePhone \triangleright \triangleleft_{StorePhone.store_id=STA.store_id}STA))$

The *StoresStaff Pair* relation contains the cross product of the *StoreStaff* relation with itself.

```
\rho(StoresStaffPair(1 \rightarrow store\_id1, 2 \rightarrow phone1, 3 \rightarrow staff\_id1, 4 \rightarrow store\_id2, 5 \rightarrow phone2, 6 \rightarrow staff\_id2), StoreStaff \times StoreStaff)
```

The *PluralStaf f Stores* relation contains stores ids and associated phone numbers of stores with at least two staff members.

```
\rho(PluralStaffStores, \pi_{store\_id1, phone1}(\sigma_{store\_id1=store\_id2} \land staff\_id1 \neq staff\_id2}StoresStaffPair))
```

The *StoresWithOneStaff* relation is the difference between all stores and stores with at least two staff members. It will thus have the store id and associated phone numbers for stores having only one staff member.

```
\rho(StoresWithOneStaff,StorePhone-PluralStaffStores)
```

c) Find every pair of customer and staff who are from the same city, by listing their customer id and staff id.

This can be done by joining customer address and staff address on city id. Let *CST*, *STF* and *ADR* be instances of the customer, staff and address relations.

```
\rho(CustomerCity, \pi_{customer\ id, city\ id}(CST) \bowtie_{CST.address\ id=ADR.address\ id}ADR)
```

```
\rho(StaffCity, \pi_{staff\_id,city\_id}(STF \bowtie_{STF.address\_id=ADR.address\_id}ADR)
```

The relations *CustomerCity* and *StaffCity* will contain the customer and staff ids respectively along with the city id of their address. If we do a natural join on these two relations, we can obtain a relation with customer id and staff id that are from the same city.

```
\pi_{customer\_id,staff\_id}(CustomerCity {\,\vartriangleright\,} StaffCity)
```

- d) Find the customer ids who *have rented all and only those* films acted by Emily Dee.
- 2. **Problem 2.** [28pts] Answer following queries using Tuple Relational Calculus.
 - a) Find the customer ids who have rented all films.
 - b) Print the store id and the phone number for each store that has only one staff member.
 - c) Find every pair of customer and staff who are from the same city, by listing their customer id and staff id.
 - d) Find the customer ids who *have rented all* films acted by Emily Dee (slightly modified from Problem 1).