

CS 7910 Computational Complexity

Assignment 5

Gopal Menon

March 1, 2016

1. (20 points) In this exercise, we consider the following *Partition Problem*.
Given a set A of n numbers, the *partition problem* is to decide whether A can be partitioned into two subsets A_1 and A_2 (i.e., $A = A_1 \cup A_2$ and $A_1 \cap A_2 = \phi$) such that the sum of the numbers in A_1 is equal to the sum of the numbers in A_2 .
For example, suppose $A = \{5, 9, 3, 12, 11\}$; then we can partition A into $A_1 = \{5, 3, 12\}$ and $A_2 = \{9, 11\}$.
Prove that the partition problem is NP-Complete (Hint: by a reduction from the subset-sum problem).

We know that the subset sum problem is NP-Complete. Let an instance of the subset sum problem be (U, M) , where U is a set of numbers and we need to find a set of numbers S such that $S \subseteq U$ and $\sum S = M$.

Let us construct an instance of the partition problem (A) , where A is a set of numbers that can be partitioned into two sets such that the elements of each set add up to the same number. Construct the set A in such a manner that it consists of all the elements in set U with the addition of an element that has a value $\sum U - 2 * M$. So the set A will have one more element than the set U .

This construction can clearly be done in polynomial time. Also, given a certificate of the partition problem, which will be a subset of A , we can verify in $O(|A|)$, or polynomial time, that the certificate is correct. This shows that the partition problem is in the set NP.

Consider the case where the instance of the subset sum problem is true. In the case of the partition problem that was constructed, we can create a subset S' such that $S' = S \cup (\sum U - 2 * M)$, where $\sum U - 2 * M$ is the value of the additional element in set A , when compared with set U . The elements in set S' will sum up to $\sum S + \sum U - 2 * M$ or $M + \sum U - 2 * M$, since $\sum S = M$. The sum of the elements in S' will reduce to $\sum U - M$. The sum of all the elements in set A will be $\sum U + (\sum U - 2 * M)$ or $2 * \sum U - 2 * M$. This can be simplified as $2 * (\sum U - M)$. This means that the sum of the elements in set A is twice the sum of the elements in set S' . Or in other words, the set A can be partitioned into two sets that sum up to the same number.

Consider the case where the set A can be partitioned into two sets that sum up to the same number. We know that the set A has been constructed by doing $U \cup (\sum U - 2 * M)$. We do not know whether there exists a subset of U that adds up to M . Each partition of set A adds up to the same amount $\sum U - M$. Consider the case of the subset sum instance. The two partitions of the set A , will be mapped to two portions of the set U , which will sum up to $\sum U - M$ and $(\sum U - M) - (\sum U - 2 * M)$. The sum of the second portion of the set U may be simplified to M . This shows that if the partition problem instance is true, the subset sum instance is also true.

So we have shown that subset sum can be reduced to the partition problem in polynomial time. This means that the partition problem is in NP-Hard. Since we already know that the partition problem is in NP, it follows that the partition problem is in NP-Complete.