

# A hazards approach to the biometric analysis of infant mortality

Göran Broström<sup>1</sup> Tommy Bengtsson<sup>2</sup>

<sup>2</sup>CED, Lund University (tommy.bengtsson@ekh.lu.se)

<sup>1</sup>CEDAR, Umeå University (gb@ehar.se, <https://github.com/goranbrostrom>)

# Outline

- Bourgeois-Pichat (1951): Biometric analysis of IM (background).
- Hazard-based alternative.
- Theoretical considerations
- Real-world examples.

# Bourgeois-Pichat and causes of death

- Two categories:
  - Endogenous: *inherited, delivery*, etc.
  - Exogenous: *accidental, infectious diseases*, etc.
- How to differentiate between the two categories without information of causes of death?

Bourgeois-Pichat's Biometric model.

# The biometric model

Two postulates:

- ① Endogenous deaths only occur during the neonatal period (0–28 days).
- ② On a specific time scale, exogenous infant mortality is uniformly distributed.


# The log-cube transform

$$g(t) = C \log^3(t + 1), \quad 0 < t \leq 365.$$

where  $C$  is a normalizing constant:


$$C = \frac{365}{\log^3(365 + 1)}$$

*Note:*  $C$  is not part of B-P's original definition.



# Demonstration of the B-P plot


Assume 100 births, of which 90 survives infancy, ten death ages:  
2, 6, 10, 15, 25, 41, 90, 120, 180, 309.



Calculation (no ties):  $\frac{1}{100}, \frac{2}{100}, \frac{3}{100}, \dots, \frac{10}{100}$ .

# Demonstration of the B-P plot II


Assume 100 births, of which 90 survives infancy, ten death ages:  
2, 6, 10, 15, 25, 41, 90, 120, 180, 309.



Calculation (no ties):  $\frac{1}{100}, \frac{2}{100}, \frac{3}{100}, \dots, \frac{10}{100}$ .

# The hazards plot


Assume 100 births, of which 90 survives infancy, ten death ages:  
2, 6, 10, 15, 25, 41, 90, 120, 180, 309 (same as before).



Calculation (no ties):  $\frac{1}{100}, \frac{2}{99}, \frac{3}{98}, \dots, \frac{10}{91}$ .




# The hazards plot II



$$\text{Exogenous slope} = \frac{\#(\text{post deaths})}{\text{post exposure}} = \frac{5}{27626} \approx 0.00018$$

$$\text{Post exposure} = 90(365 - 68) + (93 - 68) + (163 - 68) + \dots + (335 - 68) \approx 27626$$


# Adding the endogenous mortality




- Knodel & Kintner (1977).
  - Discusses effects of breastfeeding habits.
- Wrigley (1977).
  - Problems with data recording.
- Lynch, Greenhouse & Brändström (1998).
  - Mentions “constant hazard”.
  - Suggest linear regression methods.
- Bengtsson (1999).
  - Data quality.
- Manfredini (2004).
  - Deals with effect of climate variation.

## Study areas

## Skåne 1711–1800




Västerbotten 1801–1950




# Västerbotten 1861-1890

Hazards method




B-P method




# Västerbotten 1921–1950


Hazards method



B-P method




# Cohort effect of birth month, Västerbotten



Conclusion:

Post-neonatal linearity is a population property, not an individual.

# Period effect of month, Västerbotten




Looks quite proportional!




# Scania 1711–1800


Hazards method




B-P method



### Hazards method



### B-P method



# Conclusions

- Good