

A biometric analysis of infant mortality and temperature,
northern Sweden 1895-1950

Göran Broström and Tommy Bengtsson

2022-01-25 00:43:55

Abstract

The effect of extreme temperatures on infant mortality in the Umeå region 1895-1950 is studied in a biometric analysis setting. More precisely, the effect of climate and weather, measured by temperature, on infant mortality is investigated. It turns out that climate is more important than weather, low average temperatures are more important than temporary dips in temperature. The investigated geographical region is situated too far north for bad effects of high temperatures to show.

Contents

1	Introduction	1
2	Data	2
2.1	Infant mortality	2
2.2	Temperature	3
2.3	Mortality and temperature as a communal covariate	4
3	Statistical modelling	4
4	Postneonatal mortality	4
5	Neonatal mortality	5
5.1	Girls	5
5.2	Boys	5
6	Postneonatal mortality, winter and spring	6
7	Neonatal mortality, winter and spring	6
7.1	Girls	6
7.2	Boys	6
8	Conclusion	7

1 Introduction

The impact of ambient temperature variations on infant mortality is studied for a northern Sweden coastal area, the Umeå and Skellefteå regions, during the first half of the twentieth century. Two recent papers (Junkka et al., 2021; Karlsson et al., 2021) studied neonatal mortality and temperature variations in a larger geographical area containing the present one during the years 1880–1950. Climate and mortality in general is a research area that has generated great interest over the last years, see Bengtsson and Broström (2010).

The effect of seasonal variation and the occurrence of extreme monthly temperatures is studied and interacted with sex, social class, and legitimacy. Studies are performed separately for neonatal and postneonatal mortality, and for winter and summer seasons, and the classification into endogenous and exogenous factors will be discussed.

One important reason for studying neonatal and postneonatal mortality separately is the empirical findings by Bourgeois-Pichat (Bourgeois-Pichat, 1951a,b) about endogenous and exogenous mortality and the log-cube transform.

2 Data

We have two sources of data which we combine into one data set suitable for our purpose. The first is demographic data obtained from the *Centre for Demographic and Ageing Research* (CEDAR, <https://cedar.umu.se>), the second is daily temperature measurements obtained from the *Swedish Meteorological and Hydrological Institute* (SMHI, <https://www.smhi.se>).

2.1 Infant mortality

Individual data with all births between 1 January 1895 and 31 December 1950 in two coastal areas of north Sweden, Skellefteå (51560 births) and Umeå (31213 births). They were followed until death or age 365 days, whichever came first. The following *static* characteristics were observed on each child:

birthdate Date of birth.

sex Girl or boy.

exit Number of days under observation.

event Logical, *TRUE* if a death is observed.

socBranch Working branch of father (if any).

socStatus Social status of family, based on HISCLASS.

illeg Mother unmarried?

parity Order among siblings.

Some crude statistics about infant, neonatal, and postneonatal mortality are shown in Figures.

Figure 1 shows the average monthly crude infant mortality, and a clear seasonal pattern is visible.

[Figure 1 about here.]

The average monthly neonatal mortality is shown in Figure 2.

[Figure 2 about here.]

The seasonal pattern is similar to the one we found above for infant mortality.

The average monthly postneonatal mortality is shown in Figure 3.

[Figure 3 about here.]

The seasonal pattern is once again similar to the one we found for infant mortality. Next, the decline over the years in Figures 4 and 5.

[Figure 4 about here.]

[Figure 5 about here.]

Table 1: Raw temperature data from first week of 1923, Umeå.

Date	Time	Temperature	Quality
1923-01-01	07:00:00	0.4	G
1923-01-01	13:00:00	0.6	G
1923-01-01	20:00:00	0.0	G
1923-01-02	07:00:00	-1.4	G
1923-01-02	13:00:00	-1.4	G
1923-01-02	20:00:00	-1.2	G
1923-01-03	07:00:00	0.4	G
1923-01-03	13:00:00	0.8	G
1923-01-03	20:00:00	1.2	G
1923-01-04	07:00:00	1.4	G
1923-01-04	13:00:00	1.2	G
1923-01-04	20:00:00	1.0	G
1923-01-05	07:00:00	-1.4	G
1923-01-05	13:00:00	-3.2	G
1923-01-05	20:00:00	-3.4	G
1923-01-06	07:00:00	1.0	G
1923-01-06	13:00:00	0.4	G
1923-01-06	20:00:00	0.4	G
1923-01-07	07:00:00	0.6	G
1923-01-07	13:00:00	0.4	G
1923-01-07	20:00:00	0.4	G

Table 2: Weekly summarized temperature data: Umeå 1923, first week.

week	year	mintemp	maxtemp	meantemp	emintemp	emaxtemp	emeantemp
1	1923	-3.4	1.4	-0.1	-17.73	-0.36	-7.54

2.2 Temperature


Temperature data are collected from two weather stations, *Umeå* and *Bjuröklubb* (used with population data from Skellefteå coastal area). Both stations deliver daily temperature data covering our time period, usually three measures per day, morning, noon, and evening. In Table 1, the Umeå data from the week 1–7 January, 1923 is shown.

There are three measurements per day, or 21 per week. In the forthcoming analyses, the weekly data are summarized in a few measurements, see Table 2.

Weekly averages are calculated year by year, and deviations from the averages of the weekly averages are used as a time-varying *communal covariate*. See Figure 6 for the monthly distributions.

[Figure 6 about here.]

Time trends of yearly average temperatures, see Figure



2.3 Mortality and temperature as a communal covariate

The two data sets, mortality and weather, are combined into one by treating temperature data as a communal covariate and incorporate it as such in the mortality data set.

3 Statistical modelling

The analyses are performed on the log-cube scale with proportional hazards modelling. Note that the property of proportional hazards are preserved under a strictly monotone increasing time transform. However, the estimates of baseline distribution characteristics will change, of course.

We start by performing separate analyses for neonatal and postneonatal mortality. Then we make comparisons.

4 Postneonatal mortality

This is the simplest part, because $B-P$ predicts that the baseline distribution (given the log-cube transformation) is *exponential*, that is, the hazard function is *constant*.

Table 3: Postneonatal mortality, Skellefteå. Adjusted for social status and branch, illegitimacy, parity, time period, season, and sub region.

Covariate	Mean	Coef	H.R.	S.E.	L-R p
excessCold	1.443	0.037	1.037	0.008	0.000
emintemp	-1.887	-0.019	0.981	0.006	0.001
Events	1835	TTR	8020541		
Max. logLik.	-16974				

The result in Table 3 shows that *climate* (**emintemp**) is more important than *weather* (**cold**). Moreover, no signs of interaction between weather or climate and the rest of covariates (not shown).

Table 4: Postneonatal mortality, Umeå. Adjusted for social status and branch, illegitimacy, parity, time period, season, and subregion.

Covariate	Mean	Coef	H.R.	S.E.	L-R p
excessCold	1.628	0.008	1.008	0.009	0.405
emintemp	-2.866	-0.016	0.984	0.006	0.005
Events	1449	TTR	4759837		
Max. logLik.	-12875				

5 Neonatal mortality

It turns out that here we have some significant interactions, with **sex** involved. Maybe surprising. So we run separate analyses for the sexes.

5.1 Girls

Table 5: Neonatal mortality, girls. Adjusted for social status and branch, illegitimacy, parity, time period, and sub region.

Covariate	Mean	Coef	H.R.	S.E.	L-R p
cold	-0.002	0.003	1.003	0.007	0.715
emintemp	-2.129	-0.007	0.993	0.003	0.015
Events	1143	TTR	1461345		
Max. logLik.	-12007				

Same here, *climate* is more important than *weather*.

5.2 Boys

Table 6: Neonatal mortality, boys. Adjusted for social status and branch, illegitimacy, parity, time period, and sub region.

Covariate	Mean	Coef	H.R.	S.E.	L-R p
cold	0.022	-0.010	0.990	0.006	0.091
emintemp	-2.263	-0.010	0.990	0.003	0.000
Events	1494	TTR	1539122		
Max. logLik.	-15792				

Here, however, boys are sensitive to temperatures below average, in addition to the sensitivity for cold climate.

6 Postneonatal mortality, winter and spring

This is the simplest part, because $B-P$ predicts that the baseline distribution (given the log-cube transformation) is *exponential*, that is, the hazard function is *constant*.

Table 7: Postneonatal mortality, winter and spring. Adjusted for social status and branch, illegitimacy, parity, time period, and sub region.

Covariate	Mean	Coef	H.R.	S.E.	L-R p
cold	0.020	-0.015	0.985	0.004	0.001
emintemp	-9.099	-0.013	0.987	0.003	0.000
Events	1936	TTR	6362768		
Max. logLik.	-17324				

The result in Table 7 shows that *climate* (**emintemp**) is more important than *weather* (**cold**). Moreover, no signs of interaction between weather or climate and the rest of covariates (not shown).

7 Neonatal mortality, winter and spring

It turns out that here we have some significant interactions, with **sex** involved. Maybe surprising. So we run separate analyses for the sexes.

7.1 Girls

Table 8: Neonatal mortality winter and spring, girls. Adjusted for social status and branch, illegitimacy, parity, time period, and sub region.

Covariate	Mean	Coef	H.R.	S.E.	L-R p
cold	-0.021	0.005	1.005	0.008	0.505
emintemp	-9.036	-0.011	0.989	0.006	0.073
Events	613	TTR	731148		
Max. logLik.	-6012				

Same here, *climate* is more important than *weather*.

7.2 Boys

Here, however, boys are sensitive to temperatures below average, in addition to the sensitivity for cold climate.

Table 9: Neonatal mortality winter and spring, boys. Adjusted for social status and branch, illegitimacy, parity, time period, and sub region.

Covariate	Mean	Coef	H.R.	S.E.	L-R p
cold	0.016	−0.009	0.991	0.007	0.194
emintemp	−9.021	−0.007	0.993	0.005	0.165
Events	822	TTR	777881		
Max. logLik.	−8128				

8 Conclusion

Remains to be written. Feels like two papers, one about Bourgeois-Pichat and one about climate and weather. Maybe discuss exogeneity/endogeneity?

References

- Bengtsson, T. and Broström, G. (2010). Mortality crisis in rural southern Sweden 1766–1860. In Kurosu, T., Bengtsson, T., and Campbell, C., editors, *Demographic Response to Economic and Environmental Crisis*, pages 1–16. Reitaku University Press, Kashiwa.
- Bourgeois-Pichat, J. (1951a). La mesure de la mortalité infantile. I. Principes et méthodes. *Population (French Edition)*, 6:233–248.
- Bourgeois-Pichat, J. (1951b). La mesure de la mortalité infantile. II. Les causes de décès. *Population (French Edition)*, 6:459–480.
- Junkka, J., Karlsson, L., Lundevaller, E., and Schumann, B. (2021). Climate vulnerability of Swedish newborns: Gender differences and time trends of temperature-related neonatal mortality, 1880–1950. *Environmental Research*, 192. article id 110400.
- Karlsson, L., Junkka, J., Schumann, B., and Häggström Lundevaller, E. (2021). Socioeconomic disparities in climate vulnerability: neonatal mortality in northern Sweden, 1880–1950. *Population and Environment*, 10.1007.

List of Figures

1	Crude infant mortality by week of year, Umeå/Skellefteå 1895–1950.	9
2	Crude neonatal mortality by week of year, Umeå/Skellefteå 1895–1950.	9
3	Crude postneonatal mortality by week of year, Umeå/Skellefteå 1895–1950. .	10
4	Crude IMR by year, Umeå-Skellefteå 1895–1950.	10
5	Crude NMR by year, Umeå-Skellefteå 1895–1950.	11
6	Weekly max and min temperature averages, 1895–1950.	11




Figure 1: Crude infant mortality by week of year, Umeå/Skellefteå 1895–1950.




Figure 2: Crude neonatal mortality by week of year, Umeå/Skellefteå 1895–1950.




Figure 3: Crude postneonatal mortality by week of year, Umeå/Skellefteå 1895–1950.




Figure 4: Crude IMR by year, Umeå-Skellefteå 1895–1950.




Figure 5: Crude NMR by year, Umeå-Skellefteå 1895–1950.




Figure 6: Weekly max and min temperature averages, 1895–1950.