



Please be sure to mark all answers for **all** problems on this final on your scantron sheet!! Every single question! There is not one single long answer question!

1. [5 Points] An observer sees a system moving past her at a relativistic speed that consists of a mass attached to the end of a spring, oscillating. She measures a period T_1 . A second observer, who is moving with the spring-mass system, measures a period T_2 . Which of the following is true:

- A. $T_1 > T_2$
- B. $T_1 < T_2$
- C. $T_1 = T_2$

D. Not enough information to tell.

$$T_2 - \text{proper time.}$$

$$\therefore T_1 > T_2$$

2. [5 Points] In frame S, event B occurs $5.5 \mu\text{s}$ after event A ($5.5 \times 10^{-6} \text{ s}$). Event A occurs at the origin, and event B occurs along the positive x axis at position L. An observer traveling at a speed $v = 0.9c$ observes the two events to be simultaneous. Determine the value of L in frame S.

- A. $1.3 \times 10^{-8} \text{ km}$
- B. 0.80 km
- C. 1.8 km
- D. 4.2 km
- E. $5.5 \times 10^8 \text{ km}$

$$t_1' = \gamma(t_1 + \frac{x_1 v}{c^2})$$

$$t_2' = \gamma(t_2 + \frac{x_2 v}{c^2})$$

$$t_1' = t_2'$$

$$\Rightarrow \gamma t = L \frac{\vec{v}}{c^2}$$

$$L = \frac{c^2}{v} \gamma t = \frac{c}{0.9} \Delta t = 1.8 \text{ km}$$

3. [5 Points] At what speed is a particle moving when its kinetic energy is 2.5 times its rest energy?

- A. $0.85 c$
- B. $0.91 c$
- C. $0.92 c$
- D. $0.96 c$

E. Not enough information to determine.

4. [4 Points] Homer remains on earth while Ulysses travels 6 light years to planet P and turns around and comes back. Assume Ulysses spends almost all his time traveling at $0.6c$. How much does Homer predict that his clock will advance while Ulysses flies to planet P?

- A. 1 yr.
- B. 5 yr.
- C. 10 yr.
- D. 15 yr.
- E. 20 yr.

$$L = vt \Rightarrow t = 10 \text{ yr} \quad \text{There is no ref frame.}$$

5. [4 Points] Ulysses, in the spaceship reference frame, calculates how much Homer's clock will advance while he makes the trip from earth to planet P. How long does he think Homer will think it takes?

- A. 6.4 yr.
- B. 8.0 yr.
- C. 8.5 yr.
- D. 10 yr.
- E. 13 yr.

U is still - so expects Homer's clock to go slower

$$\Delta t_H = \gamma \Delta t_U$$

Δt_H is how long it takes U .

Homer knows that
 $\Delta t_H = \frac{\Delta t_{H \text{ real}}}{\gamma}$

$$(10 \text{ yr}) (\sqrt{1 - 0.6^2}) = 8 \text{ yr}$$

$$\Delta t_H = (8 \text{ yr})(0.8) = 6.4 \text{ yr}$$

Name _____


last first

Student ID _____

Score _____



6. [4 Points] Which of the following features correctly explains how the Photoelectric Effect experiment/demo done in class makes the case that light is quantized (as photons)? *(8 to make off e = > 80)*
 A. Coherent light shown through two slits shows an interference pattern.
 B. White light shown through a diffraction grating shows as a rainbow on a screen.
 C. Charge is lost from the experiment's plate faster when the light is brighter
 D. Light longer than a certain wavelength will not discharge the plate
 E. Light shorter than a certain wavelength will not discharge the plate.
7. [5 Points] For a particle in a box, the energy levels are denoted E_n . Approximately, what is $(E_{n+1} - E_n)/E_n$ dependence on n for very large n ?
 A. n^2 $E_n = n^2 E_1$ $\frac{E_{n+1} - E_n}{E_n} = \frac{(n+1)^2 - n^2}{n^2} = \frac{n^2 + 2n + 1 - n^2}{n^2} = \frac{2n + 1}{n^2}$
 B. n
 C. 1
 D. n^{-1}
 E. n^{-2}
8. [4 Points] Standing waves result from the superposition of two traveling waves that have:
 A. The same amplitude, frequency, and direction of propagation
 B. The same amplitude, frequency, and opposite directions of propagation
 C. The same amplitude, slightly different frequencies, and the same propagation direction
 D. The same amplitude, slightly different frequencies, and the opposite directions of propagation
 E. None of the above
9. [5 Points] The fundamental frequency of a pipe that has both ends closed is 230 Hz. The speed of sound for this problem is $v_s = 340$ m/s. When both ends of the same pipe are opened, the fundamental frequency is:
 A. 58 Hz
 B. 230 Hz
 C. 460 Hz
 D. 920 Hz
 E. 1400 Hz
10. [5 Points] Tritium is an unstable isotope of hydrogen with two neutrons and one proton. It has a half-life of 12.3 years. Suppose 1.0×10^{10} atoms of tritium were stored in your freshman year at UW. What is the decay rate per minute four years later?
 A. 856 per minute
 B. 1070 per minute
 C. 1120 per minute
 D. 15200 per minute
 E. 813000000 per minute



$$\frac{1}{2}\lambda = L \Rightarrow \text{freq is same}$$

$$\frac{1}{2}\lambda = L$$

$$R = R_0 e^{-\lambda t}$$

$$R_0 = \lambda N_0 \quad N_0 = 10^{10}$$

$$\lambda = \frac{\ln 2}{t_{1/2}} = \frac{\ln 2}{12.3 \text{ yr}} = 0.05633$$

$$R_0 = \lambda N_0 = 563539273$$

$$\therefore R = R_0 e^{-\lambda \cdot 4 \text{ yr}} = 449805152/\text{yr}$$

convert to /min $\Rightarrow 75.8/\text{min}$



For the next three questions, consider a traveling wave has the form:

$$f(x, t) = A \sin(kx - \omega t + \phi)$$

The wave's frequency is $f = 1300$ Hz and its speed is 340 m/s.

11. [5 Points] What is the value of k ?

- A. 1.64 per m
- B. 3.82 per m
- C. 11.2 per m
- D. 24.0 per m
- E. 41.8 per m

$$k = \frac{2\pi}{\lambda} = \frac{2\pi f}{v} = \frac{2\pi \cdot 1300 \text{ Hz}}{340 \text{ m/sec}} = 24.0 \text{ 1/m}$$


12. [5 Points] What is the value of ω ?

- A. 559 rad per sec
- B. 1300 rad per sec
- C. 3820 rad per sec
- D. 8170 rad per sec
- E. 14200 rad per sec

$$\omega = kv = 2\pi \cdot 1300 \text{ Hz} = 8168 \text{ rad/sec}$$

13. [5 Points] Assuming both ω and k are positive, which direction is the wave traveling?

- A. The wave moves perpendicular to the x -direction.
- B. The wave moves first to the right, and then to the left.
- C. The wave moves in the negative x -direction.
- D. The wave moves in the positive x -direction.
- E. There is not enough information to tell.



14. [4 Points] A mass is hung from a spring. You pull the mass down, displacing it from its equilibrium position by 30 cm. The mass then oscillates with a frequency $f = 5$ Hz. You repeat the experiment with the same mass and same spring, only you displace it from its equilibrium position by 60 cm. Which of the following is true:

- A. The new frequency is half of f .
- B. The new frequency is the same as f .
- C. The new frequency is double f .
- D. The new frequency is $\sqrt{2}f$.
- E. Not enough information to determine.

$$\omega = \sqrt{\frac{k}{m}} - \text{not depend on } A!$$

15. [5 Points] A bat flying at 5.0 m/s towards a wall emits a chirp at 50.0 kHz. If the wall reflects this sound pulse, what is the frequency of the echo received by the bat? For this problem, $v_{\text{sound}} = 340$ m/s and that the wall perfectly reflects the frequency it is hit with. Fly towards source \Rightarrow freq higher!


- A. 48.6 kHz
- B. 49.3 kHz
- C. 50.0 kHz
- D. 51.5 kHz
- E. None of the above.

Doppler effect $\times 2$

$$f_r = \frac{v + v_r}{v + v_s} f_s$$

$$f_{\text{wall}} = \frac{v}{v - v_{\text{bat}}} f_{\text{bat}}$$

$$\begin{aligned} f_{\text{ret}} &= \frac{v + v_{\text{bat}}}{v} f_{\text{wall}} = \frac{v + v_{\text{bat}}}{v - v_{\text{bat}}} f_{\text{bat}} \\ &= \frac{345}{335} 50 \text{ kHz} = 51.5 \text{ kHz} \end{aligned}$$



last first

16. [5 Points] Three plane polarizers are arranged in sequence. The first and third are rotated 90 degrees apart from each other, the first polarizer being lined up with the vertical. The second polarizer is rotated 30 degrees from the vertical. What is the intensity of the light as it leaves the third polarizer relative to unpolarized light that is shone on the first polarizer (I_0)?

- A. $0.09 I_0$
 B. $0.11 I_0$
 C. $0.19 I_0$
 D. $0.25 I_0$
 E. $0.50 I_0$

$$I_1 = \frac{1}{2} I_0$$

$$I_2 = I_1 \cos^2(30^\circ)$$


$$I_3 = I_2 \cos^2(90^\circ - 30^\circ) \rightarrow I_3 = \frac{1}{2} I_0 \cos^2(30^\circ) \cos^2(60^\circ) = 0.09 I_0$$



17. [5 Points] The expression for the electric field of a certain EM wave is $E(y, t) = E_0 \sin(ky + \omega t) \hat{z}$. What is the proper expression for the magnetic field?

- A. $B(y, t) = +B_0 \sin(ky + \omega t) \hat{x}$
 B. $B(y, t) = B_0 \sin(ky - \omega t) \hat{x}$
 C. $B(y, t) = -B_0 \sin(ky + \omega t) \hat{y}$
 D. $B(y, t) = B_0 \sin(ky - \omega t) \hat{y}$
 E. $B(y, t) = B_0 \sin(ky - \omega t) \hat{z}$

dir of prop = $E \times B$
 prop is $-\hat{y}$ (+wt)
 E is $+\hat{z}$... $\hat{z} \times \hat{y} = -\hat{y}$




Consider the reflecting optical telescope shown at right for the next two questions. The concave mirror at the bottom has $f = +1.5$ m with a diameter of 0.6 m as its objective. The eyepiece lens has a focal length of $f = +20$ cm, and it is at a distance d from the objective mirror. It is mainly used to observe blue light with $\lambda = 475$ nm.

18. [5 Points] Which is d , the approximate location of the eyepiece from the objective mirror?

- A. 0.2 m
 B. 1.3 m
 C. 1.5 m
 D. 1.7 m
 E. 3.0 m

$$f_1 + f_2 = 1.7 \text{ m}$$



19. [5 Points] What is the smallest angle between two distant stars that can be resolved by the telescope (diffraction limit)?

- A. 3.86×10^{-7} rad
 B. 7.92×10^{-7} rad
 C. 8.12×10^{-7} rad
 D. 9.66×10^{-7} rad
 E. 2.90×10^{-6} rad

$R = \text{first min - for circular aperture}$

$$R = 1.22 \frac{\lambda}{D} = 1.22 \frac{475 \times 10^{-9} \text{ m}}{0.6 \text{ m}} = 9.66 \times 10^{-7} \text{ m}$$



20 [5 points] Light from a source P travels through air ($n_{\text{air}}=1.0$) and reflects from a glass plate ($n_g=1.5$), as shown in figure 1.

In order to see the reflected light from source P, the viewer must place his eye at an angle ω equal to:

- (a) $\theta_i/2t$
- (b) $\pi + \theta_i$
- (c) θ_i
- (d) correct answer not given




Figure 1

21 [5 points] How does the phase of light ray 1 compare to the phase of the incident light ray from source P?

- (a) equal to
- (b) differs by 2π
- (c) differs by π
- (d) differs by $\pi/2$

22 [10 points] The wave length of the light from source P is 550 nm. The observer sees constructive interference and concludes that the thickness t is equal to:

- (a) 183 nm
- (b) 92 nm
- (c) 370 nm
- (d) 122 nm
- (e) 137.5 nm

23 [5 points] The image formed by a compound microscope is:

- (a) upright and magnified
- (b) inverted and magnified
- (c) neither (a) nor (b) is correct.

24 [5 points] Which of the following statements is correct for a simple camera:


- (a) a real image is created at the focal plane of lens
- (b) the image is inverted
- (c) the angular sizes of image and object are the same
- (d) a, b, and c are all correct
- (e) c is not correct



Name _____ Student ID _____
last first

III. [30 points total]

Students A and B stand at rest on the ground. Student A is exactly halfway between mountain 1 and mountain 2; student B's location on the ground is not known. A shuttle flies from right to left with speed $v = 0.7c$ relative to the ground. Assume all motion is along one dimension (*i.e.*, ignore vertical distances), and that all observers are intelligent observers.



Firecrackers 1 and 2 explode at the tops of mountains 1 and 2. Each emits a flash of light. The firecrackers explode *simultaneously* in student A's reference frame.

The figure shows the locations of student A and the shuttle in the student A's reference frame at the instant both firecrackers explode. Call event 1 "firecracker 1 explodes" and event 2 "firecracker 2 explodes."

25. [5 pts] In student B's reference frame:

- A. Event 1 occurs *before* event 2.
- B. Event 1 occurs *after* event 2.
- C. Event 1 occurs *at the same time* that event 2 occurs.
- D. It is impossible to determine the order of events without knowing student B's location.

26. [5 pts] Suppose student C stands in the middle of the shuttle (moving from right to left with a speed of $v = 0.7c$ relative to the ground). In student C's frame:

- A. Event 1 occurs *before* event 2.
- B. Event 1 occurs *after* event 2.
- C. Event 1 occurs *at the same time* that event 2 occurs.
- D. More information is needed to answer.

27. [5 pts] The location of event 1 in student C's frame is correctly determined to be x_1 . In student C's frame, the wavefront of the light from event 1 is:

- A. spherical and centered at location x_1 (which is stationary in student C's frame)
- B. spherical and centered at firecracker 1 (which moves to the left in student C's frame)
- C. non-spherical and centered at location x_1 (which is stationary in student C's frame)
- D. non-spherical and centered at firecracker 1 (which moves to the left in student C's frame).

28. [5 pts] Let Δx_A be the distance between events in student A's reference frame, and Δx_C be the distance between events in student C's reference frame. How does Δx_A compare to Δx_C ?

- A. Δx_c is not constant, and so cannot be compared to Δx_A
- B. $\Delta x_A > \Delta x_C$
- C. $\Delta x_A < \Delta x_C$
- D. $\Delta x_A = \Delta x_C$


Name _____ Student ID _____
last *first*



Monochromatic light is normally incident on a mask containing four very narrow slits. Multiple points of zero intensity are observed on a very distant screen (not shown).


29. [5 pts] Consider point X on the screen, which is the first point of zero intensity to the right of the central maximum. What is the path length difference from slits 1 and 3 to point X (*i.e.*, ΔD_{13} to point X)?

- A. $\lambda/4$
- B. $\lambda/3$
- C. $\lambda/2$
- D. λ
- E. $3\lambda/2$



30. [5 pts] A student in your class draws the diagram shown at right, and notes that the line segment M has a length of $\lambda/2$. (This is the same four-slit mask from the previous question.) Which of the following **best** describes the angle θ shown in the diagram?

- A. θ describes a line of zero intensity.
- B. θ describes a principle maximum.
- C. θ describes either a line of zero intensity or a principle maximum.
- D. θ describes neither a line of zero intensity nor a principle maximum.
- E. There is not enough information to determine what type of line is described by θ .






Problem 22 solution

Light from a source P travels through air ($n_{\text{air}}=1.0$) and reflects from a glass plate ($n_g=1.5$), as shown in figure 1.

22 [10 points] The wave length of the light from source P is 550 nm. The observer sees constructive interference and concludes that the thickness t is equal to:

- (a) 183 nm
- (b) 92 nm
- (c) 370 nm
- (d) 122 nm
- (e) 137.5 nm



Solution

1. Constructive interference occurs when the phase difference between rays 1 and 2 is a multiple of 2π .
2. Solving this requires taking into account the phase change on reflection plus the phase change due to the extra path length of ray 2.
3. Ray 1 has a phase change of π because the light is reflecting from a higher index material than it is traveling in
4. Ray 2 reflects from the bottom surface and has zero phase change because it is “bouncing off” a lower index material than it is traveling in.
5. The path length in the glass plate is given by $2\pi/\lambda'$ where $\lambda' = \lambda/n_g = \lambda/1.5$.
6. The total phase difference between two rays “bouncing off” the front and the back of the thin glass plate of thickness t and index of refraction $n_g=1.5$ is given by equation 3, page 7-7 of the lab manual
$$\phi = \pi + 2t[2\pi/\lambda']$$
7. So $\phi = 2\pi$ when $2t[2\pi/\lambda'] = \pi$ and solving for t gives $t = \lambda'/4 = \lambda/6 = 92 \text{ nm}$
8. If λ rather than the optical path λ' is used one gets 137.5 nm, which is incorrect.