

1.1a) Min $Z = 4x_1 + \sqrt{2}(x_2^+ - x_2^-) - 0.35x_3$
s.t.

$$\begin{aligned} -0.001x_1 + 200(x_2^+ - x_2^-) - c_1 &= 7\sqrt{2}61 \\ -7.07(x_2^+ - x_2^-) + 2.62x_3 - c_2 &= 4 \\ x_1, x_2^+, x_2^-, x_3, c_1, c_2 &\geq 0 \end{aligned}$$

1.1b) Max $Z = -3.1x_1 + 2\sqrt{2}x_2 - (x_3^+ - x_3^-)$
s.t.

$$\begin{aligned} 100x_1 - 20x_2 &= 7 \\ -11x_1 - 7x_2 - 2(x_3^+ - x_3^-) + s_2 &= 400 \\ (x_3^- - x_3^+) + s_3 &= 15 \\ x_1 \geq 0, x_2 \geq 0, x_3^+ \geq 0, x_3^- \geq 0, s_2 \geq 0, s_3 \geq 0 \end{aligned}$$

1.1c) Max $(x_1^+ - x_1^-) + 3x_2 - 2(x_3^+ - x_3^-)$
s.t.

$$\begin{aligned} -3(x_1^+ - x_1^-) + 5x_2 + s_1 &= 2 \\ 3(x_1^+ - x_1^-) - 5x_2 + s_2 &= 15 \\ -5(x_1^+ - x_1^-) + 20x_2 - c_3 &= 11 \\ -5(x_1^+ - x_1^-) + 20x_2 + s_4 &= 40 \\ (x_3^+ - x_3^-) + s_5 &= 10 \\ x_1^+, x_1^-, x_2, x_3^+, x_3^-, s_1, s_2, c_3, s_4, s_5 &\geq 0 \end{aligned}$$

1.2) Min $2x_1 + 6x_2 + 8x_3 = Z$
s.t.


$$\begin{aligned} x_1 + 2x_2 + x_3 &= 5 \\ 4x_1 + 6x_2 + 2x_3 &= 12 \\ x_2, x_3 &\geq 0 \end{aligned}$$

a) Min $2(x_1^+ - x_1^-) + 6x_2 + 8x_3 = Z$
s.t. $(x_1^+ - x_1^-) + 2x_2 + x_3 = 5$
 $4(x_1^+ - x_1^-) + 6x_2 + 2x_3 = 12$
 $x_1^+, x_1^-, x_2, x_3 \geq 0$

$$\begin{aligned}
 & \text{b.) Min } 2(5 - 2x_2 - x_3) + 6x_2 + 8x_3 = z \\
 & \text{s.t. } (5 - 2x_2 - x_3) + 2x_2 + x_3 = 5 \quad 5=5 \\
 & \quad 4(5 - 2x_2 - x_3) + 6x_2 + 2x_3 = 12 \quad 20 - 2x_2 - 2x_3 = 12 \\
 & \quad \quad \quad 20 - 8x_2 - 4x_3 \quad \quad \quad x_2, x_3 \geq 0
 \end{aligned}$$

$$\begin{aligned}
 & \text{Min } 2x_2 + 6x_3 + 10 = z \\
 & \text{s.t. } -2x_2 - 2x_3 = -8 \\
 & \quad \quad \quad x_2, x_3 \geq 0
 \end{aligned}$$

$$\begin{aligned}
 & \text{c.) Min } z = 2x_2 + 6x_3 + 10 \\
 & \text{s.t. } 2x_2 + 2x_3 = 8 \\
 & \quad \quad \quad x_2, x_3 \geq 0
 \end{aligned}$$




1.3)
a) No

b) Yes, plug in x_2 for x_1 .

$$\begin{aligned}
 & \text{Min } 2x_2 + 4x_3 = z \\
 & \text{s.t. } 2x_2 + 4x_3 \geq 4 \\
 & \quad \quad \quad x_2 \geq 2, x_3 \geq 0
 \end{aligned}$$

c.) standard

$$\begin{aligned}
 & \text{Min } 2x_2 + 4x_3 = z \\
 & \text{s.t. } 2x_2 + 4x_3 - e_1 = 4 \\
 & \quad \quad \quad x_2 \geq 2, x_3 \geq 0, e_1 \geq 0
 \end{aligned}$$



1.4

~~Q. Yes~~

s.t.

$$(x_1^+ - x_1^-) - 3(x_2^+ - x_2^-) + 2(x_3^+ - x_3^-) = 12$$

$$\text{Min } |x_3| + |x_4|$$

S.t.

$$x_3 + x_4 \leq 9$$

$$x_3 - 3x_4 \geq -15$$

Standard Form

$$\text{Min } x_3^+ + x_3^- + x_4^+ + x_4^-$$

S.t.

$$(x_3^+ - x_3^-) + (x_4^+ - x_4^-) + s_1 = 9 \Rightarrow 9$$

$$-(x_3^+ - x_3^-) + 3(x_4^+ - x_4^-) + S_2 = 15$$

$$x_3^+, x_3^-, x_4^+, x_4^-, s_1, s_2 \geq 0$$

$$x_1 - 5 - 3x_2 \geq 3$$

$$x_3 - 3(x_2 + 4) \geq -15$$

$$X_3 - 3X_4 \geq -15$$

1.5) x_1 : Quantity of chip-1 produced

x_2 : Quantity of chip-2 produced

$$\text{Max } Z = 15x_1 + 25x_2$$

s.t.

$$3x_1 + 4x_2 \leq 100$$

$$2x_1 + 3x_2 \leq 70$$

$$x_1 + 2x_2 \leq 30$$

$$x_1 \geq 0, x_2 \geq 0$$

standard Form

$$\text{Max } Z = 15x_1 + 25x_2$$

s.t.

$$3x_1 + 4x_2 + s_1 = 100$$

$$2x_1 + 3x_2 + s_2 = 70$$

$$x_1 + 2x_2 + s_3 = 30$$

$$x_1, s_1, s_2, s_3 \geq 0$$

$$x_2 \geq 0$$

1.6) $\text{Min } \sum_{i=1}^5 \sum_{j=1}^5 c_{ij} x_{ij}$

s.t.

$$\sum_{j=1}^5 x_{ij} = 1 \quad (i=1,2,3,4,5)$$

$$\sum_{i=1}^5 x_{ij} = 1 \quad (j=1,2,3,4,5)$$

$$x_{ij} = 0 \text{ or } 1$$

$$(i=1,2,3,4,5)$$

$$(j=1,2,3,4,5)$$

$$C = \begin{bmatrix} 5 & 5 & 7 & 4 & 8 \\ 6 & 5 & 8 & 3 & 7 \\ 6 & 8 & 9 & 5 & 10 \\ 7 & 6 & 6 & 3 & 6 \\ 6 & 7 & 10 & 6 & 11 \end{bmatrix}$$

$$1.7) \quad a = \begin{bmatrix} 60 \\ 45 \\ 30 \end{bmatrix} \quad b = [80 \ 55] \quad c = \begin{bmatrix} 190 & 230 \\ 225 & 270 \\ 180 & 250 \end{bmatrix}$$

Take each lesser cost route from source to destination:

China \rightarrow USA = 200 or 190

China \rightarrow France = 230 or 260

India \rightarrow USA = 240 or 225

India \rightarrow France = 270 or 295

Phillipines \rightarrow USA = 220 or 180

Phillipines \rightarrow France = 250

$$\text{Min } z = \sum_{i=1}^3 \sum_{j=1}^2 c_{ij} x_{ij}$$

s.t.

x_{ij} : Quantity shipped from each source to each destination

$$\sum_{j=1}^2 x_{ij} = a_i \quad (i=1,2,3)$$

$$\sum_{i=1}^3 x_{ij} = b_j \quad (j=1,2)$$

$$x_{ij} \geq 0 \quad ; (i=1,2,3 ; j=1,2)$$

b) Nothing, based on previous formulation, Hong Kong only packages textiles shipping to France, which is already limited to 55 tons.

c) It will force the model to split into multiple because the transportation model can't account for intermediate locations. The destination countries (USA and France) may also not receive their full demands since total demand is 135 tons and packaging capacity is 110 tons.

d) Source to Packaging Facilities

$$a = \begin{bmatrix} 60 \\ 45 \\ 30 \end{bmatrix} \quad b = [60 \ 50] \quad c = \begin{bmatrix} 50 & 60 \\ 90 & 95 \\ 70 & 50 \end{bmatrix}$$

$$\text{Min } z = \sum_{i=1}^3 \sum_{j=1}^2 c_{ij} x_{ij}$$

s.t.

$$\sum_{j=1}^2 x_{ij} = a_i \quad (i=1,2,3)$$

$$\sum_{i=1}^3 x_{ij} = b_j \quad (j=1,2)$$

$$x_{ij} \geq 0; (i=1,2,3; j=1,2)$$

Packaging Facilities to Destinations

$$a = \begin{bmatrix} 60 \\ 50 \end{bmatrix} \quad b = [80 \ 55] \quad c = \begin{bmatrix} 150 & 180 \\ 130 & 200 \end{bmatrix}$$

$$\text{Min } z = \sum_{i=1}^2 \sum_{j=1}^2 c_{ij} x_{ij}$$

s.t.

$$\sum_{j=1}^2 x_{ij} = a_i \quad (i=1,2)$$

$$\sum_{i=1}^2 x_{ij} \leq b_j \quad (j=1,2)$$

$$x_{ij} \geq 0; (i=1,2; j=1,2)$$

Please check the solution

96