

Homework 5 Solutions

Zheming Gao

September 27, 2017

Problem 1 (3.1)

Suppose $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$. Add slack variables $s_1, \dots, s_m \geq 0$ on each row and make it a standard form.

Let starting feasible solution be $(x_1, \dots, x_n, s_1, \dots, s_m)^T = (0, \dots, 0, b_1, \dots, b_m)^T$.

Problem 2 (3.2)

- (a) *Proof.* Let d be a feasible direction at point $x \in P$. Then, there exists $\lambda > 0$ such that $x + \lambda d \in P$, which implies $A(x + \lambda d) = b$. Since $Ax = b$, we know that $\lambda Ad = 0$ and this implies $Ad = 0$. □

- (b) *Proof.* let $d = (d_1, \dots, d_n)^T$ be a feasible direction at x . Let $\alpha = \min\{\frac{x_i}{-d_i} | d_i < 0, i = 1, \dots, n\}$. If $d \geq 0$, then let $\alpha = 1$.

It is clear that $\alpha > 0$ and $x + \alpha d \geq 0$. □

Problem 3 (3.3)

$$\begin{array}{ll} \text{Minimize} & -2x_1 - x_2 + x_3 + x_4 + 2x_5 \\ \text{subject to} & -2x_1 + x_2 + x_3 + x_4 + x_5 = 12 \\ & -x_1 + 2x_2 + x_4 - x_5 = 5 \\ & x_1 - 3x_2 + x_3 + 4x_5 = 11 \\ & x_1, x_2, x_3, x_4, x_5 \geq 0 \end{array}$$

$$\text{Here, } A = \begin{pmatrix} -2 & 1 & 1 & 1 & 1 \\ -1 & 2 & 0 & 1 & -1 \\ 1 & -3 & 1 & 0 & 4 \end{pmatrix} \text{ and } b = \begin{pmatrix} 12 \\ 5 \\ 11 \end{pmatrix}$$

(a) $B = [A_3, A_4, A_5] = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & -1 \\ 1 & 0 & 4 \end{pmatrix}$, and $N = [A_1, A_2] = \begin{pmatrix} -2 & 1 \\ -1 & 2 \\ 1 & -3 \end{pmatrix}$.

The fundamental matrix $M = \begin{bmatrix} B & N \\ \mathbf{0} & I \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & -2 & 1 \\ 0 & 1 & -1 & -1 & 2 \\ 1 & 0 & 4 & 1 & -3 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$ and $M^{-1} = \begin{bmatrix} B^{-1} & -B^{-1}N \\ \mathbf{0} & I \end{bmatrix} =$

$$\begin{bmatrix} 2 & -2 & -1 & 3 & -1 \\ -1/2 & 3/2 & 1/2 & 0 & -1 \\ -1/2 & 1/2 & 1/2 & -1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

(b) Apply Gaussian elimination on matrix $[B, N]$ and get reduced row echelon form

$$\begin{bmatrix} 1 & 0 & 0 & -3 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & -1 \end{bmatrix}$$


Hence, $x_3 = 3x_1 - x_2 + 3$, $x_4 = -x_2 + 7$, $x_5 = -x_1 + x_2 + 2$. Reform the LP problem using only two variables as the following:

$$\begin{array}{ll} \text{Minimize} & -x_1 - x_2 (+14) \\ \text{subject to} & 3x_1 - x_2 + 3 \geq 0 \\ & -x_2 + 7 \geq 0 \\ & -x_1 + x_2 + 2 \geq 0 \\ & x_1, x_2 \geq 0 \end{array}$$

(c) The feasible domain is part of the intersection of three hyperplanes on \mathbb{R}^5 , hence, its dimension is reduced by 3 and can be represented in \mathbb{R}^2 .

We plot the region of P .

Figure 1: Region P .



- (d) Basic feasible solution $\mathbf{x} = (0, 0, 3, 7, 2)^T$. And it is corresponding to point $D = (0, 0)^T$ on fig.1.

(e)

$$B^{-1}A = \begin{bmatrix} -3 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 0 & 1 \end{bmatrix}, \quad B^{-1}b = \begin{bmatrix} 3 \\ 7 \\ 2 \end{bmatrix}$$

One explanation of this:

We see that $B^{-1}A$ is the same with the reduced row echelon form of A . And $B^{-1}b$ is exactly the basic feasible solution (positive entries). This is always true since

$$Ax = b \Leftrightarrow [B|N] \begin{bmatrix} x_B \\ x_N \end{bmatrix} = b.$$

Since $x_N = 0$, we get $Bx_B = b$ so that $x_B = B^{-1}b$. This implies that $B^{-1}A$ is the reduced row echelon form of A for basic variable, since $B^{-1}Ax = B^{-1}b = x_B$.

(Other proper answers will be also acceptable).

- (f) From M^{-1} we know that $\mathbf{d}^1 = (3, 0, -1, 1, 0)^T$ and $\mathbf{d}^2 = (-1, -1, 1, 0, 1)^T$. Reduced costs:

$$r^1 = [c_B^T | c_N^T] \mathbf{d}^1 = (1, 1, 2, -2, -1) \mathbf{d}^1 = -1, \quad r^2 = [c_B^T | c_N^T] \mathbf{d}^2 = (1, 1, 2, -2, -1) \mathbf{d}^2 = -1$$

- (g) From above, either direction leads to a potential reduction in the objective value, since r^1 and r^2 are both negative. Consider the nonnegativity constraint ($x + \alpha \mathbf{d} \geq 0$), we get the step length for \mathbf{d}^1 is $\alpha_1 = 2$ and for \mathbf{d}^2 is $\alpha_2 = 3$.

- (h) 1) If we take \mathbf{d}^1 , then the new solution will be $\bar{x} = x + \alpha_1 \mathbf{d}^1 = (9, 7, 0, 2, 0)^T \geq 0$. The basis now is $\bar{B} = [A_3, A_4, A_1]$ and $\bar{N} = [A_5, A_2]$. It is easy to check that $\bar{B}\bar{x} = b$. Hence \bar{x} is a basic feasible solution(BFS). \bar{x} is also an adjacent extreme point of x . (On fig.1, \bar{x} is the point C)
- 2) If we take \mathbf{d}^2 , then the new solution will be $\bar{x} = x + \alpha_2 \mathbf{d}^2 = (0, 4, 5, 0, 3)^T \geq 0$. The basis now is $\bar{B} = [A_4, A_5, A_2]$ and $\bar{N} = [A_3, A_1]$. It is easy to check that $\bar{B}\bar{x} = b$. Hence \bar{x} is a basic feasible solution(BFS). \bar{x} is also an adjacent extreme point of x . (On fig.1, \bar{x} is the point E)
- (i) 1) If we take \mathbf{d}^1 , update $\tilde{M} = \begin{bmatrix} \tilde{B} & \tilde{N} \\ \mathbf{0} & I \end{bmatrix}$.

$$\tilde{M} = \begin{bmatrix} 1 & 1 & -2 & 1 & 1 \\ 0 & 1 & -1 & -1 & 2 \\ 1 & 0 & 1 & 4 & -3 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad \tilde{M}^{-1} = \begin{bmatrix} 1/2 & -1/2 & 1/2 & -3 & 2 \\ -1/2 & 3/2 & 1/2 & 0 & -1 \\ -1/2 & 1/2 & 1/2 & -1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

From the last two columns of \tilde{M}^{-1} we get $\tilde{\mathbf{d}}^5 = \begin{bmatrix} -3 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix}$, $\tilde{\mathbf{d}}^2 = \begin{bmatrix} 2 \\ -1 \\ 1 \\ 0 \\ 1 \end{bmatrix}$.

Let $\tilde{c}^T = [\tilde{c}_B^T | \tilde{c}_N^T]$. We get $\tilde{c}^T \tilde{\mathbf{d}}^5 = 1 > 0$, but $\tilde{c}^T \tilde{\mathbf{d}}^2 = -2 < 0$. So \tilde{x} is not an optimal solution since $\tilde{\mathbf{d}}^2$ is a good direction of translation.

- 2) If we take \mathbf{d}^2 , update $\bar{M} = \begin{bmatrix} \bar{B} & \bar{N} \\ \mathbf{0} & I \end{bmatrix}$.

$$\bar{M} = \begin{bmatrix} 1 & 1 & 1 & 1 & -2 \\ 1 & -1 & 2 & 0 & -1 \\ 0 & 4 & -3 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad \bar{M}^{-1} = \begin{bmatrix} -5/2 & 7/2 & 3/2 & 1 & -3 \\ 3/2 & -3/2 & -1/2 & -1 & 2 \\ 2 & -2 & -1 & -1 & 3 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

From the last two columns of \bar{M}^{-1} we get $\bar{\mathbf{d}}^5 = \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \\ 0 \end{bmatrix}$, $\bar{\mathbf{d}}^2 = \begin{bmatrix} -3 \\ 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}$.

Let $\bar{c}^T = [\bar{c}_B^T | \bar{c}_N^T]$. We get $\bar{c}^T \bar{\mathbf{d}}^3 = 1 > 0$, but $\bar{c}^T \bar{\mathbf{d}}^1 = -4 < 0$. So \bar{x} is not an optimal solution since $\bar{\mathbf{d}}^1$ is a good direction of translation.

- (j) 1) If basic variables are x_3, x_4, x_1 , then the reduced row echelon form (RREF) of $[\tilde{B}, \tilde{N}, -b]$ is


$$\begin{bmatrix} 1 & 0 & 0 & 3 & -2 & -9 \\ 0 & 1 & 0 & 0 & 1 & -7 \\ 0 & 0 & 1 & 1 & -1 & -2 \end{bmatrix}$$

and $x_3 = 2x_2 - 3x_5 + 9$, $x_4 = -x_2 + 7$, $x_1 = x_2 - x_5 + 2$. Reform the LP problem using only two variables as the following:

$$\begin{array}{llll} \text{Minimize} & -2x_2 + x_5(+12) \\ \text{subject to} & 2x_2 - 3x_5 + 9 & \geq 0 \\ & -x_2 + 7 & \geq 0 \\ & x_2 - x_5 + 2 & \geq 0 \\ & x_2, x_5 & \geq 0 \end{array}$$

We plot the region.

Figure 2: Region on x_2, x_5 .



- 2) If basic variables are x_4, x_5, x_2 , then the reduced row echelon form (RREF) of $[\overline{B}, \overline{N}, -b]$ is


$$\begin{bmatrix} 1 & 0 & 0 & -1 & 3 & -4 \\ 0 & 1 & 0 & 1 & -2 & -5 \\ 0 & 0 & 1 & 1 & -3 & -3 \end{bmatrix}$$

and $x_4 = x_3 - 3x_1 + 4$, $x_5 = -x_3 + 2x_1 + 5$, $x_2 = -x_3 + 3x_1 + 3$. Reform the LP problem using only two variables as the following:

$$\begin{array}{llll} \text{Minimize} & -4x_1 + x_3(+11) \\ \text{subject to} & x_3 - 3x_1 + 4 & \geq 0 \\ & -x_3 + 2x_1 + 5 & \geq 0 \\ & -x_3 + 3x_1 + 3 & \geq 0 \\ & x_1, x_3 & \geq 0 \end{array}$$

We plot the region.

Figure 3: Region on x_1, x_3 .



(k) (Any proper answer will be acceptable).

(l) We can always express $n - 2$ variables by using the rest 2 variables. Then, the LP problem can be reformed as an LP problem on \mathbb{R}^2 .

Problem 4 (3.4)

From point $x = [x_B | x_N]^T = [x_5, x_6, x_1 | x_2, x_3, x_4]^T$, we know that $B = [A_5, A_6, A_1]$ and $N = [A_2, A_3, A_4]$. Construct matrix $M = \begin{bmatrix} B & N \\ \mathbf{0} & I \end{bmatrix}$ and reduced cost $r = c_N^T - c_B^T B^{-1} N = [-1, -1, 1/2]$.

Note that r_2, r_3 are negative, and figure out the step length $\alpha_2 = \alpha_3 = 1/2$. Hence, we can pick either one from \mathbf{d}^2 or \mathbf{d}^3 . Let's pick \mathbf{d}^2 . $x_{\text{new}} = x + \alpha_2 \mathbf{d}^2 = (0, 1, 1/2, 1/2, 0, 0)^T$.

Next step:

From point $x = [x_2, x_6, x_1 | x_5, x_3, x_4] = (1/2, 1, 1/2, 0, 0, 0)^T$, we know $B = [A_2, A_6, A_1]$ and $N = [A_5, A_3, A_4]$. Construct matrix M and reduced cost $r = c_N^T - c_B^T B^{-1} N = [-1/2, -1, -1/2]$. Find the most negative direction, pick \mathbf{d}^3 with $\alpha_3 = 1/2$. $x_{\text{new}} = x + \alpha_3 \mathbf{d}^3 = (1/2, 0, 1/2, 1/2, 0, 0)^T$.

Next step:

From point $x = [x_1, x_2, x_3 | x_4, x_5, x_6] = (1/2, 1, 1/2, 0, 0, 0)^T$, we know $B = [A_1, A_2, A_3]$ and $N = [A_4, A_5, A_6]$. Compute the reduced cost $r = c_N^T - c_B^T B^{-1} N = [1/2, 1/2, 1/2] \geq 0$. Hence, $x = [x_1, x_2, x_3 | x_4, x_5, x_6] = (1/2, 1, 1/2, 0, 0, 0)^T$ is the optimal solution.

Problem 5 (3.8)

(1) We plot the graph of F_3 .

(2)

$$B = \{(x_1, x_2, x_3)^T \in \mathbb{R}^3 \mid |x_1| + |x_3| = x_2\}.$$

(3)

$$I = \{(x_1, x_2, x_3)^T \in \mathbb{R}^3 \mid |x_1| + |x_3| < x_2\}.$$

(4) Extreme point: $(0, 0, 0)^T$.

Vertex: $(0, 0, 0)^T$.

(5) *Proof.* First, we show that F_3 is a cone. Take $x = (x_1, x_2, x_3)^T \in F_3$ and $\forall \lambda \geq 0$, check if $\lambda x \in F_3$.

$$|\lambda x_1| + |\lambda x_3| = \lambda |x_1| + \lambda |x_3| = \lambda(|x_1| + |x_3|) \leq \lambda x_2.$$

Hence, λx is in F_3 . In conclusion, F_3 is a cone.

Next, we need to show F_3 is convex. Take $x, y \in F_3$ and $\forall \eta \in (0, 1)$, $\eta x + (1 - \eta)y = (\eta x_1 + (1 - \eta)y_1, \eta x_2 + (1 - \eta)y_2, \eta x_3 + (1 - \eta)y_3)$. Use triangle inequality of absolute value and we get

$$\begin{aligned} |\eta x_1 + (1 - \eta)y_1| + |\eta x_3 + (1 - \eta)y_3| &\leq \eta |x_1| + (1 - \eta)|y_1| + \eta |x_3| + (1 - \eta)|y_3| \\ &= \eta(|x_1| + |x_3|) + (1 - \eta)(|y_1| + |y_3|) \\ &\leq \eta x_2 + (1 - \eta)y_2. \end{aligned}$$

Hence, $\eta x + (1 - \eta)y \in F_3$. This implies that F_3 is convex.

□

(6) (Any reasonable answers will be fine for this question.)

Example answer:

\mathbb{R}_+^3 are different from F_3 and $\mathbb{R}_+^3 \cap F_3 \neq \phi$.

$(-1, 2, -1)^T \notin \mathbb{R}_+^3$ but in F_3 . $(2, 1, 2)^T \notin F_3$, but in \mathbb{R}_+^3 .

Problem 6

We plot the region of P_1 .

- (a) Convert P_1 to standard equality form.

$$\begin{cases} 2x_1 - 4x_2 + a_1 & = 1 \\ 3x_1 - x_2 & -a_2 = -3 \\ x_1, x_2, a_1, a_2 & \geq 0. \end{cases}$$

- (b) Basic solutions (in the form of (x_1, x_2, a_1, a_2)):

$$\begin{aligned} &(-13/10, -9/10, 0, 0) \\ &(-1, 0, 3, 0) \\ &(1/2, 0, 0, 9/2)\star \\ &(0, 3, 13, 0)\star \\ &(0, -1/4, 0, 13/4) \\ &(0, 0, 1, 3)\star \end{aligned}$$

- (c) Basic feasible solutions are those basic solutions with " \star ".

- (d) Let V be the set of all extremal directions.

$$V = \{v \in \mathbb{R}^2 | v = (1, d)^T, d \in [1/2, 3]\}.$$

- (e) From the figure, we can see that there are two moving directions to the adjacent points.
 u_1 is to point $(1/2, 0)^T$ and u_2 is to point $(0, 3)^T$.

$$\begin{aligned} u_1 &= (1, 0)^T \\ u_2 &= (0, 1)^T \end{aligned}$$

Problem 7

We plot the region of P_2 .

- (a) Convert P_2 to standard equality form.

$$\begin{cases} 2x_1 - 2x_2^+ + 2x_2^- + a_1 & = 3 \\ 8x_1 - x_2^+ + x_2^- & -a_2 = -4 \\ x_1, x_2^+, x_2^-, a_1, a_2 & \geq 0. \end{cases}$$

(b) Basic solutions (in the form of $(x_1, x_2^+, x_2^-, a_1, a_2)$):

$$\begin{aligned}
&(-1/2, -1, 0, 0, 0) \\
&(-1/2, 1, 0, 0, 0) \\
&(-1/2, 0, 0, 4, 0) \\
&(3/2, 0, 0, 0, 16)\star \\
&(0, 4, 0, 11, 0)\star \\
&(0, -3/2, 0, 0, 11/2) \\
&(0, 0, -4, 11, 0) \\
&(0, 0, 3/2, 0, 11/2)\star \\
&(0, 0, 0, 3, 4)\star
\end{aligned}$$

(c) Basic feasible solutions are those basic solutions with "★".

(d) Let V be the set of all extremal directions.

$$V = \{v \in \mathbb{R}^2 | v = (1, d)^T, d \in [1, 8]\}.$$

(e) From the figure, we can see that there is only one moving directions to the adjacent point $(0, -3/2)^T$.

$$u = (0, -1)^T$$