Using R to find the most central points in object data

Vida Zamani University of Turku, Finland


useR! 2024 conference, Salzburg, Austria

July 9, 2024

Table of Contents

- Background and overview of the subject
 - Object data
 - Extending to Metric Space
- Case Study
 - Canadian Weather Data
 - 4 Different regions
- Results

Object Data



Importance of Analyzing Object Data


$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \qquad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \qquad \begin{bmatrix} 0 & 1 & 0 \\ 1 & -1 & 1 \\ 0 & 1 & 0 \end{bmatrix} \qquad \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \qquad \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$


Table: The path of Descriptive Analysis

	Euclidean Space	Object data
Depth Functions	Tukey's depth	Metric halfspace depth
Mean	Mean sample	Fréchet mean
Dimension Reduction	PCA	MDS




Case Study


Canadian Weather Data







Deepest Stations in each groups



Permutation Distribution

