

Solution for "Quantum Computation and Quantum Information:
10th Anniversary Edition" by Nielsen and Chuang

goropikari

March 18, 2020

Copyright Notice:



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

Repository

The latest version and source \LaTeX code are located in

<https://github.com/goropikari/SolutionForQuantumComputationAndQuantumInformation>.

For readers

This is an unofficial solution manual for "Quantum Computation and Quantum Information: 10th Anniversary Edition" (ISBN-13: 978-1107002173) by Michael A. Nielsen and Isaac L. Chuang.

I have studied quantum information theory as a hobby. And I'm not a researcher. So there is no guarantee that these solutions are correct. Especially because I'm not good at mathematics, proofs are often wrong. Don't trust me. Verify yourself!

If you find some mistake or have some comments, please feel free to open an issue or a PR.

goropikari

Contents

Errata list	i
2 Introduction to quantum mechanics	1
5 The quantum Fourier transform and it's application	39
8 Quantum noise and quantum operations	41
9 Distance measures for quantum information	43
11 Entropy and information	53

Errata list

- p.101. eq (2.150) $\rho = \sum_m p(m)\rho_m$ should be $\rho' = \sum_m p(m)\rho_m$.
- p.408. eq (9.49) $\sum_i p_i D(\rho_i, \sigma_i) + D(p_i, q_i)$ should be $\sum_i p_i D(\rho_i, \sigma_i) + 2D(p_i, q_i)$.

$$\begin{aligned}
 \text{eqn (9.48)} &= \sum_i p_i \text{Tr}(P(\rho_i - \sigma_i)) + \sum_i (p_i - q_i) \text{Tr}(P\sigma_i) \\
 &\leq \sum_i p_i \text{Tr}(P(\rho_i - \sigma_i)) + \sum_i |p_i - q_i| \text{Tr}(P\sigma_i) \quad (\because p_i - q_i \leq |p_i - q_i|) \\
 &\leq \sum_i p_i \text{Tr}(P(\rho_i - \sigma_i)) + \sum_i |p_i - q_i| \quad (\because \text{Tr}(P\sigma_i) \leq 1) \\
 &= \sum_i p_i \text{Tr}(P(\rho_i - \sigma_i)) + 2 \frac{\sum_i |p_i - q_i|}{2} \\
 &= \sum_i p_i \text{Tr}(P(\rho_i - \sigma_i)) + 2D(p_i, q_i)
 \end{aligned}$$

- p.409. Exercise 9.12. If $\rho = \sigma$, then $D(\rho, \sigma) = 0$. Furthermore trace distance is non-negative. Therefore $0 \leq D(\mathcal{E}(\rho), \mathcal{E}(\sigma)) \leq 0 \Rightarrow D(\mathcal{E}(\rho), \mathcal{E}(\sigma)) = 0$. So I think the map \mathcal{E} is not strictly contractive. If $p \neq 1$ and $\rho \neq \sigma$, then $D(\mathcal{E}(\rho), \mathcal{E}(\sigma)) < D(\rho, \sigma)$ is satisfied.
- p.411. Exercise 9.16. eqn(9.73) $\text{Tr}(A^\dagger B) = \langle m|A \otimes B|m\rangle$ should be $\text{Tr}(A^T B) = \langle m|A \otimes B|m\rangle$.

Simple counter example is the case that $A = \begin{bmatrix} i & 0 \\ 0 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, In this case,

$$\begin{aligned}
 A^\dagger B &= \begin{bmatrix} -i & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} -i & 0 \\ 0 & 0 \end{bmatrix}, \\
 \text{Tr}(A^\dagger B) &= -i, \\
 A \otimes B &= \begin{bmatrix} i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \\
 \langle m|A \otimes B|m\rangle &= (\langle 00| + \langle 11|)(A \otimes B)(|00\rangle + |11\rangle) = i.
 \end{aligned}$$

Thus $\text{Tr}(A^\dagger B) \neq \langle m|A \otimes B|m\rangle$.

By using following relation, we can prove.

$$\begin{aligned}
 (I \otimes A) |m\rangle &= (A^T \otimes I) |m\rangle \\
 \text{Tr}(A) &= \langle m|I \otimes A|m\rangle
 \end{aligned}$$

$$\begin{aligned}
\mathrm{Tr}(A^T B) &= \mathrm{Tr}(BA^T) = \langle m | I \otimes BA^T | m \rangle \\
&= \langle m | (I \otimes B)(I \otimes A^T) | m \rangle \\
&= \langle m | (I \otimes B)(A \otimes I) | m \rangle \\
&= \langle m | A \otimes B | m \rangle.
\end{aligned}$$

- p.515. eqn (11.67) $S(\rho' || \rho)$ should be $S(\rho || \rho')$.

Chapter 2

Introduction to quantum mechanics

2.1

$$\begin{bmatrix} 1 \\ -1 \end{bmatrix} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

2.2

$$\begin{aligned} A|0\rangle &= A_{11}|0\rangle + A_{21}|1\rangle = |1\rangle \Rightarrow A_{11} = 0, A_{21} = 1 \\ A|1\rangle &= A_{12}|0\rangle + A_{22}|1\rangle = |0\rangle \Rightarrow A_{12} = 1, A_{22} = 0 \\ \therefore A &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \end{aligned}$$

input: $\{|0\rangle, |1\rangle\}$, output: $\{|1\rangle, |0\rangle\}$

$$\begin{aligned} A|0\rangle &= A_{11}|1\rangle + A_{21}|0\rangle = |1\rangle \Rightarrow A_{11} = 1, A_{21} = 0 \\ A|1\rangle &= A_{12}|1\rangle + A_{22}|0\rangle = |0\rangle \Rightarrow A_{12} = 0, A_{22} = 1 \\ A &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

2.3

From eq (2.12)

$$\begin{aligned} A|v_i\rangle &= \sum_j A_{ji}|w_j\rangle \\ B|w_j\rangle &= \sum_k B_{kj}|x_k\rangle \end{aligned}$$

Thus

$$\begin{aligned}
 BA|v_i\rangle &= B\left(\sum_j A_{ji}|w_j\rangle\right) \\
 &= \sum_j A_{ji}B|w_j\rangle \\
 &= \sum_{j,k} A_{ji}B_{kj}|x_k\rangle \\
 &= \sum_k \left(\sum_j B_{kj}A_{ji}\right)|x_k\rangle \\
 &= \sum_k (BA)_{ki}|x_k\rangle \\
 \therefore (BA)_{ki} &= \sum_j B_{kj}A_{ji}
 \end{aligned}$$

2.4

$$\begin{aligned}
 I|v_j\rangle &= \sum_i I_{ij}|v_i\rangle = |v_j\rangle, \quad \forall j. \\
 &\Rightarrow I_{ij} = \delta_{ij}
 \end{aligned}$$

2.5

Defined inner product on \mathcal{C}^n is

$$((y_1, \dots, y_n), (z_1, \dots, z_n)) = \sum_i y_i^* z_i.$$

Verify (1) of eq (2.13).

$$\begin{aligned}
 \left((y_1, \dots, y_n), \sum_i \lambda_i (z_{i1}, \dots, z_{in})\right) &= \sum_i y_i^* \left(\sum_j \lambda_j z_{ji}\right) \\
 &= \sum_{i,j} y_i^* \lambda_j z_{ji} \\
 &= \sum_j \lambda_j \left(\sum_i y_i^* z_{ji}\right) \\
 &= \sum_j \lambda_j ((y_1, \dots, y_n), (z_{j1}, \dots, z_{jn})) \\
 &= \sum_i \lambda_i ((y_1, \dots, y_n), (z_{i1}, \dots, z_{in})).
 \end{aligned}$$

Verify (2) of eq (2.13),

$$((y_1, \dots, y_n), (z_1, \dots, z_n))^* = \left(\sum_i y_i^* z_i \right)^* \quad (2.1)$$

$$= \left(\sum_i y_i z_i^* \right) \quad (2.2)$$

$$= \left(\sum_i z_i^* y_i \right) \quad (2.3)$$

$$= ((z_1, \dots, z_n), (y_1, \dots, y_n)) \quad (2.4)$$

Verify (3) of eq (2.13),

$$\begin{aligned} ((y_1, \dots, y_n), (y_1, \dots, y_n)) &= \sum_i y_i^* y_i \\ &= \sum_i |y_i|^2 \end{aligned}$$

Since $|y_i|^2 \geq 0$ for all i . Thus $\sum_i |y_i|^2 = ((y_1, \dots, y_n), (y_1, \dots, y_n)) \geq 0$.

From now on, I will show the following statement,

$$((y_1, \dots, y_n), (y_1, \dots, y_n)) = 0 \text{ iff } (y_1, \dots, y_n) = 0.$$

(\Leftarrow) This is obvious.

(\Rightarrow) Suppose $((y_1, \dots, y_n), (y_1, \dots, y_n)) = 0$. Then $\sum_i |y_i|^2 = 0$.

Since $|y_i|^2 \geq 0$ for all i , if $\sum_i |y_i|^2 = 0$, then $|y_i|^2 = 0$ for all i . Therefore $|y_i|^2 = 0 \Leftrightarrow y_i = 0$ for all i . Thus,

$$(y_1, \dots, y_n) = 0.$$

2.6

$$\begin{aligned} \left(\sum_i \lambda_i |w_i\rangle, |v\rangle \right) &= \left(|v\rangle, \sum_i \lambda_i |w_i\rangle \right)^* \\ &= \left[\sum_i \lambda_i (|v\rangle, |w_i\rangle) \right]^* (\because \text{linearity in the 2nd arg.}) \\ &= \sum_i \lambda_i^* (|v\rangle, |w_i\rangle)^* \\ &= \sum_i \lambda_i^* (|w_i\rangle, |v\rangle) \end{aligned}$$

2.7

$$\begin{aligned}\langle w|v\rangle &= \begin{bmatrix} 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 1 - 1 = 0 \\ \frac{|w\rangle}{\| |w\rangle \|} &= \frac{|w\rangle}{\sqrt{\langle w|w\rangle}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \\ \frac{|v\rangle}{\| |v\rangle \|} &= \frac{|v\rangle}{\sqrt{\langle v|v\rangle}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}\end{aligned}$$

2.8

If $k = 1$,

$$\begin{aligned}|v_2\rangle &= \frac{|w_2\rangle - \langle v_1|w_2\rangle |v_1\rangle}{\| |w_2\rangle - \langle v_1|w_2\rangle |v_1\rangle \|} \\ \langle v_1|v_2\rangle &= \langle v_1| \left(\frac{|w_2\rangle - \langle v_1|w_2\rangle |v_1\rangle}{\| |w_2\rangle - \langle v_1|w_2\rangle |v_1\rangle \|} \right) \\ &= \frac{\langle v_1|w_2\rangle - \langle v_1|w_2\rangle \langle v_1|v_1\rangle}{\| |w_2\rangle - \langle v_1|w_2\rangle |v_1\rangle \|} \\ &= 0.\end{aligned}$$

Suppose $\{v_1, \dots, v_n\}$ ($n \leq d-1$) is a orthonormal basis. Then

$$\begin{aligned}\langle v_j|v_{n+1}\rangle &= \langle v_j| \left(\frac{|w_{n+1}\rangle - \sum_{i=1}^n \langle v_i|w_{n+1}\rangle |v_i\rangle}{\| |w_{n+1}\rangle - \sum_{i=1}^n \langle v_i|w_{n+1}\rangle |v_i\rangle \|} \right) \quad (j \leq n) \\ &= \frac{\langle v_j|w_{n+1}\rangle - \sum_{i=1}^n \langle v_i|w_{n+1}\rangle \langle v_j|v_i\rangle}{\| |w_{n+1}\rangle - \sum_{i=1}^n \langle v_i|w_{n+1}\rangle |v_i\rangle \|} \\ &= \frac{\langle v_j|w_{n+1}\rangle - \sum_{i=1}^n \langle v_i|w_{n+1}\rangle \delta_{ij}}{\| |w_{n+1}\rangle - \sum_{i=1}^n \langle v_i|w_{n+1}\rangle |v_i\rangle \|} \\ &= \frac{\langle v_j|w_{n+1}\rangle - \langle v_j|w_{n+1}\rangle}{\| |w_{n+1}\rangle - \sum_{i=1}^n \langle v_i|w_{n+1}\rangle |v_i\rangle \|} \\ &= 0.\end{aligned}$$

Thus Gram-Schmidt procedure produces an orthonormal basis.

2.9

$$\begin{aligned}\sigma_0 &= I = |0\rangle \langle 0| + |1\rangle \langle 1| \\ \sigma_1 &= X = |0\rangle \langle 1| + |1\rangle \langle 0| \\ \sigma_2 &= Y = -i |0\rangle \langle 1| + i |1\rangle \langle 0| \\ \sigma_3 &= Z = |0\rangle \langle 0| - |1\rangle \langle 1|\end{aligned}$$

2.10

$$\begin{aligned}
|v_j\rangle \langle v_k| &= I_V |v_j\rangle \langle v_k| I_V \\
&= \left(\sum_p |v_p\rangle \langle v_p| \right) |v_j\rangle \langle v_k| \left(\sum_q |v_q\rangle \langle v_q| \right) \\
&= \sum_{p,q} |v_p\rangle \langle v_p| v_j\rangle \langle v_k| v_q\rangle \langle v_q| \\
&= \sum_{p,q} \delta_{pj} \delta_{kq} |v_p\rangle \langle v_q|
\end{aligned}$$

Thus

$$(|v_j\rangle \langle v_k|)_{pq} = \delta_{pj} \delta_{kq}$$

2.11

$$X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \det(X - \lambda I) = \det \left(\begin{bmatrix} -\lambda & 1 \\ 1 & -\lambda \end{bmatrix} \right) = 0 \Rightarrow \lambda = \pm 1$$

If $\lambda = -1$,

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Thus

$$|\lambda = -1\rangle = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

If $\lambda = 1$

$$|\lambda = 1\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$X = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \text{ w.r.t. } \{|\lambda = -1\rangle, |\lambda = 1\rangle\}$$

2.12

$$\det \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} - \lambda I \right) = (1 - \lambda)^2 = 0 \Rightarrow \lambda = 1$$

Therefore the eigenvector associated with eigenvalue $\lambda = 1$ is

$$|\lambda = 1\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Because $|\lambda = 1\rangle \langle \lambda = 1| = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix},$

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \neq c |\lambda = 1\rangle \langle \lambda = 1| = \begin{bmatrix} 0 & 0 \\ 0 & c \end{bmatrix}$$

2.13

Suppose $|\psi\rangle, |\phi\rangle$ are arbitrary vectors in V .

$$\begin{aligned} (|\psi\rangle, (|w\rangle\langle v|)|\phi\rangle)^* &= \left((|w\rangle\langle v|)^\dagger|\psi\rangle, |\phi\rangle\right)^* \\ &= \left(|\phi\rangle, (|w\rangle\langle v|)^\dagger|\psi\rangle\right) \\ &= \langle\phi|(|w\rangle\langle v|)^\dagger|\psi\rangle. \end{aligned}$$

On the other hand,

$$\begin{aligned} (|\psi\rangle, (|w\rangle\langle v|)|\phi\rangle)^* &= (\langle\psi|w\rangle\langle v|\phi\rangle)^* \\ &= \langle\phi|v\rangle\langle w|\psi\rangle. \end{aligned}$$

Thus

$$\begin{aligned} \langle\phi|(|w\rangle\langle v|)^\dagger|\psi\rangle &= \langle\phi|v\rangle\langle w|\psi\rangle \text{ for arbitrary vectors } |\psi\rangle, |\phi\rangle \\ \therefore (|w\rangle\langle v|)^\dagger &= |v\rangle\langle w| \end{aligned}$$

2.14

$$\begin{aligned} ((a_i A_i)^\dagger|\phi\rangle, |\psi\rangle) &= (|\phi\rangle, a_i A_i|\psi\rangle) \\ &= a_i(|\phi\rangle, A_i|\psi\rangle) \\ &= a_i(A_i^\dagger|\phi\rangle, |\psi\rangle) \\ &= (a_i^* A_i^\dagger|\phi\rangle, |\psi\rangle) \\ \therefore (a_i A_i)^\dagger &= a_i^* A_i^\dagger \end{aligned}$$

2.15

$$\begin{aligned} ((A^\dagger)^\dagger|\psi\rangle, |\phi\rangle) &= (|\psi\rangle, A^\dagger|\phi\rangle) \\ &= (A^\dagger|\phi\rangle, |\psi\rangle)^* \\ &= (|\phi\rangle, A|\psi\rangle)^* \\ &= (A|\psi\rangle, |\phi\rangle) \\ \therefore (A^\dagger)^\dagger &= A \end{aligned}$$

2.16

$$\begin{aligned}
P &= \sum_i |i\rangle \langle i|. \\
P^2 &= \left(\sum_i |i\rangle \langle i| \right) \left(\sum_j |j\rangle \langle j| \right) \\
&= \sum_{i,j} |i\rangle \langle i|j\rangle \langle j| \\
&= \sum_{i,j} |i\rangle \langle j| \delta_{ij} \\
&= \sum_i |i\rangle \langle i| \\
&= P
\end{aligned}$$

2.17

Proof. (\Rightarrow) Suppose A is Hermitian. Then $A = A^\dagger$. Let $|\lambda\rangle$ be eigenvectors of A with eigenvalues λ , that is,

$$A|\lambda\rangle = \lambda|\lambda\rangle.$$

Therefore

$$\langle\lambda|A|\lambda\rangle = \lambda\langle\lambda|\lambda\rangle = \lambda.$$

On the other hand,

$$\lambda^* = \langle\lambda|A|\lambda\rangle^* = \langle\lambda|A^\dagger|\lambda\rangle = \langle\lambda|A|\lambda\rangle = \lambda\langle\lambda|\lambda\rangle = \lambda.$$

Hence eigenvalues of Hermitian matrix are real.

(\Leftarrow) Suppose eigenvalues of A are real. From spectral theorem, normal matrix A can be written by

$$A = \sum_i \lambda_i |\lambda_i\rangle \langle \lambda_i| \tag{2.5}$$

where λ_i are real eigenvalues with eigenvectors $|\lambda_i\rangle$. By taking adjoint, we get

$$\begin{aligned}
A^\dagger &= \sum_i \lambda_i^* |\lambda_i\rangle \langle \lambda_i| \\
&= \sum_i \lambda_i |\lambda_i\rangle \langle \lambda_i| \quad (\because \lambda_i \text{ are real}) \\
&= A
\end{aligned}$$

Thus A is Hermitian. □

2.18

Suppose $|v\rangle$ is a eigenvector with corresponding eigenvalue λ .

$$\begin{aligned}
 U|v\rangle &= \lambda|v\rangle. \\
 1 &= \langle v|v\rangle \\
 &= \langle v|I|v\rangle \\
 &= \langle v|U^\dagger U|v\rangle \\
 &= \lambda\lambda^* \langle v|v\rangle \\
 &= \|\lambda\|^2 \\
 \therefore \lambda &= e^{i\theta}
 \end{aligned}$$

2.19

$$X^2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

2.20

$$\begin{aligned}
 U &\equiv \sum_i |w_i\rangle \langle v_i| \\
 A'_{ij} &= \langle v_i|A|v_j\rangle \\
 &= \langle v_i|UU^\dagger AUU^\dagger|v_j\rangle \\
 &= \sum_{p,q,r,s} \langle v_i|w_p\rangle \langle v_p|v_q\rangle \langle w_q|A|w_r\rangle \langle v_r|v_s\rangle \langle w_s|v_j\rangle \\
 &= \sum_{p,q,r,s} \langle v_i|w_p\rangle \delta_{pq} A''_{qr} \delta_{rs} \langle w_s|v_j\rangle \\
 &= \sum_{p,r} \langle v_i|w_p\rangle \langle w_r|v_j\rangle A''_{pr}
 \end{aligned}$$

2.21

Suppose M be Hermitian. Then $M = M^\dagger$.

$$\begin{aligned}
 M &= IMI \\
 &= (P+Q)M(P+Q) \\
 &= PMP + QMP + PMQ + QMQ
 \end{aligned}$$

Now $PMP = \lambda P$, $QMP = 0$, $PMQ = PM^\dagger Q = (QMP)^* = 0$. Thus $M = PMP + QMQ$. Next prove QMQ is normal.

$$\begin{aligned}
 QMQ(QMQ)^\dagger &= QMQQM^\dagger Q \\
 &= QM^\dagger QMQ \quad (M = M^\dagger) \\
 &= (QM^\dagger Q)QMQ
 \end{aligned}$$

Therefore QMQ is normal. By induction, QMQ is diagonal ... (following is same as Box 2.2)

2.22

Suppose A is a Hermitian operator and $|v_i\rangle$ are eigenvectors of A with eigenvalues λ_i . Then

$$\langle v_i|A|v_j\rangle = \lambda_j \langle v_i|v_j\rangle.$$

On the other hand,

$$\langle v_i|A|v_j\rangle = \langle v_i|A^\dagger|v_j\rangle = \langle v_j|A|v_i\rangle^* = \lambda_i^* \langle v_j|v_i\rangle^* = \lambda_i^* \langle v_i|v_j\rangle = \lambda_i \langle v_i|v_j\rangle$$

Thus

$$(\lambda_i - \lambda_j) \langle v_i|v_j\rangle = 0.$$

If $\lambda_i \neq \lambda_j$, then $\langle v_i|v_j\rangle = 0$.

2.23

Suppose P is projector and $|\lambda\rangle$ are eigenvectors of P with eigenvalues λ . Then $P^2 = P$.

$$P|\lambda\rangle = \lambda|\lambda\rangle \text{ and } P|\lambda\rangle = P^2|\lambda\rangle = \lambda P|\lambda\rangle = \lambda^2|\lambda\rangle.$$

Therefore

$$\begin{aligned} \lambda &= \lambda^2 \\ \lambda(\lambda - 1) &= 0 \\ \lambda &= 0 \text{ or } 1. \end{aligned}$$

2.24

Def of positive $\langle v|A|v\rangle \geq 0$ for all $|v\rangle$.

Suppose A is a positive operator. A can be decomposed as follows.

$$\begin{aligned} A &= \frac{A + A^\dagger}{2} + i \frac{A - A^\dagger}{2i} \\ &= B + iC \quad \text{where } B = \frac{A + A^\dagger}{2}, \quad C = \frac{A - A^\dagger}{2i}. \end{aligned}$$

Now operators B and C are Hermitian.

$$\begin{aligned} \langle v|A|v\rangle &= \langle v|B + iC|v\rangle \\ &= \langle v|B|v\rangle + i \langle v|C|v\rangle \\ &= \alpha + i\beta \quad \text{where } \alpha = \langle v|B|v\rangle, \quad \beta = \langle v|C|v\rangle. \end{aligned}$$

Since B and C are Hermitian, $\alpha, \beta \in \mathbb{R}$. From def of positive operator, β should be vanished because $\langle v|A|v\rangle$ is real. Hence $\beta = \langle v|C|v\rangle = 0$ for all $|v\rangle$, i.e. $C = 0$.

Therefore $A = A^\dagger$.

Reference: MIT 8.05 Lecture note by Prof. Barton Zwiebach.

https://ocw.mit.edu/courses/physics/8-05-quantum-physics-ii-fall-2013/lecture-notes/MIT8_05F13_Chap_03.pdf

Proposition. 2.0.1. *Let T be a linear operator in a complex vector space V .*

If $(u, Tv) = 0$ for all $u, v \in V$, then $T = 0$.

Proof. Suppose $u = Tv$. Then $(Tv, Tv) = 0$ for all v implies that $Tv = 0$ for all v . Therefore $T = 0$. \square

Theorem. 2.0.1. *If $(v, Av) = 0$ for all $v \in V$, then $A = 0$.*

Proof. First, we show that $(u, Tv) = 0$ if $(v, Av) = 0$. Then apply proposition 2.0.1. Suppose $u, v \in V$. Then (u, Tv) is decomposed as

$$(u, Tv) = \frac{1}{4} \left[(u+v, T(u+v)) - (u-v, T(u-v)) + \frac{1}{i}(u+iv, T(u+iv)) - \frac{1}{i}(u-iv, T(u-iv)) \right].$$

If $(v, Tv) = 0$ for all $v \in V$, the right hand side of above eqn vanishes. Thus $(u, Tv) = 0$ for all $u, v \in V$. Then $T = 0$. \square

2.25

$$\langle \psi | A^\dagger A | \psi \rangle = \|A | \psi \rangle\|^2 \geq 0 \text{ for all } | \psi \rangle.$$

Thus $A^\dagger A$ is positive.

2.26

$$\begin{aligned} |\psi\rangle^{\otimes 2} &= \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \otimes \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \\ &= \frac{1}{2}(|00\rangle + |01\rangle + |10\rangle + |11\rangle) \\ &= \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned}
|\psi\rangle^{\otimes 3} &= \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \otimes \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \otimes \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \\
&= \frac{1}{2\sqrt{2}}(|000\rangle + |001\rangle + |010\rangle + |011\rangle + |100\rangle + |101\rangle + |110\rangle + |111\rangle) \\
&= \frac{1}{2\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}
\end{aligned}$$

2.27

$$\begin{aligned}
X \otimes Z &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \otimes \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \\
&= \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix}
\end{aligned}$$

$$\begin{aligned}
I \otimes X &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\
&= \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}
\end{aligned}$$

$$\begin{aligned}
X \otimes I &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \otimes \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\
&= \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}
\end{aligned}$$

In general, tensor product is not commutable.

2.28

$$\begin{aligned}
(A \otimes B)^* &= \begin{bmatrix} A_{11}B & \cdots & A_{1n}B \\ \vdots & \ddots & \vdots \\ A_{m1}B & \cdots & A_{mn}B \end{bmatrix}^* \\
&= \begin{bmatrix} A_{11}^*B^* & \cdots & A_{1n}^*B^* \\ \vdots & \ddots & \vdots \\ A_{m1}^*B^* & \cdots & A_{mn}^*B^* \end{bmatrix} \\
&= A^* \otimes B^*.
\end{aligned}$$

$$\begin{aligned}
(A \otimes B)^T &= \begin{bmatrix} A_{11}B & \cdots & A_{1n}B \\ \vdots & \ddots & \vdots \\ A_{m1}B & \cdots & A_{mn}B \end{bmatrix}^T \\
&= \begin{bmatrix} A_{11}B^T & \cdots & A_{m1}B^T \\ \vdots & \ddots & \vdots \\ A_{1n}B^T & \cdots & A_{mn}B^T \end{bmatrix} \\
&= \begin{bmatrix} A_{11}B^T & \cdots & A_{1m}B^T \\ \vdots & \ddots & \vdots \\ A_{n1}B^T & \cdots & A_{nm}B^T \end{bmatrix} \\
&= A^T \otimes B^T.
\end{aligned}$$

$$\begin{aligned}
(A \otimes B)^\dagger &= ((A \otimes B)^*)^T \\
&= (A^* \otimes B^*)^T \\
&= (A^*)^T \otimes (B^*)^T \\
&= A^\dagger \otimes B^\dagger.
\end{aligned}$$

2.29

Suppose U_1 and U_2 are unitary operators. Then

$$\begin{aligned}
(U_1 \otimes U_2)(U_1 \otimes U_2)^\dagger &= U_1 U_1^\dagger \otimes U_2 U_2^\dagger \\
&= I \otimes I.
\end{aligned}$$

Similarly,

$$(U_1 \otimes U_2)^\dagger (U_1 \otimes U_2) = I \otimes I.$$

2.30

Suppose A and B are Hermitian operators. Then

$$(A \otimes B)^\dagger = A^\dagger \otimes B^\dagger = A \otimes B. \quad (2.6)$$

Thus $A \otimes B$ is Hermitian.

2.31

Suppose A and B are positive operators. Then

$$\langle \psi | \otimes \langle \phi | (A \otimes B) | \psi \rangle \otimes | \phi \rangle = \langle \psi | A | \psi \rangle \langle \phi | B | \phi \rangle.$$

Since A and B are positive operators, $\langle \psi | A | \psi \rangle \geq 0$ and $\langle \phi | B | \phi \rangle \geq 0$ for all $|\psi\rangle, |\phi\rangle$. Then $\langle \psi | A | \psi \rangle \langle \phi | B | \phi \rangle \geq 0$. Thus $A \otimes B$ is positive if A and B are positive.

2.32

Suppose P_1 and P_2 are projectors. Then

$$\begin{aligned}(P_1 \otimes P_2)^2 &= P_1^2 \otimes P_2^2 \\ &= P_1 \otimes P_2.\end{aligned}$$

Thus $P_1 \otimes P_2$ is also projector.

2.33

$$H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \quad (2.7)$$

$$H^{\otimes 2} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \otimes \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

2.34

Suppose $A = \begin{bmatrix} 4 & 3 \\ 3 & 4 \end{bmatrix}$.

$$\begin{aligned}\det(A - \lambda I) &= (4 - \lambda)^2 - 3^2 \\ &= \lambda^2 - 8\lambda + 7 \\ &= (\lambda - 1)(\lambda - 7)\end{aligned}$$

Eigenvalues of A are $\lambda = 1, 7$. Corresponding eigenvectors are $|\lambda = 1\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$, $|\lambda = 7\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

Thus

$$A = |\lambda = 1\rangle\langle\lambda = 1| + 7|\lambda = 7\rangle\langle\lambda = 7|.$$

$$\begin{aligned}\sqrt{A} &= |\lambda = 1\rangle\langle\lambda = 1| + \sqrt{7}|\lambda = 7\rangle\langle\lambda = 7| \\ &= \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} + \frac{\sqrt{7}}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \\ &= \frac{1}{2} \begin{bmatrix} 1 + \sqrt{7} & -1 + \sqrt{7} \\ -1 + \sqrt{7} & 1 + \sqrt{7} \end{bmatrix}\end{aligned}$$

$$\begin{aligned}\log(A) &= \log(1) |\lambda = 1\rangle\langle\lambda = 1| + \log(7) |\lambda = 7\rangle\langle\lambda = 7| \\ &= \frac{\log(7)}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}\end{aligned}$$

2.35

$$\begin{aligned}
\vec{v} \cdot \vec{\sigma} &= \sum_{i=1}^3 v_i \sigma_i \\
&= v_1 \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + v_2 \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} + v_3 \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \\
&= \begin{bmatrix} v_3 & v_1 - iv_2 \\ v_1 + iv_2 & -v_3 \end{bmatrix}
\end{aligned}$$

$$\begin{aligned}
\det(\vec{v} \cdot \vec{\sigma} - \lambda I) &= (v_3 - \lambda)(-v_3 - \lambda) - (v_1 - iv_2)(v_1 + iv_2) \\
&= \lambda^2 - (v_1^2 + v_2^2 + v_3^2) \\
&= \lambda^2 - 1 \quad (\because |\vec{v}| = 1)
\end{aligned}$$

Eigenvalues are $\lambda = \pm 1$. Let $|\lambda_{\pm 1}\rangle$ be eigenvectors with eigenvalues ± 1 .

Since $\vec{v} \cdot \vec{\sigma}$ is Hermitian, $\vec{v} \cdot \vec{\sigma}$ is diagonalizable. Then

$$\vec{v} \cdot \vec{\sigma} = |\lambda_1\rangle\langle\lambda_1| - |\lambda_{-1}\rangle\langle\lambda_{-1}|$$

Thus

$$\begin{aligned}
\exp(i\theta \vec{v} \cdot \vec{\sigma}) &= e^{i\theta} |\lambda_1\rangle\langle\lambda_1| + e^{-i\theta} |\lambda_{-1}\rangle\langle\lambda_{-1}| \\
&= (\cos \theta + i \sin \theta) |\lambda_1\rangle\langle\lambda_1| + (\cos \theta - i \sin \theta) |\lambda_{-1}\rangle\langle\lambda_{-1}| \\
&= \cos \theta (|\lambda_1\rangle\langle\lambda_1| + |\lambda_{-1}\rangle\langle\lambda_{-1}|) + i \sin \theta (|\lambda_1\rangle\langle\lambda_1| - |\lambda_{-1}\rangle\langle\lambda_{-1}|) \\
&= \cos(\theta) I + i \sin(\theta) \vec{v} \cdot \vec{\sigma}.
\end{aligned}$$

\therefore Since $\vec{v} \cdot \vec{\sigma}$ is Hermitian, $|\lambda_1\rangle$ and $|\lambda_{-1}\rangle$ are orthogonal. Thus

$$|\lambda_1\rangle\langle\lambda_1| + |\lambda_{-1}\rangle\langle\lambda_{-1}| = I.$$

2.36

$$\begin{aligned}
\text{Tr}(\sigma_1) &= \text{Tr} \left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right) = 0 \\
\text{Tr}(\sigma_2) &= \text{Tr} \left(\begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \right) = 0 \\
\text{Tr}(\sigma_3) &= \text{Tr} \left(\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \right) = 1 - 1 = 0
\end{aligned}$$

2.37

$$\begin{aligned}
\text{Tr}(AB) &= \sum_i \langle i|AB|i\rangle \\
&= \sum_i \langle i|AIB|i\rangle \\
&= \sum_{i,j} \langle i|A|j\rangle \langle j|B|i\rangle \\
&= \sum_{i,j} \langle j|B|i\rangle \langle i|A|j\rangle \\
&= \sum_j \langle j|BA|j\rangle \\
&= \text{Tr}(BA)
\end{aligned}$$

2.38

$$\begin{aligned}
\text{Tr}(A+B) &= \sum_i \langle i|A+B|i\rangle \\
&= \sum_i (\langle i|A|i\rangle + \langle i|B|i\rangle) \\
&= \sum_i \langle i|A|i\rangle + \sum_i \langle i|B|i\rangle \\
&= \text{Tr}(A) + \text{Tr}(B).
\end{aligned}$$

$$\begin{aligned}
\text{Tr}(zA) &= \sum_i \langle i|zA|i\rangle \\
&= \sum_i z \langle i|A|i\rangle \\
&= z \sum_i \langle i|A|i\rangle \\
&= z \text{Tr}(A).
\end{aligned}$$

2.39

$$(1) (A, B) \equiv \text{Tr}(A^\dagger B).$$

(i)

$$\begin{aligned}
\left(A, \sum_i \lambda_i B_i\right) &= \text{Tr} \left[A^\dagger \left(\sum_i \lambda_i B_i \right) \right] \\
&= \text{Tr}(A^\dagger \lambda_1 B_1) + \cdots + \text{Tr}(A^\dagger \lambda_n B_n) \quad (\because \text{Exercise 2.38}) \\
&= \lambda_1 \text{Tr}(A^\dagger B_1) + \cdots + \lambda_n \text{Tr}(A^\dagger B_n) \\
&= \sum_i \lambda_i \text{Tr}(A^\dagger B_i)
\end{aligned}$$

(ii)

$$\begin{aligned}
(A, B)^* &= \left(\text{Tr}(A^\dagger B) \right)^* \\
&= \left(\sum_{i,j} \langle i|A^\dagger|j\rangle \langle j|B|i\rangle \right)^* \\
&= \sum_{i,j} \langle i|A^\dagger|j\rangle^* \langle j|B|i\rangle^* \\
&= \sum_{i,j} \langle j|B|i\rangle^* \langle i|A^\dagger|j\rangle^* \\
&= \sum_{i,j} \langle i|B^\dagger|j\rangle \langle j|A|i\rangle \\
&= \sum_i \langle i|B^\dagger A|i\rangle \\
&= \text{Tr}(B^\dagger A) \\
&= (B, A).
\end{aligned}$$

(iii)

$$\begin{aligned}
(A, A) &= \text{Tr}(A^\dagger A) \\
&= \sum_i \langle i|A^\dagger A|i\rangle
\end{aligned}$$

Since $A^\dagger A$ is positive, $\langle i|A^\dagger A|i\rangle \geq 0$ for all $|i\rangle$.

Let a_i be i -th column of A . If $\langle i|A^\dagger A|i\rangle = 0$, then

$$\langle i|A^\dagger A|i\rangle = a_i^\dagger a_i = \|a_i\|^2 = 0 \text{ iff } a_i = \mathbf{0}.$$

Therefore $(A, A) = 0$ iff $A = \mathbf{0}$.

(2)

(3)

2.40

$$\begin{aligned}
[X, Y] &= XY - YX \\
&= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} - \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\
&= \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} - \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix} \\
&= \begin{bmatrix} 2i & 0 \\ 0 & -2i \end{bmatrix} \\
&= 2iZ
\end{aligned}$$

$$\begin{aligned}
[Y, Z] &= \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \\
&= \begin{bmatrix} 0 & 2i \\ 2i & 0 \end{bmatrix} \\
&= 2iX
\end{aligned}$$

$$\begin{aligned}
[Z, X] &= \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} - \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \\
&= 2i \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \\
&= 2iY
\end{aligned}$$

2.41

$$\begin{aligned}
\{\sigma_1, \sigma_2\} &= \sigma_1\sigma_2 + \sigma_2\sigma_1 \\
&= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} + \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\
&= \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} + \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix} \\
&= 0
\end{aligned}$$

$$\begin{aligned}
\{\sigma_2, \sigma_3\} &= \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \\
&= 0
\end{aligned}$$

$$\begin{aligned}
\{\sigma_3, \sigma_1\} &= \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \\
&= 0
\end{aligned}$$

$$\begin{aligned}
\sigma_0^2 &= I^2 = I \\
\sigma_1^2 &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}^2 = I \\
\sigma_2^2 &= \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}^2 = I \\
\sigma_3^2 &= \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}^2 = I
\end{aligned}$$

2.42

$$\frac{[A, B] + \{A, B\}}{2} = \frac{AB - BA + AB + BA}{2} = AB$$

2.43

From eq (2.75) and eq (2.76), $\{\sigma_j, \sigma_k\} = 2\delta_{jk}I$. From eq (2.77),

$$\begin{aligned}\sigma_j \sigma_k &= \frac{[\sigma_j, \sigma_k] + \{\sigma_j, \sigma_k\}}{2} \\ &= \frac{2i \sum_{l=1}^3 \epsilon_{jkl} \sigma_l + 2\delta_{jk}I}{2} \\ &= \delta_{jk}I + i \sum_{l=1}^3 \epsilon_{jkl} \sigma_l\end{aligned}$$

2.44

By assumption, $[A, B] = 0$ and $\{A, B\} = 0$, then $AB = 0$. Since A is invertible, multiply by A^{-1} from left, then

$$\begin{aligned}A^{-1}AB &= 0 \\ IB &= 0 \\ B &= 0.\end{aligned}$$

2.45

$$\begin{aligned}[A, B]^\dagger &= (AB - BA)^\dagger \\ &= B^\dagger A^\dagger - A^\dagger B^\dagger \\ &= [B^\dagger, A^\dagger]\end{aligned}$$

2.46

$$\begin{aligned}[A, B] &= AB - BA \\ &= -(BA - AB) \\ &= -[B, A]\end{aligned}$$

2.47

$$\begin{aligned}(i[A, B])^\dagger &= -i[A, B]^\dagger \\ &= -i[B^\dagger, A^\dagger] \\ &= -i[B, A] \\ &= i[A, B]\end{aligned}$$

2.48

(Positive)

Since P is positive, it is diagonalizable. Then $P = \sum_i \lambda_i |i\rangle\langle i|$, ($\lambda_i \geq 0$).

$$J = \sqrt{P^\dagger P} = \sqrt{PP} = \sqrt{P^2} = \sum_i \sqrt{\lambda_i^2} |i\rangle\langle i| = \sum_i \lambda_i |i\rangle\langle i| = P.$$

Therefore polar decomposition of P is $P = UP$ for all P . Thus $U = I$, then $P = P$.

(Unitary)

Suppose unitary U is decomposed by $U = WJ$ where W is unitary and J is positive, $J = \sqrt{U^\dagger U}$.

$$J = \sqrt{U^\dagger U} = \sqrt{I} = I$$

Since unitary operators are invertible, $W = UJ^{-1} = UI^{-1} = UI = U$. Thus polar decomposition of U is $U = U$.

(Hermitian)

Suppose $H = UJ$.

$$J = \sqrt{H^\dagger H} = \sqrt{HH} = \sqrt{H^2}.$$

Thus $H = U\sqrt{H^2}$.

In general, $H \neq \sqrt{H^2}$.

From spectral decomposition, $H = \sum_i \lambda_i |i\rangle\langle i|$, $\lambda_i \in \mathbb{R}$.

$$\sqrt{H^2} = \sqrt{\sum_i \lambda_i^2 |i\rangle\langle i|} = \sum_i \sqrt{\lambda_i^2} |i\rangle\langle i| = \sum_i |\lambda_i| |i\rangle\langle i| \neq H$$

2.49

Normal matrix is diagonalizable, $A = \sum_i \lambda_i |i\rangle\langle i|$.

$$J = \sqrt{A^\dagger A} = \sum_i |\lambda_i| |i\rangle\langle i|.$$

$$U = \sum_i |e_i\rangle\langle i|$$

$$A = UJ = \sum_i |\lambda_i| |e_i\rangle\langle i|.$$

2.50

Define $A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$. $A^\dagger A = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$.

Characteristic equation of $A^\dagger A$ is $\det(A^\dagger A - \lambda I) = \lambda^2 - 3\lambda + 1 = 0$. Eigenvalues of $A^\dagger A$ are $\lambda_{\pm} = \frac{3 \pm \sqrt{5}}{2}$ and associated eigenvectors are $|\lambda_{\pm}\rangle = \frac{1}{\sqrt{10 \mp 2\sqrt{5}}} \begin{bmatrix} 2 \\ -1 \pm \sqrt{5} \end{bmatrix}$.

$$A^\dagger A = \lambda_+ |\lambda_+\rangle\langle\lambda_+| + \lambda_- |\lambda_-\rangle\langle\lambda_-|.$$

$$\begin{aligned} J = \sqrt{A^\dagger A} &= \sqrt{\lambda_+} |\lambda_+\rangle\langle\lambda_+| + \sqrt{\lambda_-} |\lambda_-\rangle\langle\lambda_-| \\ &= \sqrt{\frac{3+\sqrt{5}}{2}} \cdot \frac{5-\sqrt{5}}{40} \begin{bmatrix} 4 & 2\sqrt{5}-2 \\ 2\sqrt{5}-2 & 6-2\sqrt{5} \end{bmatrix} + \sqrt{\frac{3-\sqrt{5}}{2}} \cdot \frac{5+\sqrt{5}}{40} \begin{bmatrix} 4 & -2\sqrt{5}-2 \\ -2\sqrt{5}-2 & 6+2\sqrt{5} \end{bmatrix} \end{aligned}$$

$$J^{-1} = \frac{1}{\sqrt{\lambda_+}} |\lambda_+\rangle\langle\lambda_+| + \frac{1}{\sqrt{\lambda_-}} |\lambda_-\rangle\langle\lambda_-|.$$

$$U = AJ^{-1}$$

I'm tired.

2.51

$$H^\dagger H = \left(\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right)^\dagger \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = I.$$

2.52

$$H^\dagger = \left(\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right)^\dagger = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = H.$$

Thus

$$H^2 = I.$$

2.53

$$\begin{aligned} \det(H - \lambda I) &= \left(\frac{1}{\sqrt{2}} - \lambda \right) \left(-\frac{1}{\sqrt{2}} - \lambda \right) - \frac{1}{2} \\ &= \lambda^2 - \frac{1}{2} - \frac{1}{2} \\ &= \lambda^2 - 1 \end{aligned}$$

Eigenvalues are $\lambda_\pm = \pm 1$ and associated eigenvectors are $|\lambda_\pm\rangle = \frac{1}{\sqrt{4 \mp 2\sqrt{2}}} \begin{bmatrix} 1 \\ -1 \pm \sqrt{2} \end{bmatrix}$.

2.54

Since $[A, B] = 0$, A and B are simultaneously diagonalize, $A = \sum_i a_i |i\rangle\langle i|$, $B = \sum_i b_i |i\rangle\langle i|$.

$$\begin{aligned}
 \exp(A) \exp(B) &= \left(\sum_i \exp(a_i) |i\rangle\langle i| \right) \left(\sum_i \exp(b_i) |i\rangle\langle i| \right) \\
 &= \sum_{i,j} \exp(a_i + b_j) |i\rangle\langle i| j\rangle\langle j| \\
 &= \sum_{i,j} \exp(a_i + b_j) |i\rangle\langle j| \delta_{i,j} \\
 &= \sum_i \exp(a_i + b_i) |i\rangle\langle i| \\
 &= \exp(A + B)
 \end{aligned}$$

2.55

$$H = \sum_E E |E\rangle\langle E|$$

$$\begin{aligned}
 U(t_2 - t_1) U^\dagger(t_2 - t_1) &= \exp\left(-\frac{iH(t_2 - t_1)}{\hbar}\right) \exp\left(\frac{iH(t_2 - t_1)}{\hbar}\right) \\
 &= \sum_{E, E'} \left(\exp\left(-\frac{iE(t_2 - t_1)}{\hbar}\right) |E\rangle\langle E| \right) \left(\exp\left(-\frac{iE'(t_2 - t_1)}{\hbar}\right) |E'\rangle\langle E'| \right) \\
 &= \sum_{E, E'} \left(\exp\left(-\frac{i(E - E')(t_2 - t_1)}{\hbar}\right) |E\rangle\langle E'| \delta_{E, E'} \right) \\
 &= \sum_E \exp(0) |E\rangle\langle E| \\
 &= \sum_E |E\rangle\langle E| \\
 &= I
 \end{aligned}$$

Similarly, $U^\dagger(t_2 - t_1) U(t_2 - t_1) = I$.

2.56

$$U = \sum_i \lambda_i |\lambda_i\rangle\langle \lambda_i| \quad (|\lambda_i| = 1).$$

$$\log(U) = \sum_j \log(\lambda_j) |\lambda_j\rangle\langle \lambda_j| = \sum_j i\theta_j |\lambda_j\rangle\langle \lambda_j| \quad \text{where } \theta_j = \arg(\lambda_j)$$

$$K = -i \log(U) = \sum_j \theta_j |\lambda_j\rangle\langle \lambda_j|.$$

$$K^\dagger = (-i \log U)^\dagger = \left(\sum_j \theta_j |\lambda_j\rangle\langle \lambda_j| \right)^\dagger = \sum_j \theta_j^* |\lambda_j\rangle\langle \lambda_j| = \sum_j \theta_j |\lambda_j\rangle\langle \lambda_j| = K$$

2.57

$$\begin{aligned}
|\phi\rangle &\equiv \frac{L_l |\psi\rangle}{\sqrt{\langle\psi|L_l^\dagger L_l|\psi\rangle}} \\
\langle\phi|M_m^\dagger M_m|\phi\rangle &= \frac{\langle\psi|L_l^\dagger M_m^\dagger M_m L_l|\psi\rangle}{\langle\psi|L_l^\dagger L_l|\psi\rangle} \\
\frac{M_m |\phi\rangle}{\sqrt{\langle\phi|M_m^\dagger M_m|\phi\rangle}} &= \frac{M_m L_l |\psi\rangle}{\sqrt{\langle\psi|L_l^\dagger L_l|\psi\rangle}} \cdot \frac{\sqrt{\langle\psi|L_l^\dagger L_l|\psi\rangle}}{\sqrt{\langle\psi|L_l^\dagger M_m^\dagger M_m L_l|\psi\rangle}} = \frac{M_m L_l |\psi\rangle}{\sqrt{\langle\psi|L_l^\dagger M_m^\dagger M_m L_l|\psi\rangle}} = \frac{N_{lm} |\psi\rangle}{\sqrt{\langle\psi|N_{lm}^\dagger N_{lm}|\psi\rangle}}
\end{aligned}$$

2.58

$$\begin{aligned}
\langle M \rangle &= \langle\psi|M|\psi\rangle = \langle\psi|m|\psi\rangle = m \langle\psi|\psi\rangle = m \\
\langle M^2 \rangle &= \langle\psi|M^2|\psi\rangle = \langle\psi|m^2|\psi\rangle = m^2 \langle\psi|\psi\rangle = m^2 \\
\text{deviation} &= \langle M^2 \rangle - \langle M \rangle^2 = m^2 - m^2 = 0.
\end{aligned}$$

2.59

$$\begin{aligned}
\langle X \rangle &= \langle 0|X|0\rangle = \langle 0|1\rangle = 0 \\
\langle X^2 \rangle &= \langle 0|X^2|0\rangle = \langle 0|X|1\rangle = \langle 0|0\rangle = 1 \\
\text{standard deviation} &= \sqrt{\langle X^2 \rangle - \langle X \rangle^2} = 1
\end{aligned}$$

2.60

$$\begin{aligned}
\vec{v} \cdot \vec{\sigma} &= \sum_{i=1}^3 v_i \sigma_i \\
&= v_1 \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + v_2 \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} + v_3 \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \\
&= \begin{bmatrix} v_3 & v_1 - iv_2 \\ v_1 + iv_2 & -v_3 \end{bmatrix}
\end{aligned}$$

$$\begin{aligned}
\det(\vec{v} \cdot \vec{\sigma} - \lambda I) &= (v_3 - \lambda)(-v_3 - \lambda) - (v_1 - iv_2)(v_1 + iv_2) \\
&= \lambda^2 - (v_1^2 + v_2^2 + v_3^2) \\
&= \lambda^2 - 1 \quad (\because |\vec{v}| = 1)
\end{aligned}$$

Eigenvalues are $\lambda = \pm 1$.(i) if $\lambda = 1$

$$\begin{aligned}
\vec{v} \cdot \vec{\sigma} - \lambda I &= \vec{v} \cdot \vec{\sigma} - I \\
&= \begin{bmatrix} v_3 - 1 & v_1 - iv_2 \\ v_1 + iv_2 & -v_3 - 1 \end{bmatrix}
\end{aligned}$$

Normalized eigenvector is $|\lambda_1\rangle = \sqrt{\frac{1+v_3}{2}} \begin{bmatrix} 1 \\ \frac{1-v_3}{v_1-iv_2} \end{bmatrix}$.

$$\begin{aligned} |\lambda_1\rangle\langle\lambda_1| &= \frac{1+v_3}{2} \begin{bmatrix} 1 \\ \frac{1-v_3}{v_1-iv_2} \end{bmatrix} \begin{bmatrix} 1 & \frac{1-v_3}{v_1+iv_2} \end{bmatrix} \\ &= \frac{1+v_3}{2} \begin{bmatrix} 1 & \frac{v_1-iv_2}{1+v_3} \\ \frac{v_1+iv_2}{1+v_3} & \frac{1-v_3}{1+v_3} \end{bmatrix} \\ &= \frac{1}{2} \begin{bmatrix} 1+v_3 & v_1-iv_2 \\ v_1+iv_2 & 1-v_3 \end{bmatrix} \\ &= \frac{1}{2} \left(I + \begin{bmatrix} v_3 & v_1-iv_2 \\ v_1+iv_2 & -v_3 \end{bmatrix} \right) \\ &= \frac{1}{2} (I + \vec{v} \cdot \vec{\sigma}) \end{aligned}$$

(ii) If $\lambda = -1$.

$$\begin{aligned} \vec{v} \cdot \vec{\sigma} - \lambda I &= \vec{v} \cdot \vec{\sigma} + I \\ &= \begin{bmatrix} v_3+1 & v_1-iv_2 \\ v_1+iv_2 & -v_3+1 \end{bmatrix} \end{aligned}$$

Normalized eigenvalue is $|\lambda_{-1}\rangle = \sqrt{\frac{1-v_3}{2}} \begin{bmatrix} 1 \\ -\frac{1+v_3}{v_1-iv_2} \end{bmatrix}$.

$$\begin{aligned} |\lambda_{-1}\rangle\langle\lambda_{-1}| &= \frac{1-v_3}{2} \begin{bmatrix} 1 \\ -\frac{1+v_3}{v_1-iv_2} \end{bmatrix} \begin{bmatrix} 1 & -\frac{1+v_3}{v_1+iv_2} \end{bmatrix} \\ &= \frac{1-v_3}{2} \begin{bmatrix} 1 & -\frac{v_1-iv_2}{1-v_3} \\ -\frac{v_1+iv_2}{1-v_3} & \frac{1+v_3}{1-v_3} \end{bmatrix} \\ &= \frac{1}{2} \begin{bmatrix} 1-v_3 & -(v_1-iv_2) \\ -(v_1+iv_2) & 1+v_3 \end{bmatrix} \\ &= \frac{1}{2} \left(I - \begin{bmatrix} v_3 & v_1-iv_2 \\ v_1+iv_2 & -v_3 \end{bmatrix} \right) \\ &= \frac{1}{2} (I - \vec{v} \cdot \vec{\sigma}). \end{aligned}$$

While I review my proof, I notice that my proof has a defect. The case $(v_1, v_2, v_3) = (0, 0, 1)$, second component of eigenstate, $\frac{1-v_3}{v_1-iv_2}$, diverges. So I implicitly assume $v_1 - iv_2 \neq 0$. Hence my proof is incomplete.

Since the exercise doesn't require explicit form of projector, we should prove the problem more abstractly. In order to prove, we use the following properties of $\vec{v} \cdot \vec{\sigma}$

- $\vec{v} \cdot \vec{\sigma}$ is Hermitian
- $(\vec{v} \cdot \vec{\sigma})^2 = I$ where \vec{v} is a real unit vector.

We can easily check above conditions.

$$\begin{aligned} (\vec{v} \cdot \vec{\sigma})^\dagger &= (v_1\sigma_1 + v_2\sigma_2 + v_3\sigma_3)^\dagger \\ &= v_1\sigma_1^\dagger + v_2\sigma_2^\dagger + v_3\sigma_3^\dagger \\ &= v_1\sigma_1 + v_2\sigma_2 + v_3\sigma_3 \quad (\because \text{Pauli matrices are Hermitian.}) \\ &= \vec{v} \cdot \vec{\sigma} \end{aligned}$$

$$\begin{aligned}
(\vec{v} \cdot \vec{\sigma})^2 &= \sum_{j,k=1}^3 (v_j \sigma_j)(v_k \sigma_k) \\
&= \sum_{j,k=1}^3 v_j v_k \sigma_j \sigma_k \\
&= \sum_{j,k=1}^3 v_j v_k \left(\delta_{jk} I + i \sum_{l=1}^3 \epsilon_{jkl} \sigma_l \right) \quad (\because \text{eqn(2.78) page78}) \\
&= \sum_{j,k=1}^3 v_j v_k \delta_{jk} I + i \sum_{j,k,l=1}^3 \epsilon_{jkl} v_j v_k \sigma_l \\
&= \sum_{j=1}^3 v_j^2 I \\
&= I \quad \left(\because \sum_j v_j^2 = 1 \right)
\end{aligned}$$

Proof. Suppose $|\lambda\rangle$ is an eigenstate of $\vec{v} \cdot \vec{\sigma}$ with eigenvalue λ . Then

$$\begin{aligned}
\vec{v} \cdot \vec{\sigma} |\lambda\rangle &= \lambda |\lambda\rangle \\
(\vec{v} \cdot \vec{\sigma})^2 |\lambda\rangle &= \lambda^2 |\lambda\rangle
\end{aligned}$$

On the other hand $(\vec{v} \cdot \vec{\sigma})^2 = I$,

$$\begin{aligned}
(\vec{v} \cdot \vec{\sigma})^2 |\lambda\rangle &= I |\lambda\rangle = |\lambda\rangle \\
\therefore \lambda^2 |\lambda\rangle &= |\lambda\rangle.
\end{aligned}$$

Thus $\lambda^2 = 1 \Rightarrow \lambda = \pm 1$. Therefore $\vec{v} \cdot \vec{\sigma}$ has eigenvalues ± 1 .

Let $|\lambda_1\rangle$ and $|\lambda_{-1}\rangle$ are eigenvectors with eigenvalues 1 and -1 , respectively. I will prove that $P_{\pm} = |\lambda_{\pm 1}\rangle\langle\lambda_{\pm 1}|$.

In order to prove above equation, all we have to do is prove following condition. (see Theorem 2.0.1)

$$\langle\psi|(P_{\pm} - |\lambda_{\pm 1}\rangle\langle\lambda_{\pm 1}|)|\psi\rangle = 0 \text{ for all } |\psi\rangle \in \mathbb{C}^2. \quad (2.8)$$

Since $\vec{v} \cdot \vec{\sigma}$ is Hermitian, $|\lambda_1\rangle$ and $|\lambda_{-1}\rangle$ are orthonormal vector (\because Exercise 2.22). Let $|\psi\rangle \in \mathbb{C}^2$ be an arbitrary state. $|\psi\rangle$ can be written as

$$|\psi\rangle = \alpha |\lambda_1\rangle + \beta |\lambda_{-1}\rangle \quad (|\alpha|^2 + |\beta|^2 = 1, \alpha, \beta \in \mathbb{C}).$$

$$\begin{aligned}
\langle \psi | (P_{\pm} - |\lambda_{\pm}\rangle\langle\lambda_{\pm}|) | \psi \rangle &= \langle \psi | P_{\pm} | \psi \rangle - \langle \psi | \lambda_{\pm} \rangle \langle \lambda_{\pm} | \psi \rangle . \\
\langle \psi | P_{\pm} | \psi \rangle &= \langle \psi | \frac{1}{2} (I \pm \vec{v} \cdot \vec{\sigma}) | \psi \rangle \\
&= \frac{1}{2} \pm \frac{1}{2} \langle \psi | \vec{v} \cdot \vec{\sigma} | \psi \rangle \\
&= \frac{1}{2} \pm \frac{1}{2} (|\alpha|^2 - |\beta|^2) \\
&= \frac{1}{2} \pm \frac{1}{2} (2|\alpha|^2 - 1) \quad (\because |\alpha|^2 + |\beta|^2 = 1) \\
\langle \psi | \lambda_1 \rangle \langle \lambda_1 | \psi \rangle &= |\alpha|^2 \\
\langle \psi | \lambda_{-1} \rangle \langle \lambda_{-1} | \psi \rangle &= |\beta|^2 = 1 - |\alpha|^2
\end{aligned}$$

Therefore $\langle \psi | (P_{\pm} - |\lambda_{\pm 1}\rangle\langle\lambda_{\pm 1}|) | \psi \rangle = 0$ for all $|\psi\rangle \in \mathbb{C}^2$. Thus $P_{\pm} = |\lambda_{\pm 1}\rangle\langle\lambda_{\pm 1}|$. \square

2.61

$$\begin{aligned}
\langle \lambda_1 | 0 \rangle \langle 0 | \lambda_1 \rangle &= \langle 0 | \lambda_1 \rangle \langle \lambda_1 | 0 \rangle \\
&= \langle 0 | \frac{1}{2} (I + \vec{v} \cdot \vec{\sigma}) | 0 \rangle \\
&= \frac{1}{2} (1 + v_3)
\end{aligned}$$

Post-measurement state is

$$\begin{aligned}
\frac{|\lambda_1\rangle \langle \lambda_1 | 0 \rangle}{\sqrt{\langle 0 | \lambda_1 \rangle \langle \lambda_1 | 0 \rangle}} &= \frac{1}{\sqrt{\frac{1}{2}(1 + v_3)}} \cdot \frac{1}{2} \begin{bmatrix} 1 + v_3 \\ v_1 + iv_2 \end{bmatrix} \\
&= \sqrt{\frac{1}{2}(1 + v_3)} \begin{bmatrix} 1 \\ \frac{v_1 + iv_2}{1 + v_3} \end{bmatrix} \\
&= \sqrt{\frac{1 + v_3}{2}} \begin{bmatrix} 1 \\ \frac{1 - v_3}{v_1 - iv_2} \end{bmatrix} \\
&= |\lambda_1\rangle .
\end{aligned}$$

2.62

Suppose M_m is a measurement operator. From the assumption, $E_m = M_m^\dagger M_m = M_m$. Then

$$\langle \psi | E_m | \psi \rangle = \langle \psi | M_m | \psi \rangle \geq 0.$$

for all $|\psi\rangle$.

Since M_m is positive operator, M_m is Hermitian. Therefore,

$$E_m = M_m^\dagger M_m = M_m M_m = M_m^2 = M_m.$$

Thus the measurement is a projective measurement.

2.63

$$\begin{aligned}
M_m^\dagger M_m &= \sqrt{E_m} U_m^\dagger U_m \sqrt{E_m} \\
&= \sqrt{E_m} I \sqrt{E_m} \\
&= E_m.
\end{aligned}$$

Since E_m is POVM, for arbitrary unitary U , $M_m^\dagger M_m$ is POVM.

2.64

Read following paper:

- Lu-Ming Duan, Guang-Can Guo. Probabilistic cloning and identification of linearly independent quantum states. Phys. Rev. Lett., 80:4999-5002, 1998. arXiv:quant-ph/9804064
<https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.80.4999>
<https://arxiv.org/abs/quant-ph/9804064>
- Stephen M. Barnett, Sarah Croke, Quantum state discrimination, arXiv:0810.1970 [quant-ph]
<https://arxiv.org/abs/0810.1970>
https://www.osapublishing.org/DirectPDFAccess/67EF4200-CBD2-8E68-1979E37886263936_176580/aop-1-2-238.pdf

2.65

$$|+\rangle \equiv \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |-\rangle \equiv \frac{|0\rangle - |1\rangle}{\sqrt{2}}$$

2.66

$$\begin{aligned}
X_1 Z_2 \left(\frac{|00\rangle + |11\rangle}{\sqrt{2}} \right) &= \frac{|10\rangle - |01\rangle}{\sqrt{2}} \\
\langle X_1 Z_2 \rangle &= \left(\frac{\langle 00| + \langle 11|}{\sqrt{2}} \right) X_1 Z_2 \left(\frac{|00\rangle + |11\rangle}{\sqrt{2}} \right) = \frac{\langle 00| + \langle 11|}{\sqrt{2}} \cdot \frac{|10\rangle - |01\rangle}{\sqrt{2}} = 0
\end{aligned}$$

2.67

Suppose W^\perp is the orthogonal complement of W . Then $V = W \oplus W^\perp$. Let $|w_i\rangle, |w'_j\rangle, |u'_j\rangle$ be orthonormal bases for $W, W^\perp, (\text{image}(U))^\perp$, respectively.

Define $U' : V \rightarrow V$ as $U' = \sum_i |u_i\rangle\langle w_i| + \sum_j |u'_j\rangle\langle w'_j|$, where $|u_i\rangle = U|w_i\rangle$.

Now

$$\begin{aligned}
(U')^\dagger U' &= \left(\sum_{i=1}^{\dim W} |w_i\rangle\langle u_i| + \sum_{j=1}^{\dim W^\perp} |w'_j\rangle\langle u'_j| \right) \left(\sum_i |u_i\rangle\langle w_i| + \sum_j |u'_j\rangle\langle w'_j| \right) \\
&= \sum_i |w_i\rangle\langle w_i| + \sum_j |w'_j\rangle\langle w'_j| = I
\end{aligned}$$

and

$$\begin{aligned} U'(U')^\dagger &= \left(\sum_i |u_i\rangle\langle w_i| + \sum_j |u'_j\rangle\langle w'_j| \right) \left(\sum_i |w_i\rangle\langle u_i| + \sum_j |w'_j\rangle\langle u'_j| \right) \\ &= \sum_i |u_i\rangle\langle u_i| + \sum_j |u'_j\rangle\langle u'_j| = I. \end{aligned}$$

Thus U' is a unitary operator. Moreover, for all $|w\rangle \in W$,

$$\begin{aligned} U'|w\rangle &= \left(\sum_i |u_i\rangle\langle w_i| + \sum_j |u'_j\rangle\langle w'_j| \right) |w\rangle \\ &= \sum_i |u_i\rangle \langle w_i|w\rangle + \sum_j |u'_j\rangle \langle w'_j|w\rangle \\ &= \sum_i |u_i\rangle \langle w_i|w\rangle \quad (\because |w'_j\rangle \perp |w\rangle) \\ &= \sum_i U |w_i\rangle \langle w_i|w\rangle \\ &= U |w\rangle. \end{aligned}$$

Therefore U' is an extension of U .

2.68

$$|\psi\rangle = \frac{|00\rangle + |11\rangle}{\sqrt{2}}.$$

Suppose $|a\rangle = a_0|0\rangle + a_1|1\rangle$ and $|b\rangle = b_0|0\rangle + b_1|1\rangle$.

$$|a\rangle|b\rangle = a_0b_0|00\rangle + a_0b_1|01\rangle + a_1b_0|10\rangle + a_1b_1|11\rangle.$$

If $|\psi\rangle = |a\rangle|b\rangle$, then $a_0b_0 = 1$, $a_0b_1 = 0$, $a_1b_0 = 0$, $a_1b_1 = 1$ since $\{|ij\rangle\}$ is an orthonormal basis.

If $a_0b_1 = 0$, then $a_0 = 0$ or $b_1 = 0$.

When $a_0 = 0$, this is contradiction to $a_0b_0 = 1$. When $b_1 = 0$, this is contradiction to $a_1b_1 = 1$.

Thus $|\psi\rangle \neq |a\rangle|b\rangle$.

2.69

Define Bell states as follows.

$$\begin{aligned}
 |\psi_1\rangle &\equiv \frac{|00\rangle + |11\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} \\
 |\psi_2\rangle &\equiv \frac{|00\rangle - |11\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix} \\
 |\psi_3\rangle &\equiv \frac{|01\rangle + |10\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} \\
 |\psi_4\rangle &\equiv \frac{|01\rangle - |10\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \end{bmatrix}
 \end{aligned}$$

First, we prove $\{|\psi_i\rangle\}$ is a linearly independent basis.

$$\begin{aligned}
 a_1 |\psi_1\rangle + a_2 |\psi_2\rangle + a_3 |\psi_3\rangle + a_4 |\psi_4\rangle &= 0 \\
 \therefore \frac{1}{\sqrt{2}} \begin{bmatrix} a_1 + a_2 \\ a_3 + a_4 \\ a_3 - a_4 \\ a_1 - a_2 \end{bmatrix} &= 0 \\
 \therefore \begin{cases} a_1 + a_2 = 0 \\ a_3 + a_4 = 0 \\ a_3 - a_4 = 0 \\ a_1 - a_2 = 0 \end{cases} \\
 \therefore a_1 = a_2 = a_3 = a_4 &= 0
 \end{aligned}$$

Thus $\{|\psi_i\rangle\}$ is a linearly independent basis.

Moreover $\| |\psi_i\rangle \| = 1$ and $\langle \psi_i | \psi_j \rangle = \delta_{ij}$ for $i, j = 1, 2, 3, 4$. Therefore $\{|\psi_i\rangle\}$ forms an orthonormal basis.

2.70

For any Bell states we get $\langle \psi_i | E \otimes I | \psi_i \rangle = \frac{1}{2}(\langle 0 | E | 0 \rangle + \langle 1 | E | 1 \rangle)$.

Suppose Eve measures the qubit Alice sent by measurement operators M_m . The probability that Eve gets result m is $p_i(m) = \langle \psi_i | M_m^\dagger M_m \otimes I | \psi_i \rangle$. Since $M_m^\dagger M_m$ is positive, $p_i(m)$ are same values for all $|\psi_i\rangle$. Thus Eve can't distinguish Bell states.

2.71

From spectral decomposition,

$$\begin{aligned}\rho &= \sum_i p_i |\psi_i\rangle\langle\psi_i|, \quad p_i \geq 0, \quad \sum_i p_i = 1. \\ \rho^2 &= \sum_{i,j} p_i p_j |i\rangle\langle i|j\rangle\langle j| \\ &= \sum_{i,j} p_i p_j |i\rangle\langle j| \delta_{ij} \\ &= \sum_i p_i^2 |i\rangle\langle i|\end{aligned}$$

$$\text{Tr}(\rho^2) = \text{Tr}\left(\sum_i p_i^2 |i\rangle\langle i|\right) = \sum_i p_i^2 \text{Tr}(|i\rangle\langle i|) = \sum_i p_i^2 \langle i|i\rangle = \sum_i p_i^2 \leq \sum_i p_i = 1 \quad (\because p_i^2 \leq p_i)$$

Suppose $\text{Tr}(\rho^2) = 1$. Then $\sum_i p_i^2 = 1$. Since $p_i^2 < p_i$ for $0 < p_i < 1$, only single p_i should be 1 and otherwise have to vanish. Therefore $\rho = |\psi_i\rangle\langle\psi_i|$. It is a pure state.

Conversely if ρ is pure, then $\rho = |\psi\rangle\langle\psi|$.

$$\text{Tr}(\rho^2) = \text{Tr}(|\psi\rangle\langle\psi| |\psi\rangle\langle\psi|) = \text{Tr}(|\psi\rangle\langle\psi|) = \langle\psi|\psi\rangle = 1.$$

2.72

(1) Since density matrix is Hermitian, matrix representation is $\rho = \begin{bmatrix} a & b \\ b^* & d \end{bmatrix}$, $a, d \in \mathbb{R}$ and $b \in \mathbb{C}$ w.r.t. standard basis. Because ρ is density matrix, $\text{Tr}(\rho) = a + d = 1$. Define $a = (1 + r_3)/2$, $d = (1 - r_3)/2$ and $b = (r_1 - ir_2)/2$, ($r_i \in \mathbb{R}$).

In this case,

$$\rho = \begin{bmatrix} a & b \\ b^* & d \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 + r_3 & r_1 - ir_2 \\ r_1 + ir_2 & 1 - r_3 \end{bmatrix} = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma}).$$

Thus for arbitrary density matrix ρ can be written as $\rho = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma})$.

Next, we derive the condition that ρ is positive.

If ρ is positive, all eigenvalues of ρ should be non-negative.

$$\begin{aligned}\det(\rho - \lambda I) &= (a - \lambda)(b - \lambda) - |b|^2 = \lambda^2 - (a + d)\lambda + ad - |b|^2 = 0 \\ \lambda &= \frac{(a + d) \pm \sqrt{(a + d)^2 - 4(ad - |b|^2)}}{2} \\ &= \frac{1 \pm \sqrt{1 - 4\left(\frac{1-r_3^2}{4} - \frac{r_1^2+r_2^2}{4}\right)}}{2} \\ &= \frac{1 \pm \sqrt{1 - (1 - r_1^2 - r_2^2 - r_3^2)}}{2} \\ &= \frac{1 \pm \sqrt{|\vec{r}|^2}}{2} \\ &= \frac{1 \pm |\vec{r}|}{2}\end{aligned}$$

Since ρ is positive, $\frac{1-|\vec{r}|}{2} \geq 0 \rightarrow |\vec{r}| \leq 1$.

Therefore an arbitrary density matrix for a mixed state qubit is written as $\rho = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma})$.

(2)

$\rho = I/2 \rightarrow \vec{r} = 0$. Thus $\rho = I/2$ corresponds to the origin of Bloch sphere.

(3)

$$\begin{aligned}
 \rho^2 &= \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma}) \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma}) \\
 &= \frac{1}{4} \left[I + 2\vec{r} \cdot \vec{\sigma} + \sum_{j,k} r_j r_k \left(\delta_{jk} I + i \sum_{l=1}^3 \epsilon_{jkl} \sigma_l \right) \right] \\
 &= \frac{1}{4} (I + 2\vec{r} \cdot \vec{\sigma} + |\vec{r}|^2 I) \\
 \text{Tr}(\rho^2) &= \frac{1}{4} (2 + 2|\vec{r}|^2)
 \end{aligned}$$

If ρ is pure, then $\text{Tr}(\rho^2) = 1$.

$$\begin{aligned}
 1 = \text{Tr}(\rho^2) &= \frac{1}{4} (2 + 2|\vec{r}|^2) \\
 \therefore |\vec{r}| &= 1.
 \end{aligned}$$

Conversely, if $|\vec{r}| = 1$, then $\text{Tr}(\rho^2) = \frac{1}{4} (2 + 2|\vec{r}|^2) = 1$. Therefore ρ is pure.

2.73

Theorem 2.6

$$\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i| = \sum_i |\tilde{\psi}_i\rangle\langle\tilde{\psi}_i| = \sum_j |\tilde{\varphi}_j\rangle\langle\tilde{\varphi}_j| = \sum_j q_j |\varphi_j\rangle\langle\varphi_j| \Leftrightarrow |\tilde{\psi}_i\rangle = \sum_j u_{ij} |\tilde{\varphi}_j\rangle$$

where u is unitary.

The transformation in theorem 2.6, $|\tilde{\psi}_i\rangle = \sum_j u_{ij} |\tilde{\varphi}_j\rangle$, corresponds to

$$\left[|\tilde{\psi}_1\rangle \cdots |\tilde{\psi}_k\rangle \right] = \left[|\tilde{\varphi}_1\rangle \cdots |\tilde{\varphi}_k\rangle \right] U^T$$

where $k = \text{rank}(\rho)$.

$$\sum_i |\tilde{\psi}_i\rangle\langle\tilde{\psi}_i| = \left[|\tilde{\psi}_1\rangle \cdots |\tilde{\psi}_k\rangle \right] \begin{bmatrix} \langle\tilde{\psi}_1| \\ \vdots \\ \langle\tilde{\psi}_k| \end{bmatrix} \quad (2.10)$$

$$= \left[|\tilde{\varphi}_1\rangle \cdots |\tilde{\varphi}_k\rangle \right] U^T U^* \begin{bmatrix} \langle\tilde{\varphi}_1| \\ \vdots \\ \langle\tilde{\varphi}_k| \end{bmatrix} \quad (2.11)$$

$$= \left[|\tilde{\varphi}_1\rangle \cdots |\tilde{\varphi}_k\rangle \right] \begin{bmatrix} \langle\tilde{\varphi}_1| \\ \vdots \\ \langle\tilde{\varphi}_k| \end{bmatrix} \quad (2.12)$$

$$= \sum_j |\tilde{\varphi}_j\rangle\langle\tilde{\varphi}_j|. \quad (2.13)$$

From spectral theorem, density matrix ρ is decomposed as $\rho = \sum_{k=1}^d \lambda_k |k\rangle\langle k|$ where $d = \dim \mathcal{H}$. Without loss of generality, we can assume $p_k > 0$ for $k = 1 \cdots l$ where $l = \text{rank}(\rho)$ and $p_k = 0$ for $k = l+1, \dots, d$. Thus $\rho = \sum_{k=1}^l p_k |k\rangle\langle k| = \sum_{k=1}^l |\tilde{k}\rangle\langle\tilde{k}|$, where $|\tilde{k}\rangle = \sqrt{\lambda_k} |k\rangle$.

Suppose $|\psi_i\rangle$ is a state in support ρ . Then

$$|\psi_i\rangle = \sum_{k=1}^l c_{ik} |k\rangle, \quad \sum_k |c_{ik}|^2 = 1.$$

Define $p_i = \frac{1}{\sum_k \frac{|c_{ik}|^2}{\lambda_k}}$ and $u_{ik} = \frac{\sqrt{p_i} c_{ik}}{\sqrt{\lambda_k}}$.

Now

$$\sum_k |u_{ik}|^2 = \sum_k \frac{p_i |c_{ik}|^2}{\lambda_k} = p_i \sum_k \frac{|c_{ik}|^2}{\lambda_k} = 1.$$

Next prepare a unitary operator ¹ such that i th row of U is $[u_{i1} \cdots u_{ik} \cdots u_{il}]$. Then we can

¹By Gram-Schmidt procedure construct an orthonormal basis $\{\mathbf{u}_j\}$ (row vector) with $\mathbf{u}_i = [u_{i1} \cdots u_{ik} \cdots u_{il}]$.

Then define unitary $U = \begin{bmatrix} \mathbf{u}_1 \\ \vdots \\ \mathbf{u}_i \\ \vdots \\ \mathbf{u}_l \end{bmatrix}$.

define another ensemble such that

$$\left[|\tilde{\psi}_1\rangle \cdots |\tilde{\psi}_i\rangle \cdots |\tilde{\psi}_l\rangle \right] = \left[|\tilde{k}_1\rangle \cdots |\tilde{k}_l\rangle \right] U^T$$

where $|\tilde{\psi}_i\rangle = \sqrt{p_i} |\psi_i\rangle$. From theorem 2.6,

$$\rho = \sum_k |\tilde{k}\rangle \langle \tilde{k}| = \sum_k |\tilde{\psi}_k\rangle \langle \tilde{\psi}_k|.$$

Therefore we can obtain a minimal ensemble for ρ that contains $|\psi_i\rangle$.

Moreover since $\rho^{-1} = \sum_k \frac{1}{\lambda_k} |k\rangle \langle k|$,

$$\langle \psi_i | \rho^{-1} | \psi_i \rangle = \sum_k \frac{1}{\lambda_k} \langle \psi_i | k \rangle \langle k | \psi_i \rangle = \sum_k \frac{|c_{ik}|^2}{\lambda_k} = \frac{1}{p_i}.$$

Hence, $\frac{1}{\langle \psi_i | \rho^{-1} | \psi_i \rangle} = p_i$.

2.74

$$\begin{aligned} \rho_{AB} &= |a\rangle \langle a|_A \otimes |b\rangle \langle b|_B \\ \rho_A &= \text{Tr}_B \rho_{AB} = |a\rangle \langle a| \text{Tr}(|b\rangle \langle b|) = |a\rangle \langle a| \\ \text{Tr}(\rho_A^2) &= 1 \end{aligned}$$

Thus ρ_A is pure.

2.75

Define $|\Phi_{\pm}\rangle = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle)$ and $|\Psi_{\pm}\rangle = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle)$.

$$\begin{aligned} |\Phi_{\pm}\rangle \langle \Phi_{\pm}|_{AB} &= \frac{1}{2}(|00\rangle \langle 00| \pm |00\rangle \langle 11| \pm |11\rangle \langle 00| + |11\rangle \langle 11|) \\ \text{Tr}_B(|\Phi_{\pm}\rangle \langle \Phi_{\pm}|_{AB}) &= \frac{1}{2}(|0\rangle \langle 0| + |1\rangle \langle 1|) = \frac{I}{2} \\ |\Psi_{\pm}\rangle \langle \Psi_{\pm}| &= \frac{1}{2}(|01\rangle \langle 01| \pm |01\rangle \langle 10| \pm |10\rangle \langle 01| + |10\rangle \langle 10|) \\ \text{Tr}_B(|\Psi_{\pm}\rangle \langle \Psi_{\pm}|) &= \frac{1}{2}(|0\rangle \langle 0| + |1\rangle \langle 1|) = \frac{I}{2} \end{aligned}$$

2.76

Unsolved. I think the polar decomposition can only apply to square matrix A , not arbitrary linear operators. Suppose A is $m \times n$ matrix. Then size of $A^\dagger A$ is $n \times n$. Thus the size of U should be $m \times n$. Maybe U is isometry, but I think it is not unitary.

I misunderstand linear operator.

Quoted from "Advanced Linear Algebra" by Steven Roman, ISBN 0387247661.

A linear transformation $\tau : V \rightarrow V$ is called a **linear operator** on V .²

²According to Roman, some authors use the term linear operator for any linear transformation from V to W .

Thus coordinate matrices of linear operator are square matrices. And Nielsen and Chaung say at Theorem 2.3, "Let A be a linear operator on a vector space V ." Therefore A is a linear transformation such that $A : V \rightarrow V$.

2.77

$$\begin{aligned} |\psi\rangle &= |0\rangle |\Phi_+\rangle \\ &= |0\rangle \left[\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \right] \\ &= (\alpha |\phi_0\rangle + \beta |\phi_1\rangle) \left[\frac{1}{\sqrt{2}}(|\phi_0\phi_0\rangle + |\phi_1\phi_1\rangle) \right] \end{aligned}$$

where $|\phi_i\rangle$ are arbitrary orthonormal states and $\alpha, \beta \in \mathbb{C}$. We cannot vanish cross term. Therefore $|\psi\rangle$ cannot be written as $|\psi\rangle = \sum_i \lambda_i |i\rangle_A |i\rangle_B$.

2.78

Proof. Former part.

If $|\psi\rangle$ is product, then there exist a state $|\phi_A\rangle$ for system A , and a state $|\phi_B\rangle$ for system B such that $|\psi\rangle = |\phi_A\rangle |\phi_B\rangle$.

Obviously, this Schmidt number is 1.

Conversely, if Schmidt number is 1, the state is written as $|\psi\rangle = |\phi_A\rangle |\phi_B\rangle$. Hence this is a product state. \square

Proof. Later part.

(\Rightarrow) Proved by exercise 2.74.

(\Leftarrow) Let a pure state be $|\psi\rangle = \sum_i \lambda_i |i_A\rangle |i_B\rangle$. Then $\rho_A = \text{Tr}_B(|\psi\rangle\langle\psi|) = \sum_i \lambda_i^2 |i\rangle\langle i|$. If ρ_A is a pure state, then $\lambda_j = 1$ and otherwise 0 for some j . It follows that $|\psi_j\rangle = |j_A\rangle |j_B\rangle$. Thus $|\psi\rangle$ is a product state. \square

2.79

Procedure of Schmidt decomposition.

Goal: $|\psi\rangle = \sum_i \sqrt{\lambda_i} |i_A\rangle |i_B\rangle$

- Diagonalize reduced density matrix $\rho_A = \sum_i \lambda_i |i_A\rangle\langle i_A|$.
- Derive $|i_B\rangle$, $|i_B\rangle = \frac{(I \otimes \langle i_A|) |\psi\rangle}{\sqrt{\lambda_i}}$
- Construct $|\psi\rangle$.

(i)

$$\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \text{ This is already decomposed.}$$

(ii)

$$\frac{|00\rangle + |01\rangle + |10\rangle + |11\rangle}{2} = \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right) \otimes \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right) = |\psi\rangle |\psi\rangle \text{ where } |\psi\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}$$

(iii)

$$|\psi\rangle_{AB} = \frac{1}{\sqrt{3}}(|00\rangle + |01\rangle + |10\rangle)$$

$$\rho_{AB} = |\psi\rangle\langle\psi|_{AB}$$

$$\rho_A = \text{Tr}_B(\rho_{AB}) = \frac{1}{3} (2|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| + |1\rangle\langle 1|)$$

$$\det(\rho_A - \lambda I) = \left(\frac{2}{3} - \lambda \right) \left(\frac{1}{3} - \lambda \right) - \frac{1}{9} = 0$$

$$\lambda^2 - \lambda + \frac{1}{9} = 0$$

$$\lambda = \frac{1 \pm \sqrt{5}/3}{2} = \frac{3 \pm \sqrt{5}}{6}$$

Eigenvector with eigenvalue $\lambda_0 \equiv \frac{3 + \sqrt{5}}{6}$ is $|\lambda_0\rangle \equiv \frac{1}{\sqrt{\frac{5+\sqrt{5}}{2}}} \begin{bmatrix} \frac{1+\sqrt{5}}{2} \\ 1 \end{bmatrix}$.

Eigenvector with eigenvalue $\lambda_1 \equiv \frac{3 - \sqrt{5}}{6}$ is $|\lambda_1\rangle \equiv \frac{1}{\sqrt{\frac{5-\sqrt{5}}{2}}} \begin{bmatrix} \frac{1-\sqrt{5}}{2} \\ 1 \end{bmatrix}$.

$$\rho_A = \lambda_0 |\lambda_0\rangle\langle\lambda_0| + \lambda_1 |\lambda_1\rangle\langle\lambda_1|.$$

$$|a_0\rangle \equiv \frac{(I \otimes \langle\lambda_0|) |\psi\rangle}{\sqrt{\lambda_0}}$$

$$|a_1\rangle \equiv \frac{(I \otimes \langle\lambda_1|) |\psi\rangle}{\sqrt{\lambda_1}}$$

Then

$$|\psi\rangle = \sum_{i=0}^1 \sqrt{\lambda_i} |a_i\rangle |\lambda_i\rangle.$$

(It's too tiresome to calculate $|a_i\rangle$)**2.80**Let $|\psi\rangle = \sum_i \lambda_i |\psi_i\rangle_A |\psi_i\rangle_B$ and $|\varphi\rangle = \sum_i \lambda_i |\varphi_i\rangle_A |\varphi_i\rangle_B$.Define $U = \sum_i |\psi_j\rangle\langle\varphi_j|_A$ and $V = \sum_j |\psi_j\rangle\langle\varphi_j|_B$.

Then

$$\begin{aligned} (U \otimes V) |\varphi\rangle &= \sum_i \lambda_i U |\varphi_i\rangle_A V |\varphi_i\rangle_B \\ &= \sum_i \lambda_i |\psi_i\rangle_A |\psi_i\rangle_B \\ &= |\psi\rangle. \end{aligned}$$

2.81

Let the Schmidt decomposition of $|AR_1\rangle$ be $|AR_1\rangle = \sum_i \sqrt{p_i} |\psi_i^A\rangle |\psi_i^R\rangle$ and let $|AR_2\rangle = \sum_i \sqrt{q_i} |\phi_i^A\rangle |\phi_i^R\rangle$.

Suppose ρ^A has orthonormal decomposition $\rho^A = \sum_i p_i |i\rangle\langle i|$.

Since $|AR_1\rangle$ and $|AR_2\rangle$ are purifications of the ρ^A , we have

$$\begin{aligned} \text{Tr}_R(|AR_1\rangle\langle AR_1|) &= \text{Tr}_R(|AR_2\rangle\langle AR_2|) = \rho^A \\ \therefore \sum_i p_i |\psi_i^A\rangle\langle\psi_i^A| &= \sum_i q_i |\phi_i^A\rangle\langle\phi_i^A| = \sum_i \lambda_i |i\rangle\langle i|. \end{aligned}$$

The $|i\rangle$, $|\psi_i^A\rangle$, and $|\phi_i^A\rangle$ are orthonormal bases and they are eigenvectors of ρ^A . Hence without loss of generality, we can consider

$$\lambda_i = p_i = q_i \text{ and } |i\rangle = |\psi_i^A\rangle = |\phi_i^A\rangle.$$

Then

$$\begin{aligned} |AR_1\rangle &= \sum_i \lambda_i |i\rangle |\psi_i^R\rangle \\ |AR_2\rangle &= \sum_i \lambda_i |i\rangle |\phi_i^R\rangle \end{aligned}$$

Since $|AR_1\rangle$ and $|AR_2\rangle$ have same Schmidt numbers, there are two unitary operators U and V such that $|AR_1\rangle = (U \otimes V) |AR_2\rangle$ from exercise 2.80.

Suppose $U = I$ and $V = \sum_i |\psi_i^R\rangle\langle\phi_i^R|$. Then

$$\begin{aligned} \left(I \otimes \sum_j |\psi_j^R\rangle\langle\phi_j^R| \right) |AR_2\rangle &= \sum_i \lambda_i |i\rangle \left(\sum_j |\psi_j^R\rangle\langle\phi_j^R| \phi_i^R \right) \\ &= \sum_i \lambda_i |i\rangle |\psi_i^R\rangle \\ &= |AR_1\rangle. \end{aligned}$$

Therefore there exists a unitary transformation U_R acting on system R such that $|AR_1\rangle = (I \otimes U_R) |AR_2\rangle$.

2.82

(1)

Let $|\psi\rangle = \sum_i \sqrt{p_i} |\psi_i\rangle |i\rangle$.

$$\begin{aligned} \text{Tr}_R(|\psi\rangle\langle\psi|) &= \sum_{i,j} \sqrt{p_i} \sqrt{p_j} |\psi_i\rangle\langle\psi_j| \text{Tr}_R(|i\rangle\langle j|) \\ &= \sum_{i,j} \sqrt{p_i} \sqrt{p_j} |\psi_i\rangle\langle\psi_j| \delta_{ij} \\ &= \sum_i p_i |\psi_i\rangle\langle\psi_i| = \rho. \end{aligned}$$

Thus $|\psi\rangle$ is a purification of ρ .

(2)

Define the projector P by $P = I \otimes |i\rangle\langle i|$. The probability we get the result i is

$$\text{Tr}[P|\psi\rangle\langle\psi|] = \langle\psi|P|\psi\rangle = \langle\psi|(I \otimes |i\rangle\langle i|)|\psi\rangle = p_i \langle\psi_i|\psi_i\rangle = p_i.$$

The post-measurement state is

$$\frac{P|\psi\rangle}{\sqrt{p_i}} = \frac{(I \otimes |i\rangle\langle i|)|\psi\rangle}{\sqrt{p_i}} = \frac{\sqrt{p_i}|\psi_i\rangle|i\rangle}{\sqrt{p_i}} = |\psi_i\rangle|i\rangle.$$

If we only focus on the state on system A ,

$$\text{Tr}_R(|\psi_i\rangle|i\rangle) = |\psi_i\rangle.$$

(3)

($\{|\psi_i\rangle\}$ is not necessary an orthonormal basis.)

Suppose $|AR\rangle$ is a purification of ρ and its Schmidt decomposition is $|AR\rangle = \sum_i \sqrt{\lambda_i} |\phi_i^A\rangle |\phi_i^R\rangle$.

From assumption

$$\text{Tr}_R(|AR\rangle\langle AR|) = \sum_i \lambda_i |\phi_i^A\rangle\langle\phi_i^A| = \sum_i p_i |\psi_i\rangle\langle\psi_i|.$$

By theorem 2.6, there exists a unitary matrix u_{ij} such that $\sqrt{\lambda_i} |\phi_i^A\rangle = \sum_j u_{ij} \sqrt{p_j} |\psi_j\rangle$. Then

$$\begin{aligned} |AR\rangle &= \sum_i \left(\sum_j u_{ij} \sqrt{p_j} |\psi_j\rangle \right) |\phi_i^R\rangle \\ &= \sum_j \sqrt{p_j} |\psi_j\rangle \otimes \left(\sum_i u_{ij} |\phi_i^R\rangle \right) \\ &= \sum_j \sqrt{p_j} |\psi_j\rangle |j\rangle \\ &= \sum_i \sqrt{p_i} |\psi_i\rangle |i\rangle \end{aligned}$$

where $|i\rangle = \sum_k u_{ki} |\phi_k^R\rangle$.

About $|i\rangle$,

$$\begin{aligned} \langle k|l\rangle &= \sum_{m,n} u_{mk}^* u_{nl} \langle\phi_m^R|\phi_n^R\rangle \\ &= \sum_{m,n} u_{mk}^* u_{nl} \delta_{mn} \\ &= \sum_m u_{mk}^* u_{ml} \\ &= \delta_{kl}, \quad (\because u_{ij} \text{ is unitary.}) \end{aligned}$$

which implies $|j\rangle$ is an orthonormal basis for system R .

Therefore if we measure system R w.r.t $|j\rangle$, we obtain j with probability p_j and post-measurement state for A is $|\psi_j\rangle$ from (2). Thus for any purification $|AR\rangle$, there exists an orthonormal basis $|i\rangle$ which satisfies the assertion.

Problem 2.1

From Exercise 2.35, $\vec{n} \cdot \vec{\sigma}$ is decomposed as

$$\vec{n} \cdot \vec{\sigma} = |\lambda_1\rangle\langle\lambda_1| - |\lambda_{-1}\rangle\langle\lambda_{-1}|$$

where $|\lambda_{\pm 1}\rangle$ are eigenvector of $\vec{n} \cdot \vec{\sigma}$ with eigenvalues ± 1 .

Thus

$$\begin{aligned} f(\theta \vec{n} \cdot \vec{\sigma}) &= f(\theta) |\lambda_1\rangle\langle\lambda_1| + f(-\theta) |\lambda_{-1}\rangle\langle\lambda_{-1}| \\ &= \left(\frac{f(\theta) + f(-\theta)}{2} + \frac{f(\theta) - f(-\theta)}{2} \right) |\lambda_1\rangle\langle\lambda_1| + \left(\frac{f(\theta) + f(-\theta)}{2} - \frac{f(\theta) - f(-\theta)}{2} \right) |\lambda_{-1}\rangle\langle\lambda_{-1}| \\ &= \frac{f(\theta) + f(-\theta)}{2} (|\lambda_1\rangle\langle\lambda_1| + |\lambda_{-1}\rangle\langle\lambda_{-1}|) + \frac{f(\theta) - f(-\theta)}{2} (|\lambda_1\rangle\langle\lambda_1| - |\lambda_{-1}\rangle\langle\lambda_{-1}|) \\ &= \frac{f(\theta) + f(-\theta)}{2} I + \frac{f(\theta) - f(-\theta)}{2} \vec{n} \cdot \vec{\sigma} \end{aligned}$$

Problem 2.2

Unsolved

Problem 2.3

Unsolved

Chapter 5

The quantum Fourier transform and its application

5.1

We can write the Fourier quantum transformation as a matrix U . By definition, for an arbitrary ket $|x\rangle = \sum_{j=0}^{N-1} x_j |j\rangle$, the quantum fourier transform maps $|x\rangle$ to $|y\rangle$ as such, noting $w = e^{2i\pi/N}$:

$$F(|x\rangle) = |y\rangle = \frac{1}{\sqrt{N}} \sum_{j=0}^{N-1} \sum_{n=0}^{N-1} x_n w^{kn} |j\rangle$$

the Fourier quantum transformation can then be written as a matrix equation, giving the Fourier transform's matrix, $F|x\rangle = |y\rangle$, with :

$$F = \frac{1}{\sqrt{N}} \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & w & \dots & w^{N-1} \\ 1 & w^2 & \dots & w^{2(N-1)} \\ \vdots & \vdots & \vdots & \vdots \\ 1 & w^{N-1} & \dots & w^{(N-1)(N-1)} \end{bmatrix}.$$

Let's now show that this matrix is indeed unitary. If we note $A = FF^\dagger$, then

$$A_{j,k} = \sum_{n=1}^N F_{j,n} F_{n,k}^\dagger = \frac{1}{N} (1 + w^{j-n} + w^{2(j-n)} + \dots + w^{(N-1)(j-n)})$$

If $j = n$ then $A_{j,k} = \frac{N}{N} = 1$.

If $j \neq n$ then $A_{j,k} = 0$ according to the following orthogonal property over the roots of unity

$$\sum_{k=0}^{N-1} (z^{jk})^\dagger z^{j'k} = 0 \quad \text{if } j \neq j'$$

We then see that $FF^\dagger = I$. It is straightforward to show that $FF^\dagger = F^\dagger F$, we can then conclude that the quantum Fourier transform is unitary.

5.2

the N qubit state $|00\dots 0\rangle$ is a computational basis state, we can use directly the provided formula 5.2 :

$$|j\rangle \rightarrow \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} e^{2i\pi jk/N} |k\rangle$$

applying it to $|00\dots 0\rangle$ we get :

$$|00\dots 0\rangle \rightarrow \frac{1}{2^{N/2}} \sum_{k=0}^{2^N-1} |k\rangle$$

with $|k\rangle$ the 2^N computation basis states for N qubits.

Chapter 8

Quantum noise and quantum operations

8.1

Density operator of initial state is written by $|\psi\rangle\langle\psi|$ and final state is written by $U |\psi\rangle\langle\psi| U^\dagger$. Thus time development of $\rho = |\psi\rangle\langle\psi|$ can be written by $\mathcal{E}(\rho) = U\rho U^\dagger$.

8.2

From eqn (2.147) (on page 100),

$$\rho_m = \frac{M_m \rho M_m^\dagger}{\text{Tr}(M_m^\dagger M_m \rho)} = \frac{M_m \rho M_m^\dagger}{\text{Tr}(M_m \rho M_m^\dagger)} = \frac{\mathcal{E}_m(\rho)}{\text{Tr} \mathcal{E}_m(\rho)}.$$

And from eqn (2.143) (on page 99), $p(m) = \text{Tr}(M_m^\dagger M_m \rho) = \text{Tr}(M_m \rho M_m^\dagger) = \text{Tr} \mathcal{E}_m(\rho)$.

8.3

8.4

8.5

8.6

8.7

8.8

8.9

8.10

8.11

8.12

8.13

8.14

8.15

8.16

8.17

8.18

8.19

8.20

8.21

8.22

8.23

8.24

8.25

8.26

8.27

8.28

8.29

8.30

8.31

8.32

8.33

8.34

8.35

Chapter 9

Distance measures for quantum information

9.1

$$\begin{aligned} D((1, 0), (1/2, 1/2)) &= \frac{1}{2} (|1 - 1/2| + |0 - 1/2|) \\ &= \frac{1}{2} \left(\frac{1}{2} + \frac{1}{2} \right) \\ &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} D((1/2, 1/3, 1/6), (3/4, 1/8, 1/8)) &= \frac{1}{2} (|1/2 - 3/4| + |1/3 - 1/8| + |1/6 - 1/8|) \\ &= \frac{1}{2} (1/4 + 5/24 + 1/24) \\ &= \frac{1}{4} \end{aligned}$$

9.2

$$\begin{aligned} D((p, 1 - p), (q, 1 - q)) &= \frac{1}{2} (|p - q| + |(1 - p) - (1 - q)|) \\ &= \frac{1}{2} (|p - q| + |-p + q|) \\ &= |p - q| \end{aligned}$$

9.3

$$F((1, 0), (1/2, 1/2)) = \sqrt{1 \cdot 1/2} + \sqrt{0 \cdot 1/2} = \frac{1}{\sqrt{2}}$$

$$\begin{aligned} F((1/2, 1/3, 1/6), (3/4, 1/8, 1/8)) &= \sqrt{1/2 \cdot 3/4} + \sqrt{1/3 \cdot 1/8} + \sqrt{1/6 \cdot 1/8} \\ &= \frac{4\sqrt{6} + \sqrt{3}}{12} \end{aligned}$$

9.4

Define $r_x = p_x - q_x$. Let U be the whole index set.

$$\begin{aligned} \max_S |p(S) - q(S)| &= \max_S \left| \sum_{x \in S} p_x - \sum_{x \in S} q_x \right| \\ &= \max_S \left| \sum_{x \in S} (p_x - q_x) \right| \\ &= \max_S \left| \sum_{x \in S} r_x \right| \end{aligned}$$

Since $\sum_{x \in S} r_x$ is written as

$$\sum_{x \in S} r_x = \sum_{\substack{x \in S \\ r_x \geq 0}} r_x + \sum_{\substack{x \in S \\ r_x < 0}} r_x, \quad (9.1)$$

$|\sum_{x \in S} r_x|$ is maximized when $S = \{x \in U | r_x \geq 0\}$ or $S = \{x \in U | r_x < 0\}$.

Define $S_+ = \{x \in U | r_x \geq 0\}$ and $S_- = \{x \in U | r_x < 0\}$.

Now the sum of all r_x is 0,

$$\begin{aligned} \sum_{x \in U} r_x &= \sum_{x \in S_+} r_x + \sum_{x \in S_-} r_x = 0 \\ \therefore \sum_{x \in S_+} r_x &= - \sum_{x \in S_-} r_x. \end{aligned}$$

Thus

$$\max_S \left| \sum_{x \in S} r_x \right| = \sum_{x \in S_+} r_x = - \sum_{x \in S_-} r_x. \quad (9.2)$$

On the other hand,

$$\begin{aligned} D(p_x, q_x) &= \frac{1}{2} \sum_{x \in U} |p_x - q_x| \\ &= \frac{1}{2} \sum_{x \in U} |r_x| \\ &= \frac{1}{2} \sum_{x \in S_+} |r_x| + \frac{1}{2} \sum_{x \in S_-} |r_x| \\ &= \frac{1}{2} \sum_{x \in S_+} r_x - \frac{1}{2} \sum_{x \in S_-} r_x \\ &= \frac{1}{2} \sum_{x \in S_+} r_x + \frac{1}{2} \sum_{x \in S_+} r_x \quad (\because \text{eqn(9.2)}) \\ &= \sum_{x \in S_+} r_x \\ &= \max_S \left| \sum_{x \in S} r_x \right|. \end{aligned}$$

Therefore $D(p_x, q_x) = \max_S |\sum_{x \in S} p_x - \sum_{x \in S} q_x| = \max_S |p(S) - q(S)|$.

9.5

From eqn (9.1) and (9.2), maximizing $|\sum_{x \in S} r_x|$ is equivalent to maximizing $\sum_{x \in S} r_x$.

Hence

$$D(p_x, q_x) = \max_S (p(S) - q(S)) = \max_S \left(\sum_{x \in S} p_x - \sum_{x \in S} q_x \right).$$

9.6

Define $\rho = \frac{3}{4} |0\rangle\langle 0| + \frac{1}{4} |1\rangle\langle 1|$, $\sigma = \frac{2}{3} |1\rangle\langle 1| + \frac{1}{3} |1\rangle\langle 1|$.

$$\begin{aligned} D(\rho, \sigma) &= \frac{1}{2} \text{Tr} |\rho - \sigma| \\ &= D((3/4, 1/4), (2/3, 1/3)) \\ &= \frac{1}{2} \left(\left| \frac{3}{4} - \frac{2}{3} \right| + \left| \frac{1}{4} - \frac{1}{3} \right| \right) \\ &= \frac{1}{2} \left(\frac{1}{12} + \frac{1}{12} \right) \\ &= \frac{1}{12} \end{aligned}$$

Define $\rho = \frac{3}{4} |0\rangle\langle 0| + \frac{1}{4} |1\rangle\langle 1|$, $\sigma = \frac{2}{3} |+\rangle\langle +| + \frac{1}{3} |-\rangle\langle -|$.

$$\begin{aligned} |+\rangle\langle +| &= \frac{1}{2} (|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| + |1\rangle\langle 1|) \\ |-\rangle\langle -| &= \frac{1}{2} (|0\rangle\langle 0| - |0\rangle\langle 1| - |1\rangle\langle 0| + |1\rangle\langle 1|) \end{aligned}$$

$$\begin{aligned} \rho - \sigma &= \left(\frac{3}{4} - \frac{1}{2} \right) |0\rangle\langle 0| - \frac{1}{6} (|0\rangle\langle 1| + |1\rangle\langle 0|) + \left(\frac{1}{4} - \frac{1}{2} \right) |1\rangle\langle 1| \\ &= \frac{1}{4} |0\rangle\langle 0| - \frac{1}{6} (|0\rangle\langle 1| + |1\rangle\langle 0|) - \frac{1}{4} |1\rangle\langle 1| \end{aligned}$$

$$\begin{aligned} (\rho - \sigma)^\dagger (\rho - \sigma) &= \frac{1}{4^2} |0\rangle\langle 0| - \frac{1}{4 \cdot 6} |0\rangle\langle 1| + \frac{1}{6^2} |0\rangle\langle 0| + \frac{1}{6 \cdot 4} |0\rangle\langle 1| - \frac{1}{4 \cdot 6} |1\rangle\langle 0| + \frac{1}{6^2} |1\rangle\langle 1| + \frac{1}{4 \cdot 6} |1\rangle\langle 0| + \frac{1}{4^2} |1\rangle\langle 1| \\ &= \left(\frac{1}{4^2} + \frac{1}{6^2} \right) (|0\rangle\langle 0| + |1\rangle\langle 1|) \end{aligned}$$

$$\begin{aligned} D(\rho, \sigma) &= \frac{1}{2} \text{Tr} |\rho - \sigma| \\ &= \sqrt{\frac{1}{4^2} + \frac{1}{6^2}} \end{aligned}$$

9.7

Since $\rho - \sigma$ is Hermitian, we can apply spectral decomposition. Then $\rho - \sigma$ is written as

$$\rho - \sigma = \sum_{i=1}^k \lambda_i |i\rangle\langle i| + \sum_{i=k+1}^n \lambda_i |i\rangle\langle i|$$

where λ_i are positive eigenvalues for $i = 1, \dots, k$ and negative eigenvalues for $i = k+1, \dots, n$.

Define $Q = \sum_{i=1}^k \lambda_i |i\rangle\langle i|$ and $S = -\sum_{i=k+1}^n \lambda_i |i\rangle\langle i|$. Then P and S are positive operator. Therefore $\rho - \sigma = P - S$.

Proof of $|\rho - \sigma| = Q + S$.

$$\begin{aligned} |\rho - \sigma| &= |Q - S| \\ &= \sqrt{(Q - S)^\dagger (Q - S)} \\ &= \sqrt{(Q - S)^2} \\ &= \sqrt{Q^2 - QS - SQ + S^2} \\ &= \sqrt{Q^2 + S^2} \\ &= \sqrt{\sum_i \lambda_i^2 |i\rangle\langle i|} \\ &= \sum_i |\lambda_i| |i\rangle\langle i| \\ &= Q + S \end{aligned}$$

9.8

Suppose $\sigma = \sigma_i$. Then $\sigma = \sum_i p_i \sigma_i$.

$$D\left(\sum_i p_i \rho_i, \sigma\right) = D\left(\sum_i p_i \rho_i, \sum_i p_i \sigma_i\right) \quad (9.3)$$

$$\leq \sum_i p_i D(\rho_i, \sigma_i) \quad (\because \text{eqn(9.50)}) \quad (9.4)$$

$$= \sum_i p_i D(\rho_i, \sigma). \quad (\because \text{assumption}). \quad (9.5)$$

9.9

9.10

9.11

9.12

Suppose $\rho = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma})$ and $\sigma = \frac{1}{2}(I + \vec{s} \cdot \vec{\sigma})$ where \vec{r} and \vec{s} are real vectors s.t. $|\vec{r}|, |\vec{s}| \leq 1$.

$$\mathcal{E}(\rho) = p \frac{I}{2} + (1-p)\rho, \quad \mathcal{E}(\sigma) = p \frac{I}{2} + (1-p)\sigma.$$

$$\begin{aligned}
D(\mathcal{E}(\rho), \mathcal{E}(\sigma)) &= \frac{1}{2} \text{Tr} |\mathcal{E}(\rho) - \mathcal{E}(\sigma)| \\
&= \frac{1}{2} \text{Tr} |(1-p)(\rho - \sigma)| \\
&= \frac{1}{2} (1-p) \text{Tr} |\rho - \sigma| \\
&= (1-p) D(\rho, \sigma) \\
&= (1-p) \frac{|\vec{r} - \vec{s}|}{2}
\end{aligned}$$

Is this strictly contractive?

9.13

Bit flip channel $E_0 = \sqrt{p}I$, $E_1 = \sqrt{1-p}\sigma_x$.

$$\begin{aligned}
\mathcal{E}(\rho) &= E_0 \rho E_0^\dagger + E_1 \rho E_1^\dagger \\
&= p\rho + (1-p)\sigma_x \rho \sigma_x.
\end{aligned}$$

Since $\sigma_x \sigma_x \sigma_x = \sigma_x$, $\sigma_x \sigma_y \sigma_x = -\sigma_y$ and $\sigma_x \sigma_z \sigma_x = -\sigma_z$, then $\sigma_x(\vec{r} \cdot \vec{\sigma}) = r_1 \sigma_x - r_2 \sigma_y - r_3 \sigma_z$. Thus

$$\begin{aligned}
D(\mathcal{E}(\rho), \mathcal{E}(\sigma)) &= \frac{1}{2} \text{Tr} |\mathcal{E}(\rho) - \mathcal{E}(\sigma)| \\
&= \frac{1}{2} \text{Tr} |p(\rho - \sigma) + (1-p)(\sigma_x \rho \sigma_x - \sigma_x \sigma \sigma_x)| \\
&\leq \frac{1}{2} p \text{Tr} |\rho - \sigma| + \frac{1}{2} (1-p) \text{Tr} |\sigma_x(\rho - \sigma)\sigma_x| \\
&= p D(\rho, \sigma) + (1-p) D(\sigma_x \rho \sigma_x, \sigma_x \sigma \sigma_x) \\
&= D(\rho, \sigma) \quad (\because \text{eqn(9.21)}).
\end{aligned}$$

Suppose $\rho_0 = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma})$ is a fixed point. Then

$$\begin{aligned}
\rho_0 &= \mathcal{E}(\rho_0) = p\rho_0 + (1-p)\sigma_x \rho_0 \sigma_x \\
\therefore (1-p)\rho_0 - (1-p)\sigma_x \rho_0 \sigma_x &= 0 \\
\therefore (1-p)(\rho - \sigma_x \rho_0 \sigma_x) &= 0 \\
\therefore \rho_0 &= \sigma_x \rho_0 \sigma_x \\
\therefore \frac{1}{2}(I + r_1 \sigma_x + r_2 \sigma_y + r_3 \sigma_z) &= \frac{1}{2}(I + r_1 \sigma_x - r_2 \sigma_y - r_3 \sigma_z)
\end{aligned}$$

Since $\{I, \sigma_x, \sigma_y, \sigma_z\}$ are linearly independent, $r_2 = -r_2$ and $r_3 = -r_3$. Thus $r_2 = r_3 = 0$.

Therefore the set of fixed points for the bit flip channel is $\{\rho \mid \rho = \frac{1}{2}(I + r\sigma_x), |r| \leq 1, r \in \mathbb{R}\}$

9.14

$$\begin{aligned}
F(U\rho U^\dagger, U\sigma U^\dagger) &= \text{Tr} \sqrt{(U\rho U^\dagger)^{1/2} \sigma (U\rho U^\dagger)} \\
&= \text{Tr} \sqrt{U\rho^{1/2} \sigma \rho^{1/2} U^\dagger} \\
&= \text{Tr}(U \sqrt{\rho^{1/2} \sigma \rho^{1/2}} U^\dagger) \\
&= \text{Tr}(\sqrt{\rho^{1/2} \sigma \rho^{1/2}} U^\dagger U) \\
&= \text{Tr} \sqrt{\rho^{1/2} \sigma \rho^{1/2}} \\
&= F(\rho, \sigma)
\end{aligned}$$

I think the fact $\sqrt{UAU^\dagger} = U\sqrt{A}U^\dagger$ is not restricted for positive operator. Suppose A is a normal matrix. From spectral theorem, it is decomposed as

$$A = \sum_i a_i |i\rangle\langle i|.$$

Let f be a function. Then

$$\begin{aligned}
f(UAU^\dagger) &= f\left(\sum_i a_i U|i\rangle\langle i|U^\dagger\right) \\
&= \sum_i f(a_i) U|i\rangle\langle i|U^\dagger \\
&= U\left(\sum_i f(a_i) |i\rangle\langle i|\right)U^\dagger \\
&= Uf(A)U^\dagger
\end{aligned}$$

9.15

$|\psi\rangle = (U_R \otimes \sqrt{\rho}U_Q)|m\rangle$ is any fixed purification of ρ , and $|\phi\rangle = (V_R \otimes \sqrt{\sigma}V_Q)|m\rangle$ is purification of σ . Suppose $\sqrt{\rho}\sqrt{\sigma} = |\sqrt{\rho}\sqrt{\sigma}|V$ is the polar decomposition of $\sqrt{\rho}\sqrt{\sigma}$. Then

$$\begin{aligned}
|\langle\psi|\phi\rangle| &= \left| \langle m| \left(U_R^\dagger V_R \otimes U_Q^\dagger \sqrt{\rho}\sqrt{\sigma} V_Q \right) |m\rangle \right| \\
&= \left| \text{Tr} \left((U_R^\dagger V_R)^T U_Q^\dagger \sqrt{\rho}\sqrt{\sigma} V_Q \right) \right| \\
&= \left| \text{Tr} \left(V_R^T U_R^* U_Q^\dagger \sqrt{\rho}\sqrt{\sigma} V_Q \right) \right| \\
&= \left| \text{Tr} \left(V_Q V_R^T U_R^* U_Q^\dagger \sqrt{\rho}\sqrt{\sigma} \right) \right| \\
&= \left| \text{Tr} \left(V_Q V_R^T U_R^* U_Q^\dagger |\sqrt{\rho}\sqrt{\sigma}| V \right) \right| \\
&= \left| \text{Tr} \left(V V_Q V_R^T U_R^* U_Q^\dagger |\sqrt{\rho}\sqrt{\sigma}| \right) \right| \\
&\leq \text{Tr} |\sqrt{\rho}\sqrt{\sigma}| \\
&= F(\rho, \sigma)
\end{aligned}$$

Choosing $V_Q = V^\dagger$, $V_R^T = (U_Q^* U_R^\dagger)^\dagger$ we see that equality is attained.

9.16

I think eq (9.73) has a typo. $\text{Tr}(A^\dagger B) = \langle m|A \otimes B|m\rangle$ should be $\text{Tr}(A^T B) = \langle m|A \otimes B|m\rangle$. See errata list.

In order to show that this exercise, I will prove following two properties,

$$\text{Tr}(A) = \langle m|(I \otimes A)|m\rangle, \quad (I \otimes A)|m\rangle = (A^T \otimes I)|m\rangle$$

where A is a linear operator and $|m\rangle$ is unnormalized maximally entangled state, $|m\rangle = \sum_i |ii\rangle$.

$$\begin{aligned} \langle m|I \otimes A|m\rangle &= \sum_{ij} \langle ii|(I \otimes A)|jj\rangle \\ &= \sum_{ij} \langle i|I|j\rangle \langle i|A|j\rangle \\ &= \sum_{ij} \delta_{ij} \langle i|A|j\rangle \\ &= \sum_i \langle i|A|i\rangle \\ &= \text{Tr}(A) \end{aligned}$$

Suppose $A = \sum_{ij} a_{ij} |i\rangle\langle j|$.

$$\begin{aligned} (I \otimes A)|m\rangle &= \left(I \otimes \sum_{ij} a_{ij} |i\rangle\langle j| \right) \sum_k |kk\rangle \\ &= \sum_{ijk} a_{ij} |k\rangle \otimes |i\rangle \langle j|k\rangle \\ &= \sum_{ijk} a_{ij} |k\rangle \otimes |i\rangle \delta_{jk} \\ &= \sum_{ij} a_{ij} |j\rangle \otimes |i\rangle \\ &= \sum_{ij} a_{ji} |i\rangle \otimes |j\rangle \end{aligned}$$

$$\begin{aligned} (A^T \otimes I)|m\rangle &= \left(\sum_{ij} a_{ji} |i\rangle\langle j| \otimes I \right) \sum_k |kk\rangle \\ &= \sum_{ij} a_{ji} |i\rangle \langle j|k\rangle \otimes |k\rangle \\ &= \sum_{ij} a_{ji} |i\rangle \delta_{jk} \otimes |k\rangle \\ &= \sum_{ij} a_{ji} |ij\rangle \\ &= (I \otimes A)|m\rangle \end{aligned}$$

Thus

$$\begin{aligned} \text{Tr}(A^T B) &= \text{Tr}(BA^T) = \langle m|I \otimes BA^T|m\rangle \\ &= \langle m|(I \otimes B)(I \otimes A^T)|m\rangle \\ &= \langle m|(I \otimes B)(A \otimes I)|m\rangle \\ &= \langle m|A \otimes B|m\rangle. \end{aligned}$$

9.17

If $\rho = \sigma$, then $F(\rho, \sigma) = 1$. Thus $A(\rho, \sigma) = \arccos F(\rho, \sigma) = \arccos 1 = 0$.

If $A(\rho, \sigma) = 0$, then $\arccos F(\rho, \sigma) = 0 \Rightarrow \cos(\arccos F(\rho, \sigma)) = \cos(0) \Rightarrow F(\rho, \sigma) = 1$ (\because text p.411, the fifth line from bottom).

9.18

For $0 \leq x \leq y \leq 1$, $\arccos(x) \geq \arccos(y)$. From $F(\mathcal{E}(\rho), \mathcal{E}(\sigma)) \geq F(\rho, \sigma)$ and $0 \leq F(\mathcal{E}(\rho), \mathcal{E}(\sigma)), F(\rho, \sigma) \leq 1$,

$$\begin{aligned} \arccos F(\mathcal{E}(\rho), \mathcal{E}(\sigma)) &\geq \arccos F(\rho, \sigma) \\ \therefore A(\mathcal{E}(\rho), \mathcal{E}(\sigma)) &\geq A(\rho, \sigma) \end{aligned}$$

9.19

From eq (9.92)

$$\begin{aligned} F\left(\sum_i p_i \rho_i, \sum_i p_i \sigma_i\right) &\geq \sum_i \sqrt{p_i p_i} F(\rho_i, \sigma_i) \\ &= \sum_i p_i F(\rho_i, \sigma_i). \end{aligned}$$

9.20

Suppose $\sigma_i = \sigma$. Then

$$\begin{aligned} F\left(\sum_i p_i \rho_i, \sigma\right) &= F\left(\sum_i p_i \rho_i, \sum_i p_i \sigma\right) \\ &= F\left(\sum_i p_i \rho_i, \sum_i p_i \sigma_i\right) \\ &\geq \sum_i p_i F(\rho_i, \sigma_i) \quad (\because \text{Exercise 9.19}) \\ &= \sum_i p_i F(\rho_i, \sigma) \end{aligned}$$

9.21

$$1 - F(|\psi\rangle, \sigma)^2 = 1 - \langle\psi|\sigma|\psi\rangle \quad (\because \text{eq(9.60)})$$

$$\begin{aligned} D(|\psi\rangle, \sigma) &= \max_P \text{Tr}(P(\rho - \sigma)) \quad (\text{where } P \text{ is projector.}) \\ &\geq \text{Tr}(|\psi\rangle\langle\psi|(\rho - \sigma)) \\ &= \langle\psi|(|\psi\rangle\langle\psi| - \sigma)|\psi\rangle \\ &= 1 - \langle\psi|\sigma|\psi\rangle \\ &= 1 - F(|\psi\rangle, \sigma)^2. \end{aligned}$$

9.22

(ref: QCQI Exercise Solutions (Chapter 9) - めもめも

<http://enakai00.hatenablog.com/entry/2018/04/12/134722>)

For all ρ , following inequality is satisfied,

$$\begin{aligned} d(VU\rho U^\dagger V^\dagger, \mathcal{F} \circ \mathcal{E}(\rho)) &\leq d(VU\rho U^\dagger V^\dagger, \mathcal{F}(U\rho U^\dagger)) + d(\mathcal{F}(U\rho U^\dagger), \mathcal{F} \circ \mathcal{E}(\rho)) \\ &\leq d(VU\rho U^\dagger V^\dagger) + d(U\rho U^\dagger, \mathcal{E}(\rho)) \\ &\leq E(V, \mathcal{F}) + E(U, \mathcal{E}). \end{aligned}$$

First inequality is triangular inequality, second is contractivity of the metric¹ and third is from definition of E .

Above inequality is hold for all ρ . Thus $E(VU, \mathcal{F} \circ \mathcal{E}) \leq E(V, \mathcal{F}) + E(U, \mathcal{E})$.

9.23

(\Leftarrow) If $\mathcal{E}(\rho_j) = \rho_j$ for all j such that $p_j > 0$, then

$$\bar{F} = \sum_j p_j F(\rho_j, \mathcal{E}(\rho_j))^2 = \sum_j p_j F(\rho_j, \rho_j)^2 = \sum_j p_j 1^2 = \sum_j p_j = 1.$$

(\Rightarrow) Suppose $\mathcal{E}(\rho_j) \neq \rho_j$. Then $F(\rho_j, \mathcal{E}(\rho_j)) < 1$ (\because text p.411, the fifth line from bottom). Thus

$$\bar{F} = \sum_j p_j F(\rho_j, \mathcal{E}(\rho_j))^2 < \sum_j p_j = 1.$$

Therefore if $\bar{F} = 1$, then $\mathcal{E}(\rho_j) = \rho_j$.

Problem 1

Problem 2

Problem 3

Theorem 5.3 of "Theory of Quantum Error Correction for General Noise", Emanuel Knill, Raymond Laflamme, and Lorenza Viola, Phys. Rev. Lett. 84, 2525 – Published 13 March 2000. arXiv:quant-ph/9604034 <https://arxiv.org/abs/quant-ph/9604034>

¹Trace distance and angle are satisfied with contractive (eq (9.35), eq (9.91)), but I don't assure that arbitrary metric satisfied with contractive.

Chapter 11

Entropy and information

11.1

Fair coin:

$$H(1/2, 1/2) = \left(-\frac{1}{2} \log \frac{1}{2}\right) \times 2 = 1 \quad (11.1)$$

Fair die:

$$H(p) = \left(-\frac{1}{6} \log \frac{1}{6}\right) \times 6 = \log 6. \quad (11.2)$$

The entropy decreases if the coin or die is unfair.

11.2

From assumption $I(pq) = I(p) + I(q)$.

$$\frac{\partial I(pq)}{\partial p} = \frac{\partial I(p)}{\partial p} + 0 = \frac{\partial I(p)}{\partial p} \quad (11.3)$$

$$\frac{\partial I(pq)}{\partial q} = 0 + \frac{\partial I(q)}{\partial q} = \frac{\partial I(q)}{\partial q} \quad (11.4)$$

$$\frac{\partial I(pq)}{\partial p} = \frac{\partial I(pq)}{\partial(pq)} \frac{\partial(pq)}{\partial p} = q \frac{\partial I(pq)}{\partial(pq)} \Rightarrow \frac{\partial I(pq)}{\partial(pq)} = \frac{1}{q} \frac{\partial I(p)}{\partial p} \quad (11.5)$$

$$\frac{\partial I(pq)}{\partial q} = \frac{\partial I(pq)}{\partial(pq)} \frac{\partial(pq)}{\partial q} = p \frac{\partial I(pq)}{\partial(pq)} \Rightarrow \frac{\partial I(pq)}{\partial(pq)} = \frac{1}{p} \frac{\partial I(q)}{\partial q} \quad (11.6)$$

Thus

$$\frac{1}{q} \frac{\partial I(p)}{\partial p} = \frac{1}{p} \frac{\partial I(q)}{\partial q} \quad (11.7)$$

$$\therefore p \frac{dI(p)}{dp} = q \frac{dI(q)}{dq} \quad \text{for all } p, q \in [0, 1]. \quad (11.8)$$

$$(11.9)$$

Then $p(dI(p)/dp)$ is constant.

If $p(dI(p)/dp) = k$, $k \in \mathbb{R}$. Then $I(p) = k \ln p = k' \log p$ where $k' = k/\log e$.

11.3

$$H_{\text{bin}}(p) = -p \log p - (1-p) \log(1-p).$$

$$\frac{dH_{\text{bin}}(p)}{dp} = \frac{1}{\ln 2} (-\log p - 1 + \log(1-p) + 1) \quad (11.10)$$

$$= \frac{1}{\ln 2} \ln \frac{1-p}{p} = 0 \quad (11.11)$$

$$\Rightarrow \frac{1-p}{p} = 1 \quad (11.12)$$

$$\Rightarrow p = 1/2. \quad (11.13)$$

11.4

11.5

$$H(p(x, y) || p(x)p(y)) = \sum_{x, y} p(x, y) \log \frac{p(x)p(y)}{p(x, y)} \quad (11.14)$$

$$= -H(p(x, y)) - \sum_{x, y} p(x, y) \log [p(x)p(y)] \quad (11.15)$$

$$= -H(p(x, y)) - \sum_{x, y} p(x, y) [\log p(x) + \log p(y)] \quad (11.16)$$

$$= -H(p(x, y)) - \sum_{x, y} p(x, y) \log p(x) - \sum_{x, y} p(x, y) \log p(y) \quad (11.17)$$

$$= -H(p(x, y)) - \sum_x p(x) \log p(x) - \sum_y p(y) \log p(y) \quad (11.18)$$

$$= -H(p(x, y)) + H(p(x)) + H(p(y)) \quad (11.19)$$

$$= -H(X, Y) + H(X) + H(Y). \quad (11.20)$$

From the non-negativity of the relative entropy,

$$H(X) + H(Y) - H(X, Y) \geq 0 \quad (11.21)$$

$$\therefore H(X) + H(Y) \geq H(X, Y). \quad (11.22)$$

11.6

11.7

11.8

11.9

11.10

11.11

11.12

11.13

11.14

11.15

11.16

11.17

11.18

11.19

11.20

11.21

11.22

11.23

11.24

11.25

11.26

Problem 11.1

Problem 11.2

Problem 11.3

Problem 11.4

Problem 11.5