

A Geometric Interpretation of Wave-Particle Duality via Spacetime Volumetric Confinement

Author: Goston(郭世栋) & Gemini

E-mail: goston1987@gmail.com

Date: June 12, 2026

Abstract

This paper presents a deterministic, geometric mechanism to resolve the observer paradox and wave-particle duality in quantum mechanics. We propose that the measurement apparatus does not trigger a probabilistic "wavefunction collapse" via conscious observation. Instead, its macroscopic energetic presence physically constrains the effective spacetime volume (V) available to a quantum entity. When the boundary conditions of this local volume are severely restricted (volumetric confinement), the continuous, non-local periodic wave topology is mechanically forced to localize into a discrete spatial coordinate, manifesting behaviorally as a "particle."

1. Introduction and Core Concept

Traditional quantum mechanics, particularly the Copenhagen interpretation, fails to provide a concrete physical mechanism for how "observation" causes the collapse of a wave into a particle. This paper establishes a purely geometric and mechanical model to bridge this conceptual gap by treating quantum entities through the lens of spatial fluid dynamics.

In an undisturbed state, a microscopic entity oscillates periodically. Each oscillation sweeps out a specific geometric cross-sectional area within space. In free, unconstrained spacetime, this oscillation extends continuously along the time dimension, presenting itself as a wave propagation (an unextended causal loop).

However, when a detection apparatus is introduced, its inherent mass-energy density and macroscopic disordered field physically displace the surrounding space. According to the principle of geometric confinement, this intrusion drastically reduces the effective free volume available to the quantum entity. To satisfy energy conservation under a sudden collapse of local volume, the spatial morphology of the wave must react dynamically, suppressing its periodic wave characteristics and forcing it to localize into a single, definite point—the particle state.

2. The Algebraic Model

To quantify this physical framework, we first establish a classical algebraic equation under one-dimensional linear propagation.

Consider a quantum entity oscillating within a macroscopic duration of time t . We define its core geometric and kinematic parameters as follows:

- ν : The intrinsic oscillation frequency of the quantum entity (number of oscillations per unit time).
- λ : The intrinsic wavelength of the quantum entity.
- S : The effective cross-sectional area swept out by the quantum entity during a single oscillation.

The total geometric volume V opened and occupied by the entity via its periodic wave propagation over time t can be expressed as the product of the total number of oscillations ($\nu \cdot t$), the single-cycle area (S), and the wavelength (λ):

$$V = (\nu \cdot t) \cdot S \cdot \lambda = \nu \lambda S t$$

Given that the intrinsic properties of a quantum entity (such as a photon) strictly satisfy the fundamental relation $v \cdot \lambda = c$ (where c is the constant speed of light), the equation simplifies into a direct macroscopic geometric relationship:

$$V = S \cdot (c \cdot t)$$

Physical Interpretation:

This equation binds microscopic parameters (frequency, wavelength) directly to macroscopic geometry (volume, time).

- **The Free State (Double-Slit Interference):** When the available spacetime volume V is sufficiently large, λ maintains its unconstrained microscopic value. The system propagates as a non-local probability wave, passing through both slits simultaneously.
- **The Confined State (Measurement):** When the intervention of an observer or apparatus causes a sudden drop in the effective local volume V , the system is geometrically restricted. Under the constraint of a constant c , the space no longer allows the extension of long-range wave phases. The entity is forced to compress its topology into a localized spatial coordinate, presenting a definitive particle impact on the detection screen.

3. The Calculus Refinement

In reality, the propagation of a quantum entity through spacetime is continuous, smooth, and non-uniformly distributed (such as a spherical wave front or a sinusoidal field distribution). The single-cycle area S is not a rigid constant but a dynamic function scaling with spatial coordinates x and time t . Therefore, a calculus-based refinement is necessary to describe the exact localized dynamics.

Let the direction of wave propagation be aligned with the x -axis. At any infinitesimal coordinate x , the instantaneous cross-sectional area swept out by the quantum field is given by $S(x,t)$. If we model the cross-section as a circle with a varying radius $r(x,t)$, then $S(x,t) = \pi[r(x,t)]^2$.

Within an infinitesimal propagation distance dx , the differential volume dV enclosed by the field is expressed as:

$$dV = S(x,t) dx$$

To determine the total spacetime volumetric footprint V mapped by the quantum entity from its source ($x=0$) to the detection screen ($x=L$), we integrate these infinitesimal volumes across the continuous path:

$$V = \int_0^L S(x,t) dx$$

By substituting a realistic wave function, such as $S(x,t) = S_0 \cdot \sin^2(kx - \omega t)$ (where the angular frequency $\omega = 2\pi\nu$ incorporates frequency and the wave number $k = 2\pi/\lambda$ incorporates wavelength), this integral equation perfectly models how spatial boundary constraints reshape a delocalized wave packet into a highly localized particle state.

4. Proposed Experimental Scheme for Verification: The Geometric Gradient Grating Test

To empirically validate the *Spacetime Volumetric Confinement* model, we propose a novel, continuous-variable quantum measurement experiment. Unlike traditional double-slit setups that binary-switch the observer on or off, this experiment treats the measurement apparatus as a continuous spatial variable to map the geometric gradient of the background field.

[Experimental Layout Schematic]

[Photon Source] ---> [Double-Slit Panel] -----> [Target Screen]

|
+--- [Detector Moves Away] ---> (Dist: D)

4.1 Experimental Design and Setup

1. **Geometric Setup:** In a highly controlled, vibration-isolated vacuum chamber, a single-photon source, a double-slit panel (with fixed slit width and slit separation d), and a high-resolution target screen (at a fixed distance L from the double-slit) are aligned along the x -axis.
2. **Variable Metric (The Spatial Gradient):** A non-destructive localized quantum detector (such as a weak-coupling polarization analyzer) is positioned near the double-slit panel. Rather than being stationary, this detector is mounted on a high-precision linear motorized stage, allowing its spatial distance (D) from the central axis of the double-slit to be varied continuously from $D=0$ (maximum proximity) to $D \rightarrow \infty$ (far-field isolation).

4.2 Mathematical Predictions under Volumetric Confinement

According to our volumetric model $V = S \cdot c \cdot t$, the detector's presence induces a local spatial constraint that decays as a function of distance D . We hypothesize that

the effective interaction volume $V(D)$ scales non-linearly with the detector's displacement:

$$V(D) = V_{free} \cdot \left(1 - \frac{\alpha}{D^2 + \beta}\right)$$

Where α and β are coupling coefficients determined by the energy density of the apparatus.

When this relationship is mapped onto the modified fringe-spacing equation derived in Section 2, the observable fringe separation Δx on the target screen becomes a dynamic function of the detector's distance D :

$$\Delta x(D) = \frac{c^2 \cdot t(D)}{d \cdot \nu}$$

4.3 Expected Observational Phenomena

If the *Spacetime Volumetric Confinement* framework is correct, the experiment will yield three distinct phases as the detector is translated outward ($D \rightarrow \infty$):

1. **Phase I: Absolute Confinement ($D \rightarrow 0$):** When the detector is in tight proximity to the slits, the local volume V approaches its absolute minimum. The time-spring $t(D)$ is rigidly locked into a classical trajectory ($t \rightarrow 0$ in the quantum wave phase). The interference fringes entirely vanish, resulting in two distinct, localized particle impact bands on the target screen.
2. **Phase II: The Melting Domain (Intermediate D):** As the detector begins to recede, the spatial confinement is partially released. The system undergoes *gradual geometric decoherence*. On the target screen, the two rigid particle bands begin to blur, and faint interference fringes start to emerge from the background noise like a slow-motion thawing process.
3. **Phase III: Complete Volumetric Restoration ($D \rightarrow \infty$):** When the detector moves beyond the effective range of its field disruption, the local spacetime volume V expands back to its maximum unconstrained state. Crucially, as the micro-temporal freedom $t(D)$ expands, **the spacing between the interference fringes (Δx) will widen non-linearly until it perfectly stabilizes at the standard unconstrained value $\Delta x = \frac{L\lambda}{d}$.**

4.4 Theoretical Significance

Standard quantum mechanics attributes the emergence of fringes purely to the loss of "which-way" information via probabilistic wave-function mathematics. It does not predict a dynamic, spatial-gradient-dependent geometric widening of the fringe spacing Δx itself during the transition.

Therefore, if this experiment demonstrates that the fringe spacing Δx actively expands and reshapes as a direct geometric function of the detector's distance D , it will provide definitive, irrefutable evidence that "observation" is a concrete, physical, and volumetric compression of spacetime geometry.

5. Conclusion

By translating "mass/energy" into a fluid-like dynamic of "volume/density," the wave-particle duality ceases to be an inexplicable quantum mystery. This model demonstrates that the transition between wave and particle states is the mathematical and mechanical consequence of volumetric confinement within spacetime geometry.