

Building a better metaweb: predicting spatiotemporally explicit plant-pollinator networks

Michael D. Catchen^{1,2}, Paul CaraDonna^{3,4}, Jane E. Ogilvie³, Francis Banville³, Dominique Caron^{1,2},
Philippe Desjardins-Proulx^{5,2}, Norma R. Forero-Muñoz^{5,2}, Dominique Gravel^{6,2}, Laura Pollock^{1,2},
Tanya Strydom^{5,2}, Timothée Poisot^{5,2}, Julian Resasco⁷, Andrew Gonzalez^{1,2}

¹ McGill University; ² Québec Centre for Biodiversity Sciences; ³ Rocky Mountain Biological Laboratory;
⁴ Chicago Botanic Garden; ⁵ Université de Montréal; ⁶ Université de Sherbrooke; ⁷ University of Colorado
Boulder

Correspondance to:

Michael D. Catchen — michael.catchen@mail.mcgill.ca

Purpose: This template provides a series of scripts to render a markdown document into an interactive website and a series of PDFs.

Motivation: It makes collaborating on text with GitHub easier, and means that we never need to think about the output.

Internals: GitHub actions and a series of python scritpts. The markdown is handled with pandoc.

Keywords:
species interactions
ecological forecasting
pollinators
bumblebees
network ecology

0.1. Abstract Using a data set of [DESCRIBE EACH DATASET IN A NICE WAY], we predict a spatiotemporally explicit metaweb of interactions between bumblebees (*Bombus*) and wildflowers (within *find clade*). We integrate this data with crowdsourced occurrence data and climate data to [best paint the picture of the Colorado bumblebee-plant metaweb]. Using temporal climate data, we forecast how the spatiotemporal overlap of interacting species will change under proposed climate scenarios. We use this to estimate what interactions between bees and plants need the most attention to prevent the spatiotemporal decoupling of an interactions from threatening ecosystem functioning or the persistence of a species.

1

Introduction

Species interactions are important. It is ultimately interactions between individuals of different species that drive the structure, dynamics, and persistence of ecosystems, and the abundance and diversity of the species within them. Plant-pollinator interactions specifically drive the function and persistence of “architecture of biodiversity” (Bascompte & Jordano 2007). However, we are far from a robust understanding of plant-pollinator networks. This is because sampling interactions is costly. Interactions vary in space and time (Poisot *et al.* 2015)—particularlyly relevent in this system (CaraDonna *et al.* 2014). This is why there is interest in using models to predict interactions from sparse data (Strydom2021?). In this paper, we combine several datasets, each spanning several years, to produce spatially and temporally explicit predictions of the bumblebee (genus *Bombus*) and wildflower pollination network across the state of Colorado.

We do this in two parts: (1) metaweb prediction and (2) conditioning our metaweb prediction on co-occurrence probability. First, we build a model to predict the metaweb—the network of *all* interactions, aggregated across all times and spatial locations—of *Bombus* and wildflower species across Colorado. (Why do this? The metaweb is more predictable than local interactions.) We do this using network

Metaweb Prediction




Figure 1 todo

embedding (**cite?**). Network embedding takes each node in the network (either a bumblebee or a wild-flower) and represents it in a latent n dimensional space. Combination of running models on Temporal niche (T), Phylogenetic niche (P), Environmental niche (E), and relative abundance in community (RA).

Second, we then use this metaweb to predict the structure of networks at specific locations and times of year (Gravel *et al.* 2019). Finally we suggest a map of sampling priority, which suggests the locations to sample that will best improve our understanding of the Colorado *Bombus* pollination metaweb.

Why is this good for science, what does this contribute to our understanding of plant-pollinator ints, networks, *Bombus*, predictive models, etc., and how can these results be useful.

2 _____

Data

We use three separate field datasets to estimate the Colorado *Bombus* metaweb.

3 _____

Methods

4 _____

Metaweb Model

4.1. Phylogeny Construction

4.2. Feature Embedding




Figure 2 todo

4.2.1 Relative Abundance

4.2.2 Phylogenetic features

4.2.3 Environmental niche features

4.2.4 Temporal niche features

4.3. Metaweb Model Fitting and Validation

5

Spatiotemporally Explicit Networks

Now that we have a metaweb.....

Figure 3: Maps over time figure and Prob(Connectance) vs. Month figure

Sampling Prioritization

Figure 4: Uncertainty and sampling priority map

Discussion

- Bascompte, J. & Jordano, P. (2007). Plant-Animal Mutualistic Networks: The Architecture of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 38, 567–593.
- CaraDonna, P.J., Iler, A.M. & Inouye, D.W. (2014). Shifts in flowering phenology reshape a subalpine plant community. *Proceedings of the National Academy of Sciences*, 111, 4916–4921.
- Gravel, D., Baiser, B., Dunne, J.A., Kopelke, J.-P., Martinez, N.D., Nyman, T., *et al.* (2019). Bringing Elton and Grinnell together: A quantitative framework to represent the biogeography of ecological interaction networks. *Ecography*, 42, 401–415.
- Poisot, T., Stouffer, D.B. & Gravel, D. (2015). Beyond species: Why ecological interaction networks vary through space and time. *Oikos*, 124, 243–251.