

The missing link: discerning true from false negatives when sampling species interaction networks

[Michael D. Catchen](#)^{1,2} [Timothée Poisot](#)^{3,2} [Laura J. Pollock](#)^{1,2} [Andrew Gonzalez](#)^{1,4}

¹ McGill University ² Québec Centre for Biodiversity Sciences ³ Université de Montréal ⁴ Québec Centre for Biodiversity Science

Correspondance to:

Michael D. Catchen — michael.catchen@mcgill.ca

Abstract: Ecosystems are composed of networks of interacting species. These interactions allow communities of species to persist through time through both neutral and adaptive processes. Despite their importance, a robust understanding of (and ability to predict and forecast) interactions among species remains elusive. This knowledge-gap is largely driven by a shortfall of data—although species occurrence data has rapidly increased in the last decade, species interaction data has not kept pace, largely due to the intrinsic difficulty and effort required to sample interactions. This means there are many interactions between species that occur in nature, but we do not think this interaction occurs because we have no record of it. These so-called “false-negatives” bias data and hinder inference about the structure and dynamics of interaction networks. Here, we demonstrate the realized rate of false-negatives in data can be quite high, even in thoroughly sampled systems, due to the intrinsic variation in abundances across species in a community. We illustrate how a null model of occurrence detection can be used to estimate the false-negative rate in a given dataset. We also show how to directly incorporate uncertainty due to observation error into model-based predictions of interaction probabilities between species. One hypothesis is that interactions between “rare” species are themselves rare because these species are less likely to encounter one-another than species of higher relative abundance, and that this can (in part) explain the common pattern of nestedness in bipartite interaction networks. However, we demonstrate that across several datasets of spatial or temporally replicated networks, there are positive associations between species co-occurrence and interactions, which suggests these interactions among “rare” species actually exist but simply are not observed. Finally, we assess how false negatives influence various models of network prediction, and recommend directly accounting for observation error in predictive models. We conclude by discussing how the understanding of false-negatives can inform how we design monitoring schemes for species interactions.

1 Introduction

2 Species interactions drive many processes in evolution and ecology. A better understanding of species
3 interactions is an imperative to understand the evolution of life on Earth, to mitigate the impacts of
4 anthropogenic change on biodiversity (Makiola *et al.* 2020), and for predicting zoonotic spillover of
5 disease to prevent future pandemics (Becker *et al.* 2021). At the moment we lack sufficient data to meet
6 these challenges (Poisot *et al.* 2021), largely because species interactions are hard to sample (Jordano
7 2016). Over the past few decades biodiversity data has become increasingly available through remotely
8 collected data and adoption of open data practices (Kenall *et al.* 2014; Stephenson 2020). Still, interaction
9 data remains relatively scarce because sampling typically requires human observation. This induces a
10 constraint on the amount, spatial scale, and temporal frequency of resulting data that it is feasible to
11 collect by humans. Many crowdsourced methods for biodiversity data aggregation (e.g. GBIF, eBird) still
12 rely on automated identification of species, which does not easily generalize to interaction sampling.
13 There is interest in using remote methods for interaction sampling, which primarily detect co-occurrence
14 and derive properties like species avoidance from this data (Niedballa *et al.* 2019). However, co-occurrence
15 itself is not necessarily indicative of an interaction (Blanchet *et al.* 2020). This is an example of semantic
16 confusion around the word “interaction”—for example one might consider competition a type of species
17 interaction, even though it is marked by a lack of co-occurrence between species, unlike other types of
18 interactions, like predation or parasitism, which require both species to be together at the same place and
19 time. Here we consider interaction in the latter sense, where two species have fitness consequences on
20 one-another if (and only if) they are in the sample place at the same time. In addition, here we only
21 consider direct (not higher-order) interactions.

22 We cannot feasibly observe all (or even most) of the interactions that occur in an ecosystem. This means
23 we can be confident two species actually interact if we have a record of it (assuming they are correctly
24 identified), but not at all confident that a pair of species *do not* interact if we have *no record* of those
25 species observed together. In other words, it is difficult to distinguish *true-negatives* (two species never
26 interact) from *false-negatives* (two species interact sometimes, but we do not have a record of this
27 interaction). For a concrete example of a false-negative in a food web, see fig. 1. Because even the most
28 highly sampled systems will still contain false-negatives, there is increasing interest in combining
29 species-level data (e.g. traits, abundance, range, phylogenetic relatedness, etc.) to build models to predict

30 interactions between species we haven't observed together before (Strydom *et al.* 2021). However, the
31 noise of false-negatives could impact the efficacy of our predictive models and have practical
32 consequences for answering questions about interactions (de Aguiar *et al.* 2019). This data constraint is
33 amplified as the interaction data we have is geographically biased toward the usual suspects (Poisot *et al.*
34 2021). We therefore need a statistical approach to assessing these biases in the observation process and
35 their consequences for our understanding of interaction networks.

36 The importance of *sampling effort* and its impact on resulting ecological data has produced a rich body of
37 literature. The recorded number of species in a dataset or sample depends on the total number of
38 observations (Walther *et al.* 1995; Willott 2001), as do estimates of population abundance (Griffiths 1998).
39 This relationship between sampling effort and spatial coverage and species detectability has motivated
40 more quantitatively robust approaches to account for error in sampling data in many contexts: to
41 determine if a given species is extinct (Boakes *et al.* 2015), to determine sampling design (Moore &
42 McCarthy 2016), and to measure species richness across large scales (Carlson *et al.* 2020). In the context of
43 interactions, an initial concern was the compounding effects of limited sampling effort combined with the
44 amalgamation of data (across both study sites, time of year, and taxonomic scales) could lead any
45 empirical set of observations to inadequately reflect the reality of how species interact (Paine 1988) or the
46 structure of the network as a whole (Martinez *et al.* 1999; McLeod *et al.* 2021). Martinez *et al.* (1999)
47 showed that in a plant-endophyte trophic network, network connectance is robust to sampling effort, but
48 this was done in the context of a system for which observation of 62,000 total interactions derived from
49 164,000 plant-stems was feasible. In some systems (e.g. megafauna food-webs) this many observations is
50 either impractical or infeasible due to the absolute abundance of the species in question.

51 The intrinsic properties of ecological communities create several challenges for sampling: first, species are
52 not observed with equal probability—we are much more likely to observe a species of high abundance
53 than one of very low abundance (Poisot *et al.* 2015). Canard *et al.* (2012) presents a null model of food-web
54 structure where species encounter one-another in proportion to each species' relative-abundance. This
55 assumes that there are no associations in species co-occurrence due to an interaction (perhaps because
56 this interaction is "important" for both species; Cazelles *et al.* (2016)), but in this paper we later show
57 increasing strength of associations leads to increasing probability of false-negatives in interaction data,
58 and that these positive associations are common in existing network data. Second, observed co-occurrence
59 is often equated with meaningful interaction strength, but this is not necessarily the case (Blanchet *et al.*

2020)—a true “non-interaction” would require that neither of two species, regardless of whether they co-occur, ever exhibit any meaningful effect on the fitness of the other. So, although co-occurrence is not directly indicative of an interaction, it is a precondition for an interaction.

Here, we illustrate how our confidence that a pair of species never interacts highly depends on sampling effort. We suggest that surveys of species interactions can benefit from simulation modeling of the sampling process. We demonstrate how the realized false-negative-rate of interactions is related to the relative abundance of the species pool, and introduce a method to produce a null estimate of the false-negative-rate given total sampling effort (the total count of all interactions seen among all species-pairs) and a method for including uncertainty into model predictions of interaction probabilities to account for observation error. We then show that positive associations in co-occurrence data can increase the realized number of false-negatives, and demonstrate these positive associations are rampant in network datasets, and conclude by recommending that the simulation of sampling effort and species occurrence can and should be used to help design surveys of species interaction diversity (Moore & McCarthy 2016), and by advocating use of null models like those presented here as a tool for both guiding design of surveys of species interactions and for modeling detection error in predictive models.

Accounting for false-negatives in species interactions

In this section, we demonstrate how difference in relative-abundance can lead to many false-negatives in interaction data. We also introduce a method for producing a null estimate of the false-negative-rate in datasets via simulation, and a method for incorporating uncertainty directly into predictions of species interactions to account for observation error.

How many observations of a non-interaction do we need to be confident it's a true negative?

We start with a naive model of interaction detection: we assume that every interacting pair of species is incorrectly observed as not-interacting with an independent and fixed probability, which we denote p_{fn} and subsequently refer to as the False-Negative-Rate (FNR). If we observe the same species not-interacting N times, then the probability of a true-negative (denoted p_{tn}) is given by $p_{tn} = 1 - (p_{fn})^N$. This relation




Figure 1. This conceptual example considers a sample of the trophic community of bears, wolves, salmon (pink fish), pike (yellow fish), berries, trees, and aspen trees. The true metaweb (all realized interactions across the entire spatial extent) is shown on the left. In the center is what a hypothetical ecologist samples at each site. Notice that although bears and wolves are observed co-occurring with both salmon and pike, there was never a direct observation of bears eating pike, even though they actually do. Therefore, this interaction between bears and pike

86 (the probability-mass-function of geometric distribution, a special case of the negative-binomial
 87 distribution) is shown in fig. 2(A) for varying values of p_{fn} and illustrates a fundamental link between our
 88 ability to reliably say an interaction doesn't exist— p_{tn} —and the number of times N we have observed a
 89 given species. In addition, note that there is no non-zero p_{fn} for which we can ever *prove* that an
 90 interaction does not exist—no matter how many observations of non-interactions N we have, $p_{tn} < 1$.
 91 From fig. 2(A) it is clear that the more often we see two species co-occurring, but *not interacting*, the more
 92 likely the interaction is a true-negative. This has several practical consequences: first it means negatives
 93 taken outside the overlap of the range of each species aren't informative because co-occurrence was not
 94 possible, and therefore neither was an interaction. Second, we can use this relation to compute the
 95 expected number of total observations needed to obtain a “goal” number of observations of a particular
 96 pair of species (fig. 2(B)). As an example, if we hypothesize that A and B do not interact, and we want to see
 97 species A and B both co-occurring and *not interacting* 10 times to be confident this is a true negative, then
 98 we need an expected 1000 observations of all species if the relative abundances of A and B are both 0.1.
 99 Because the true FNR is latent, we can never actually be sure what the actual number of false-negatives in
 100 our data—however, we can use simulation to estimate the FNR for datasets of a given size using neutral
 101 models of observation. If some of the “worst-case” FNRs presented in fig. 2(A) seem unrealistically high,
 102 considering that species are observed in proportion to their relative abundance. In the next section we
 103 demonstrate that the distribution of abundance in ecosystems can lead to very high realized values of FNR
 104 (p_{fn}) simply as an artifact of sampling effort.

105 **False-negatives as a product of relative abundance**

106 We now show that the realized FNR changes drastically with sampling effort due to the intrinsic variation
 107 of the abundance of individuals of each species within a community. We do this by simulating the process
 108 of observation of species interactions, applied both to 243 empirical food webs from the Mangal database
 109 (Banville *et al.* 2021) and random food-webs generated using the niche model, a simple generative model
 110 of food-web structure that accounts for allometric scaling (Williams & Martinez 2000). Our neutral model
 111 of observation assumes each observed species is drawn in proportion to each species' abundance at that
 112 place and time. The abundance distribution of a community can be reasonably-well described by a
 113 log-normal distribution (Volkov *et al.* 2003). In addition to the log-normal distribution, we also tested the





Figure 2
(A) The probability that an observed interaction is a true-negative (y-axis) given how many times it has been sampled as a non-interaction (x-axis). Each color reflects different value of p_{fi} the false-negative-rate (FNR)—this is effectively the cdf of the geometric distribution.
(B) The expected number of total observations needed (colors) to observe 10 co-occurrences between a species with relative abundance $P(A)$ (x-axis) and a second species with relative abundance $P(Y)$. **(C)** false-negative-rate (y-axis) as a function of total sampling

114 case where the abundance distribution is derived from power-law scaling $Z^{(\log(T_i)-1)}$ where T_i is the
 115 trophic level of species i and Z is a scaling coefficient (Savage *et al.* 2004), which yields the same
 116 qualitative behavior. The practical consequence of abundance distributions spanning many orders of
 117 magnitude of abundance is that observing two “rare” species interacting requires two low probability
 118 events: observing two rare species *at the same time*.

119 To simulate the process of observation, for an ecological network M with S species, we sample abundances
 120 for each species from a standard-log-normal distribution. For each true interaction in the adjacency matrix
 121 M (i.e. $M_{ij} = 1$) we estimate the probability of observing both species i and j at a given place and time by
 122 simulating n observations of all individuals of any a species, where the species of the individual observed
 123 at the $\{1, 2, \dots, n\}$ -th observation is drawn from the generated log-normal distribution of abundances. For
 124 each pair of species (i, j) , if both i and j are observed within the n -observations, the interaction is tallied as
 125 a true positive if $M_{ij} = 1$. If only one of i or j are observed—but not both—in these n observations, but
 126 $M_{ij} = 1$, this is counted as a false-negative, and a true-negative otherwise. For each pair of species (i, j) , if
 127 both i and j are observed within the n -observations, the interaction is tallied as a true positive if $M_{ij} = 1$.
 128 If only one of i or j are observed—but not both—in these n observations, but $M_{ij} = 1$, this is counted as a
 129 false-negative, and a true-negative otherwise ($M_{ij} = 0$). This process is illustrated conceptually in fig. 3(A).
 130 In fig. 2(C) we see this model of observation applied to niche model networks across varying levels of
 131 species richness, and in fig. 2(D) the observation model applied to Mangal food webs. For all niche model
 132 simulations in this manuscript, for a given number of species S the number of interactions is drawn from
 133 the flexible-links model fit to Mangal data (MacDonald *et al.* 2020), effectively drawing the number of
 134 interactions L for a random niche model food-web as

$$L \sim \text{BetaBinomial}(S^2 - S + 1, \mu\phi, 1 - \mu\phi)$$

135 where the maximum *a posteriori* (MAP) estimate of (μ, ϕ) applied to Mangal data from (MacDonald *et al.*
 136 2020) is $(\mu = 0.086, \phi = 24.3)$. All simulations were done with 500 independent replicates of unique niche
 137 model networks per unique number of observations n . All analyses presented here are done in Julia v1.8
 138 (Bezanson *et al.* 2015) using both EcologicalNetworks.jl v0.5 and Mangal.jl v0.4 (Banville *et al.* 2021) and
 139 are hosted on [Github](#)). Note that the empirical data, for the reasons described above, very likely already
 140 contains many false-negatives, we’ll revisit this issue in the final section.

141 From fig. 2(C) it is evident that the number of species considered in a study is inseparable from the
142 false-negative-rate in that study, and this effect should be taken into account when designing samples of
143 ecological networks in the future. We see a similar qualitative pattern in fig. 2(D) where the FNR drops off
144 quickly as a function of observation effort, mediated by total richness. The practical consequence of the
145 bottom row of fig. 2 is whether the total number of observations of all species (the x-axis) for the threshold
146 FNR we deem acceptable (the y-axis) is feasible. This raises two points: first, empirical data on
147 interactions are subject to the practical limitations of funding and human-work hours, and therefore
148 existing data tend to fall on the order of hundreds or thousands observations of individuals per site. Clear
149 aggregation of data on sampling effort has proven difficult to find and a meta-analysis of network data and
150 sampling effort seems both pertinent and necessary, in addition to the effects of aggregation of interactions
151 across taxonomic scales (Gauzens *et al.* 2013; Giacomuzzo & Jordán 2021). This inherent limitation on
152 in-situ sampling means we should optimize where we sample across space so that for a given number of
153 samples, we obtain the maximum information possible. Second, what is meant by “acceptable” FNR? This
154 raises the question: does a shifting FNR lead to rapid transitions in our ability inference and predictions
155 about the structure and dynamics of networks, or does it produce a roughly linear decay in model efficacy?
156 We explore this in the next section.

157 We conclude this section by advocating for the use of neutral models similar to above to generate
158 expectations about the number of false-negatives in a data set of a given size. This could prove fruitful
159 both for designing surveys of interactions but also because we may want to incorporate models of
160 imperfect detection error into predictive interactions models, as Joseph (2020) does for species occurrence
161 modeling. Additionally, we emphasize that one must consider the context for sampling—is the goal to
162 detect a particular species (as in fig. 2(C)), or to get a representative sample of interactions across the
163 species pool? These arguments are well-considered when sampling individual species (Willott 2001), but
164 have not yet been adopted for designing samples of communities.

165 **Including observation error in interaction predictions**

166 Here we show how to incorporate uncertainty into model predictions of interaction probability to account
167 for imperfect observation (both false-negatives and false-positives). Models for interaction prediction
168 typically yield a probability of interaction between each pair of species, p_{ij} . When these are considered
169 with uncertainty, it is usually model-uncertainty, e.g. the variance in the interaction probability prediction

across several cross-validation folds, where the data is split into training and test sets several times. The method we introduce adjusts the value of a model's predictions to produce a distribution of interaction probabilities, which are adjusted by a given false-negative-rate p_{fn} and false-positive-rate p_{fp} (outlined in figure fig. 3). We describe first how to sample from this distribution of adjusted interaction probabilities via simulation, and show that this distribution can be well-approximated analytically.

We then consider the output prediction from an arbitrary prediction model, which is the probability p_{ij} that two species i and j interact. To get an estimate of p_{ij} that accounts for observation error, we resample the probability of each interaction p_{ij} by simulating a set of several 'particles,' where each particle is a realization of an interaction occurring (either true or false with probabilities p_{ij} and $1 - p_{ij}$ respectively) and then being correctly observed with probabilities given by p_{fp} and p_{fn} to yield a single boolean outcome for each particle. ("Resampling" within fig. 3 (B)). Over many samples of particles, the resulting frequency of 'true' outcomes is a single resample of the interaction probability p_{ij}^* . Across several samples each of several particles, this forms a distribution of probabilities which are adjusted by the true and false negative rates.

There is also an analytic way to approximate this distribution using the normal approximation to binomial. As a reminder, as the total number of samples n from a binomial distribution with success probability p from approaches infinity, the sum of total successes across all samples approaches a normal distribution with mean np and variance $np(1 - p)$. We can use this to correct the estimate p_{ij} based on the expected false-negative-rate p_{fn} and false-positive rate p_{fp} to obtain the limiting distribution as the number of resamples approaches infinity for the resampled p_{ij}^* for a given number of particles n_p . We do this by first adjusting for the rates of observation error to get the mean resampled probability, $\mathbb{E}[p_{ij}^*]$, as

$$\mathbb{E}[p_{ij}^*] = p_{ij}(1 - p_{fp}) + (1 - p_{ij})p_{fn}$$

which yields the normal approximation

$$\sum_{i=1}^{n_p} p_{ij}^* \sim \mathcal{N}\left(n_p \cdot \mathbb{E}[p_{ij}^*], \sqrt{n_p \mathbb{E}[p_{ij}^*](1 - \mathbb{E}[p_{ij}^*])}\right)$$

which then can be converted back to a distribution of frequency of successes to yield the final approximation




Figure 1: Workflow for estimating interaction probabilities with uncertainty. (A) The process for estimating the false negative-rate (FNR) for a given interaction dataset consisting of N total observed interactions. (B) The method for resampling interaction probability based on estimates of false negative and false positive rates. (C) The method for interaction probability resampling applied to three mammals and three parasites from the Hadfield *et al.* (2014) dataset.

$$p_{ij}^* \sim \mathcal{N}\left(\mathbb{E}[p_{ij}^*], \sqrt{\frac{\mathbb{E}[p_{ij}^*](1 - \mathbb{E}[p_{ij}^*])}{n_p}}\right) \quad (1)$$

We can then further truncate to remain on the interval $(0, 1)$ (as the output is a probability, although in practice often the probability mass outside $(0, 1)$ is extremely low. As an example case study, we use a boosted-regression-tree to predict interactions in a host-parasite network (Hadfield *et al.* 2014) (with features derived in the same manner as Strydom *et al.* (2021) derives features on this data) to produce a set of interaction predictions. We then applied this method to a set of a few resampled interaction probabilities between mammals and parasite species shown in figure fig. 3(C).

Why is this useful? For one, this analytic method avoids the extra computation required by simulating samples from this distribution directly. Further, it enables continuous examination of the number of particles n_p as a uncertainty width. The natural analogue for the number of particles sampled is the number of observations of co-occurrence for a given pair of species—the fewer the particles, the higher the variance of the resulting approximation. The normal approximation is undefined for 0 particles (i.e. 0 observations co-occurrence), although as n_p approaches 0 the approxated normal (once truncated) approaches a uniform distribution on the interval $(0, 1)$, the maximum entropy distribution where we have no information about the possibility of an interaction.

This also has implications for what we mean by ‘uncertainty’ in interaction predictions. A model’s prediction can be ‘uncertain’ in two different ways: (1) the model’s predictions may have high variance, or (2) the model’s predictions may be centered around a probability of interaction of 0.5, where we are the most unsure about whether this interaction exists. Improving the incorporation of different forms of uncertainty in probabilistic interaction predictions seems a necessary next step toward understanding what pairs of species we know the least about, in order to prioritize sampling to provide the most new information possible.

Positive associations in co-occurrence increase the false-negative-rate

The model above doesn’t consider the possibility that there are positive or negative associations which shift the probability of species cooccurrence away from what is expected based on their relative abundances due to their interaction (Cazelles *et al.* 2016). However, here we demonstrate that the probability of having a

219 false-negative can be higher if there is some positive association in the occurrence of species A and B . If
 220 we denote the probability that we observe the co-occurrence of two species A and B as $P(AB)$ and if there
 221 is no association between the marginal probabilities of observing A and observing B , denoted $P(A)$ and
 222 $P(B)$ respectively, then the probability of observing their co-occurrence is the product of the marginal
 223 probabilities for each species, $P(AB) = P(A)P(B)$. In the other case where there is some positive strength
 224 of association between observing both A and B because this interaction is “important” for each species,
 225 then the probability of observation both A and B , $P(AB)$, is greater than $P(A)P(B)$ as $P(A)$ and $P(B)$ are
 226 not independent and instead are positively correlated, i.e. $P(AB) > P(A)P(B)$. In this case, the probability
 227 of observing a single false-negative in our naive model from fig. 2(A) is $p_{fn} = 1 - P(AB)$, which due to the
 228 above inequality implies $p_{fn} > 1 - P(A)P(B)$. This indicates an increasingly greater probability of a false
 229 negative as the strength of association gets stronger, $P(AB) \rightarrow P(AB) \gg P(A)P(B)$. However, this still does
 230 not consider variation in species abundance in space and time (Poisot *et al.* 2015). If positive or negative
 231 associations between species structure variation in the distribution of $P(AB)$ across space/time, then the
 232 spatial/temporal biases induced by data collection would further impact the realized false-negative-rate, as
 233 the probability of false negative would not be constant for each pair of species across sites.

234 To test for these positive associations in data we scoured Mangal for datasets with many spatial or temporal
 235 replicates of the same system, which led to the resulting seven datasets set in figure fig. 4. For each
 236 dataset, we compute the marginal probability $P(A)$ of occurrence of each species A across all networks in
 237 the dataset. For each pair of interacting species A and B , we then compute and compare the probability of
 238 co-occurrence if each species occurs independently, $P(A)P(B)$, to the empirical joint probability of
 239 co-occurrence, $P(AB)$. Following our analysis above, if $P(AB)$ is greater than $P(A)P(B)$, then we expect
 240 our neutral estimates of the FNR above to underestimate the realized FNR. In fig. 4, we see the difference
 241 between $P(AB)$ and $P(A)P(B)$ for the seven suitable datasets with enough spatio-temporal replicates and a
 242 shared taxonomic backbone (meaning all individual networks use common species identifiers) found on
 243 Mangal to perform this analysis. Further details about each dataset are reported in tbl. 1.

244 In each of these datasets, the joint probability of co-occurrence $P(AB)$ is decisively greater than our
 245 expectation if species co-occur in proportion to their relative abundance $P(A)P(B)$. This suggests that
 246 there may not be as many “neutrally forbidden links” (Canard *et al.* 2012) as we might think, and that the
 247 reason we do not have records of interactions between rare species is probably due to observation error.
 248 This has serious ramifications for the widely observed property of nestedness seen in bipartite networks

(Bascompte & Jordano 2007)—perhaps the reason we have lots of observations between generalists is because they are more abundant, and this is particularly relevant as we have strong evidence that generalism drives abundance (Song *et al.* 2022a), not vice-versa.

Table 1: The datasets used in the above analysis (Fig 2). The table reports the type of each dataset, the total number of networks in each dataset (N), the total species richness in each dataset (S), the connectance of each metaweb (all interactions across the entire spatial-temporal extent) (C), the mean species richness across each local network \bar{S} , the mean connectance of each local network \bar{C} , the mean β -diversity among overlapping species across all pairs of network species ($\bar{\beta}_{OS}$), and the mean β -diversity among all species in the metaweb ($\bar{\beta}_{WN}$). Both metrics are computed using KGL β -diversity (Koleff *et al.* 2003)

Network	Type	N	S	C	\bar{S}	\bar{C}	$\bar{\beta}_{OS}$	$\bar{\beta}_{WN}$
Kopelke <i>et al.</i> (2017)	Food Web	100	98	0.037	7.87	0.142	1.383	1.972
Thompson & Townsend (2000)	Food Web	18	566	0.014	80.67	0.049	1.617	1.594
Havens (1992)	Food Web	50	188	0.065	33.58	0.099	1.468	1.881
Ponisio <i>et al.</i> (2017)	Pollinator	100	226	0.079	23.0	0.056	1.436	1.870
Hadfield <i>et al.</i> (2014)	Host-Parasite	51	327	0.085	32.71	0.337	1.477	1.952
Closs & Lake (1994)	Food Web	12	61	0.14	29.09	0.080	1.736	1.864
CaraDonna <i>et al.</i> (2017)	Pollinator	86	122	0.18	21.42	0.312	1.527	1.907

The impact of false-negatives on network properties and prediction

Here, we assess the effect of false-negatives on our ability to make predictions about interactions, as well as their effect on network structure. The prevalence of false-negatives in data is the catalyst for interaction prediction in the first place, and as a result methods have been proposed to counteract this bias (Stock *et al.* 2017; Poisot *et al.* 2022). However, it is feasible that the FNR in a given dataset is so high that it could induce too much noise for an interaction prediction model to detect the signal of possible interaction between species.

To test this we use the dataset from Hadfield *et al.* (2014) that describes host-parasite interaction networks sampled across 51 sites, and the same method as Strydom *et al.* (2021) to extract latent features for each species in this dataset based on applying PCA to the co-occurrence matrix. We then predict a metaweb (equivalent to predicting true or false for an interaction between each species pair, effectively a binary classification problem) from these species-level features using four candidate models for binary




Figure 4: The difference between joint-probability of co-occurrence ($P(AB)$) and expected probability of co-occurrence under independence ($P(A)P(B)$) for interacting species for each dataset. The red dashed line indicates 0 (no association). Each histogram represents a density estimate meaning the area of the entire curve sums to 1. The continuous density estimate (computed using local smoothing) is shown in grey. The p-value of each plot is the result of a one-sided t-test comparing the mean of each distribution to 0.

classification—three often used machine-learning (ML) methods (Boosted Regression Tree (BRT), Random Forest (RF), Decision Tree (DT)), and one naive model from classic statistics (Logistic Regression (LR)). Each of the ML models are bootstrap aggregated (or bagged) with 100 replicates each. We partition the data into 80-20 training-test split, and then seed the training data with false negatives at varying rates, but crucially do nothing to the test data. We fit all of these models using MLJ.jl, a high-level Julia framework for a wide-variety of ML models (Blaom *et al.* 2020). We evaluate the efficacy of these models using two common measures of binary classifier performance: the area under the receiver-operator curve (ROC-AUC) and the area under the precision-recall curve (PR-AUC), for more details see Poisot (2022). Here, PR-AUC is slightly more relevant as it is a better indicator of prediction of false-negatives. The results of these simulations are shown in fig. 5(A&B).

One interesting result seen in fig. 5(A&B) is that the ROC-AUC value does not approach random in the same way the PR-AUC curve does as we increase the added FNR. The reason for this is that ROC-AUC is fundamentally not as useful a metric in assessing predictive capacity as PR-AUC. As we keep adding more false-negatives, the network eventually becomes a zeros matrix, and these models can still learn to predict “no-interaction” for all possible species pairs, which does far better than random guessing (ROC-AUC = 0.5) in terms of the false positive rate (one of the components of ROC-AUC). This highlights a more broad issue of label class imbalance, meaning there are far more non-interactions than interactions in data. A full treatment of the importance of class-balance is outside the scope of this paper, but is explored in-depth in Poisot (2022).

Although these ML models are surprisingly performant at link prediction given their simplicity, there have been several major developments in applying deep-learning methods to many tasks in network inference and prediction—namely graph-representation learning (GRL, Khoshraftar & An (2022)) and graph convolutional networks (Zhang *et al.* 2019). At this time, these advances can not yet be applied to ecological networks because they require far more data than we currently have. We already have lots of features that could be used as inputs into these models (i.e. species level data about occurrence, genomes, abundance, etc.), but our network datasets barely get into the hundreds of local networks sampled across space and time (tbl. 1). Once we start to get into the thousands, these models will become more useful, but this can only be done with systematic monitoring of interactions. This again highlights the need to optimize our sampling effort to maximize the amount of information contained in our data given the expense of sampling interactions.




Figure 5. Performance of different predictive models across species richness and added FNR. (A) The area-under the receiver-operator curve (ROC-AUC) and (B) The area-under the precision-recall curve (PR-AUC; right) for each different predictive model (colors/shapes) across spectrum of the proportion of added false-negatives (x-axis). (C) The mean trophic-level of all species in a network generated with the niche model across different species richnesses (colors). For each value of the FNR, the mean trophic level was computed across 5 replicates. The shaded region for each line is one standard-deviation across those

294 We also consider how the FNR affects network properties. In fig. 5(C) we see the mean trophic level across
295 networks simulated using the niche model (as above), across a spectrum of FNR values. In addition to the
296 clear dependence on richness, we see that mean trophic level, despite varying widely between niche model
297 simulations, tends to be relatively robust to false-negatives and does not deviate widely from the true value
298 until very large FNRs, i.e. $p_{fn} > 0.7$. This is not entirely unsurprising. Removing links randomly from a
299 food-web is effectively the inverse problem of the emergence of a giant component (more than half of the
300 nodes are in a connected network) in random graphs (see Li *et al.* (2021) for a thorough review). The
301 primary difference being that we are removing edges, not adding them, and thus we are witnessing the
302 dissolution of a giant component, rather than the emergence of one. Further applications of percolation
303 theory (Li *et al.* 2021) to the topology of sampled ecological networks could improve our understanding of
304 how false-negatives impact the inferences about the structure and dynamics on these networks.

305 Discussion

306 Species interactions enable the persistence and functioning of ecosystems, but our understanding of
307 interactions is limited due to the intrinsic difficulty of sampling them. Here we have provided a null
308 model for the expected number of false-negatives in an interaction dataset. We demonstrated that we
309 expect many false-negatives in species interaction datasets purely due to the intrinsic variation of
310 abundances within a community. We also, for the first time to our knowledge, measured the strength of
311 association between co-occurrence and interactions (Cazelles *et al.* 2016) across many empirical systems,
312 and found that these positive associations are both very common, and showed algebraically that they
313 increase the realized FNR. We have also shown that false-negatives could further impact our ability to
314 both predict interactions and infer properties of the networks, which highlights the need for further
315 research into methods for correcting this bias in existing data.

316 A better understanding of how false-negatives impact species interaction data is a practical
317 necessity—both for inference of network structure and dynamics, but also for prediction of interactions by
318 using species level information. False-negatives could pose a problem for many forms of inference in
319 network ecology. For example, inferring the dynamic stability of a network could be prone to error if the
320 observed network is not sampled “enough.” What exactly “enough” means is then specific to the
321 application, and should be assessed via methods like those here when designing samples. Further,

322 predictions about network rewiring (Thompson & Gonzalez 2017) due to range shifts in response to
323 climate change could be error-prone without accounting for interactions that have not been observed but
324 that still may become climatically infeasible. As is evident from fig. 2(A), we can never guarantee there are
325 no false-negatives in data. In recent years, there has been interest toward explicitly accounting for
326 false-negatives in models (Stock *et al.* 2017; Young *et al.* 2021), and a predictive approach to
327 networks—rather than expecting our samples to fully capture all interactions (Strydom *et al.* 2021). As a
328 result, better models for predicting interactions are needed for interaction networks. This includes
329 explicitly accounting for observation error (Johnson & Larremore 2021)—certain classes of models have
330 been used to reflect hidden states which account for detection error in occupancy modeling (Joseph 2020),
331 and could be integrated in the predictive models of interactions in the future.

332 This work has several practical consequences for the design of surveys for species' interactions.
333 Simulating the process of observation could be a powerful tool for estimating the sampling effort required
334 by a study that takes relative abundance into account, and provides a null baseline for expected FNR. It is
335 necessary to take the size of the species pool into account when deciding how many total samples is
336 sufficient for an “acceptable” FNR (fig. 2(C & D)). Further the spatial and temporal turnover of
337 interactions means any approach to sampling prioritization must be spatiotemporal. We demonstrated
338 earlier that observed negatives outside of the range of both species aren't informative, and therefore using
339 species distribution models could aid in this spatial prioritization of sampling sites.

340 We also should address the impact of false-negatives on the inference of process and causality in
341 community ecology. We demonstrated that in model food webs, false-negatives do not impact the measure
342 of total trophic levels until very high FNR (figure fig. 5(C)), although we cannot generalize this further to
343 other properties. This has immediate practical concern for how we design what taxa to sample—does it
344 matter if the sampled network is fully connected? It has been shown that the stability of subnetworks can
345 be used to infer the stability of the metaweb paper beyond a threshold of samples (Song *et al.* 2022b). But
346 does this extend to other network properties? And how can we be sure we are at the threshold at which we
347 can be confident our sample characterizes the whole system? We suggest that modeling observation error
348 as we have done here can address these questions and aid in the design of samples of species interactions.
349 To try to survey to avoid all false-negatives is a fool's errand. Species ranges overlap to form mosaics,
350 which themselves are often changing in time. Communities and networks don't end in space, and the
351 interactions that connect species on the ‘periphery’ of a given network to species outside the spatial extent

352 of a given sample will inevitably appear as false-negatives in practical samples. The goal should instead be
353 to sample a system enough to have a statistically robust estimate of the current state and empirical change
354 over time of an ecological community at a given spatial extent and temporal resolution, and to determine
355 what the sampling effort required should be prior to sampling.

356 Our work highlights the need for a quantitatively robust approach to sampling design, both for
357 interactions (Jordano 2016) and all other aspects of biodiversity (Carlson *et al.* 2020). As anthropogenic
358 forces create rapid shifts in our planet's climate and biosphere, this is an imperative to maximize the
359 amount of ecological information we get in our finite samples, and make our inferences and decisions
360 based on this data as robust as possible. Where we choose to sample, and how often we choose to sample
361 there, has strong impacts on the inferences we make from data. Incorporating a better understanding of
362 sampling effort and bias to the design of biodiversity monitoring systems, and the inference and predictive
363 models we apply to this data, is imperative in understanding how biodiversity is changing, and making
364 forecasts that can guide conservation action.

365 **Acknowledgements**

366 AG & MDC acknowledge the support of the Liber Ero Chair for Biodiversity conservation and NSERC.

367 **References**

- 368 Banville, F., Vissault, S. & Poisot, T. (2021). Mangal.jl and EcologicalNetworks.jl: Two complementary
369 packages for analyzing ecological networks in Julia. *Journal of Open Source Software*, 6, 2721.
- 370 Bascompte, J. & Jordano, P. (2007). Plant-Animal Mutualistic Networks: The Architecture of Biodiversity.
371 *Annual Review of Ecology, Evolution, and Systematics*, 38, 567–593.
- 372 Becker, D.J., Albery, G.F., Sjodin, A.R., Poisot, T., Bergner, L.M., Dallas, T.A., *et al.* (2021). Optimizing
373 predictive models to prioritize viral discovery in zoonotic reservoirs.
- 374 Bezanson, J., Edelman, A., Karpinski, S. & Shah, V.B. (2015). Julia: A Fresh Approach to Numerical
375 Computing.

Blanchet, F.G., Cazelles, K. & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions. *Ecology Letters*, 23, 1050–1063.

Blaom, A.D., Kiraly, F., Lienart, T., Simillides, Y., Arenas, D. & Vollmer, S.J. (2020). MLJ: A Julia package for composable machine learning. *Journal of Open Source Software*, 5, 2704.

Boakes, E.H., Rout, T.M. & Collen, B. (2015). Inferring species extinction: The use of sighting records. *Methods in Ecology and Evolution*, 6, 678–687.

Canard, E., Mouquet, N., Marescot, L., Gaston, K.J., Gravel, D. & Mouillot, D. (2012). Emergence of Structural Patterns in Neutral Trophic Networks. *PLOS ONE*, 7, e38295.

CaraDonna, P.J., Petry, W.K., Brennan, R.M., Cunningham, J.L., Bronstein, J.L., Waser, N.M., *et al.* (2017). Interaction rewiring and the rapid turnover of plantpollinator networks. *Ecology Letters*, 20, 385–394.

Carlson, C.J., Dallas, T.A., Alexander, L.W., Phelan, A.L. & Phillips, A.J. (2020). What would it take to describe the global diversity of parasites? *Proceedings of the Royal Society B: Biological Sciences*, 287, 20201841.

Cazelles, K., Araújo, M.B., Mouquet, N. & Gravel, D. (2016). A theory for species co-occurrence in interaction networks. *Theoretical Ecology*, 9, 39–48.

Closs, G.P. & Lake, P.S. (1994). Spatial and Temporal Variation in the Structure of an Intermittent-Stream Food Web. *Ecological Monographs*, 64, 1–21.

de Aguiar, M.A.M., Newman, E.A., Pires, M.M., Yeakel, J.D., Boettiger, C., Burkle, L.A., *et al.* (2019). Revealing biases in the sampling of ecological interaction networks. *PeerJ*, 7, e7566.

Gauzens, B., Legendre, S., Lazzaro, X. & Lacroix, G. (2013). Food-web aggregation, methodological and functional issues. *Oikos*, 122, 1606–1615.

Giacomuzzo, E. & Jordán, F. (2021). Food web aggregation: Effects on key positions. *Oikos*, 130, 2170–2181.

Griffiths, D. (1998). Sampling effort, regression method, and the shape and slope of sizeabundance relations. *Journal of Animal Ecology*, 67, 795–804.

Hadfield, J.D., Krasnov, B.R., Poulin, R. & Nakagawa, S. (2014). A Tale of Two Phylogenies: Comparative Analyses of Ecological Interactions. *The American Naturalist*, 183, 174–187.

403 Havens, K. (1992). Scale and Structure in Natural Food Webs. *Science*, 257, 1107–1109.

404 Johnson, E.K. & Larremore, D.B. (2021). Bayesian estimation of population size and overlap from random
405 subsamples.

406 Jordano, P. (2016). Sampling networks of ecological interactions. *Functional Ecology*, 30, 1883–1893.

407 Joseph, M.B. (2020). Neural hierarchical models of ecological populations. *Ecology Letters*, 23, 734–747.

408 Kenall, A., Harold, S. & Foote, C. (2014). An open future for ecological and evolutionary data? *BMC*
409 *Evolutionary Biology*, 14, 66.

410 Khoshraftar, S. & An, A. (2022). A Survey on Graph Representation Learning Methods.

411 Koleff, P., Gaston, K.J. & Lennon, J.J. (2003). Measuring beta diversity for presenceabsence data. *Journal*
412 *of Animal Ecology*, 72, 367–382.

413 Kopelke, J.-P., Nyman, T., Cazelles, K., Gravel, D., Vissault, S. & Roslin, T. (2017). Food-web structure of
414 willow-galling sawflies and their natural enemies across Europe. *Ecology*, 98, 1730–1730.

415 Li, M., Liu, R.-R., Lü, L., Hu, M.-B., Xu, S. & Zhang, Y.-C. (2021). Percolation on complex networks:
416 Theory and application. *Physics Reports*, Percolation on complex networks: Theory and application,
417 907, 1–68.

418 MacDonald, A.A.M., Banville, F. & Poisot, T. (2020). Revisiting the Links-Species Scaling Relationship in
419 Food Webs. *Patterns*, 1.

420 Makiola, A., Compson, Z.G., Baird, D.J., Barnes, M.A., Boerlijst, S.P., Bouchez, A., *et al.* (2020). Key
421 Questions for Next-Generation Biomonitoring. *Frontiers in Environmental Science*, 7.

422 Martinez, N.D., Hawkins, B.A., Dawah, H.A. & Feifarek, B.P. (1999). Effects of Sampling Effort on
423 Characterization of Food-Web Structure. *Ecology*, 80, 1044–1055.

424 McLeod, A., Leroux, S.J., Gravel, D., Chu, C., Cirtwill, A.R., Fortin, M.-J., *et al.* (2021). Sampling and
425 asymptotic network properties of spatial multi-trophic networks. *Oikos*, 130, 2250–2259.

426 Moore, A.L. & McCarthy, M.A. (2016). Optimizing ecological survey effort over space and time. *Methods*
427 *in Ecology and Evolution*, 7, 891–899.

428 Niedballa, J., Wilting, A., Sollmann, R., Hofer, H. & Courtiol, A. (2019). Assessing analytical methods for
429 detecting spatiotemporal interactions between species from camera trapping data. *Remote Sensing in*

430 *Ecology and Conservation*, 5, 272–285.

431 Paine, R.T. (1988). Road Maps of Interactions or Grist for Theoretical Development? *Ecology*, 69,
432 1648–1654.

433 Poisot, T. (2022). Guidelines for the prediction of species interactions through binary classification.

434 Poisot, T., Bergeron, G., Cazelles, K., Dallas, T., Gravel, D., MacDonald, A., *et al.* (2021). Global knowledge
435 gaps in species interaction networks data. *Journal of Biogeography*, 48, 1552–1563.

436 Poisot, T., Ouellet, M.-A., Mollentze, N., Farrell, M.J., Becker, D.J., Brierly, L., *et al.* (2022). Network
437 embedding unveils the hidden interactions in the mammalian virome.

438 Poisot, T., Stouffer, D.B. & Gravel, D. (2015). Beyond species: Why ecological interaction networks vary
439 through space and time. *Oikos*, 124, 243–251.

440 Ponisio, L.C., Gaiarsa, M.P. & Kremen, C. (2017). Opportunistic attachment assembles plantpollinator
441 networks. *Ecology Letters*, 20, 1261–1272.

442 Savage, V.M., Gillooly, J.F., Brown, J.H., West, G.B. & Charnov, E.L. (2004). Effects of Body Size and
443 Temperature on Population Growth. *The American Naturalist*, 163, 429–441.

444 Song, C., Simmons, B.I., Fortin, M.-J. & Gonzalez, A. (2022a). Generalism drives abundance: A
445 computational causal discovery approach. *PLOS Computational Biology*, 18, e1010302.

446 Song, C., Simmons, B.I., Fortin, M.-J., Gonzalez, A., Kaiser-Bunbury, C.N. & Saavedra, S. (2022b). Rapid
447 monitoring for ecological persistence.

448 Stephenson, P. (2020). Technological advances in biodiversity monitoring: Applicability, opportunities
449 and challenges. *Current Opinion in Environmental Sustainability*, Open issue 2020 part A: Technology
450 Innovations and Environmental Sustainability in the Anthropocene, 45, 36–41.

451 Stock, M., Poisot, T., Waegeman, W. & De Baets, B. (2017). Linear filtering reveals false negatives in
452 species interaction data. *Scientific Reports*, 7, 45908.

453 Strydom, T., Catchen, M.D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., *et al.* (2021). A
454 roadmap towards predicting species interaction networks (across space and time). *Philosophical
455 Transactions of the Royal Society B: Biological Sciences*, 376, 20210063.

456 Thompson, P.L. & Gonzalez, A. (2017). Dispersal governs the reorganization of ecological networks under
 457 environmental change. *Nature Ecology & Evolution*, 1, 1–8.

458 Thompson, R.M. & Townsend, C.R. (2000). Is resolution the solution?: The effect of taxonomic resolution
 459 on the calculated properties of three stream food webs. *Freshwater Biology*, 44, 413–422.

460 Volkov, I., Banavar, J.R., Hubbell, S.P. & Maritan, A. (2003). Neutral theory and relative species abundance
 461 in ecology. *Nature*, 424, 1035–1037.

462 Walther, B.A., Cotgreave, P., Price, R.D., Gregory, R.D. & Clayton, D.H. (1995). Sampling Effort and
 463 Parasite Species Richness. *Parasitology Today*, 11, 306–310.

464 Williams, R.J. & Martinez, N.D. (2000). Simple rules yield complex food webs. *Nature*, 404, 180–183.

465 Willott, S.j. (2001). Species accumulation curves and the measure of sampling effort. *Journal of Applied*
 466 *Ecology*, 38, 484–486.

467 Young, J.-G., Valdovinos, F.S. & Newman, M.E.J. (2021). Reconstruction of plantpollinator networks from
 468 observational data. *Nature Communications*, 12, 3911.

469 Zhang, S., Tong, H., Xu, J. & Maciejewski, R. (2019). Graph convolutional networks: A comprehensive
 470 review. *Computational Social Networks*, 6, 11.