

RADAR MIDTERM - Joshua Gould

Contents

- [SNR at 20km](#)
- [CNR at 20km](#)
- [Generate S/\(C+N\) at 20 km.](#)
- [Mean sea clutter reflectivity at S/\(C+N\) of exactly 15 dB?](#)

```
clc
f = 5; %GHz (C-Band - vertical polarization)
PKP = 5; %kW
CPA = 1.00; %meters
NoiseFigure = 3;
TotalSystemLosses = 10; %dB
Dutyf = .05; %
DwellL = 1.0; %ms
B = 1; %MHz.
```

SNR at 20km

```
Ar = (0.5);
c = 299792458;
R_g = (Ar*4*pi)^2 / (WL^2);
R_g_db = 10*log10(R_g);
SNR = 7.2724;
fprintf('The SNR at 20km = %f db. ', SNR);
fprintf('\nThe SNR is NOT detectable as %f is less than 15.', SNR);
```

The SNR at 20km = 7.272400 db.

The SNR is NOT detectable as 7.272400 is less than 15.

CNR at 20km

```
clc
D = 1.00; %diameter
PKP = 10*log10(5000); %peak power

Tpross = 10* log10(.001); %processing time, ms
Gt = 34.38;
Gr = 34.38;
DF = 10*log10(0.05);
WL = (c/5000000000); %wavelength lamda
WLdb = 10*log10(90000 / (5000)^2);
pi4 = 10 * log10(64*pi*pi*pi);
Bwidth = WL/D; %beamwidth
PulseWidth = 1 / (B*1000000);
km = 40*log10(20000); %m
Sstate = 3; %sea state
```

```

AngleG = 0.125; %grazing angle
L = 1 / B;
Pol = 'V';
NRL = NRL_SigmaSea(f,Sstate,'V',AngleG);
WCC = 20 * 1000 * Bwidth; %Width of clutter cell, m
RCC = PulseWidth * (1.5*(10)^8);%Range of clutter cell, m =pulsewidth x 1.5E8 m/s
CLUTA = WCC * RCC; %Clutter Area
RCSdb = 10* log10(CLUTA * 10^(NRL / 10));
RCS = CLUTA * 10^(NRL / 10);
k = 228.60; % Boltzman Constant
T0 = 24.62; %temp
Nfig = 3;
Eloss = 10;
CNR = Gt+ Gr + PKP + RCSdb + DF + k - pi4 + Tpross - T0 + WLdb - Nfig - Eloss - km;
fprintf('The CNR at 20km = %f db. ', CNR);
fprintf('\nThe Clutter Cross Sectional area = %f dbm^2. ', RCS);

```

The CNR at 20km = 15.898734 db.
The Clutter Cross Sectional area = 0.145673 dbm^2.

Generate S/(C+N) at 20 km.

```

clc
SCPN = 1/(1/(SNR) + 1/(CNR));
fprintf('The S/(C+N) at 20km = %f db. ', SCPN);
fprintf('\nThe target is NOT detectable as %f is less than 15 db.', SCPN);

```

The S/(C+N) at 20km = 4.989913 db.
The target is NOT detectable as 4.989913 is less than 15 db.

Mean sea clutter reflectivity at S/(C+N) of exactly 15 dB?

```

clc;
NRLn = -90.931063;
RCSn = 10*log10(CLUTA * 10^(NRLn / 10));
CNRn = Gt + Gr + PKP + RCSn + DF + k - pi4 + Tpross - T0 + WLdb - Nfig - Eloss - km;
fprintf('Mean sea clutter reflectivity = %f dbm^2 / m^2',NRLn);
fprintf('\nNew CNR = %f db',CNRn);
SCPNn = 1/(1/(SNR) + 1/(CNRn));
fprintf('\nNew S/(C+N) = %f db',SCPNn);

```

Mean sea clutter reflectivity = -90.931063 dbm^2 / m^2
New CNR = -14.116414 db
New S/(C+N) = 15.000000 db

