

# Tanaka Certificates

July 15, 2026

## Abstract

In these notes, I am building up the intuition and theory behind supermartingale certificates, with relaxed assumptions on the certificate function  $V$ . In particular, it is explored what happens when  $V$  is not twice continuously-differentiable, but instead piecewise twice continuously-differentiable.

## Contents

|          |                                                                             |           |
|----------|-----------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Piecewise linear certificates in dimension one</b>                       | <b>2</b>  |
| 1.1      | Proof of the supermartingale property . . . . .                             | 3         |
| <b>2</b> | <b>Piecewise <math>\mathcal{C}^2</math> certificates in dimension one</b>   | <b>3</b>  |
| <b>3</b> | <b>Studying piecewise certificates on concrete one-dimensional examples</b> | <b>5</b>  |
| 3.1      | Ornstein–Uhlenbeck process in three parameter regimes . . . . .             | 5         |
| 3.2      | Failure of piecewise linearity and success of curvature . . . . .           | 5         |
| 3.3      | A piecewise $\mathcal{C}^2$ certificate with curvature and a kink . . . . . | 7         |
| <b>4</b> | <b>Piecewise quadratic certificates in dimension one</b>                    | <b>8</b>  |
| 4.1      | Failure of piecewise linear certificates in dimension one . . . . .         | 8         |
| 4.2      | Sufficiency of piecewise quadratic certificates in dimension one . . . . .  | 10        |
| 4.3      | Connection to Taylor expansion of the best possible certificate . . . . .   | 11        |
| <b>5</b> | <b>Piecewise quadratic supermartingales in multiple dimensions</b>          | <b>11</b> |
| 5.1      | Proving the supermartingale property . . . . .                              | 11        |
| 5.1.1    | Proof of the supermartingale property . . . . .                             | 14        |
| 5.1.2    | Interpretation of the multidimensional concavity condition . . . . .        | 15        |
| 5.1.3    | Summary of sufficient conditions . . . . .                                  | 15        |
| 5.2      | Reach–avoid stopping convention . . . . .                                   | 16        |
| 5.3      | Asymptotic completeness of piecewise quadratic certificates . . . . .       | 16        |
| <b>6</b> | <b>The local-time condition by construction</b>                             | <b>18</b> |
| 6.1      | The deep ReLU ICNN . . . . .                                                | 18        |
| 6.2      | Convexity determines the interface sign . . . . .                           | 19        |
| 6.3      | The residual certificate . . . . .                                          | 19        |

# 1 Piecewise linear certificates in dimension one

Consider the following setup:

## Setup

Let:

1.  $dX_t = f(X_t)dt + g(X_t)dW_t$  be a one-dimensional SDE
2.  $V(x)$  be a piecewise linear function such that  $V(x) = a_i x + b_i$ , where  $x \in (x_i, x_{i+1})$ .
3.  $V(x)$  is continuous at breakpoints:  $V(x_i^+) = V(x_i^-) = V(x_i)$
4. as a consequence,  $V'_-(x_i) = a_{i-1}$ ,  $V'_+(x_i) = a_i$

Now we will consider what conditions on  $V$  are sufficient to guarantee that  $V(X_t)$  is a supermartingale.

## Hypothesis

Does concavity of  $V(x)$  (along with an interior / drift term condition) guarantee that  $V(X_t)$  is a supermartingale?

To answer this, let's apply the Itô-Tanaka formula to  $V(X_t)$ :

$$V(X_t) = V(X_0) + \int_0^t V'(X_s) f(X_s) ds \quad (1)$$

$$+ \int_0^t V'(X_s) g(X_s) dW_s \quad (2)$$

$$+ \frac{1}{2} \sum_{i=1}^{n-1} (V'_+(x_i) - V'_-(x_i)) L_t^{x_i}(X) \quad (3)$$

$$(4)$$

We will focus on these terms:

1.  $D_t = \int_0^t V'(X_s) f(X_s) ds$ , the drift term
2.  $M_t = \int_0^t V'(X_s) g(X_s) dW_s$ , the diffusion term (stochastic integral)
3.  $K_t = \frac{1}{2} \sum_{i=1}^{n-1} (V'_+(x_i) - V'_-(x_i)) L_t^{x_i}(X)$ , the kink term

we need to prove that  $D_t + M_t + K_t$  is a supermartingale.

Some natural assumptions, which are sufficient to guarantee that  $V(X_t)$  is a supermartingale, are the following:

1.  $f(x)V'(x) \leq 0$  for all  $x \in (x_i, x_{i+1})$ . This is because of the following:

## Fact

If  $f(X)V'(X) \leq 0$ , then the drift term  $D_t$  is a supermartingale.

**Proof:** We want to prove  $\mathbb{E}[D_t | \mathcal{F}_s] \leq D_s$  for  $s \leq t$ .

$$\mathbb{E}[D_t | \mathcal{F}_s] = \mathbb{E} \left[ \int_0^t V'(X_u) f(X_u) du \middle| \mathcal{F}_s \right] \quad (5)$$

$$= \int_0^s V'(X_u) f(X_u) du + \mathbb{E} \left[ \int_s^t V'(X_u) f(X_u) du \middle| \mathcal{F}_s \right] \quad (6)$$

$$\leq D_s + 0 = D_s \quad (7)$$

2.  $M_t$  is a martingale. Assuming  $V(x)$  has finitely many pieces (as is the case for a neural network with ReLU activation functions), it is enough to impose a square-integrability condition on  $g(x)$ <sup>1</sup>
3.  $V'_+(x_i) - V'_-(x_i) \leq 0 \iff a_{i-1} \geq a_i$ , i.e. the **concavity condition**. This is because of the following:

<sup>1</sup>For example,  $\mathbb{E} \left[ \int_0^t g(X_s)^2 ds \right] < \infty$ .

**Fact**

If  $V'_+(x_i) - V'_-(x_i) \leq 0$  and  $L_t^{x_i}(X) \geq 0$ , then the kink term  $K_t$  is a supermartingale.

**Proof:** We want to prove  $\mathbb{E}[K_t|\mathcal{F}_s] \leq K_s$  for  $s \leq t$ .

$$\mathbb{E}[K_t|\mathcal{F}_s] = \mathbb{E}\left[\frac{1}{2} \sum_{i=1}^{n-1} (V'_+(x_i) - V'_-(x_i)) L_t^{x_i}(X) \middle| \mathcal{F}_s\right] \quad (8)$$

$$= \frac{1}{2} \sum_{i=1}^{n-1} (V'_+(x_i) - V'_-(x_i)) \mathbb{E}[L_t^{x_i}(X)|\mathcal{F}_s] \quad (9)$$

$$\leq \frac{1}{2} \sum_{i=1}^{n-1} (V'_+(x_i) - V'_-(x_i)) L_s^{x_i}(X) = K_s \quad (10)$$

Note that the inequality follows from the fact that  $L_t^{x_i}(X)$  is non-decreasing in  $t$ , hence  $\mathbb{E}[L_t^{x_i}(X)|\mathcal{F}_s] \geq \mathbb{E}[L_s^{x_i}(X)|\mathcal{F}_s] = L_s^{x_i}(X)$ .

We will consider what happens in different practical cases to these terms, including:

1. A simple one-dimensional SDE with a piecewise linear certificate.
2. A pendulum with a stochastic torque, and  $X_t$  measuring the angle of the pendulum.
3. A network  $b$  black and  $w$  white nodes, with given transition probabilities, and  $X_t$  measuring the amount of mass in the white nodes.

to ensure our assumptions are reasonable.

## 1.1 Proof of the supermartingale property

First, let's note the following:

**Fact**

The sum of a martingale and a supermartingale is a supermartingale.

**Proof:** By definition, given a martingale  $M_t$  and a supermartingale  $S_t$ , for  $r < t$  we have  $\mathbb{E}[M_t|\mathcal{F}_r] = M_r$  and  $\mathbb{E}[M_t|\mathcal{F}_r] \leq S_r$ . Then, for  $r < t$ :

$$\mathbb{E}[M_t + S_t|\mathcal{F}_r] = \mathbb{E}[M_t|\mathcal{F}_r] + \mathbb{E}[S_t|\mathcal{F}_r] \quad (11)$$

$$= M_r + \mathbb{E}[S_t|\mathcal{F}_r] \quad (12)$$

$$\leq M_r + S_r = (M + S)_r \quad (13)$$

hence  $M_t + S_t$  is a supermartingale.

The drift term  $D_t$  is a supermartingale by assumption 1. The kink term  $K_t$  is a supermartingale by assumption 3. The stochastic integral  $M_t$  is a martingale by assumption 2. Therefore, the sum of these three terms is a supermartingale (as a consequence of the previous fact), and hence  $V(X_t)$  is a supermartingale.

## 2 Piecewise $\mathcal{C}^2$ certificates in dimension one

The setup is pretty similiar, with the exception that  $V$  is now piecewise  $\mathcal{C}^2$  instead of piecewise linear.

**Setup**

Let:

1.  $dX_t = f(X_t)dt + g(X_t)dW_t$  be a one-dimensional SDE.
2.  $V(x)$  be a piecewise  $\mathcal{C}^2$  function such that  $V(x) \in \mathcal{C}^2$ , where  $x \in (x_i, x_{i+1})$ .
3.  $V(x)$  is continuous at breakpoints:  $V(x_i^+) = V(x_i^-) = V(x_i)$ .

We will again consider what conditions on  $V$  are sufficient to guarantee that  $V(X_t)$  is a supermartingale. Again, the natural hypothesis is:

**Hypothesis**

Does concavity of  $V(x)$  (along with an interior / drift term condition) guarantee that  $V(X_t)$  is a supermartingale?

We apply the Itô-Tanaka formula to  $V(X_t)$ , and we get similar terms, with the addition of

$$C_t = \frac{1}{2} \int_0^t V''(X_s) g^2(X_s) ds \quad (14)$$

Since the integral in the  $C_t$  term is over the same intervals as the  $D_t$  term, we can combine them to recover the generator / Lie derivative term:

$$D_t + C_t = \int_0^t \left( V'(X_s) f(X_s) + \frac{1}{2} V''(X_s) g^2(X_s) \right) ds = \int_0^t \mathcal{L}V(X_s) ds \quad (15)$$

The proof that  $V(X_t)$  is a supermartingale is similar to the piecewise linear case, but now we need to impose the condition:

$$\mathcal{L}V(x) = V'(x)f(x) + \frac{1}{2}V''(x)g^2(x) \leq 0 \quad (16)$$

Table 1: Piecewise linear certificates examples.

| SDE              | $V(x)$ | $D_t$                | $M_t$                            | $K_t$       |
|------------------|--------|----------------------|----------------------------------|-------------|
| $dX_t = dW_t$    | $- x $ | 0                    | $\int_0^t -\text{sgn}(X_s) dW_s$ | $-L_t^0(X)$ |
| $dX_t = -X_t dt$ | $ x $  | $-\int_0^t  X_s  ds$ | 0                                | 0           |

Table 2: Piecewise  $\mathcal{C}^2$  certificates examples.

| SDE                                    | $V(x)$                                                                  | $D_t$                         | $M_t$                              | $C_t$                                               | $K_t$                   |
|----------------------------------------|-------------------------------------------------------------------------|-------------------------------|------------------------------------|-----------------------------------------------------|-------------------------|
| $dX_t = -\lambda X_t dt + \sigma dW_t$ | $x^2$                                                                   | $-2\lambda \int_0^t X_s^2 ds$ | $\int_0^t 2\sigma X_s dW_s$        | $\sigma^2 t$                                        | 0                       |
| $dX_t = dt + \sqrt{2} dW_t$            | $x - \frac{x^2}{2}$                                                     | $\int_0^t (1 - X_s) ds$       | $\int_0^t (1 - X_s) \sqrt{2} dW_s$ | $-t$                                                | 0                       |
| $dX_t = dW_t$                          | $\begin{cases} -\frac{1}{2}x^2, & x < 0, \\ -x, & x \geq 0 \end{cases}$ | 0                             | $\int_0^t V'(X_s) dW_s$            | $-\frac{1}{2} \int_0^t \mathbf{1}_{\{X_s < 0\}} ds$ | $-\frac{1}{2} L_t^0(X)$ |

### 3 Studying piecewise certificates on concrete one-dimensional examples

Some useful examples to study are given in Tables 1 and 2.

#### 3.1 Ornstein–Uhlenbeck process in three parameter regimes

Consider the Ornstein–Uhlenbeck process

$$dX_t = -\lambda X_t dt + \sigma dW_t$$

with the quadratic certificate  $V(x) = x^2$ . Its generator is

$$\mathcal{L}V(x) = -2\lambda x^2 + \sigma^2 = -2\lambda \left( x^2 - \frac{\sigma^2}{2\lambda} \right).$$

Consequently, if

$$r = \frac{\sigma}{\sqrt{2\lambda}},$$

then the generator is negative for  $|x| > r$ , zero for  $|x| = r$ , and positive for  $|x| < r$ . The decomposition is

$$X_t^2 - X_0^2 = \underbrace{-2\lambda \int_0^t X_s^2 ds}_{D_t} + \underbrace{2\sigma \int_0^t X_s dW_s}_{M_t} + \underbrace{\sigma^2 t}_{C_t},$$

with  $K_t = 0$ .

Figure 1 fixes  $\lambda = 1$  and  $X_0 = 1$ , and uses  $r = 0.5$ ,  $r = 1$ , and  $r = 1.5$ . These choices make the initial generator negative, zero, and positive, respectively. Moreover,

$$\mathbb{E}[X_t^2] = r^2 + (X_0^2 - r^2)e^{-2\lambda t},$$

so the expected generator retains the corresponding sign. In the middle case,  $X_t^2$  is constant in expectation, but it is not a martingale: the pointwise generator vanishes only when  $|X_t| = r$ .

The stochastic integral  $M_t$  need not visibly oscillate on every individual sample path; the martingale statement is instead that its conditional drift is zero. Figure 1 therefore plots Monte Carlo expectations. The mean of  $M_t$  remains statistically consistent with zero, while  $D_t$  and  $C_t$  determine whether  $\mathbb{E}[X_t^2 - X_0^2]$  decreases, remains zero, or increases.

#### 3.2 Failure of piecewise linearity and success of curvature

For  $f = 1$  and  $g = \sqrt{2}$  on  $[0, 1]$ , consider a case where want to find a certificate  $V$  such that  $V(0) = 0$  and  $V(1) = 1$ .

OU process with  $V(x) = x^2$ ,  $\lambda = 1$ , and  $X_0 = 1$




Figure 1: Monte Carlo expectations from 8000 Ornstein-Uhlenbeck paths for  $V(x) = x^2$ . The three columns use  $\sigma/\sqrt{2\lambda} = 0.5, 1, 1.5$ . The shaded region is a pointwise approximate 95% Monte Carlo band for  $\mathbb{E}[M_t]$ .

A smooth candidate could be:

$$V_{\mathcal{C}^2}(x) = x - \frac{x^2}{2},$$

the curvature contribution compensates for the positive drift:

$$\mathcal{L}V_{\mathcal{C}^2}(x) = (1 - x) + \frac{1}{2}(-1)(2) = -x \leq 0.$$

and indeed  $V_{\mathcal{C}^2}$  is a valid supermartingale certificate.

On the other hand, let's assume  $V$  is piecewise linear. Then there exists an interval  $(x_i, x_{i+1}) \subset [0, 1]$  where  $V'(x) = a_i > 0$ . On that interval, given an  $x \in (x_i, x_{i+1})$ , the curvature term vanishes, and the generator is

$$\mathcal{L}V(x) = V'(x)f(x) + \frac{1}{2}V''(x)g^2(x) = V'(x)f(x) = V'(x) = a_i > 0.$$

This example motivates the following hypothesis, which we will study in more detail in [Section 4](#).

### Hypothesis

While piecewise linear certificates  $V_{\text{PL}}$  are not sufficient to guarantee that  $V_{\text{PL}}$  is a supermartingale, **piecewise quadratic certificates** are sufficient, as the curvature term can compensate for positive drift.




Figure 2: An increasing piecewise-linear candidate fails the generator condition for  $f = 1$ ,  $g = \sqrt{2}$ , whereas negative curvature makes the smooth candidate satisfy it throughout  $[0, 1]$ .

### 3.3 A piecewise $\mathcal{C}^2$ certificate with curvature and a kink

Finally, let  $dX_t = dW_t$  and

$$V(x) = \begin{cases} -\frac{1}{2}x^2, & x < 0, \\ -x, & x \geq 0. \end{cases}$$

Here  $D_t = 0$ , while

$$M_t = \int_0^t V'(X_s) dW_s, \quad C_t = -\frac{1}{2} \int_0^t \mathbf{1}_{\{X_s < 0\}} ds, \quad K_t = -\frac{1}{2} L_t^0(X).$$

Thus  $C_t$  decreases while the path lies below zero, and  $K_t$  decreases only when local time accumulates at the kink. Figure 3 shows these different mechanisms on the same Brownian realization. The plotted local time uses an occupation-density approximation with bandwidth 0.06.




Figure 3: A Brownian realization and the terms for the hybrid piecewise  $\mathcal{C}^2$  certificate. The curvature and kink terms are both nonincreasing, while the martingale term fluctuates.

## 4 Piecewise quadratic certificates in dimension one

In this section, we will study a failure regime of the piecewise linear certificates in dimension one and see how it leads to sufficiency of piecewise quadratic certificates.

Additionally, we observe an interesting connection to the *Taylor expansion* of the best possible certificate, namely a martingale certificate for which the generator is zero.

### 4.1 Failure of piecewise linear certificates in dimension one

We will consider the setup from [Section 1](#) again, but with additional assumptions on absorption at the boundary, an increase condition on the piecewise linear function  $V(x)$ , and a non-degeneracy condition on the diffusion term  $g(x)$ .

The key point is that the piecewise linear setup is not flexible enough when the certificate is required to increase in the same direction as the drift. In that case, a piecewise linear function has no curvature term which could compensate for the positive drift contribution.

So essentially, we will show that:

#### Fact

(*Informally.*) There is a case where no piecewise linear certificate can make  $V(X_t)$  a supermartingale.

Although it should be clear from previous sections, for completeness, the setup is as follows:

#### Setup

Let:

1.  $dX_t = f(X_t)dt + g(X_t)dW_t$  be a one-dimensional SDE.
2.  $V(x)$  be a piecewise linear function such that  $V(x) = a_i x + b_i$ , where  $x \in (x_i, x_{i+1})$ .
3.  $V(x)$  is continuous at breakpoints:  $V(x_i^+) = V(x_i^-) = V(x_i)$ .
4. as a consequence,  $V'_-(x_i) = a_{i-1}$ ,  $V'_+(x_i) = a_i$
5. The  $X_t$  process is **absorbed** at the boundary of  $I = [a, b]$ .
6.  $V(x)$  has the **increase condition** at the boundary:

$$V(a) < V(b). \tag{17}$$

7.  $g(x)^2 > 0$  for all  $x \in (a, b)$ , i.e. the **diffusion term is non-degenerate**.
8.  $M_t$  is a martingale, i.e. the **martingale condition**.

Motivated by the example given in [Section 3.2](#), we will now show that there is a broad class of SDEs for which increasing piecewise linear certificates cannot be supermartingales.



### Fact

Assume that for all  $\varepsilon > 0$ , there exists  $x \in (a, a + \varepsilon)$  such that

$$f(x) > 0. \quad (18)$$

Then there is no continuous piecewise linear  $V$  satisfying the increase condition

$$V(a) < V(b) \quad (19)$$

and the concavity condition

$$V'_+(x_i) - V'_-(x_i) \leq 0 \quad (20)$$

at every breakpoint, such that  $V(X_t)$  is a supermartingale.

**Proof.** Assume, for contradiction, that such a piecewise linear function  $V$  exists and that  $V(X_t)$  is a supermartingale.

Let  $p(x) = V'(x)$  on each linear region. Since  $V$  is piecewise linear, the Itô–Tanaka formula gives

$$V(X_t) = V(X_0) + \int_0^t p(X_s) f(X_s) ds \quad (21)$$

$$+ \int_0^t p(X_s) g(X_s) dW_s \quad (22)$$

$$+ \frac{1}{2} \sum_i (V'_+(x_i) - V'_-(x_i)) L_t^{x_i}(X). \quad (23)$$

The stochastic integral is a martingale by assumption. Therefore, if  $V(X_t)$  is a supermartingale, the finite-variation part cannot create positive drift on any linear region. Hence the necessary interior condition is

$$p(x)f(x) \leq 0 \quad \text{on every linear region.} \quad (24)$$

This also follows from [Fact 1](#) in [Section 1](#).

We now use the concavity condition. Since

$$V'_+(x_i) - V'_-(x_i) \leq 0 \quad (25)$$

at every breakpoint, the slopes of  $V$  are nonincreasing from left to right. Therefore  $V$  is concave.

Let  $(a, x_1)$  be the first linear region. We claim that the slope of  $V$  on this first region must be positive. Indeed, suppose instead that the first slope were nonpositive. Since the slopes are nonincreasing, all later slopes would also be nonpositive. Thus,

$$V(b) - V(a) = \int_a^b V'(x) dx \quad (26)$$

$$\leq 0. \quad (27)$$

This contradicts the increase condition  $V(a) < V(b)$ . Hence the first slope must be positive, so

$$p(x) > 0 \quad \text{for all } x \in (a, x_1). \quad (28)$$

Now use the assumption on the drift. Since for every  $\varepsilon > 0$  there exists  $x \in (a, a + \varepsilon)$  such that  $f(x) > 0$ , we may choose  $\varepsilon = x_1 - a$ . Then there exists  $x^* \in (a, x_1)$  such that

$$f(x^*) > 0. \quad (29)$$

Combining this with (28), we get

$$p(x^*)f(x^*) > 0. \quad (30)$$

This contradicts the necessary interior condition (24). Therefore no such piecewise linear  $V$  can make  $V(X_t)$  a supermartingale.

The conclusion is that piecewise linear certificates fail when the certificate is forced to increase in the same

direction as the drift. The obstruction comes from the fact that piecewise linear functions have no interior curvature term. Indeed, away from the kinks,

$$\mathcal{L}V(x) = V'(x)f(x), \quad (31)$$

so a positive drift contribution cannot be compensated. This motivates the piecewise quadratic case, where the additional curvature term

$$\frac{1}{2}V''(x)g(x)^2 \quad (32)$$

can offset the positive drift contribution.

## 4.2 Sufficiency of piecewise quadratic certificates in dimension one

In this subsection we will show that piecewise quadratic certificates are sufficient to guarantee that  $V(X_t)$  is a supermartingale.

Let  $V$  be piecewise quadratic on  $I = [a, b]$ , so that on every interval  $(x_i, x_{i+1})$  we have

$$V(x) = a_i x^2 + b_i x + c_i. \quad (33)$$

Assume also that  $V$  is continuous at the breakpoints. Then

$$V'(x) = 2a_i x + b_i, \quad V''(x) = 2a_i, \quad x \in (x_i, x_{i+1}). \quad (34)$$

Therefore the generator on the interior of a quadratic piece is

$$\mathcal{L}V(x) = (2a_i x + b_i) f(x) + a_i g(x)^2. \quad (35)$$

Here, there always exists a solution  $(a_i, b_i)$  to the interior condition  $\mathcal{L}V(x) \leq 0$  on each piece. This gives the following sufficient condition.

### Fact

Let  $V$  be continuous and piecewise quadratic on  $I = [a, b]$ . Suppose that:

1. on every open piece  $(x_i, x_{i+1})$ ,

$$\mathcal{L}V(x) = V'(x)f(x) + \frac{1}{2}V''(x)g(x)^2 \leq 0; \quad (36)$$

2. at every breakpoint,

$$V'_+(x_i) - V'_-(x_i) \leq 0; \quad (37)$$

3. the stochastic integral

$$M_t = \int_0^t V'(X_s)g(X_s) dW_s \quad (38)$$

is a martingale.

Then  $V(X_t)$  is a supermartingale.

**Proof.** By the Itô–Tanaka formula for a piecewise  $C^2$  function,

$$V(X_t) = V(X_0) + \int_0^t \left( V'(X_s)f(X_s) + \frac{1}{2}V''(X_s)g(X_s)^2 \right) ds \quad (39)$$

$$+ \int_0^t V'(X_s)g(X_s) dW_s \quad (40)$$

$$+ \frac{1}{2} \sum_i (V'_+(x_i) - V'_-(x_i)) L_t^{x_i}(X). \quad (41)$$

The first term is nonincreasing by (36). The stochastic integral is a martingale by assumption. Finally, since local time is nondecreasing in  $t$ , the kink term is nonincreasing by (37). Therefore  $V(X_t)$  is a supermartingale.

### 4.3 Connection to Taylor expansion of the best possible certificate

Let us now again consider the SDE

$$dX_t = dt + \sqrt{2} dW_t, \quad (42)$$

the generator is

$$\mathcal{L}V(x) = V'(x) + V''(x). \quad (43)$$

A martingale certificate is obtained by asking for the generator to vanish:

$$V'(x) + V''(x) = 0. \quad (44)$$

Solving this ODE gives

$$V(x) = C_1 + C_2 e^{-x}. \quad (45)$$

If we impose the normalization  $V(0) = 0$  and  $V(1) = 1$ , then

$$V_\star(x) = \frac{1 - e^{-x}}{1 - e^{-1}}. \quad (46)$$

This is the martingale certificate:  $\mathcal{L}V_\star(x) = 0$  on the interior.

Now expand the numerator of (46) around 0:

$$1 - e^{-x} = x - \frac{x^2}{2} + \frac{x^3}{6} - \dots. \quad (47)$$

The quadratic certificate used above is exactly the second-order truncation of this numerator:

$$V_{\text{quad}}(x) = x - \frac{x^2}{2}. \quad (48)$$

So, up to the constant  $(1 - e^{-1})^{-1}$  and up to truncation error, the quadratic certificate is the Taylor approximation of the martingale certificate.

## 5 Piecewise quadratic supermartingales in multiple dimensions

Now we will finally consider a very general setup.

First, we prove that when a piecewise-quadratic function  $V : \mathbb{R}^d \rightarrow \mathbb{R}$  satisfies similar concavity and smoothness conditions as in the one-dimensional case, then  $V(X_t)$  is a supermartingale.

Second, we will prove that if a smooth certificate exists, then we can construct a piecewise quadratic certificate that is a supermartingale, essentially showing how quadratic certificates are in some sense sufficient, while piecewise linear certificates are not.

### 5.1 Proving the supermartingale property

We will show that a piecewise quadratic function gives a supermartingale if:

1. the generator is nonpositive inside each region,
2. the stochastic integral is a martingale,
3. the normal derivative jumps in the correct direction across each interface.

### Setup

Let:

1.  $dX_t = f(X_t)dt + \sigma(X_t)dW_t$  be a  $d$ -dimensional SDE, where  $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  and  $\sigma : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$ .
2.  $W_t$  is an  $m$ -dimensional Brownian motion.
3.  $a(x) = \sigma(x)\sigma(x)^\top$ .
4.  $K \subseteq \mathbb{R}^d$  is a compact set partitioned into finitely many polyhedral cells:

$$K = \bigcup_{i=1}^N K_i.$$

5.  $V : K \rightarrow \mathbb{R}$  is a continuous piecewise quadratic function such that

$$V(x) = V_i(x) = c_i + p_i^\top x + \frac{1}{2}x^\top Q_i x, \quad x \in K_i,$$

where  $Q_i = Q_i^\top$ .

6.  $V$  is continuous across interfaces:

$$V_i(x) = V_j(x), \quad x \in K_i \cap K_j.$$

Now we will consider what conditions on  $V$  are sufficient to guarantee that  $V(X_t)$  is a supermartingale, at least up to the first exit time from  $K$ .

Define the stopping time

$$\tau_K = \inf\{t \geq 0 : X_t \notin K\}.$$

### Hypothesis

Does a multidimensional analogue of concavity, together with an interior generator condition, guarantee that

$$V(X_{t \wedge \tau_K})$$

is a supermartingale?

To answer this, we apply the multidimensional Itô–Tanaka formula to  $V(X_{t \wedge \tau_K})$ .

On each cell  $K_i$ , we have

$$\nabla V_i(x) = p_i + Q_i x, \quad \nabla^2 V_i(x) = Q_i.$$

Therefore, the infinitesimal generator of  $V_i$  is

$$\mathcal{L}V_i(x) = \nabla V_i(x) \cdot f(x) + \frac{1}{2} \text{tr}(a(x) \nabla^2 V_i(x)).$$

Substituting the quadratic form gives

$$\mathcal{L}V_i(x) = (p_i + Q_i x) \cdot f(x) + \frac{1}{2} \text{tr}(a(x) Q_i).$$

For  $i \neq j$ , let

$$\mathcal{F}_{ij} = \{F : F \text{ is a connected } (d-1)\text{-dimensional face of } K_i \cap K_j\}.$$

### Sidenote

Each cell  $K_i$  carries its own quadratic piece  $V_i$ ; the function need not be given by one quadratic expression on  $K_i \cup K_j$ . Two such cells can touch along more than one  $(d-1)$ -dimensional interface, or facet. The set  $\mathcal{F}_{ij}$  records these connected faces separately so that the normal and the Tanaka condition can be specified on each one.

For each  $F \in \mathcal{F}_{ij}$ , choose a unit normal  $n_{i \rightarrow j, F}$  pointing from  $K_i$  into  $K_j$ . If

$$F \subseteq \{x : h_F^\top x = \gamma_F\},$$

then one may take

$$n_{i \rightarrow j, F} = \text{sign}(h_F^\top (x_j - x_i)) \frac{h_F}{\|h_F\|},$$

where  $x_i \in \text{int}(K_i)$  and  $x_j \in \text{int}(K_j)$ .

The multidimensional Itô–Tanaka formula gives

$$V(X_{t \wedge \tau_K}) = V(X_0) + \int_0^{t \wedge \tau_K} \mathcal{L}V(X_s) ds \quad (49)$$

$$+ \int_0^{t \wedge \tau_K} \nabla V(X_s)^\top \sigma(X_s) dW_s \quad (50)$$

$$+ \frac{1}{2} \sum_{i < j} \sum_{F \in \mathcal{F}_{ij}} \int_0^{t \wedge \tau_K} (\nabla V_j(X_s) - \nabla V_i(X_s)) \cdot n_{i \rightarrow j, F} dL_s^F. \quad (51)$$

Here  $L_t^F$  denotes the surface local time accumulated by  $X_t$  on the face  $F$ .

We will focus on these terms:

1.

$$D_t = \int_0^{t \wedge \tau_K} \mathcal{L}V(X_s) ds,$$

the interior drift term.

2.

$$M_t = \int_0^{t \wedge \tau_K} \nabla V(X_s)^\top \sigma(X_s) dW_s,$$

the diffusion term.

3.

$$K_t = \frac{1}{2} \sum_{i < j} \sum_{F \in \mathcal{F}_{ij}} \int_0^{t \wedge \tau_K} (\nabla V_j(X_s) - \nabla V_i(X_s)) \cdot n_{i \rightarrow j, F} dL_s^F,$$

the interface term.

We need to prove that

$$D_t + M_t + K_t$$

is a supermartingale.

Some natural assumptions, which are sufficient to guarantee that  $V(X_{t \wedge \tau_K})$  is a supermartingale, are the following.

1. The interior generator is nonpositive on every cell:

$$\mathcal{L}V_i(x) \leq 0, \quad x \in \text{int}(K_i).$$

Equivalently,

$$(p_i + Q_i x) \cdot f(x) + \frac{1}{2} \text{tr}(a(x) Q_i) \leq 0, \quad x \in \text{int}(K_i).$$

This is the multidimensional analogue of the condition

$$f(x)V'(x) \leq 0$$

from the one-dimensional piecewise linear case.

#### Fact

If  $\mathcal{L}V(X_s) \leq 0$  for all  $s \leq \tau_K$ , then the drift term  $D_t$  is a supermartingale.

**Proof:** We want to prove

$$\mathbb{E}[D_t \mid \mathcal{F}_r] \leq D_r, \quad r \leq t.$$

By definition,

$$\mathbb{E}[D_t \mid \mathcal{F}_r] = \mathbb{E} \left[ \int_0^{t \wedge \tau_K} \mathcal{L}V(X_s) ds \mid \mathcal{F}_r \right] \quad (52)$$

$$= \int_0^{r \wedge \tau_K} \mathcal{L}V(X_s) ds + \mathbb{E} \left[ \int_{r \wedge \tau_K}^{t \wedge \tau_K} \mathcal{L}V(X_s) ds \mid \mathcal{F}_r \right]. \quad (53)$$

Since  $\mathcal{L}V(X_s) \leq 0$ , the second term is nonpositive. Hence

$$\mathbb{E}[D_t \mid \mathcal{F}_r] \leq D_r.$$

Therefore  $D_t$  is a supermartingale.

2. The stochastic integral  $M_t$  is a martingale. A sufficient condition is

$$\mathbb{E} \left[ \int_0^{t \wedge \tau_K} \|\nabla V(X_s)^\top \sigma(X_s)\|^2 ds \right] < \infty.$$

Since  $V$  is piecewise quadratic on finitely many cells,  $\nabla V$  is piecewise linear. Therefore, on a compact set  $K$ ,  $\nabla V$  is bounded. Thus this condition is usually easy to verify if  $\sigma$  is bounded on  $K$ .

3. The interface contribution is nonpositive. The Tanaka interface condition is imposed face by face: for every  $F \in \mathcal{F}_{ij}$ , we assume

$$(\nabla V_j(x) - \nabla V_i(x)) \cdot n_{i \rightarrow j, F} \leq 0, \quad x \in F.$$

Since

$$\nabla V_i(x) = p_i + Q_i x$$

and

$$\nabla V_j(x) = p_j + Q_j x,$$

this condition can be written explicitly as

$$(p_j - p_i + (Q_j - Q_i)x) \cdot n_{i \rightarrow j, F} \leq 0, \quad x \in F, \quad F \in \mathcal{F}_{ij}.$$

This is the multidimensional version of the concavity condition

$$V'_+(x_i) - V'_-(x_i) \leq 0$$

from the one-dimensional case.

#### Fact

If

$$(\nabla V_j(x) - \nabla V_i(x)) \cdot n_{i \rightarrow j, F} \leq 0 \quad \text{on } F,$$

for every  $F \in \mathcal{F}_{ij}$ , and  $L_t^F$  is nondecreasing in  $t$ , then the interface term  $K_t$  is a supermartingale.

**Proof:** We want to prove

$$\mathbb{E}[K_t \mid \mathcal{F}_r] \leq K_r, \quad r \leq t.$$

The interface term is

$$K_t = \frac{1}{2} \sum_{i < j} \sum_{F \in \mathcal{F}_{ij}} \int_0^{t \wedge \tau_K} \gamma_{ij, F}(X_s) dL_s^F,$$

where

$$\gamma_{ij, F}(x) = (\nabla V_j(x) - \nabla V_i(x)) \cdot n_{i \rightarrow j, F}.$$

By assumption,

$$\gamma_{ij, F}(x) \leq 0 \quad x \in F, \quad F \in \mathcal{F}_{ij}.$$

Also,  $dL_s^F$  is a nonnegative measure, because surface local time is nondecreasing. Therefore

$$\int_r^t \gamma_{ij, F}(X_s) dL_s^F \leq 0.$$

Hence the future increment of  $K_t$  after time  $r$  is nonpositive, and so

$$\mathbb{E}[K_t \mid \mathcal{F}_r] \leq K_r.$$

Therefore  $K_t$  is a supermartingale.

#### 5.1.1 Proof of the supermartingale property

We now combine the three conditions.

By the multidimensional Itô–Tanaka formula,

$$V(X_{t \wedge \tau_K}) = V(X_0) + D_t + M_t + K_t. \tag{54}$$

The drift term  $D_t$  is a supermartingale by the interior generator condition:

$$\mathcal{L}V_i(x) \leq 0 \quad x \in \text{int}(K_i).$$

The stochastic integral  $M_t$  is a martingale by the square-integrability assumption:

$$\mathbb{E} \left[ \int_0^{t \wedge \tau_K} |\nabla V(X_s)^\top \sigma(X_s)|^2 ds \right] < \infty.$$

The interface term  $K_t$  is a supermartingale by the multidimensional concavity condition:

$$(\nabla V_j(x) - \nabla V_i(x)) \cdot n_{i \rightarrow j, F} \leq 0, \quad x \in F, \quad F \in \mathcal{F}_{ij}.$$

Therefore  $D_t + K_t$  is a supermartingale, and  $M_t$  is a martingale. Since the sum of a martingale and a supermartingale is a supermartingale, we conclude that

$$V(X_{t \wedge \tau_K})$$

is a supermartingale.

### 5.1.2 Interpretation of the multidimensional concavity condition

The condition

$$(\nabla V_j(x) - \nabla V_i(x)) \cdot n_{i \rightarrow j, F} \leq 0, \quad x \in F, \quad F \in \mathcal{F}_{ij},$$

says that when the process crosses from cell  $K_i$  into cell  $K_j$ , the directional derivative of  $V$  in the crossing direction cannot increase.

Indeed, the normal derivative from the  $K_i$  side is

$$\nabla V_i(x) \cdot n_{i \rightarrow j, F},$$

while the normal derivative from the  $K_j$  side is

$$\nabla V_j(x) \cdot n_{i \rightarrow j, F}.$$

Thus the condition is equivalent to

$$\nabla V_j(x) \cdot n_{i \rightarrow j, F} \leq \nabla V_i(x) \cdot n_{i \rightarrow j, F}.$$

This is exactly the analogue of the one-dimensional condition

$$V'_+(x_i) \leq V'_-(x_i).$$

So, in multiple dimensions, ordinary concavity is not the only thing we need to look at. What matters for the local time term is the jump in the normal derivative across each interface. If that jump is nonpositive, then the surface local time contribution pushes  $V(X_t)$  downwards rather than upwards.

### 5.1.3 Summary of sufficient conditions

We have shown that the following assumptions are sufficient.

1.  $V$  is continuous and piecewise quadratic:

$$V(x) = c_i + p_i^\top x + \frac{1}{2} x^\top Q_i x, \quad x \in K_i.$$

2. The generator is nonpositive inside each cell:

$$(p_i + Q_i x) \cdot f(x) + \frac{1}{2} \text{tr}(a(x) Q_i) \leq 0, \quad x \in \text{int}(K_i).$$

3. The stochastic integral is a martingale:

$$\mathbb{E} \left[ \int_0^{t \wedge \tau_K} \|\nabla V(X_s)^\top \sigma(X_s)\|^2 ds \right] < \infty.$$

4. The interface condition holds on every shared face:

$$(p_j - p_i + (Q_j - Q_i)x) \cdot n_{i \rightarrow j, F} \leq 0, \quad x \in F, \quad F \in \mathcal{F}_{ij}.$$

Under these assumptions,

$$V(X_{t \wedge \tau_K})$$

is a supermartingale.

This gives the multidimensional version of the one-dimensional piecewise linear certificate result. The interior generator condition controls the ordinary Itô drift, while the interface concavity condition controls the Tanaka local time terms.

## 5.2 Reach-avoid stopping convention

The exit-time result above is a general auxiliary result. For the reach-avoid verifier, we stop earlier. Let  $X_\tau \subseteq K$  be the target and let  $\beta$  be the upper certificate level. The stopping time used by the verifier is

$$\tau = \inf \{t \geq 0 : X_t \in X_\tau \text{ or } X_t \in \partial K \text{ or } V(X_t) \geq \beta\}.$$

Thus touching the target, the boundary of the verification domain, or the super- $\beta$  set is the stopping moment. In particular, the unsafe set is not a separate stopping event: the condition  $V \geq \beta$  on the unsafe set ensures that a continuous path reaching it has already reached the super- $\beta$  stopping set.

With this convention, the interior generator condition is needed only on

$$\{x \in \text{int}(K) \setminus X_\tau : V(x) < \beta\}.$$

For verification it is conservative to check its closure, replacing  $V < \beta$  by  $V \leq \beta$ . Likewise, the normal-derivative jump condition is needed only on the portions of internal faces outside the target and in this sub- $\beta$  basin. No generator or interface condition is required after the process has reached any of the three stopping sets.

The stopping convention also requires the domain-boundary separation condition

$$\inf_{x \in \partial K \setminus X_\tau} V(x) \geq \beta.$$

Together with

$$\sup_{x \in X_0} V(x) \leq \alpha, \quad \inf_{x \in X_u} V(x) \geq \beta, \quad \alpha < \beta,$$

these are the conditions implemented by the multidimensional piecewise-quadratic verifier. Applying the preceding Itô-Tanaka argument up to  $\tau$  shows that  $V(X_{t \wedge \tau})$  is a supermartingale.

## 5.3 Asymptotic completeness of piecewise quadratic certificates

The theorem below retains the stronger exit-time formulation: it assumes the generator inequality throughout  $K$  and therefore proves a supermartingale up to  $\tau_K$ . To use its approximants as reach-avoid certificates under the convention above, one additionally needs a strict boundary margin for the smooth certificate on  $\partial K \setminus X_\tau$ ; uniform approximation then preserves  $V_h \geq \beta$  there.

**Theorem (asymptotic completeness under a strict smooth certificate).** Let  $K \subset \mathbb{R}^2$  be compact and polygonal, and consider the stopped SDE

$$dX_t = f(X_t) dt + \sigma(X_t) dW_t, \quad a(x) = \sigma(x)\sigma(x)^\top,$$

stopped at

$$\tau_K = \inf\{t \geq 0 : X_t \notin K\}.$$

Assume that  $f$  and  $a$  are bounded on  $K$ . Suppose there exists a strict smooth certificate  $U \in C^3(K)$  such that

$$\mathcal{L}U(x) = \nabla U(x) \cdot f(x) + \frac{1}{2} \text{tr}(a(x) \nabla^2 U(x)) \leq -\varepsilon \quad x \in K,$$

for some  $\varepsilon > 0$ . Assume also that the boundary/separation conditions hold with margin: for some  $\delta > 0$ ,

$$\sup_{x \in X_0} U(x) \leq \alpha - \delta, \quad \inf_{x \in X_u} U(x) \geq \beta + \delta, \quad \alpha < \beta.$$



Let  $\{\mathcal{T}_h\}_{h \downarrow 0}$  be a shape-regular sequence of refined triangulations of  $K$  supporting  $C^1$  piecewise-quadratic spline interpolants  $I_h U$  satisfying

$$\|I_h U - U\|_{C^2(K)} \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

Then, for all sufficiently small  $h$ , the function

$$V_h := I_h U$$

is a continuous  $C^1$  piecewise-quadratic certificate satisfying

$$\mathcal{L}V_h(x) \leq -\frac{\varepsilon}{2} \quad x \in K,$$

and

$$\sup_{x \in X_0} V_h(x) \leq \alpha, \quad \inf_{x \in X_u} V_h(x) \geq \beta.$$

Consequently,

$$V_h(X_{t \wedge \tau_K})$$

is a supermartingale.

**Proof.** Since  $f$  and  $a$  are bounded on  $K$ , define

$$F := \sup_{x \in K} \|f(x)\|, \quad A := \sup_{x \in K} \|a(x)\|.$$

Let

$$E_h := V_h - U.$$

Then

$$\mathcal{L}V_h - \mathcal{L}U = \nabla E_h \cdot f + \frac{1}{2} \text{tr}(a \nabla^2 E_h).$$

Hence, for a constant  $C_K > 0$  depending only on the dimension and the chosen matrix norm,

$$|\mathcal{L}V_h(x) - \mathcal{L}U(x)| \leq F \|\nabla E_h\|_{L^\infty(K)} + C_K A \|\nabla^2 E_h\|_{L^\infty(K)}.$$

Because

$$\|E_h\|_{C^2(K)} \rightarrow 0,$$

there exists  $h_0 > 0$  such that, for all  $h < h_0$ ,

$$|\mathcal{L}V_h(x) - \mathcal{L}U(x)| \leq \frac{\varepsilon}{2} \quad x \in K.$$

Since  $\mathcal{L}U \leq -\varepsilon$ , it follows that

$$\mathcal{L}V_h \leq -\varepsilon + \frac{\varepsilon}{2} = -\frac{\varepsilon}{2}.$$

Similarly, since  $V_h \rightarrow U$  uniformly, for all sufficiently small  $h$ ,

$$\|V_h - U\|_{L^\infty(K)} \leq \delta.$$

Therefore

$$\sup_{X_0} V_h \leq \sup_{X_0} U + \delta \leq \alpha,$$

and

$$\inf_{X_u} V_h \geq \inf_{X_u} U - \delta \geq \beta.$$

It remains to check the nonsmooth interfaces. The spline  $V_h$  is  $C^1$  across every internal face of the triangulation. Therefore, if two quadratic pieces  $V_i, V_j$  meet across a face  $F_{ij}$ , then

$$\nabla V_i(x) = \nabla V_j(x) \quad x \in F_{ij}.$$

Hence the Tanaka normal-derivative jump is

$$[\nabla V_j(x) - \nabla V_i(x)] \cdot n_{i \rightarrow j} = 0,$$

so every local-time contribution is zero.

Applying the multidimensional Itô–Tanaka formula gives

$$V_h(X_{t \wedge \tau_K}) = V_h(X_0) + \int_0^{t \wedge \tau_K} \mathcal{L}V_h(X_s) ds + \int_0^{t \wedge \tau_K} \nabla V_h(X_s) \sigma(X_s) dW_s.$$

The drift term is nonpositive because

$$\mathcal{L}V_h \leq -\frac{\varepsilon}{2} < 0.$$

The stochastic integral is a true martingale after stopping, since  $K$  is compact and  $V_h, \sigma$  are bounded on  $K$ . Therefore

$$V_h(X_{t \wedge \tau_K})$$

is a martingale plus a nonincreasing finite-variation process, and is hence a supermartingale.  $\square$

## 6 The local-time condition by construction

This section proves that the deep ReLU input-convex neural network used in the certificate architecture enforces the required sign of the local-time term by construction. The relevant point is a sign convention: the network  $C_\psi$  defined below is convex, and hence its gradient jump across an activation interface has the convex sign. Therefore the supermartingale certificate uses the residual form

$$V_\theta(x) = S_\theta(x) - C_\psi(x),$$

where  $S_\theta$  is continuously differentiable. The branch  $-C_\psi$ , rather than  $C_\psi$  itself, then satisfies the nonpositive normal-derivative jump condition required by the Itô–Tanaka formula.

### 6.1 The deep ReLU ICNN

Let  $x \in \mathbb{R}^d$ . Consider a network with  $L \geq 1$  hidden layers, all of width  $m$ , defined recursively by

$$z_1(x) = \text{ReLU}(U_1 x + b_1), \tag{55}$$

$$z_\ell(x) = \text{ReLU}(U_\ell x + b_\ell + W_\ell z_{\ell-1}(x)), \quad \ell = 2, \dots, L, \tag{56}$$

and with scalar output

$$C_\psi(x) = u^\top x + c + w^\top z_L(x). \tag{57}$$

The input matrices  $U_\ell$ , biases  $b_\ell$ , output input weight  $u$ , and output bias  $c$  are unrestricted. The recurrent matrices and final hidden-to-output weights satisfy

$$W_\ell \geq 0, \quad w \geq 0, \tag{58}$$

entrywise.

In the implementation, these inequalities hold for every parameter value because

$$W_\ell = \text{softplus}(\widetilde{W}_\ell), \quad w = \text{softplus}(\widetilde{w}).$$

Since  $\text{softplus}(r) > 0$  for every finite  $r$ , the effective weights are nonnegative without requiring a constrained optimization step.

**Proposition 6.1** (Convexity and piecewise-affine structure). *The function  $C_\psi : \mathbb{R}^d \rightarrow \mathbb{R}$  defined by (57)–(58) is finite, continuous, convex, and piecewise affine.*

*Proof.* We prove convexity by induction over the hidden layers.

For the first layer, each coordinate has the form

$$(z_1)_k(x) = \max\{0, a_k^\top x + \beta_k\},$$

which is the maximum of two affine functions and is therefore convex. Consequently every coordinate of  $z_1$  is convex.

Suppose that every coordinate of  $z_{\ell-1}$  is convex. For a fixed coordinate  $k$  of the next preactivation, write

$$q_{\ell,k}(x) = (U_\ell x + b_\ell)_k + \sum_{r=1}^m (W_\ell)_{kr} (z_{\ell-1})_r(x).$$

The first term is affine. Every coefficient  $(W_\ell)_{kr}$  is nonnegative, so the second term is a nonnegative linear combination of convex functions. Thus  $q_{\ell,k}$  is convex. Because ReLU is convex and nondecreasing,

$$(z_\ell)_k(x) = \text{ReLU}(q_{\ell,k}(x))$$

is convex. This completes the induction.

Finally,

$$C_\psi(x) = u^\top x + c + \sum_{k=1}^m w_k (z_L)_k(x)$$

is the sum of an affine function and a nonnegative linear combination of convex functions, and is therefore convex.

Every operation in the network is obtained from affine maps, finite sums, and coordinatewise ReLU. Hence the network is continuous and piecewise affine, with finitely many affine pieces.  $\square$

## 6.2 Convexity determines the interface sign

Let  $K_i$  and  $K_j$  be two full-dimensional affine regions of  $C_\psi$  that share a codimension-one face  $F$ . On the two regions, write

$$C_\psi(x) = a_i^\top x + c_i, \quad x \in K_i,$$

and

$$C_\psi(x) = a_j^\top x + c_j, \quad x \in K_j.$$

Let  $n_{i \rightarrow j, F}$  be a unit normal to  $F$  pointing from  $K_i$  into  $K_j$ .

**Lemma 6.2** (Normal derivative jump of a convex CPWL function). *For every shared face  $F$ ,*

$$(a_j - a_i) \cdot n_{i \rightarrow j, F} \geq 0. \quad (59)$$

*Equivalently,*

$$(\nabla C_{\psi,j}(x) - \nabla C_{\psi,i}(x)) \cdot n_{i \rightarrow j, F} \geq 0, \quad x \in \text{relint}(F).$$

*Proof.* Choose  $x_0 \in \text{relint}(F)$ . Since the two affine pieces agree on  $F$ , their difference vanishes on every tangent direction to  $F$ . Therefore  $a_j - a_i$  is normal to  $F$ , so there exists  $\lambda \in \mathbb{R}$  such that

$$a_j - a_i = \lambda n_{i \rightarrow j, F}.$$

It remains to prove  $\lambda \geq 0$ .

For sufficiently small  $\varepsilon > 0$ , the points

$$x_- = x_0 - \varepsilon n_{i \rightarrow j, F} \in K_i, \quad x_+ = x_0 + \varepsilon n_{i \rightarrow j, F} \in K_j.$$

Restrict  $C_\psi$  to the line

$$g(t) = C_\psi(x_0 + t n_{i \rightarrow j, F}).$$

Because  $C_\psi$  is convex,  $g$  is a convex function of one variable. Its left and right derivatives at zero are

$$g'_-(0) = a_i \cdot n_{i \rightarrow j, F}, \quad g'_+(0) = a_j \cdot n_{i \rightarrow j, F}.$$

The one-dimensional derivative of a convex function is nondecreasing, hence

$$a_i \cdot n_{i \rightarrow j, F} \leq a_j \cdot n_{i \rightarrow j, F}.$$

This is exactly (59).  $\square$

## 6.3 The residual certificate

We now consider the certificate

$$V_\theta(x) = \rho(S_\theta(x) - C_\psi(x)), \quad \rho > 0, \quad (60)$$

where  $S_\theta \in C^1$ . In the implementation,  $S_\theta$  is the shallow piecewise-quadratic network

$$S_\theta(x) = c + p^\top x + \frac{1}{2} x^\top H x + \frac{1}{2} \sum_{k=1}^M v_k \text{ReLU}(a_k^\top x + b_k)^2.$$

Since the scalar function  $r \mapsto \frac{1}{2} \text{ReLU}(r)^2$  is continuously differentiable,  $S_\theta$  has no gradient jump across any of its activation hyperplanes.

The common refinement of the activation regions of  $S_\theta$  and  $C_\psi$  is a finite polyhedral partition. Let  $R_i$  and  $R_j$  be adjacent full-dimensional cells of this refinement, sharing a face  $F$ , and let  $n_{i \rightarrow j, F}$  point from  $R_i$  to  $R_j$ .

**Theorem 6.3** (Local-time condition by construction). *For every choice of the unconstrained parameters of  $S_\theta$  and  $C_\psi$ , every choice of raw recurrent and output parameters, and every adjacent pair of cells in the common refinement, the certificate (60) satisfies*

$$(\nabla V_{\theta,j}(x) - \nabla V_{\theta,i}(x)) \cdot n_{i \rightarrow j, F} \leq 0, \quad x \in \text{relint}(F). \quad (61)$$

Consequently, every codimension-one local-time contribution in the multidimensional Itô–Tanaka formula for  $V_\theta(X_t)$  is nonpositive. No activation-region enumeration or pairwise interface verification is needed to establish this sign.

*Proof.* On the two sides of  $F$ ,

$$\nabla V_{\theta,i} = \rho(\nabla S_{\theta,i} - \nabla C_{\psi,i}), \quad \nabla V_{\theta,j} = \rho(\nabla S_{\theta,j} - \nabla C_{\psi,j}).$$

Therefore

$$(\nabla V_{\theta,j} - \nabla V_{\theta,i}) \cdot n_{i \rightarrow j, F} \quad (62)$$

$$= \rho(\nabla S_{\theta,j} - \nabla S_{\theta,i}) \cdot n_{i \rightarrow j, F} - \rho(\nabla C_{\psi,j} - \nabla C_{\psi,i}) \cdot n_{i \rightarrow j, F}. \quad (63)$$

Because  $S_\theta \in C^1$ , its gradient is continuous across every face of the common refinement. Hence

$$\nabla S_{\theta,j}(x) - \nabla S_{\theta,i}(x) = 0, \quad x \in F.$$

If  $F$  is introduced only by the smooth branch, then  $C_\psi$  is affine on both sides and its gradient jump is also zero. If  $F$  lies on an activation interface of  $C_\psi$ , Lemma 6.2 gives

$$(\nabla C_{\psi,j}(x) - \nabla C_{\psi,i}(x)) \cdot n_{i \rightarrow j, F} \geq 0.$$

Substituting these facts into (63), and using  $\rho > 0$ , yields

$$(\nabla V_{\theta,j}(x) - \nabla V_{\theta,i}(x)) \cdot n_{i \rightarrow j, F} = -\rho(\nabla C_{\psi,j}(x) - \nabla C_{\psi,i}(x)) \cdot n_{i \rightarrow j, F} \leq 0.$$

This proves (61).

In the multidimensional Itô–Tanaka formula, the contribution of  $F$  is

$$\frac{1}{2} \int_0^t (\nabla V_{\theta,j}(X_s) - \nabla V_{\theta,i}(X_s)) \cdot n_{i \rightarrow j, F} dL_s^F.$$

Surface local time is nondecreasing, so  $dL_s^F$  is a nonnegative measure. The integrand is nonpositive by the preceding argument; therefore the entire integral is nonpositive. Summing over all faces preserves this sign.  $\square$

**Corollary 6.4** (Max-affine specialization). *The conclusion of Theorem 6.3 also holds when the convex branch is*

$$C_\psi(x) = \max_{1 \leq r \leq R} \{a_r^\top x + b_r\}.$$

*In particular, it applies to the residual max-affine certificate used in the initial radial Ornstein–Uhlenbeck training experiment.*

*Proof.* A finite maximum of affine functions is finite, continuous, convex, and piecewise affine. The proof of Theorem 6.3 uses no other property of  $C_\psi$ , so it applies unchanged.  $\square$

*Remark 6.5* (What is and is not guaranteed). The theorem concerns only the singular, interface part of the generalized Itô formula. It does not by itself imply that  $V_\theta(X_t)$  is a supermartingale. One must still verify the interior generator inequality, for example

$$\mathcal{L}V_{\theta,i}(x) \leq 0$$

on every relevant smooth cell, together with the required boundary, separation, stopping, and stochastic-integrability conditions. The architecture removes the combinatorial verification of the local-time sign; it does not remove the ordinary generator verification problem.

*Remark 6.6* (Distributional-Hessian interpretation). The result can be stated without referring to a particular activation-region partition. A finite convex piecewise-affine function has a positive semidefinite singular Hessian measure. Since  $S_\theta \in C^1$ , it has no codimension-one singular Hessian mass, and therefore

$$D_{\text{sing}}^2 V_\theta = -\rho D^2 C_\psi \preceq 0.$$

Thus the architecture enforces nonpositive singular curvature globally. The facewise normal-jump condition in (61) is the polyhedral expression of this measure inequality.